

Turkish Version of The Renal Inpatient Nutrition Screening Tool: Validity and Reliability for Hemodialysis Patients

Nursena ERSOY¹, Emine KARADEMİR², Ebru BAYRAK³, Muslu Kazım KÖREZ⁴, Hülya YARDIMCI⁵

¹Postal Address: Ankara University Institute of Health Sciences, Dışkapı Campus Şehit Ömer Halisdemir Boulevard 06110 Dışkapı ANKARA, ORCID: 0000-0003-4327-0775, E-mail: nersoy@ankara.edu.tr

²Postal Address: Niğde Ömer Halisdemir University Faculty of Health Sciences, Department of Nutrition and Dietetics, Bor/NIGDE/TURKEY, ORCID: 0000-0002-8319-7210, E-mail: eminekarademir@ohu.edu.tr

³Postal Address: Selçuk University Faculty of Health Sciences, Department of Nutrition and Dietetics, Alaeddin Keykubat Campus 299/1 42250 Selçuklu/KONYA/TURKEY, ORCID: 0000-0001-7279-3255, E-mail: ebayrak@selcuk.edu.tr

⁴Postal Address: Selçuk University Faculty of Health Sciences, Department of Biostatistic, Alaeddin Keykubat Campus 299/1 42250 Selçuklu/KONYA/TURKEY, ORCID: 0000-0001-9524-6115, E-mail: mkkorez@gmail.com

⁵Postal Address: Ankara University Faculty of Health Sciences, Department of Nutrition and Dietetics, Fatih Street. Tepebaşı District, No:197/A, 06300 Kecioren-ANKARA/TURKEY, ORCID: 0000-0002-2664-4176, E-mail: hulyardimci@gmail.com

Corresponding author: Nursena ERSOY (MSc), Postal Address: Ankara University Institute of Health Sciences, Dışkapı Campus Şehit Ömer Halisdemir Boulevard 06110 Dışkapı ANKARA, ORCID: 0000-0003-4327-0775, Phone Number: +90 (312) 381 23 50-6545, E-mail: nersoy@ankara.edu.tr



This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI

10.1017/S0007114524003192

The British Journal of Nutrition is published by Cambridge University Press on behalf of The Nutrition Society

Abstract

The aim of this study was to analyze the validity and reliability of the Turkish version of the Renal Inpatient Nutrition Screening Tool (Renal iNUT) for hemodialysis patients.

The Renal Inpatient Nutrition Screening Tool (Renal iNUT) and the Malnutrition Universal Screening Tool (MUST) were used in adult hemodialysis patients at two different centers to identify malnutrition. The Subjective Global Assessment (SGA), regarded as the gold standard for nutritional status assessment, was utilized for comparison. Structural validity was assessed using biochemical values and anthropometric measurements, while reliability was assessed using repeated the Renal iNUT assessment. Of the 260 patients admitted, 42.3% were malnourished (SGA score was B or C). According to the Renal iNUT, 59.6% of the patients were at increased risk for malnutrition (score ≥ 1) and 3.8% required referral to a dietitian (score ≥ 2). According to the MUST, 13.1% the of patients were at increased risk for malnutrition and 8.5% required referral to a dietitian. The Renal iNUT was found to be more sensitive in detecting increased risk of malnutrition in hemodialysis patients compared with the MUST (59.6% versus 13.1%). According to the SGA, the sensitivity of the Renal iNUT is higher compared to the MUST (89% and 45%, respectively). Kappa-assessed reliability of the Renal iNUT was 0.48 (95% CI, 0.58-0.9) and a moderate concordance was observed. The Renal iNUT is a valid and reliable nutritional screening tool for evaluating hemodialysis patients to determine their nutritional status. The use of the Renal iNUT by dietitians will contribute to the identification of malnutrition and its treatment.

Keywords: Chronic kidney disease; Hemodialysis; Malnutrition; Renal; Nutritional Assessment

Abbreviations List:

BMI	Body Mass Index
CKD	Chronic Kidney Disease
ESRF	End-Stage Renal Failure
HGS	Handgrip Strength
MUAC	Mid-Upper Arm Circumference
MUST	Malnutrition Universal Screening Tool
NPV	Negative Predictive Value
PPV	Positive Predictive Value
Renal iNUT	The Renal Inpatient Nutrition Screening Tool
ROC	Receiver Operating Characteristics
SGA	Subjective Global Assessment
TST	Triceps Skinfold Thickness

INTRODUCTION

Chronic kidney disease (CKD) is a disease characterized by the chronic and progressive impairment of renal functions, resulting from damage and dysfunction of the kidney's hemofiltration processes ⁽¹⁾. This disease, which affects approximately 10% of the global population and one in two individuals over the age of 75, represents a significant global public health concern ⁽²⁾. It was reported that the median prevalence of CKD is 9.5% ⁽³⁾. In Turkey, the prevalence is 15.7% ⁽⁴⁾.

The nature of chronic kidney disease results in malnutrition in hemodialysis patients due to a combination of factors, including dialysis-related issues, psychosocial influences, and a decline in physical capacity ^(5, 6). As a consequence of diminished appetite resulting from inflammatory processes and alterations in taste perception associated with uremia, these patients exhibit a deficiency in energy and protein intake ⁽⁷⁾. This condition, which is associated with hypercatabolism, leads to muscle wasting and malnutrition ⁽⁸⁾. Given the association between malnutrition in these patients and poor prognosis and quality of life, it is imperative to prioritize the prevention and early diagnosis of malnutrition is important ⁽⁷⁾.

It is recommended that the nutritional status of hemodialysis patients be reviewed during the initial 90-day period following the commencement of dialysis and subsequently on an annual basis, or at a frequency determined by the results of a nutritional screening tool. A comprehensive nutritional assessment should include an evaluation of appetite, dietary history, biochemical parameters, anthropometric measurements, and any physical symptoms that may be related to nutritional status. The use of screening and malnutrition assessment tools is strongly recommended for this purpose. Nevertheless, there are notable distinctions between nutritional screening and assessment tools. Nutritional screening tools are designed to identify the risk of malnutrition in a given population, whereas nutritional assessment tools are used to conduct a comprehensive examination of the nutritional status of individuals who are at risk of malnutrition ⁽⁹⁾. Accordingly, when selecting a tool for the assessment of malnutrition, it is essential to exercise caution in interpreting the results, regardless of whether the tool is intended for screening or assessment purposes. The National Kidney Foundation recommends the use of the 7-item Subjective Global Assessment (SGA) tool for dialysis patients. The SGA is a valid and reliable tool for dialysis patients, as it addresses a range of pertinent health issues, including weight loss, food intake, gastrointestinal problems, muscle loss, loss of fat stores, edema, and functional capacity over the previous 6 months ⁽¹⁰⁾.

¹¹⁾. In Turkey, the 7-P SGA is also a valid and reliable tool for use with hemodialysis patients ⁽¹²⁾.

The Malnutrition Universal Screening Tool (MUST) is another commonly used screening tool for malnutrition ^(13, 14). The MUST includes an assessment of BMI, body weight loss, and any acute illness associated with fasting for more than five days. It should be noted that the sensitivity is lower in all renal inpatients, not only those undergoing hemodialysis, than it is in patients assessed using the SGA as a reference tool ⁽¹⁵⁾. In Turkey, the MUST has also been shown to be a specific screening tool for this population, but with less sensitivity ⁽¹⁶⁾. Given the challenges associated with utilizing the SGA and the reduced sensitivity of the MUST, the Renal iNUT screening tool for malnutrition was devised with the objective of enhancing simplicity and practicality by employing a series of questions pertaining to body weight loss, BMI, appetite, food intake, and nutritional support ⁽¹⁷⁾. This study aimed to examine the validity of this nutritional screening tool, which has been demonstrated to be a valid and reliable instrument for use in renal specialist wards in the UK, in assessing the nutritional status of patients receiving hemodialysis in dialysis centers, which are common in Turkey.

METHOD

The data were gathered from two private hemodialysis centers in two distinct cities within the CCC region between February and April of 2022. A total of 260 volunteer patients, aged 20 to 65 years who had been undergoing regular hemodialysis treatment for a minimum of one year and could speak Turkish were enrolled in the study. Patients were excluded from the study if they were younger than nineteen years of age, older than sixty-five, pregnant, had hearing difficulties, an intellectual disability, or were experiencing fatigue during hemodialysis (Figure 1). The study was conducted in accordance with the ethical standards of the Ethics Board for Noninvasive Research of ... University (approval no. 2022/60) and in compliance with the STROBE checklist for observational research. Prior to their participation, all subjects provided informed consent.

Permission to translate the Renal iNUT was granted via e-mail by Helena S. Jackson. A standard translation and back translation were conducted. In the preliminary translation phase, two Turkish-speaking translators proficient in English translated the screening tool independently into Turkish. The discrepancies between the two versions were collectively addressed by the research team. Another bilingual translator, who was not familiar with the English version, translated the screening tool back into English. A total of ten experts in the field of nutrition were invited to participate in a content validity test of Renal iNUT. The

experts were provided with the expert form via email and asked to rate the simplicity, clarity, appropriateness, and necessity of each question. In light of the expert opinions, the Content Validity Index was calculated to be 0.76. This value indicates that the scale is deemed to possess content validity ⁽¹⁸⁾. The comments provided by the experts were evaluated by the research teams. Following this evaluation, a pilot study was conducted on 20 individuals without making any significant alterations to the screening tool. Once the necessary corrections had been made, the final version of the screening tool was produced.

The patients were queried regarding their age, sex, date of diagnosis of CKD, duration of dialysis treatment, and the presence of any chronic diseases. The Charlson Comorbidity Index was calculated based on the patients' medical histories ⁽¹⁹⁾. The biochemical parameters were used in the standard clinical practice. Additionally, an expert measured the following variables after dialysis: dry body weight, height, mid-upper arm circumference (MUAC), handgrip strength (HGS), and triceps skinfold thickness (TST). Patients were weighed without shoes or outer garments in order to determine their dry body weight. This measurement was taken with the use of an accurate Seca 799 scale, calibrated in 0.1-kilogram increments ⁽²⁰⁾. The patients' heights were measured with their feet bare and in an upright position, with their heads aligned with the Frankfurt plane, using a portable stadiometer (Seca 769) ⁽²¹⁾. The MUAC was measured by having the subjects adopt an upright stance and flex the arm at the elbow to achieve a 90° angle. The midpoint between the acromion and the olecranon process was marked and the mean humeral circumference was measured with a tape measure ⁽²²⁾. In the TST, the subjects were instructed to flex their elbows at a 90° angle, parallel to the floor, in an upright position. The right MUAC was determined and marked. Subcutaneous adipose tissue was then held with the thumb and index finger at the center of the right mid-dorsum of the upper arm, and measured with a Holtain brand caliper ⁽²³⁾. The HGS was measured on at least three occasions from the arm devoid of fistula with the use of a Takei digital dynamometer. The procedure was conducted with the shoulder in a state of adduction, with the elbow maintained at a flexion angle of 90°. The highest value obtained was deemed to be the patient's HGS ⁽²⁴⁾.

The Renal iNUT is a renal-specific nutrition screening tool comprising five questions developed by Jackson et al. in 2019. The total score obtained from the Renal iNUT is indicative of the necessity for further evaluation and intervention. A score of 0 indicates that weekly screening is sufficient, while a score of 1 suggests the need for follow-up of high-risk

patients. A score of ≥ 2 indicates that the patient should be referred to a dietician. In this screening tool, a score of ≥ 1 was identified as indicative of an increased risk of malnutrition, consistent with the findings of the original study ⁽¹⁷⁾. A total of 6-7 points from SGA 7P indicates that the patient is well nourished (A), 3-5 points indicate moderate malnutrition (B), and 1-2 points indicate severe malnutrition (C) ⁽¹¹⁾. In this study, SGA A was classified as indicating a low nutritional risk, SGA B as indicating an increased nutritional risk, and SGA C as indicating the need for referral to a dietician. Furthermore, SGA B+C was classified as indicating an increased risk of malnutrition. The MUST consists of three questions. A total score of 0 is indicative of low risk, 1 is suggestive of increased nutritional risk, and a score of ≥ 2 is indicative of the necessity for referral to a dietitian ⁽²⁵⁾. For MUST and Renal iNUT, a score of 1 or greater was classified as indicative of an increased risk for malnutrition. All screening and assessment tools were applied by nurses and dietitians in the centres and by experts on the study team.

All statistical analyses were performed with the R software, version 4.2.1. (The R Foundation for Statistical Computing, Vienna, Austria; <https://www.r-project.org>). The Shapiro-Wilk normality test and Q-Q plots were used to evaluate the normality of the data. In addition, Levene's test was used to check the homogeneity of variances. Numerical variables are expressed as mean \pm standard deviation or median with interquartile range (25th percentile – 75th percentile), as appropriate. One-Way ANOVA (analysis of variances) and the Kruskal-Wallis tests were performed to determine whether there was a statistically significant difference between the Renal iNUT categories, and these statistical test results were also used to examine the construct validity of the Renal iNUT. Subsequently, post-hoc comparisons were conducted using the Tukey HSD and Bonferroni-corrected Dunn tests, respectively, in cases where the variables were found to be significant following the aforementioned tests. To ascertain the diagnostic efficacy of the Renal iNUT method for the identification of malnutrition as determined by the SGA method, receiver operating characteristics (ROC) curve analysis and statistical diagnostic measures (sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV), diagnostic accuracy, and kappa value) were employed. Furthermore, we conducted a comparative analysis of the diagnostic efficacy of the Renal iNUT method and the MUST method for the identification of malnutrition risk in hemodialysis patients. The McNemar test was performed to compare the sensitivity and specificity of the Renal iNUT and MUST methods, and the Weighted Generalized score test was also applied to compare the negative and positive predictive values of these methods. We

examined the interrater reliability of the Renal iNUT method using Weighted kappa value. A *p*-value of less than 5% was considered statistically significant.

RESULTS

A total of 260 patients (52% male and 48% female) who received hemodialysis at two private dialysis centers in Ankara ($n=167$, 64.2%) and Konya ($n=93$, 35.8%) were included in the study. The mean age of the patients was 62.91 ± 13.39 years. The data regarding anthropometric measures and biochemical parameters are presented in Table 1.

According to SGA category A, screening tool Renal iNUT=0 and MUST=0, 57.7%, 40.4% and 78.5% of hemodialysis patients have low nutritional risk, respectively. According to SGA B, Renal iNUT=1 and MUST=1, respectively 38.5%, 36.9% and 13.1% of the patients who increased nutritional risk and according to SGA category C, Renal iNUT ≥ 2 and MUST ≥ 2 scores, it was found that 3.8%, 22.7% and 8.5% of the patients needed to be referred to a dietician. In addition, the number of patients found to be at increased risk for malnutrition according to Renal iNUT ($n=96$) was more than double the number of patients with MUST ($n=34$) (Figure 2).

A comparison of Renal iNUT and MUST in patients with low and severe malnutrition (SGA B+C) (42.3%) reveals a higher prevalence of increased nutritional risk with Renal iNUT ≥ 1 (37.7%) than with MUST ≥ 1 (19.2%) (Table 2).

A comparison of Renal iNUT and MUST sensitivity, as reported by SGA, revealed that Renal iNUT sensitivity was statistically significantly higher than that of MUST (89% vs 45%, *p*-value $< .001$). The Kappa coefficient of concordance, which indicates the degree of concordance between SGA and renal iNUT, was calculated to be 0.484, indicating a moderate level of concordance (Table 2). Figure 3 illustrates the results of the ROC curve analysis, which indicates that Renal iNUT and MUST are effective in predicting risk malnutrition in hemodialysis patients (AUC=0.755 [95% CI, 0.696 – 0.815] and AUC=0.707 [95% CI, 0.640 – 0.774], respectively).

To evaluate the construct validity of Renal iNUT, anthropometric measurements and biochemical parameters were evaluated in accordance with the patients' Renal iNUT scores. As the Renal iNUT score increased, there was a statistically significant decrease in BMI in each Renal iNUT category (Figure 4-A). In the group referred to a dietician (Renal iNUT score ≥ 2), there was a statistically significant decrease in MUAC (Figure 4-B). In addition,

the handgrip scores of individuals at low nutritional risk were statistically significantly higher than those who were referred to a dietician (Figure 4-D). In the biochemical parameters, a significant reduction in CRP levels was observed in subjects with Renal iNUT=0 compared to those with Renal iNUT=1 (Figure 4-E) (Table 3). The weighted kappa for the repeated Renal iNUT assessment in a subsample of 69 subjects was 0.608 (95% CI, 0.449-0.766) for scores 0 to 3, indicating good and substantial agreement between the scores.

DISCUSSION

In this study, the Renal iNUT screening tool, developed by Jackson et al. (2019), was adapted for the purpose of determining the prevalence of malnutrition among Turkish hemodialysis patients. The tool was applied to patients receiving treatment in hemodialysis centers. Anthropometric measurements and biochemical parameters were analyzed, and the Renal iNUT was found to be a valid tool for this purpose. Furthermore, this tool has been demonstrated to be reliable for the Turkish population when used for repeated Renal iNUT assessments. Additionally, the Renal iNUT has been shown to be more sensitive for this population than the MUST, a commonly used screening tool.

It is crucial to evaluate the nutritional status of patients undergoing dialysis for chronic kidney disease and to identify instances of malnutrition through the assessment of multiple parameters⁽²⁵⁾. Nutritional screening tools facilitate the assessment of nutritional status and the evaluation of the efficacy of implemented interventions⁽²⁶⁾. Malnutrition may result from deficiencies, excesses, and imbalances in energy and nutrient intake in these patients. Such imbalances can lead to alterations in body composition (e.g., changes in fat and muscle percentage), a decline in physical and mental functions, and complications⁽²⁷⁾. It is therefore of great importance to be able to identify cases of malnutrition at an early stage and to take appropriate action.

The SGA is a valid and reliable nutritional assessment tool used in hemodialysis patients⁽²⁸⁾. In our study, the prevalence of malnutrition risk was 42.3% among hemodialysis patients according to the SGA. A recent meta-analysis revealed that the prevalence of malnutrition among hemodialysis patients according to the SGA ranges between 28% and 54%⁽²⁹⁾. In this sample, the percentage in Turkey was reported to be 29.3⁽³⁰⁾. According to the Renal iNUT, which is a new screening tool for renal patients, 45.4% of hemodialysis patients are at malnutrition risk⁽³¹⁾, whereas in our study, this ratio was found to be 59.6%. The Renal iNUT

questionnaire encompasses inquiries pertaining to body mass index (BMI), weight loss, dietary intake, and the utilization of nutritional supplements. In addition to other components, the assessment of supplement use, particularly in renal patients, may be crucial for identifying the risk of malnutrition. According to the MUST, which is another screening tool used in this study, 21.6% of the patients were at risk of malnutrition. The sensitivities of the Renal iNUT and MUST were reported to be 89% and 45%, respectively, compared with the SGA gold standards. The Renal iNUT has been demonstrated to be a statistically significantly better predictor of malnutrition in hemodialysis patients than the MUST. In a study conducted by Jackson et al. (2019), Renal iNUT was found to have a sensitivity of 92.1%, while the sensitivity of MUST was determined to be 44.4%. The findings of both the original study and our own investigation indicate that the Renal iNUT is a more effective screening instrument than the MUST in patients undergoing hemodialysis⁽¹⁷⁾.

Anthropometric measures can be incorporated into a comprehensive nutritional assessment in people with kidney disease⁽³²⁾. In patients undergoing hemodialysis, edema frequently arises for a number of reasons, including low albumin levels, excessive fluid intake, diminished cardiac function, and hypotension⁽³³⁾. The presence of edema can impede the accurate assessment of malnutrition by obscuring body weight. To mitigate this potential confounding factor, this study employed a methodology wherein anthropometric measurements were obtained following the completion of hemodialysis. As per the Renal iNUT categories, the BMI was 27.94 kg/m² (Renal iNUT=0), 25.81 kg/m² (Renal iNUT=1), and 22.90 kg/m² (Renal iNUT≥2), with a notable decline in BMI as the risk of malnutrition increased. Similarly, in another study, a negative correlation was observed between BMI and the score on the MUST, with a score of 0-1 indicating a BMI of 23.7 kg/m² and a score of >2 indicating a BMI of 17.3 kg/m².¹⁶ In patients undergoing hemodialysis, both underweight and obesity are associated with an elevated risk of mortality, as indicated by BMI⁽³²⁾. Accordingly, the objective is to ascertain the precise body weight of hemodialysis patients and ensure that their BMI values remain within the normal range.

In addition to BMI, body composition, which is composed of extracellular fluid, muscle mass, and fat mass, is a significant factor in the assessment of nutritional status. In order to evaluate body fat composition, it is recommended that renal patients undergo skinfold thickness measurement, as this method is both simple and cost-effective⁽³²⁾. The present study did not identify a statistically significant correlation between the Renal iNUT classification and the TST results⁽³⁴⁾. Handgrip strength, which is a sign of adequate energy

and protein intake, is also indicative of skeletal muscle function and nutrition. In this instance, the recommended approach is to utilize the HGS method, which is a straightforward, non-invasive, and highly effective technique for evaluating muscle function⁽³⁵⁾. In our study, the mean HGS value was found to be 22.3 kg (range: 16.5-27.9 kg) in males and 10.9 kg (range: 7.5-15.3 kg) in females. While the mean HGS score in various Turkish studies is higher than that observed in the present study, the HGS score of males is comparable to that of females^(37, 38). As reported by Renal iNUT, the HGS score of the cohort who had consulted with a dietician was found to be statistically significantly lower in comparison to the cohort that was deemed to be well-nourished. These findings reinforce the notion that despite patients exhibiting normal BMI values, muscle mass should be assessed independently, as BMI is not a reliable indicator of body composition⁽³⁹⁾. MUAC, which is also an indicator of body fat, varies between Renal iNUT categories. As a result, both BMI and HGS along with MUAC values differ among Renal iNUT classifications. This shows that as the risk of malnutrition increases, HGS and MUAC levels decrease in parallel with BMI.

In patients with renal disease, biomarkers such as serum albumin are also useful additional methods for assessing nutritional status⁽³²⁾. Although albumin levels did not demonstrate significant discrepancies between categories of nutritional risk as defined by Renal iNUT, elevated CRP levels were observed in individuals at elevated nutritional risk. Moreover, there was no correlation between albumin levels and nutritional status, even when CRP levels were elevated. This may provide a novel perspective on the use of CRP in lieu of albumin in SGA or other global measures of nutritional status. The occurrence of inflammatory markers in these patients can be attributed to a number of factors, including infection of the fistula, the composition of the dialysate, filtration processes, chronic infection, and inadequate nutritional intake⁽⁴⁰⁾. Despite the absence of studies demonstrating the validity and reliability of inflammatory markers as a standalone assessment of nutritional status, there is evidence indicating a correlation between inflammatory markers and other nutritional markers⁽⁴¹⁾. However, because these markers are also influenced by factors other than diet, biomarkers should be assessed together rather than separately to evaluate nutritional status.

The principal strength of this study is that it was conducted at two different centers in Turkey, and that a number of anthropometric measurements and routine biochemical parameters were evaluated. Furthermore, the repeat test conducted five days later served to corroborate the reliability of Renal iNUT. The researchers, who are dietitians, applied the Renal iNUT. A limitation of the study is that the content and ease of use of the screening tool were not

evaluated by the nurses or the dietitian at the dialysis center. Furthermore, it should be noted that the present study, which was conducted at two hemodialysis centers, may yield different results when compared to those observed in inpatients. It is possible that alterations in the environment may result in a shift in the characteristics of the patient population. The population under investigation comprises patients who reside in their own homes and who visit the center for dialysis sessions. It would be beneficial to conduct further studies with larger samples, including inpatients from diverse regions within Turkey.

The Renal iNUT nutritional screening tool is a valid and reliable method for assessing the nutritional status of patients undergoing hemodialysis. Based on the results, it was recommended that iNUT be included in usual care by nurses or dietitians. The use of Renal iNUT allows the screening of nutritional status and allows the early treatment of malnutrition once it has been identified as a risk. The Renal iNUT is a specific screening tool developed for patients with renal diseases. It is recommended that future studies include a greater variety of samples, comprising renal patients receiving different medical treatments such as peritoneal dialysis.

Acknowledgements: None

Financial Support: None

Conflict of Interest: None

Authorship: Ersoy, Nursena: Conceptualization (Equal), Investigation (Equal), Methodology (Equal), Resources (Equal), Visualization (Equal), Writing – original draft (Equal), Writing – review & editing (Equal) **Karademir, Emine:** Conceptualization (Equal), Investigation (Equal), Methodology (Equal), Resources (Equal), Writing – original draft (Equal), Writing – review & editing (Equal) **Bayrak, Ebru:** Conceptualization (Equal), Investigation (Equal), Methodology (Equal), Supervision (Equal), Writing – review & editing (Equal) **Körez, Muslu Kazım:** Formal analysis (Lead), Methodology (Lead), Resources (Equal), Validation (Lead), Writing – review & editing (Equal)

Yardımcı, Hülya: Conceptualization (Equal), Methodology (Equal), Supervision (Equal), Writing – review & editing (Equal)

References

1. National Institute of Diabetes and Digestive and Kidney Disease, NIH. Chronic kidney disease. Accessed June 12, 2023. <https://www.niddk.nih.gov/health-information/kidney-disease/chronic-kidney-disease-ckd>.
2. National Kidney Federation. Chronic kidney disease. Accessed May 19, 2022. <https://www.kidney.org/atoz/content/about-chronic-kidney-disease>.
3. Jadoul M., Aoun, M., Imani MM. The major global burden of chronic kidney disease. *The Lancet Global Health*, 2024; **12(3)**, e342-e343.
4. Süleymanlar G, Utaş C, Arinsoy T, et al. A population-based survey of Chronic Renal Disease In Turkey—the CREDIT study. *Nephrol Dial Transplant*. 2011; **26(6)**:1862-1871.
5. Turner JM, Bauer C, Abramowitz MK, et al. Treatment of chronic kidney disease. *Kidney Int*. 2012; **81(4)**:351-362.
6. Moncef, E, Kadiri MB, Nechba RB, et al. Factors predicting malnutrition in hemodialysis patients. *Saudi J Kidney Dis Transpl*. 2011; **22 (4)**: 695-704.
7. Sahathevan S, Khor BH, Ng HM, et al. Understanding Development of Malnutrition in Hemodialysis Patients: A Narrative Review. *Nutrients*. 2020;**12(10)**.
8. Zaki DS, Mohamed RR, Mohammed NA, et al. Assessment of malnutrition status in hemodialysis patients. *Clinical Medicine and Diagnostics*. 2019; **1**: 10-20.
9. Reber E, Gomes F, Vasiloglou MF, et al. Nutritional risk screening and assessment. *Journal of clinical medicine*. 2019; **8(7)**: 1065.
10. Ikizler TA, Burrowes JD, Byham-Gray LD, et al. KDOQI clinical practice guideline for nutrition in CKD: 2020 update. *American Journal of Kidney Diseases*. 2020; **76(3)**: 1-107.
11. de Mutsert R, Grootendorst DC, Boeschoten EW, et al. Subjective global assessment of nutritional status is strongly associated with mortality in chronic dialysis patients. *Am J Clin Nutr*. 2009; **89(3)**: 787-793.
12. Olcay Eminsoy İ, Kızıltan G, Akçıl Ok M, et al. Adaptation of the “7-Points Subjective Global Assessment (SGA-7P) into Turkish and Evaluation of SGA-3P Compliance with Pilot Study on Hemodialysis Patients. *J Nutr Diet*. 2019;**1-12**.
13. Mueller C, Compher C, Ellen DM. The American Society for Parenteral and Enteral Nutrition (A.S.P.E.N.) Board of Directors. A.S.P.E.N. Clinical Guidelines: Nutrition Screening, Assessment, and Intervention in Adults. *J Parenter Enter Nutr*. 2011; **35(1)**: 16-24.
14. Russell CA, Elia M. Nutrition Screening Surveys In Hospitals In The Uk, 2007-2011. *British Association For Parenteral And Enteral Nutrition*. 2014; **60**.

15. Lawson CS, Campbell KL, Dimakopoulos I, Dockrellet ME. Assessing the Validity and Reliability of the MUST and MST Nutrition Screening Tools in Renal Inpatients. *J Ren Nutr.* 2012; **22(5)**: 499-506.
16. Uyar S, Kök M, Unsal A, et al. The Validity of Malnutrition Universal Screening Tool (MUST) for Nutritional Screening in Hemodialysis Patients. *Turk J Nephrol.* 2019; **28(2)**: 109-113.
17. Jackson HS, MacLaughlin HL, Vidal-Diez A, et al. A new renal inpatient nutrition screening tool (Renal iNUT): a multicenter validation study. *Clin Nutr.* 2019; **38(5)**: 2297-2303.
18. Yeşilyurt S, Çapraz C. A Road Map for the Content Validity Used in Scale Development Studies. *Erzincan University Journal of Education Faculty.* 2018; **20(1)**: 251-264.
19. Charlson ME, Pompei P, Ales K, et al. A new method of classifying prognostic comorbidity in longitudinal studies: Development and validation. *J Chronic Dis.* 1987; **40(5)**: 373-383.
20. WHO [Internet]. Waist Circumference and Waist–Hip Ratio. Accessed June 12, 2022. <https://www.who.int/publications/i/item/9789241501491>.
21. Pekcan G. Beslenme Durumunun Saptanması. 2008; **726**: 67-141.
22. NHANES [Internet]. Anthropometry Procedures Manual. Accessed May 23, 2022. https://www.cdc.gov/nchs/data/nhanes/nhanes_07_08/manual_an.pdf.
23. Su CY, Lin JH, Chien TH, et al. Grip strength in different positions of elbow and shoulder. *Arch Phys Med Rehabil.* 1994; **75(7)**: 812-815.
24. Mathiowetz V, Weber K, Volland G, et al. Reliability and validity of grip and pinch strength evaluations. *J Hand Surg.* 1984; **9(2)**: 222-226.
25. Rakıcioğlu N, Nergiz R, Ozkan Ş. Evaluation of nutritional status and habits in hemodialysis patients. *Journal of Nutrition and Dietetics.* 2005; **33(1)**: 13-25.
26. Kondrup J, Allison SP, Elia M, et al. ESPEN Guidelines for Nutrition Screening. 2002. *Clin Nutr.* 2003; **22(4)**: 415-421.
27. Hassanin IA, Hassanein H, Elmenshawy P, et al. Malnutrition score and Body Mass Index as nutritional screening tools for hemodialysis patients. *Clin Nutr ESPEN.* 2021; **42**: 403-406.
28. Steiber A, Leon JB, Secker D, et al. Multicenter Study of the Validity and Reliability of Subjective Global Assessment in the Hemodialysis Population. *J Ren Nutr.* 2007; **17(5)**: 336-342.
29. Carrero JJ, Thomas F, Nagy K, et al. Global Prevalence of Protein-Energy Wasting in Kidney Disease: A Meta-analysis of Contemporary Observational Studies From the International Society of Renal Nutrition and Metabolism. *J Ren Nutr.* 2018; **28(6)**: 380-392.

30. Deniz Güneş B, Köksal E. Screening for malnutrition with malnutrition inflammation score and geriatric nutritional risk index in hemodialysis patients. *Hemodialysis International*. 2022; **26(4)**: 562-568.
31. Badrasawi M, Zidan S, Sharif I, et al. Prevalence and correlates of malnutrition among hemodialysis patients at hebron governmental hospital, Palestine: cross-sectional study. *BMC Nephrol*. 2021;**22(1)**:214.
32. Ikizler TA, Burrowes JD, Byham-Gray LD, et al. KDOQI Clinical Practice Guideline for Nutrition in CKD: 2020 Update. *American Journal of Kidney Diseases* 2020; **76(3)**:1-107.
33. Jayanama K, Putadechakun S, Srisuwarn P, et al. Evaluation of Body Composition in Hemodialysis Thai Patients: Comparison between Two Models of Bioelectrical Impedance Analyzer and Dual-Energy X-Ray Absorptiometry. Zemel MB, ed. *J Nutr Metab*. 2018.
34. Deger SM, Hung AM, Gamboa JL, et al. Systemic inflammation is associated with exaggerated skeletal muscle protein catabolism in maintenance hemodialysis patients. *JCI Insight* 2017;2(22).
35. Wang AY, Sea MM, Ho ZS, et al. Evaluation of handgrip strength as a nutritional marker and prognostic indicator in peritoneal dialysis patients. *Am J Clin Nutr*. 2005;81(1):79-86.
36. Birajdar N, Anandh U, Premlatha S, et al. Hand grip strength in patients on maintenance hemodialysis: An observational cohort study from India. *Indian J Nephrol*. 2019;29(6):393.
37. Başcı S, Özkök A, Kaçar M, et al. Handgrip Strength and Serum Uric Acid Levels are Associated with Dialysis-Malnutrition Scores in Chronic Hemodialysis Patients. *Turk Nephrol Dial Transplant*. 2017;26(03):323-332.
38. Karahan Yılmaz S, Rakıcıoğlu N. Assessment of Relationship Between Nutritional Status and Handgrip Strength in Hemodialysis Patients. *Galician Med J*. 2020; 27(3):202034.
39. Köksal E, Küçükerdönmez Ö. Detection of Obesity and Current Approaches; Baysal A, Baş M, Eds. *Yetişkinlerde Ağırlık Yönetimi*. 1 th. Ed. Seyrantepe, İstanbul; 2008; 35-42
40. Heidari B. C-reactive protein and other markers of inflammation in hemodialysis patients. *Caspian Journal of Internal Medicine*. 2013;(1): 611.
41. Vannini FD, Antunes AA, Caramori JC, et al. Associations between nutritional markers and inflammation in hemodialysis patients. *Int Urol Nephrol*. 2009;41(4):1003-1009.

Table 1. Various biochemical parameters and anthropometric measurements of hemodialysis patients

Variables	Median (IQR)
Body weight (post dialysis)	69.0 (59.0-69.5)
Height (m)	1.65 (1.59-1.75)
BMI (kg/m ²)	25.5 (21,9-28,9)
TST (mm) (n=260)	11.0 (7.4-17.0)
TST, women (mm) (n=125)	17.0 (12.0-20.0)
TST, men (mm) (n=135)	8.4 (6.2-11.0)
HGS (kg) (n= 260)	16.1 (10.0-23.2)
HGS, women (n= 125)	10.9 (7.5-15.3)
HGS, men (kg) (n=135)	22.3 (16,5-27,9)
Serum albumin (g/L)	3.7 (3.6-3.9)
Total protein (g/L)	6.8 (6.5-7.1)
C-reactive protein (mg/dL)	6.6 (2.7-12.7)
Creatinine (mg/dL)	2.6 (2.0-3.3)
Potassium (mEq/L)	3.5 (3.2-3.8)
Phosphor (mg/dL)	4.8 (4.0-5.5)
Calcium (mg/dL)	8.6 (8.2-9.0)
Total cholesterol (mg/dL)	163.0 (146.0-187.0)
Triglycerides (mg/dL)	136.0 (94.0-185.0)
HDL (mg/dL)	35.5 (31.0-43.3)
LDL (mg/dL)	91.0 (68.6-114.0)
Charlson Comorbidity Index	5.0 (3.0-6.0)

Abbreviations: BMI: Body Mass Index, TST: Triceps Skinfold Thickness, HDL: High Density Lipoprotein, HGS: Handgrip Strength, LDL: Low Density Lipoprotein

Table 2. The validity of the Renal Inpatient Nutrition Screening Tool (Renal iNUT) and Malnutrition Universal Screening Tool (MUST) according to clinical outcome category, following comparison with subjective global assessment in hemodialysis patient population ($n=260$)

	Renal iNUT		MUST		<i>p</i> -value
	iNUT=0 ($n=105$)	iNUT \geq 1 ($n=155$)	MUST=0 ($n=204$)	MUST \geq 1 ($n=56$)	
SGA (Gold standard)					
SGA A ($n=150$)	93	57	144	6	
SGA (B+C) ($n=110$)	12	98	60	50	
ROC Curve analysis					
AUC (95% CI)	0.755 (0.696 – 0.815)		0.707 (0.640 – 0.744)		
<i>p</i>-value	<.001		<.001		
Youden index (<i>J</i>)	0.511		0.415		
Statistical Diagnostic Measures					
Sensitivity (95% CI)	89 (82-94)		45 (36 – 55)		<.001 ¹
Specificity (95% CI)	62 (54-70)		96 (91 – 99)		<.001 ¹
Accuracy (95% CI)	73 (68 – 79)		75 (69 – 80)		
PPV (95% CI)	63 (55 – 71)		89 (78 – 96)		<.001 ²
NPV (95% CI)	89 (81 – 94)		71 (64 – 77)		<.001 ²
κ	0.484		0.444		

Abbreviations: AUC; area under the curve, 95% CI; 95% confidence intervals, PPV; positive predictive value, NPV; negative predictive value, κ ; kappa value

¹ McNemar test

² Weighted generalized score test

Table 3. Differences between Renal iNUT scores of anthropometric measurements and parameters ($n=260$)

	Renal iNUT categories			<i>p</i> -value
	Renal iNUT=0 ($n=105$)	Renal iNUT=1 ($n=96$)	Renal iNUT \geq 2 ($n=59$)	
Anthropometric measures				
BMI	27.94 \pm 5.20 ^a	25.81 \pm 5.23 ^b	22.90 \pm 4.32 ^c	<.001 ¹
MUAC	27.70 \pm 3.25 ^a	26.52 \pm 4.06 ^a	25.08 \pm 3.66 ^b	<.001 ¹
TST	13.12 \pm 7.18	12.70 \pm 7.04	12.45 \pm 6.41	.827 ¹
HGS	19.25(11.65–5.85) ^a	15.02(10.45–23.32)	12.30(7.72–18.65) ^b	.001 ²
Biochemical parameters				
CRP	4.15(1.7 – 10.82) ^a	7.90(3.92–13.10) ^b	6.60 (2.42 – 12)	.003 ²
Serum albumin	3.74 (3.57 – 3.96)	3.71 (3.52 – 3.87)	3.69 (3.57 – 3.84)	.648 ²
CCI	5 (3 – 6)	5 (3.75 – 6)	5 (4 – 6)	.657 ²

Abbreviations: BMI; Body Mass Index, MUAC; Middle Upper Arm Circumference, TST; Triceps Skinfold Thickness, HGS; Handgrip Strength, CRP; C Reactive Protein, CCI; Charlson Comorbidity Index

Data were expressed as mean \pm standard deviation or median (interquartile range).

Different small superscripts in each row indicate that statistically significant difference between groups after multiple comparison.

¹ One-Way ANOVA followed by Tukey HSD test

² Kruskal Wallis followed by Dunn test with Bonferroni correction




Figure 1. Flow diagram of the conduct of the study. Renal iNUT, Renal Inpatient Nutrition Screening Tool




Figure 2. Subjective Global Assessment (SGA), Renal Inpatient Nutrition Screening Tool (Renal iNUT) and Malnutrition Universal Screening Universal (MUST) for patients with low nutritional risk, increased nutritional risk and referral to a dietitian.




Figure 3. The Receiver Operating Characteristics (ROC) curves for the comparison of Renal iNUT and MUST methods in diagnosing of malnutrition, which is assessed using the SGA method. The reference line was colored with black; the ROC curve of Renal iNUT was colored with blue; ROC curve of MUST was colored with blue. DeLong's test revealed that there was no statistically significant difference between AUCs for methods of malnutrition screening tools ($p=.124$).




Figure 4. The box-plot of anthropometrical measurements and biochemical parameters in patients with hemodialysis. Horizontal lines in each box indicate the median. Data were represented as median with interquartile ranges. The One-Way ANOVA followed by Tukey HSD, and Kruskal-Wallis H test followed by Dunn post-hoc test with Bonferroni correction was used for comparisons between Renal iNUT categories. Abbreviations: BMI: Body Mass Index, MUAC: Middle Upper Arm Circumference, TST: Triceps Skinfold Thickness, HGS: Handgrip Strength, CRP: C Reactive Protein, CCI: Charlson Comorbidity Index.