

significant inverse association with dementia occurrence. These results underscore the importance of promoting regular physical activity as a potential preventive measure against dementia among older Brazilian adults.

FC30: Bridging the gap: an estimate of undetected dementia in Brazil and differences between genders, age groups and regions.

Authors: Andrew Christopher Miguel¹ MD; Matheus Ghossain Barbosa¹ MD, PhD; Sonia Brucki² MD, PhD; Paulo Caramelli, MD³ PhD; Jerson Laks, MD⁴ PhD; Ricardo Nitri² MD, PhD; Claudia Kimie Suemoto⁵ MD, PhD; Lucas Martins¹ MD; Cleusa P Ferri¹ MD, PhD.

1. Department of Psychiatry. Federal University of São Paulo (UNIFESP). São Paulo. SP. Brazil.
2. Department of Neurology. University of São Paulo Medical School (USP), São Paulo. SP. Brazil.
3. Department of Neurology. Federal University of Minas Gerais (UFMG). Belo Horizonte. MG. Brazil.
4. Department of Psychiatry. Federal University of Rio de Janeiro (UFRJ). Rio de Janeiro. RJ. Brazil.
5. Division of Geriatrics. University of São Paulo Medical School (USP). São Paulo. SP. Brazil.

Objectives: Despite the increasing number of people with dementia (PWD), detection remains low worldwide. In Brazil, PWD is expected to triple by 2050, and diagnosis can be challenging, contributing to high and growing rates of underdiagnosis. At the moment, there is no national estimate of the under detection or characteristics of its distribution according to gender, age and region. We aimed to estimate the proportion of PWD not diagnosed in relation to the estimated number of PWD.

Methods: The number of diagnosed individuals were estimated based on national records of the prescription of anticholinesterases drugs (AChE) in 2022 for the treatment of mild and moderate stages of Alzheimer's Disease (AD) held by the Unified Health System (SUS). Data were obtained from <ftp://ftp.datasus.gov.br> and drugs were dispensed according to the national clinical protocol. Studies from the national literature were consulted to estimate: (i) the number of people currently diagnosed with mild and moderate AD; (ii) the proportion of those who obtain AChE from SUS; (iii) the proportion of those who do not take AChE; and (iv) the proportion of AD related to other dementias. We assumed that the under-detection rate of AD would be similar to other dementias and 70% of the diagnosed AD individuals obtain AChE from SUS.

Results: More than 80% of the PWD 60+ are undetected (88.7%, 95% CI = 88.6–88.7). The poorest regions had higher rates (94.6% 95%, CI = 94.5–94.6) than the richest (84.8%, 95% CI = 84.7–84.8). Men had higher rates (89.8%, 95% CI = 89.7–89.9) than women (87.4%, 95% CI = 87.4–87.5). The youngest age group (60-64) had the highest rate (94.6%, 95% CI = 94.5–94.7) which decreased until 85–89 (84.3%, 95% CI = 84.2–84.4), before increasing again to 91.1% (95% CI = 91.0–91.2) among 90+.

Conclusions: Dementia under detection in Brazil is among the highest in the world. Fast populational aging and the highest rates among the youngest individuals are of concern as it may be related to late diagnosis. Gender and regional disparities also need to be considered when developing health policies.

FC31: Experiences And Meanings of Alzheimer's Diagnosis In The Early Stage In Puerto Rico

Authors: Ana Gratacos; Roberto González, Zaira Mateo, Heriberto Acosta

Summary: Studies suggest that people in the early stages of Alzheimer's dementia (AD), a disease first described in 1906, do not have sufficient support to cope with the disease process after a diagnosis (Solomon, 2016). AD is