

P8: Depression, diabetes, and healthcare services use – Brazilian Longitudinal Study of Aging (ELSI-Brazil, 2015-2016)

Authors: Ana Keinert, MD, Matheus Ghossain Barbosa, PhD, Maria Fernanda Lima-Costa, Cleusa Pinheiro Ferri

Objectives: Low- and middle-income countries face increasing burden of noncommunicable chronic diseases due to rapid population aging. The Objectives of this study is to estimate the association of co-occurring depression and diabetes with healthcare services utilization in the Brazilian population aged 50 years and older.

Methods: This is an analysis using baseline data of the ELSI-Brazil study. Measurements used were self-reported previous diagnosis for diabetes and a cut-off point of 4 on the CES-D-8 score for depression. Any medical consultations, specialist consultations and hospitalizations in the previous 12 months were assessed for measuring health services use. Logistic regression was used to estimate the association of having depression, diabetes, or both, compared to having none of the conditions with healthcare services utilization.

Results: In a final sample of 8303 participants, the adjusted odds ratios (95%CI) in the depression only group (27.9%), in the diabetes only group (9.7%), and in the depression and diabetes group (5.8%), for any medical consultations were 1.2 (1.01-1.42), 3.39 (2.4-4.79) and 3.12 (1.82-5.35); for specialist consultations, 1.05 (0.91-1.21), 1.34 (1.07-1.68) and 1.13 (0.86-1.49); and for hospitalizations, 1.42 (1.1-1.84), 1.42 (1.02-1.96) and 3.1 (2.12-4.54), respectively. All models were adjusted for sex, age, education, marital status, insufficient physical activity, current drinking and smoking, obesity and number of other diseases and conditions.

Conclusions: Those with diabetes seemed more likely to have any or specialized medical consultations. However, those with the depression and diabetes comorbidity were more likely than any other group to have been admitted to a hospital in the last year, which is distressing and costly. Screening for depression could be incorporated into diabetes usual care to reduce related complications and hospitalizations.

Funding: This study did not receive any funding. Authors: Ferri and Lima-Costa are recipient of CNPq research productive fellowship.

Keywords: depression; diabetes; comorbidity; health services use; aging.

P9: Computer-based Neuropsychological Rehabilitation in Latin America

Authors: Alvarado Rodas Andrea Cristina, Psy.D., M.Sc., PhD student^{1,2}, Orellana Palacios Hernán Mateo, M.D., M.Sc., PhD student ^{1,2}

1. Universidad de Salamanca, España.

2. Centro MAIOR Ecuador. Centro de Especialidades del Adulto Mayor.

Summary: Globally, older adults grow faster than the rest of the population (United Nations et al., 2019). Ecuador predicts 2054 this group will represent 18% of the population. Cognitive and functional impairment affects quality of life of individuals and their families. Conditions such as dementia have increased, with almost 100,000 people affected in Ecuador.