

Weeds

VOLUME 8
OCTOBER 1960
NUMBER 4

Journal of the Weed Society of America

Weeds

Issued Quarterly by the Weed Society of America

- C. J. Willard, Editor, Department of Agronomy, The Ohio State University, 1827 Neil Ave., Columbus 10, Ohio
- W. C. JACOB, Business Manager, Department of Agronomy, University of Illinois, Urbana, Illinois

EDITORIAL COMMITTEE

- O. C. Lee, Department of Botany and Plant Pathology, Purdue University, Lafayette, Indiana
- R. A. Peters, Department of Plant Science, University of Connecticut, Storrs, Connecticut
- W. C. ROBOCKER, Crops Research Division, ARS, USDA, State College of Washington, Pullman, Washington
- E. G. Rodgers, Department of Agronomy, University of Florida, Gainesville, Florida
- D. W. STANIFORTH, Department of Botany and Plant Pathology, Iowa State University, Ames, Iowa

WEEDS is a quarterly journal published by the Weed Society of America. The subscription price is \$6.00 yearly for four issues; single copies \$1.50. Address all communications regarding subscriptions, advertising, and reprints to W. C. Jacob, Department of Agronomy, University of Illinois. All checks, money orders, and other remittances should be made payable to the Weed Society of America. The editorial office is located at Department of Agronomy, The Ohio State University, 1827 Neil Avenue, Columbus 10, Ohio. Inquiries concerning manuscripts and other material for publication should be addressed to the editorial office. Printing is by the W. F. Humphrey Press, Inc., Geneva, New York.

Entered as second-class matter at the post office at Urbana, Illinois with additional entry at Geneva, New York.

Table of Contents

	Page
Weed Control Research—Past, Present, and Future. A. S. Crafts	535
Practices. H. Gysin	541
Research and Organizational Needs in the Field of Weed Control. M. W. Parker	556
Weed Control—Its Status and Organization in Great Britain, E. K. Woodford	561
A Conference Was Born, Walter S. Ball.	573
Growth of Witchweed (Striga asiatica) as Affected by Soil Types and Soil and Air Temperatures, E. L. Robinson	576
The Effect of Temperature on the Herbicidal Activity and Translocation of Arsenicals. Charles B. Rumburg, Ralph E. Engel, and William F. Meggitt	582
The Effects of Four Herbicides on Microorganisms in Nine Oregon Soils. Purna Chandra, William R, Furtick, and Walter B, Bollen	589
Persistence and Movement of CDAA and CDEC in Soil and the Tolerance of Corn Seedlings to these Herbicides, R. L. Gantz and F. W. Slife	599
Simazine for Annual Weed Control in Cranberries, Malcolm N. Dana	.607
The Influence of Concentration of Fumigant and Time of Exposure on the Killing of Dormant Imbibed Seeds. Stanley J. Pieczarka and G. E. Warren	612
Weed Control in Cotton Over a Ten-year Period by Use of the More Promising Materials and Techniques. Vernon C. Harris	616
Kill of Blue Oak and Poison Oak by Aircraft Spraying with Phenoxy Herbicides. O. A. Leonard and C. E. Carlson	625
Quackgrass Control in Potato Production, D. R. Isleib	631
Factors Affecting the Competitive Ability of Foxtail Barley (Hordeum jubatum). H. P. Cords	636
The Relative Toxicity of Three Boron Compounds and Sodium Chlorate Used as Soil Sterilants. E. G. Scott	645
Histoautoradiographic Studies of Herbicidal Translocation. M. A. Radwan, C. R. Stocking, and H. B. Currier	657
Relative Toxicity of Extracts from Vegetative Organs of Quackgrass to Alfalfa. J. H. Ohman and Thor Kommedahl	666
Simazine for Controlling Weeds in Fruit Tree and Grape Plantings. R. P. Larsen and S. K. Ries	671
Brief Papers	
Where We Stand With Witchweed. W. G. Westmoreland	678
Control of Horsenettle in Pastures. W. B. Albert	680
Effect of Tansy Mustard (Descurainia intermedia) on Moisture Storage During Fallow. A. F. Wiese	683
News and Notes	686
Membership List, Weed Society of America	687
Sustaining MembersIndex to Volume 8	703
Index to Volume 8	704
Bibliography of Weed Investigations, January to March, 1960	710
Advertisers Index	

For Agriculture and Industry . . .

Du Pont WEED KILLERS

offer new economies and efficiency in killing weeds, grass and brush

KARMEX® diuron for weed control in many crops; irrigation and drainage ditches; and industrial areas.

KARMEX® DL for pre-emergence weed control in cotton.

TELVAR® monuron for weed control in many crops; irrigation and drainage ditches; and industrial areas.

KLOBEN® neburon for weed and grass control in nursery plantings and tomatoes.

DYBAR® fenuron—a pelleted product for dry application to the soil for control of brush in fence rows, drainage ditches, utility and railroad right-of-ways, and other non-cultivated areas.

TRYSBEN® 200, a new weed killer based on trichlorobenzoic acid, for control of bindweed and other noxious weeds; mixed broadleaf weeds; and certain woody vines and brush.

AMMATE® X for long-term, low-cost brush control.



Better Things for Better Living...through Chemistry





& CHEMICAL CORPORATION

630 Shatto Place, Los Angeles 5, California

= NEWS ABOUT

VAPAM®=

Summary of Application Methods

After careful soil preparation, a wide range of application methods can be used for effective control of weeds, germinating weed seed, nematodes, and soil fungi with VAPAM fumigant. Most of the methods do not require special equipment.

For Seed Beds, Plant Beds, Lawns and Other Limited Areas

—A sprinkling can, hose proportioner or rotary tiller are all effective. In some cases, it might be practical to use soil injection.

Field Application to Beds or Rows—Soil injection, beddingover or soil covering are the quickest and easiest methods for most growers. Basin flooding is the recommended method for treating tree replant sites.

For Treating Entire Fields—Soil injection, sprinkler irrigation, and check or flood irrigation are recommended.

Full details on these application methods, plus the treatment of potting soils, are on the VAPAM label.

Copy of Label Available

Write to Stauffer for a copy of the label, or for further information on VAPAM. Address your inquiry to Stauffer Chemical Company, Agricultural Chemicals Division, 380 Madison Ave., New York 17, N. Y., or 636 California St., San Francisco 8, California St., San Fr

®VAPAM is Stauffer Chemical Company's trade-mark (registered in principal countries) for sodium methyl dithiocarbamate, a soil fumigant.





ATLACIDE: Safer chlorate weed killer...widely used for non-selective eradication of bindweed, Canada thistle, quack grass, Johnson grass and other tough perennials. Kills roots...discourages regrowth. Applied dry or dissolved in water.

ATLACIDE -2,4-D: Offers dual killing action of sodium chlorate and 2,4-D.

CHLOREA: A combination of sodium chlorate, borate and monuron in powder form. Kills weeds and grasses. Combines the proven effectiveness of chlorate on deep-rooted weeds with the soil-surface action of monuron on shallow-rooted grasses and annual seedling growth. Lasting residual effect inhibits new growth. Does not create a fire hazard when used as directed. Applied dry or as a water-mixed spray.

CHLOREA GRANULAR Similar to Chlorea, but a "pelletized" material. No mixing...easy to apply by hand or with mechanical spreader.

CHLORAX "40": A composition of sodium chlorate and borate...for weed and grass control. Has lasting residual effect. Does not create a fire hazard. Applied dry or as a spray.

ATLAS "A": A 40% sodium arsenite solution (4 lbs. arsenic trioxide per gal.). Destroys vegetation and algae in ponds and lakes. Controls certain turf weeds. Used as general weed killer and to kill trees and stumps. Also used to kill potato vines prior to harvesting.

SODIUM ARSENITE: A powder containing 75% arsenic trioxide. Used for the same purposes as Atlas "A". Applied dry or as a spray.

CHIP-CAL: New low-lime calcium arsenate formulation in *granular* and spray *powder* forms. Used as crabgrass preventatives for established turf. Applied any time before crabgrass seed starts to germinate.

2,4-D & 2,4,5-T: 2,4-D Amine and 2,4-D Ester liquids; 2,4-D Ester dusts; Low Volatile 2,4,5-T and Brush Killer.

METHOXONE: Contains MCP sodium salt. Used for control in small grains, flax, rice and grass. Controls same weeds as 2,4-D; considered safer for selective spraying.

BUTOXONE: A liquid containing 4–(2,4–DB). For selective control of many broadleaf weeds in established alfalfa and certain other legumes.

Write for Weed Control Booklets

CHIPMAN CHEMICAL COMPANY, INC.

Chicago, Ill.

Palo Alto, Calif.

Pasadena, Tex.

Portland, Ore.

Pasadena, Tex.

Bessemer, Ala.

Manufacturers of Weed Killers Since 1912



Supplied in a full range of interchangeable orifice tip and strainer sizes to meet every capacity requirement. TeeJet Spray Nozzles for Weed Control by spraying make it possible to take maximum advantage of the chemical and sprayer unit. TeeJet nozzles are precision built and provide a flat spray with uniform distribution. Atomization is properly controlled to give coverage with an absolute minimum of driftage. Patented tip design, with set-back orifice opening protects precision orifice from accidental damage. TeeJet spray nozzles are built for use on spray booms and portable sprayers.

OFF-CENTER SPRAY NOZZLES

Spraying Systems Spray Nozzles with TeeJet tips are supplied in a variety of special body types to meet any unusual spraying requirement. For example, one type of off-center spray nozzle with swivel body provides a flat spray up to 35 feet wide for spraying areas with a single nozzle, that are not accessible with a boom.

SUPPLEMENTARY EQUIPMENT

Complete accessories relating to nozzle use are supplied. These include strainers, special nozzle fittings, and hand valve equipment.

TEEJET SPRAY NOZZLE male pipe connection



INTER-CHANGE-ABLE ORIFICE TIPS flat and cone spray types

TeeJet Spray Nozzles are supplied for Weed Control... as well as all other types of agricultural spraying. For complete information and reference data write for Catalog 30

PRAYING SYSTEMS CO.

Engineers and Manufacturers

3296 RANDOLPH STREET

BELLWOOD, ILLINOIS