




The Nutrition Society Irish Section Conference 2023 was held at the Technological University of the Shannon on 14th–16th June 2023

Conference on ‘Understanding the role of sex and gender in nutrition research’ Symposium two: Outcomes: Impact of sex and gender on risk of non-communicable disease

Calcium and vitamin D homoeostasis in male fertility

Sam Kafai Yahyavi^{1,2,†}, Ida Marie Boisen^{1,2,†}, Zhihui Cui^{1,2}, Mads Joon Jorsal^{1,2}, Ireen Kooij^{1,2}, Rune Holt^{1,2}, Anders Juul^{3,4,5} and Martin Blomberg Jensen^{1,2,*} 

¹Division of Translational Endocrinology, Department of Endocrinology and Internal Medicine, Copenhagen University Hospital – Herlev and Gentofte, Herlev, Denmark

²Group of Skeletal, Mineral, and Gonadal Endocrinology, Department of Growth and Reproduction, Copenhagen University Hospital – Rigshospitalet, Copenhagen, Denmark

³Department of Growth and Reproduction, Copenhagen University Hospital – Rigshospitalet, Copenhagen, Denmark

⁴International Centre for Research and Research Training in Endocrine Disruption of Male Reproduction and Child Health (EDMaRC), Copenhagen University Hospital – Rigshospitalet, Copenhagen, Denmark

⁵Department of Clinical Medicine, Copenhagen University Hospital – Rigshospitalet, Copenhagen, Denmark

Calcium and vitamin D have well-established roles in maintaining calcium balance and bone health. Decades of research in human subjects and animals have revealed that calcium and vitamin D also have effects on many other organs including male reproductive organs. The presence of calcium-sensing receptor, vitamin D receptor, vitamin D activating and inactivating enzymes and calcium channels in the testes, male reproductive tract and human spermatozoa suggests that vitamin D and calcium may modify male reproductive function. Functional animal models have shown that vitamin D deficiency in male rodents leads to a decrease in successful mating and fewer pregnancies, often caused by impaired sperm motility and poor sperm morphology. Human studies have to a lesser extent validated these findings; however, newer studies suggest a positive effect of vitamin D supplementation on semen quality in cases with vitamin D deficiency, which highlights the need for initiatives to prevent vitamin D deficiency. Calcium channels in male reproductive organs and spermatozoa contribute to the regulation of sperm motility and capacitation, both essential for successful fertilisation, which supports a need to avoid calcium deficiency. Studies have demonstrated that vitamin D, as a regulator of calcium homoeostasis, influences calcium influx in the testis and spermatozoa. Emerging evidence suggests a potential link between vitamin D deficiency and male infertility, although further investigation is needed to establish a definitive causal relationship. Understanding the interplay between vitamin D, calcium and male reproductive health may open new avenues for improving fertility outcomes in men.

Key words: Vitamin D: Calcium: Fertility: Male reproduction

Abbreviations: 1, 25(OH)₂D₃, 1, 25-dihydroxyvitamin D₃; 25OHD, 25-hydroxyvitamin D; CaSR, calcium-sensing receptor; CYP, cytochrome P450; PMCA, plasma-membrane calcium ATPase; PTH, parathyroid hormone; TRPV, transient receptor potential vanilloid; VDR, vitamin D receptor.

*Corresponding author: Martin Blomberg Jensen, email martin.blomberg.jensen@regionh.dk

†These authors contributed equally to the study.

Male reproduction is of increasing interest and concern worldwide due to the growing issues with reproductive health and low fertility rates⁽¹⁾. According to the UN, the average global fertility rate stood at 2.3 births per woman in 2021, falling from about 5 births per woman in the mid-twentieth century⁽²⁾. Male infertility contributes to the problem in almost 50% of infertile couples, with up to 30% of all cases being attributed to a male factor as the sole cause^(3–5). Numerous causes can influence male reproductive function. These include genetic mutations, environmental aspects, lifestyle choices, medical conditions or medications^(6–9). In recent years, the potential impact of nutritional factors on male reproductive health has gained significant attention. Both vitamin D deficiency and low-ionised calcium have been linked with impaired semen quality⁽¹⁰⁾. While it can be challenging to directly attribute instances of male infertility to dietary or lifestyle changes, these factors can contribute to the problem. Modern dietary habits have seen a shift towards processed foods⁽¹¹⁾, which are often low in essential nutrients and can lead to nutrient deficiencies that may have a negative influence on fertility.

Vitamin D is a steroid hormone essential for various physiological processes in many different organs beyond its classical role in maintaining calcium homeostasis and bone health⁽¹²⁾. In the past 20 years, there has been a great interest in exploring the beneficial effects of vitamin D on human health and in various organs during health and disease^(13–15). It has been suggested to be involved in several aspects of male reproductive function, such as sperm production, sperm motility and hormonal regulation^(16–18). Generally, vitamin D deficiency is defined as serum 25-hydroxyvitamin D (25OHD) levels below 25 or 30 nmol/l and vitamin D insufficiency as levels below 50 nmol/l. The active vitamin D metabolite, 1,25-dihydroxyvitamin D₃ (1,25(OH)₂D₃) (calcitriol), exerts its effects through the vitamin D receptor (VDR), which is present in various reproductive tissues including testes, epididymis and prostate⁽⁹⁾. The VDR, upon activation by 1,25(OH)₂D₃, potentially regulates the expression of genes involved in steroidogenesis, spermatogenesis and sperm maturation⁽⁸⁾.

Vitamin D regulates serum calcium concentrations by enhancing intestinal absorption, utilising bone storage, and increasing renal reabsorption⁽⁸⁾. These actions are involved in the tight regulation of calcium in the body which is essential for almost all physiological functions. Calcium is not only crucial for skeletal integrity but is also involved in intracellular signalling pathways, muscle contraction and hormonal regulation. In the male reproductive system, calcium is essential for sperm motility, sperm capacitation and the acrosome reaction, which are crucial steps for successful fertilisation⁽¹⁰⁾. Disturbances in calcium homeostasis can affect sperm function and contribute to male infertility^(10,19). Therefore, adequate levels of both vitamin D and calcium have been proposed to be important for optimal male reproductive health.

Understanding the significance of vitamin D and calcium in male reproductive health has the potential to guide clinical practice, form interventions and improve

reproductive outcomes for men. As such, this review summarises the current understanding of the effects of vitamin D and calcium on male reproductive function.

Male reproductive tract

Male reproductive organs ensure the production, storage and delivery of motile and competent spermatozoa and the synthesis and secretion of male sex steroids. The testes are responsible for the production of spermatozoa through a complicated process known as spermatogenesis, which occurs in the seminiferous tubules⁽⁸⁾. Spermatogenesis involves both mitosis and meiosis to complete the development of mature spermatozoa from spermatogonial stem cells⁽²⁰⁾. Spermatogenesis relies on the interplay between various cell types, including germ cells, Sertoli cells, peritubular cells and Leydig cells. Within the seminiferous tubules, the specialised Sertoli cells, in conjunction with peritubular cells, create the microenvironment that allows germ cells to develop from diploid spermatogonia to haploid spermatids⁽²¹⁾. They provide physical support and nutrition, and regulate the milieu required for proper sperm development. Various genetic and endocrine factors tightly regulate the functions of Sertoli and peritubular cells, with follicle-stimulating hormone and testosterone being prominent regulators necessary for optimal spermatogenesis⁽²²⁾. Leydig cells, located in the interstitial spaces between the seminiferous tubules, produce testosterone in response to luteinising hormone stimulation. This ensures a high intratesticular testosterone concentration that can be up to 100 times higher than in serum^(23,24). Testosterone has a significant impact on fetal testis development, the descent of the testes into the scrotum and the maturation of Sertoli cells during puberty. The maturation of Sertoli cells marks the initiation of spermatogenesis and coincides with the cessation of their proliferation during puberty. Mature Sertoli cells cannot proliferate, thus the quantity of Sertoli cells established during early life determines the capacity for sperm production in adulthood⁽²¹⁾. Consequently, the number of sperm present in a semen sample is, to a certain extent, determined during fetal and early life.

After successful spermatogenesis, the spermatozoa are immotile and undergo passive transportation to the epididymis through the efferent ducts. In the epididymis, the spermatozoa undergo final maturation and become motile. The epididymal fluid composition is tightly regulated and differs markedly as the spermatozoa transit through the caput, corpus and cauda of the epididymis. During ejaculation, spermatozoa meet secretions from the prostate and seminal vesicles that may also influence sperm function.

Vitamin D

Physiology and metabolism

The inactive form of vitamin D, cholecalciferol, is synthesised in the skin following the conversion of

7-dehydrocholesterol by UV B radiation from the sun^(25,26). Cholecalciferol can also be absorbed from the diet or via supplementation. In detail, cholecalciferol must undergo two hydroxylations to form the active form of vitamin D, 1,25(OH)₂D₃ that can bind and activate the VDR^(27–29). These steps involve the microsomal cytochrome P450 (CYP) enzymes. The first step is 25-hydroxylation by hepatic CYP2R1 which forms the prohormone 25OHD (calcidiol) followed by 1 α -hydroxylation by renal CYP27B1 to form 1,25(OH)₂D₃, whereas CYP24A1 inactivates 25OHD and 1,25(OH)₂D₃⁽³⁰⁾. Parathyroid hormone (PTH) is the main inducer of renal CYP27B1 expression and thus the main inducer of circulating 1,25(OH)₂D₃ and a powerful mobiliser of calcium from the skeleton⁽³¹⁾. 1,25(OH)₂D₃ binds and activates the VDR which forms a heterodimer with the retinoid X receptor. This complex recognises vitamin D response elements in the promoter region of target genes and regulates transcription⁽³⁰⁾. The DNA-binding domain of the VDR mediates the transcriptional effects, but VDR also has an alternative ligand-binding pocket that mediates rapid non-genomic effects by regulating second messenger systems. In clinical practice, vitamin D status is determined by measuring serum levels of 25OHD and not 1,25(OH)₂D₃, as 25OHD correlates with bone mass density, serum calcium level and PTH secretion^(32,33). However, cellular responsiveness to circulating vitamin D metabolites goes beyond VDR expression as the presence of activating and inactivating enzymes locally in the target organs influences the availability of substrates for VDR and thereby influences the effect of vitamin D on the target tissue^(34,35).

Expression of vitamin D regulating enzymes and vitamin D receptor in the male reproductive system

Studies have consistently found mRNA and protein expression of the VDR and the vitamin D activating and inactivating enzymes in various cell types of the gonads and male reproductive tract⁽⁹⁾. In most studies, VDR is co-expressed with the activating enzyme, CYP27B1, and the deactivating enzyme, CYP24A1. This co-expression occurs throughout the reproductive tract, including in some Sertoli cells, most germ cells, spermatozoa, Leydig cells and the epithelial cells lining the tract^(36–42). The local presence and activity of the enzymes play a crucial role in regulating the concentration of 1,25(OH)₂D₃ within cells and consequently activating the VDR.

In adult testes, the VDR and metabolising enzymes are primarily expressed in human germ cells, with a marked expression in spermatogonia, spermatocytes and spermatids in both human subjects and rodents^(18,43). Studies have also found VDR expression in both the nucleus and cytoplasm of primary cultures of immature Sertoli cells, in fetal Sertoli cells and in the immature mouse Sertoli cell line TM4 where 1,25(OH)₂D₃ is known to mediate fast non-genomic effects^(44–50). The expression levels of CYP2R1, CYP27B1 and CYP24A1 in the testes indicate that the cells are equipped to

respond to and regulate the effects of vitamin D metabolites^(8,18). Furthermore, the effects of cholecalciferol or calcitriol treatment on gonadal function may vary depending on the specific expression pattern of these enzymes. Cholecalciferol can be activated locally, potentially eliciting a response, whereas calcitriol is already active and immediately triggers responses or undergoes inactivation. This suggests that the local regulation of vitamin D is important for spermatogenesis and sperm function. The expression of VDR in germ cells suggests that calcitriol may play a role in sperm function⁽⁵¹⁾. Multiple studies have reported the presence of VDR in specific regions of mature human spermatozoa, including the post-acrosomal part of the head, the neck region and the midpiece^(18,52). Additionally, the vitamin D activating enzymes are also expressed in spermatozoa, and the inactivating enzyme CYP24A1 exhibits a distinct expression pattern at the annulus. Notably, the expression levels of VDR and CYP24A1 are higher in spermatozoa from healthy men compared with infertile men⁽⁵³⁾, which supports that vitamin D may be beneficial for sperm function.

VDR expression has also been demonstrated in Leydig cells, suggesting a potential direct impact of vitamin D on steroidogenesis and thereby also an indirect effect on spermatogenesis and male fertility. Recent studies have detected VDR at the protein level in fetal and adult Leydig cells in human subjects, roosters and mice^(18,38,54,55). Moreover, VDR, CYP27B1 and CYP24A1 are all expressed in the epididymis, prostate and seminal vesicles^(18,37,38), which highlights a potential direct effect throughout the male reproductive system.

Animal models: vitamin D deficiency and male fertility

Numerous animal models have been utilised to investigate the relationship between vitamin D and male fertility (defined as successful conception). An overview of the models is shown in Table 1. The first association was found in a study exploring vitamin D deficiency in rats⁽⁵⁶⁾. Vitamin D deficiency in male rats led to a 45% decrease in successful mating, defined as the presence of sperm in the vaginal tract, compared with rats with sufficient vitamin D levels. Additionally, rats with vitamin D deficiency had 71% fewer pregnancies compared with vitamin D-sufficient rats⁽⁵⁶⁾. When the rats were supplemented with calcium, the fertility potential was restored, implying an indirect effect of hypovitaminosis D leading to hypocalcaemia and secondarily reduced fertility⁽⁵⁷⁾. However, after reassessment of the data by adjusting for the mating period and male:female ratio, and by determining the time to pregnancy, it was evident that the rats still had lower pregnancy rates after insemination with sperm from normocalcemic vitamin D-deficient rats than normocalcemic vitamin D-replete rats⁽⁵⁷⁾. This indicates that vitamin D itself is important for semen quality and that the impaired fertility caused by vitamin D deficiency cannot be fully reversed by correcting hypocalcaemia alone⁽⁹⁾. Newer animal studies on vitamin D deficiency have supported the negative effects of inadequate vitamin D levels on semen quality and

Table 1. Six studies investigating the relationship between vitamin D deficiency and fertility in rodents

Subgroup	Fertility	Sperm count	Sperm motility	Sperm morphology	Reference
Naval Medical Research Institute (NMRI) mice*	↓	N.D.	↓	→	Shahreza <i>et al.</i> ⁽⁶²⁾
Sprague Dawley rats	N.D.	↓	↓	↓	Zamani <i>et al.</i> ⁽⁶¹⁾
Sprague Dawley rats	→	N.D.	↓	N.D.	Merino <i>et al.</i> ⁽⁶⁴⁾
Institute of Cancer Research (ICR) mice	→	↓	N.D.	N.D.	Fu <i>et al.</i> ⁽⁶⁰⁾
Sprague Dawley rats	↓	N.D.	N.D.	N.D.	Uhland <i>et al.</i> ⁽⁵⁷⁾
Sprague Dawley rats	↓	N.D.	N.D.	N.D.	Kwieceński <i>et al.</i> ⁽⁵⁶⁾

N.D., not determined.

Animal models made vitamin D-deficient either by diet, knock-out of the vitamin D receptor or the 1 α -hydroxylase (*Cyp27b1*).

*Phenotype reported for prolonged vitamin D-deficient mice; ↓, negatively associated; →, no association.

male fertility. In general, the low fertility rates in vitamin D-deficient male rodents seem to be caused by impaired sperm count, motility and occasionally poor sperm morphology^(58,59).

Dietary vitamin D-deficient mice have also been shown to have lower testicular weight and fewer spermatozoa in the cauda epididymis compared with controls⁽⁶⁰⁾. Another study revealed adverse impacts on semen quality in vitamin D-deficient rats when compared with control groups. These effects included reductions in germ cell numbers, sperm count, sperm morphology, motility and viability. Furthermore, a decrease in serum testosterone levels was also observed in vitamin D-deficient rats⁽⁶¹⁾. Duration and severity of the deficiency may be an important factor in reproductive function. Vitamin D deficiency through one spermatogenic cycle induced a significant increase in sperm DNA fragmentation but no significant effect on other semen parameters⁽⁶²⁾. In contrast, prolonged vitamin D deficiency through two spermatogenic cycles resulted in a modest but significant decrease in sperm motility. Furthermore, different degrees of vitamin D deficiency have been induced in mice, and no effect was observed on sperm concentration and sperm viability, but sperm morphology was decreased in the severe deficiency group⁽⁶³⁾. In another study, DNA fragmentation has also been shown to increase significantly in spermatozoa of vitamin D-deficient rats, which could explain the impaired fertility⁽⁶⁴⁾. Combined, these studies indicate that vitamin D is involved in the regulation of male fertility in rodents, suggesting a potential link between vitamin D status and reproductive health.

Human studies

Many of the cross-sectional studies investigating the link between vitamin D status and testicular function have been reviewed and discussed before, and the link critically depends on the fraction of men with low vitamin D status in the study population⁽⁶⁵⁾. Recent years have seen a surge in human intervention trials with vitamin D and male fertility outcomes. In this review, we have evaluated six randomised clinical intervention studies, with four of them being conducted after 2020. An overview of the trials is shown in Table 2. In the first randomised controlled trial from 2014, 86 infertile men with idiopathic oligoasthenozoospermia were included and

randomised to receive either a 3 month supplementation of cholecalciferol (5 μ g daily) and calcium (600 mg daily), or placebo⁽⁶⁶⁾. This study found a significant increase in the number of progressive motile sperm and a higher pregnancy rate in the vitamin D *v.* placebo-treated men. These findings are supported by the most recent randomised clinical trial of 120 men conducted in 2022 on Indian men⁽⁶⁷⁾. They found an increased sperm volume, progressive motility, total motility and sperm count in men treated with cholecalciferol (100 μ g daily for 10 weeks) when compared with the placebo group. The study does not report the serum 25OHD levels before or after treatment, so it is not possible to determine whether the effect was found in vitamin D insufficient, deficient or sufficient men.

In 2018, the biggest study to date was conducted and contradicted these findings⁽¹⁷⁾. In the Copenhagen bonegonadal study, 330 Danish infertile men with vitamin D insufficiency (defined as serum 25OHD <50 nm/l) were randomised to receive either a high single-dose cholecalciferol supplementation (7 500 μ g) followed by daily supplementation of cholecalciferol (35 μ g) and calcium (500 mg daily), or placebo. The study showed that supplementation did not affect semen parameters or live birth rates when comparing the groups. The spontaneous pregnancy rate tended to be higher in couples in which the men were treated with vitamin D and calcium compared with couples in which the men were in the placebo group, but the results were not statistically significant. However, subgroup analysis of oligospermic men did demonstrate an increased live birth rate in the treatment group (36 *v.* 18%). One criticism of the study is the use of a high-loading dosage that could lead to high CYP24A1 activity locally in many organs and thus inactivate vitamin D activity in the gonads. The overall negative findings of the study are consistent with three other recent randomised clinical trials conducted on Iranian men^(68–70). One trial with results published in 2020, investigated the effects of vitamin D supplementation in sixty-two infertile men with impaired semen quality and vitamin D insufficiency (defined as serum 25OHD <50 nm/l). The men were randomised to receive either cholecalciferol (1 250 μ g once weekly) for 8 weeks, followed by a maintenance dose (1 250 μ g once) lasting the final 4 weeks, or placebo treatment. The same study design was used by another Iranian research group from 2021, conducted on forty-four men with asthenozoospermia⁽⁶⁸⁾. Both studies

Table 2. Interventional studies of vitamin D and semen quality in infertile men

Population	N, treatment	N, control	Age	BMI	25OHD status	Vitamin D ₃ dose	Fertility	Sperm count	Sperm motility	Sperm morphology	Reference
Indian	60	60	34(sd 8)	N.D.	N.D.	100 µg daily for 10 weeks	N.D.	↑	↑	↑	Padmapriya <i>et al.</i> ⁽⁶⁷⁾ Gheflati <i>et al.</i> ⁽⁶⁸⁾
	20	20	33(sd 1)	24.5(sd 0.9)	<75 nw/l	1 250 µg weekly for 8 weeks, followed by 1 250 µg once for final 4 weeks	N.D.	↑	↑	↑	
Iranian	43	43	35(sd 5)	28.4(sd 3.0)	<75 nw/l	100 µg daily for 12 weeks Maghsoumi-Norouzabad <i>et al.</i> ⁽⁶⁹⁾	→	→	↑	→	Amini <i>et al.</i> ⁽⁷⁰⁾
Iranian	30	32	35(sd 5)	25.7(sd 1.9)	<75 nw/l	1 250 µg weekly for 8 weeks, followed by 1 250 µg once for final 4 weeks	→	→	→	→	
Danish	164	166	35(sd 7)	26.5(sd 4.2)	<50 nw/l	7 500 µg as initial dose, followed by 35 µg and 500 mg calcium daily for 150 days	↑	→	→	→	Blomberg Jensen <i>et al.</i> ⁽¹⁷⁾
Chinese	43	43	N.D.	N.D.	N.D.	5 µg and 600 mg calcium daily for 12 weeks	↑	↑	↑	N.D.	Deng <i>et al.</i> ⁽⁶⁶⁾

25OHD, 25-hydroxyvitamin D; N, cohort size; N.D., not determined. Six human randomised controlled studies investigating the relationship between serum vitamin D levels and semen quality in human subjects were evaluated. ↓, decreased with supplementation; ↑, increased with supplementation; →, no changes.

used vitamin D-deficient men (defined as serum levels <75 nm/l) and found no effect of cholecalciferol supplementation on semen parameters. A third Iranian study published in 2021 evaluated the effects of cholecalciferol on both endocrine markers and semen parameters in eighty-six infertile men with asthenozoospermia⁽⁶⁹⁾. Supplementation with 100 µg cholecalciferol daily for 12 weeks had no significant effects on semen volume, sperm count or sperm morphology compared with the placebo group. However, they found a positive effect on total and progressive sperm motility, to some extent supporting the results found in the Chinese and Indian study populations.

Overall, most of the interventional human data available indicate that vitamin D supplementation does not affect sperm count, volume, morphology or pregnancies. However, it must be noted that there are some contradictory reports on the presumed effect on total and progressive sperm motility, but positive data have only been generated in a few relatively small studies. Furthermore, with the diverse study populations of the mentioned studies (co-morbidities, age and BMI), design, inclusion criteria and baseline vitamin D status, it is hard to find consensus on the effect. Observational human studies and case-control studies investigating the association between vitamin D status and reproductive functions have also been conducted. These results are conflicting; however, most studies find that low vitamin D status is linked to impaired semen quality and fertility^(20,71–76). Therefore, it seems likely that the possible effect of vitamin D supplementation on semen quality and male fertility is present under conditions with vitamin D deficiency or prolonged vitamin D insufficiency. These data are in line with animal studies showing impaired fertility and poor semen quality in males with vitamin D deficiency, which by combining suggest that vitamin D deficiency should be avoided to secure optimal reproductive function.

Calcium

Systemic calcium homeostasis

Serum calcium is maintained within a narrow physiological range to secure normal function of several organs as multiple cellular processes are influenced by calcium. Virtually all cells in the body have a 20 000-fold gradient of calcium ions (Ca²⁺) from the extracellular to the intracellular environment (about 100 nm/l-Ca²⁺ intracellularly), making them capable of using Ca²⁺ as a second messenger when entry is permitted into the cell by Ca²⁺ channels⁽⁷⁷⁾. The total calcium concentration in serum is 2.20–2.55 mm/l and the non-protein-bound ionised calcium level is 1.18–1.32 mm/l although the exact reference levels vary among laboratories. A large fraction of calcium is protein-bound or in complex with anions such as citrate, lactate, phosphate or bicarbonate and the free and active fraction is therefore dependent on pH and availability of binding partners⁽⁷⁸⁾. To maintain serum calcium within this narrow range several potent regulators influence systemic calcium homeostasis by

modifying intestinal absorption, renal reabsorption and excretion and release from the skeletal reservoir. The calcium-sensing receptor (CaSR) in the parathyroid cells senses the Ca^{2+} concentration in serum and transduces an intracellular signal that inhibits the production and release of PTH into the bloodstream⁽⁷⁹⁾. PTH binds primarily to the PTH receptor 1 and mobilises calcium to the circulation. In bone, PTH mediates calcium release by promoting bone resorption, and in the kidney, it increases calcium reabsorption and up-regulates CYP27B1 activity which ensures high vitamin D activity. In mice, the CaSR is indispensable as *Casr* knockout mice die shortly after birth and suffer from hypercalcaemia and hyperparathyroidism⁽⁸⁰⁾. In human subjects, homozygous loss-of-function mutations in *CASR* cause neonatal severe hyperparathyroidism, a potentially life-threatening condition with severe hypercalcaemia, bone demineralisation and respiratory distress, whereas a heterozygous loss-of-function mutation in *CASR* causes familial hypocalciuric hypercalcaemia 1⁽⁸¹⁾, which shows that full compensation by other factors does not occur.

In the epithelia lining the intestine and kidney, calcium is transported from the lumen to the blood through the cell cytoplasm by the transcellular pathway or between intercellular spaces by the paracellular pathway. Expression of transcellular calcium transporters in the kidney and intestine is regulated mainly by $1,25(\text{OH})_2\text{D}_3$ ⁽⁸²⁾. Transcellular calcium transport involves three steps: first, Ca^{2+} is transported into the cytosol by transient receptor potential vanilloid –5 or –6 (TRPV –5/–6) Ca^{2+} channels. Secondly, Ca^{2+} binds intracellularly to calbindin- D_{9k} (or D_{28k}) to maintain a low intracellular Ca^{2+} concentration. Thirdly, Ca^{2+} is removed from the cytosol by the plasma-membrane calcium ATPase (PMCA) proteins or the $\text{Na}^+/\text{Ca}^{2+}$ exchanger 1⁽⁸³⁾. CaSR is expressed in the thick ascending limb of Henle in the kidney, and acute inhibition of CaSR increases the permeability of the paracellular Ca^{2+} pathway⁽⁸⁴⁾. More recently, it has been shown that CaSR inhibits the paracellular Ca^{2+} transport by increasing the expression of renal claudin-14 responsible for the inhibition of reabsorption in a PTH-independent manner⁽⁸⁵⁾. In addition, PTH directly increases renal Ca^{2+} reabsorption by inhibiting claudin-14⁽⁸⁶⁾.

Calcium balance in the male reproductive tract

Calcium and phosphate concentrations are very different in the proximal and the distal parts of the epididymis. The concentration of calcium decreases, whereas the concentration of phosphate increases⁽⁸⁷⁾. These variations in calcium and phosphate levels are believed to play a crucial role in sperm maturation and the initiation of sperm motility and in keeping the sperm quiescent during storage in the distal epididymis. The high calcium concentration in the seminal fluid may be of great importance as spermatozoa are transcriptionally silent and heavily rely on intracellular calcium as a signalling system⁽⁸⁸⁾. A list of observational studies exploring total and ionised calcium levels in the human seminal fluid/plasma is

summarised in Table 3. Based on the weighted average of the studies included in Table 3, the concentration of total calcium in the seminal fluid is about 7.48 mmol/l, and the concentration of ionised calcium is about 0.23 mmol/l. Citrate and phosphate concentrations are also high in the seminal fluid, ensuring competent buffer systems and making the ionised calcium concentration lower than the corresponding serum level^(89–93). Studies exploring the potential impact of the total calcium concentration in seminal fluid on sperm parameters are inconclusive – some suggest it is associated with increased motility^(94,95), another study found an inverse relationship with sperm morphology⁽⁹⁶⁾ and other studies found no impact on sperm function^(97–99). One study showed that low total calcium content in the seminal fluid (<5 mmol/l) was associated with fewer progressive and total motile sperm and fewer morphologically normal sperm compared with men with calcium levels between 5 and 10 mmol/l⁽¹⁹⁾. Regarding studies that investigated the link between ionised calcium and semen quality variables, two studies found a positive association with motility^(95,97), one showed a negative association⁽¹⁰⁰⁾ and one study found no effect⁽⁹¹⁾. The possible link between total calcium/ Ca^{2+} levels and sperm motility/morphology/concentration is interesting because these semen variables are predictors of male fertility potential. However, the spermatozoa are only briefly exposed to the seminal fluid before it is mixed with female fluids, which questions a direct effect of calcium content on the fertility potential. The total calcium concentration in seminal fluid is more than 2-fold higher than the level found in serum, and the levels are not associated (Fig. 1A)^(19,100), hence there must be an active transport of calcium from serum into the seminal fluid. The calcium concentration in the different epididymal compartments (studies have only been conducted in rodents) varies largely from low to several-fold higher than the corresponding serum concentration (Fig. 1B)^(101–103). To maintain this steep gradient the mechanism and proteins underlying calcium homeostasis must be tightly regulated. Seminal fluid comprises fluid from the epididymis (about 10%), prostate (about 20%) and seminal vesicles (about 70%), which indicates that the calcium content in the ejaculate is highly dependent on secretions occurring after the transit and storage of the spermatozoa in the epididymis. At the time of ejaculation, spermatozoa escape the millimolar concentrations of citrate and phosphate in seminal fluid and encounter about 2.2 mmol/l-total calcium and about 1.23 mmol/l- Ca^{2+} in the follicular fluid in the female reproductive tract^(91,104). This influences sperm function and facilitates sperm capacitation and hyperactivation that help the sperm swim up in the vicinity of the oocyte.

Gene expression is silenced during spermatogenesis as histones are replaced by protamines that further condense DNA and prevent transcription within the small spermatozoa (about $4.4 \times 2.8 \mu\text{m}$)^(105,106). Protein synthesis and processing are also hampered and therefore, spermatozoa critically depend on second messenger systems to transmit extracellular signals, and they mainly rely on changes in the intracellular calcium concentration

Table 3. Studies investigating ionised and total calcium levels in human seminal fluid/plasma

Seminal plasma/fluid (mm/l)				
Ionised calcium	Total calcium	Subgroup	Conclusions	Reference
0.19(SD 0.03)	5.13(SD 2.50)	Healthy young men (n 9)	CaSR is important for sensing Ca ²⁺ in spermatozoa	Boisen <i>et al.</i> ⁽¹⁰⁴⁾
0.16(SD 0.04)	5.60(SD 1.43)	Men recruited from a fertility clinic	Men with the lowest sperm motility had significantly lower Ca ²⁺ levels in seminal fluid	Kılıç <i>et al.</i> ⁽⁹⁷⁾
0.37(SD 0.28)	5.48(SD 1.37)	Sperm motility <60% (n 45)		
0.18(SD 0.01)	3.30(SD 0.12)	Sperm motility >60% (n 30)		
0.23(SD 0.01)	3.42(SD 0.13)	Men referred to a fertility clinic	Men with the lowest sperm motility had significantly lower Ca ²⁺ and total calcium levels in seminal fluid	Prien <i>et al.</i> ⁽⁹⁵⁾
0.25(SD 0.07)		Sperm motility <60% (n 15)		
		Sperm motility >60% (n 21)		
0.24(SD 0.02)	11.07(SD 0.41)	Men referred to an andrology unit for different reasons (n 27)	No association between ionised calcium concentrations and motility	Magnus <i>et al.</i> ⁽⁹¹⁾
0.17(SD 0.05)	5.70(SD 2.50)	Healthy men (n 45)	N.A.	Ford and Harrison ⁽⁹⁰⁾
		Men referred to a reproductive unit for different andrological reasons (n 32–37)	Spermatozoa from semen samples with Ca ²⁺ levels below average had higher motility than spermatozoa in semen samples with higher levels. Ca ²⁺ was not correlated with total calcium	Arver <i>et al.</i> ⁽¹⁰⁰⁾
	7.8(SD 3.90)	Subfertile men (n 165)	Men with a seminal fluid calcium concentration between 5 and 10 mm had higher sperm motility compared with men with a calcium concentration between 1 and 5 mm	Boisen <i>et al.</i> ⁽¹⁹⁾
	6.79(SD 0.43)	Healthy men (n 24)	N.A.	Valsa <i>et al.</i> ⁽¹³³⁾
	4.86(SD 1.93)	Normozoospermic (n 30)	No difference in the calcium concentrations between normospermic and azoospermic semen samples	N'Guessan <i>et al.</i> ⁽⁹³⁾
	5.32(SD 1.23)	Azoospermic (n 30)		
	9.61(SD 3.76)	Healthy men, 18–55 years old (n 515)	Calcium was positively associated with sperm concentration	Liang <i>et al.</i> ⁽¹³⁴⁾
	4.55(SD 4.28)	Men referred to a fertility clinic	In the neurologically intact group, seminal calcium was negatively correlated with sperm morphology	Salsabili <i>et al.</i> ⁽⁹⁶⁾
	5.39(SD 4.94)	Spinal cord-injured (n 93)		
	11.88	Neurologically intact (n 145)	Low calcium was associated with better CASA parameters (average path velocity, straight-line velocity and linearity)	Sørensen <i>et al.</i> ⁽⁹⁴⁾
	6.11(SD 29.19)	Healthy men (n 6)	Semen samples from spinal cord injured patients were obtained by electroejaculation	Hirsch <i>et al.</i> ⁽¹³⁵⁾
	2.35(SD 1.65)	Spinal cord injured (n 6)		
	4.17(SD 1.10)	Normozoospermic (n 19)	The total calcium concentration does not discriminate between fertile and infertile patients	Abou-Shakra <i>et al.</i> ⁽⁹⁹⁾
	5.04(SD 1.45)	Oligozoospermic (n 19)		
	4.47(SD 1.25)	Severely oligozoospermic (n 17)		
	4.87(SD 1.57)	Azoospermic (n 19)		
	6.11(SD 1.97)	Healthy men (n 22)	The total calcium concentration does not discriminate between fertile and infertile patients	Umeyama <i>et al.</i> ⁽⁹⁸⁾
	6.59(SD 3.12)	Infertile men (n 69)		
	7.50(SD 2.00)	Men at fertility clinical with normal semen parameters (n 17)	N.A.	Kavanagh ⁽¹³⁶⁾
	6.21(SD 1.55)	Normozoospermic (n 40)	N.A.	Adamopoulos and Deliyiannis ⁽⁹²⁾
	7.11(SD 0.37)	Azoospermic (n 12)		
Weighted average				
0.23	7.48			

CASA, computer-assisted semen analysis; CaSR, calcium-sensing receptor; n, cohort size; N.A., not applicable (the studies did not contain a conclusion on calcium levels and sperm function).

The ionised calcium and/or total calcium concentration measured in the seminal fluid or seminal plasma. The conclusions drawn by the authors of the papers regarding the calcium content and semen parameters are mentioned in the third column. All studies are mean(SD) except for Sørensen *et al.*⁽⁹⁴⁾, which is given as the median. The weighted averages are calculated based on the study sizes.

([Ca²⁺]_i). An increase in [Ca²⁺]_i occurs by opening Ca²⁺ channels across the cell membrane or from intracellular stores – constituting a Ca²⁺ signal. [Ca²⁺]_i regulates capacitation, hyperactivation and the acrosome reaction⁽¹⁰⁷⁾. Regulation of the [Ca²⁺]_i ensures the activation of spermatozoa at the appropriate time. The importance is

indicated by studies showing that loss-of-function mutations in the gene coding for the main Ca²⁺ channel in sperm, CatSper, lead to infertility, and spermatozoa from men in fertility treatment have lower Ca²⁺ influx in response to progesterone compared with spermatozoa from healthy men^(108,109). Studies have shown that men




Fig. 1. Calcium and vitamin D in the male reproductive tract. (A) Seminal fluid concentration relative to serum concentration of calcium, 25-hydroxyvitamin D (25OHD), 1,25-dihydroxyvitamin D₃ (1,25(OH)₂D₃) and 24,25-dihydroxyvitamin D₃ (24,25(OH)₂D₃) in men. Dotted lines represent the detection limit in seminal fluid, solid lines represent the constant concentration in seminal fluid and grey solid lines represent the undetectable levels in seminal fluid. The figure is based on data presented in^(19,138). (B) The concentration of calcium and vitamin D in the intraluminal fluid of rodent epididymis relative to the serum concentration. In caput, the 25OHD concentration is 15× lower and 11× lower in cauda compared with serum levels. The calcium concentration is 2× higher in caput and 4× lower in cauda. The figure is based on data presented in^(76,101,102).

with severely impaired semen quality and men who had unsuccessful fertility treatment have significantly lower Ca²⁺ influx upon stimulation with female secreted factors compared with spermatozoa from healthy men indicating that these Ca²⁺ signals are clinically relevant^(108,110).

Animal models with ablation of genes related to calcium homoeostasis in the testes or epididymis

Calcium homoeostasis in the male reproductive tract and its impact on male fertility has been studied in various animal models. Table 4 presents rodent models where a gene ablation has led to changes in local calcium homoeostasis in the testes or epididymis. The *Cyp27b1*^{-/-} mice exhibit hypocalcaemia, decreased sperm count and motility resulting in impaired fertility. *Cyp27b1*^{-/-} mice also have lower expression of calcium transport proteins including CaSR, CaV3.1 and TRPV5 in testes compared with wild-type mice⁽⁶³⁾. Interestingly, a rescue diet (containing high calcium, phosphate and lactose) re-established both the fertility and the expression of CaSR, CaV3.1 and TRPV5 in the testes of *Cyp27b1*^{-/-} mice⁽⁶³⁾. A study in *Vdr*^{-/-} mice also found reduced *Casr* expression in the testes compared with wild-type mice⁽¹⁹⁾, which is in accordance with the role of VDR as a regulator of calcium transporters in various organs. Previous studies have shown CaSR expression in the testes and spermatozoa from different species^(63,111–117), and a proteomic study in bulls showed that the CaSR was the most differentially expressed protein when comparing

good v. bad quality spermatozoa⁽¹¹⁸⁾. Moreover, treatment with different CaSR agonists increased sperm motility in rodents⁽¹¹¹⁾. However, a germ cell-specific *Casr* knockdown model in male mice had no major reproductive phenotype compared with tamoxifen-treated controls irrespectively of the timing of *Casr* knockdown as conducted both pre- and post-pubertally (70 and 84 % reduction, respectively)⁽¹⁹⁾, which questions the importance of CaSR in spermatozoa, at least in mice. In a recent paper investigating CaSR function in human sperm, three patients with loss-of-function mutations in *CASR*, and one patient with a gain-of-function mutation were included. Spermatozoa from all patients had aberrant Ca²⁺ signalling. Moreover, two of the patients with loss-of-function mutations in *CASR* had either low sperm motility and few morphologically normal spermatozoa, or calcifications in the efferent ducts⁽¹⁰⁴⁾.

Myotubularin-related protein 14 (MTMR14) is a phosphoinositide phosphatase and knockout of the gene results in impaired male fertility. In the *Mtmr14*^{-/-} mice, the mRNA expression of the calcium channels inositol 1,4,5-trisphosphate receptor type 1 and 2 (*Itp1-2*), and ryanodine receptor 3 (*Ryr3*) was decreased and [Ca²⁺]_i in spermatozoa derived from epididymis was lower compared with spermatozoa from wild-type mice⁽¹¹⁹⁾. Additionally, *Tmem203*-deficient male mice are sterile and exhibit a profound defect in spermatogenesis. mRNA expression profiling revealed down-regulation of the calcium channels *Trpv6* and transient receptor potential cation channel subfamily M members

Table 4. Mouse knockout models that impact local calcium homeostasis in the male reproductive tract

Approach	Subgroup	Fertility	Sperm count	Sperm motility	Sperm morphology	Impact on local calcium homeostasis	Reference
<i>Vdr</i> KO	Black Swiss	N.D.	N.D.	N.D.	N.D.	Lower <i>Casr</i> expression in testes	Boisen <i>et al.</i> ⁽¹⁹⁾
<i>Cyp27b1</i> KO	BALB/c	↓	↓	↓	↓	Low expression of CaSR, CaV3-1 and TRPV5 in testes. Rescued by calcium-containing diet	Sun <i>et al.</i> ⁽⁶³⁾
<i>Mttr14</i> KO	C57BL/6	↓	↓	↓	↓	Decreased expression of the calcium channels <i>Itpr1</i> , <i>Itpr2</i> and <i>Ryr3</i> in spermatozoa. Spermatozoa had lower [Ca ²⁺] _i	Wen <i>et al.</i> ⁽¹¹⁹⁾
<i>Tmem203</i> KO	C57BL/6J	↓	↓	N.A.	↓	<i>Trpv6</i> , <i>Trpm5</i> and <i>Trpm8</i> were down-regulated in testes. <i>Pmca1</i> and <i>Ip3r1</i> were up-regulated. Altered calcium entry kinetics in testicular cells	Shambharkar <i>et al.</i> ⁽¹²⁰⁾
<i>Sppl2c</i> KO	C57BL/6N	→	N.D.	↓	→	SPPL2c cleaves phospholamban, which interacts with SERCA2 thereby regulating intracellular calcium concentration in germ cells	Niemeyer <i>et al.</i> ⁽¹²¹⁾
<i>Trpv6</i> KO	Mixed (129/C57BL)	↓	↓*	↓	N.D.	Calcium level in cauda epididymis was 10-fold higher resulting in low sperm motility	Weissgerber <i>et al.</i> ⁽¹²⁴⁾
<i>Pmca4</i> KO	Mixed (129/black Swiss)	↓	N.D.	↓	→	PMCA4 is important in the regulation of basal calcium levels and calcium clearance in sperm	Schuh <i>et al.</i> ⁽¹³⁷⁾

CaSR, calcium-sensing receptor; *Ip3r1*, intracellular ion channel inositol 1,4,5-triphosphate receptor 1; *Itpr1/2*, inositol 1,4,5-trisphosphate receptor type 1/2; KO, knockout; *Mttr14*, myotubularin related protein 14; N.A., not applicable; N.D., not determined; *Pmca1/4*, plasma membrane Ca²⁺-ATPase 1/4; *Ryr*, ryanodine receptor; SERCA2, sarcoplasmic/endoplasmic reticulum Ca²⁺-ATPase; *Sppl2c*, signal peptide peptidase-like 2C; *Trpm5/8*, transient receptor potential cation channel subfamily M 5/8; *Trpv5/6*, transient receptor potential vanilloid sub-type 5/6; *Vdr*, vitamin D receptor.

↓, negatively associated; →, no association; *the low caudal sperm count after swim-out may be explained by impaired motility rather than impaired sperm production or storage.

5 and 8 (*Trpm* -5/-8), and an increase in *Pmca1* and intracellular ion channel inositol 1,4,5-triphosphate receptor (*Ip3r1*) in testes. Testicular cells from *Tmem203* mice also exhibited altered calcium mobilisation⁽¹²⁰⁾. Signal peptide peptidase-like 2C (*Sppl2c*) deficiency in male mice leads to dysregulation of the calcium pump sarcoplasmic/endoplasmic reticulum calcium ATPase (SERCA2) involved in intracellular Ca²⁺ handling in male germ cells. The disturbed calcium homeostasis resulted in impaired motility of spermatozoa but preserved fertility⁽¹²¹⁾. Efflux of intracellular Ca²⁺ can be mediated by the PMCA pump, which is essential for spermatozoa as *Pmca4*^{-/-} mice are infertile due to Ca²⁺ overload and inability of spermatozoa to undergo capacitation^(122,123).

The epididymis requires active calcium transport to maintain the calcium concentration gradient (Fig. 1B), which is necessary to ensure normal sperm function. In cauda epididymis in global *Trpv6* knockout mice a 10-fold higher calcium concentration impaired sperm motility⁽¹²⁴⁾. Ma *et al.*⁽¹²⁵⁾ found an excessive calcium accumulation in the epididymis of vitamin K2-deficiency rats due to dysregulation of γ -glutamyl carboxylase (GGCX) and matrix gla protein (MGP). The vitamin K2-deficiency rats had lower sperm count and sperm motility. In agreement with the results in rats, the study also identified an SNP mutation in *GGCX* in an infertile man with asthenozoospermia, indicating that the influence

of GGCX and MGP on fertility is conserved between species.

Human relevance of calcium in male fertility

The link between calcium, vitamin D, bone and gonadal function⁽¹²⁶⁻¹²⁹⁾ complicates interpretations of human and animal studies as many of the effects of calcium may be indirect. In most human studies on vitamin D and male fertility, potential calcium-mediated effects have been neglected. In a study of infertile men, serum Ca²⁺ levels were negatively associated with sperm motility indicating that Ca²⁺ may influence semen quality in infertile men⁽¹⁰⁾. This association suggests that systemic regulators of calcium homeostasis may also influence local calcium transport in the male reproductive tract, as calcium levels in serum and calcium levels in the epididymis and seminal fluid are not associated (Fig. 1A).

Significant interspecies differences exist in fertility and sperm function. Hence, extrapolating results from mice to human subjects in the fertility field is problematic and should be performed with caution⁽¹³⁰⁾. Depending on the species, fertilisation occurs under entirely different aqueous conditions with different calcium and vitamin D levels, which remains to be thoroughly studied *in vivo*. Most data have been generated by *in vitro* studies and the true levels of calcium surrounding the spermatozoa as it

moves through the female reproductive tract to the oocyte and in the moment of fertilisation are largely unknown.

Conclusions and perspectives

In conclusion, vitamin D and calcium are implicated in various facets of male fertility. Preliminary findings from intervention studies suggest that vitamin D supplementation may be beneficial for men with low vitamin D status, especially those with a serum 25OHD level below 25 nm/l. However, further investigation through larger clinical studies is required to enhance our understanding and address whether vitamin D supplementation can improve semen quality mostly in men with insufficient vitamin D levels. It seems that many of the observed effects of vitamin D are mediated indirectly through changes in local levels of factors such as calcium or phosphate, not only in the testes but also in epididymis, prostate and seminal vesicles. The crucial role of calcium in male reproductive function extends beyond sperm motility and fertilisation, as calcium signalling is involved in various processes, such as spermatogenesis, sperm maturation, capacitation, acrosome reaction and imbalances in calcium levels may disrupt these events and impair fertility. The exact mechanisms underlying the impact of calcium on male reproduction are not fully understood, although studies have suggested that maintaining optimal calcium homeostasis is essential for overall reproductive health.

More research on the specific role of vitamin D and other regulators of local calcium and phosphate signalling in the male reproductive organs is warranted. Recently, we identified the receptor activator of NF κ B ligand as a novel regulator of the production and maturation of spermatozoa⁽¹²⁸⁾. Inhibition of receptor activator of NF κ B ligand increased sperm motility and sperm count in a subgroup of men with preserved Sertoli cell capacity, and identification of predictive biomarkers for positive treatment outcomes is crucial and requires further investigation. Moreover, it is crucial to shift focus towards investigating more definitive outcomes such as conception rates and live births. Currently, there is a concerning lack of evidence-based treatments available for men with idiopathic infertility, despite the high prevalence of male infertility^(7,131). Rather than tailoring treatments to the specific causes of infertility^(17,24,129,132), most infertile couples are commonly subjected to inseminations or assisted reproductive techniques. Assisted reproductive techniques have demonstrated high success rates, but they come with substantial financial costs and impose a significant burden on the female partner. The invasive procedures and prolonged hormonal treatments often lasting several months contribute to the treatment burden faced by women undergoing assisted reproductive techniques. In this regard, supplementation of vitamin D presents a promising avenue for improving semen quality in certain cases of idiopathic male infertility and supports screening of vitamin D and mineral status in cases of idiopathic male infertility. It offers a safe and non-invasive treatment option that could potentially benefit some infertile couples.

Moreover, the devastating effect of vitamin D deficiency on semen quality in animal models and human subjects indicates that society should continue to focus on preventing people from having vitamin D deficiency.

Acknowledgements

The authors thank colleagues and peers for constructive discussions while putting this review together.

Financial Support

None.

Conflict of Interest

M. B. J. holds two patents on the use of RANKL inhibitors to treat male infertility. All other authors state no conflict of interest.

Authorship

S. K. Y. and I. M. B. reviewed the literature, wrote the initial draft and had the main responsibility for the preparation of the paper. M. B. J. was responsible for the overall paper design, critical discussions and final approval of the manuscript. In line with the mentioned authors, Z. C., M. J. J., I. K., R. H. and A. J. participated in the discussion and interpretation of the results, critically revised the manuscript for intellectual content and approved the final version.

References

1. Boivin J, Bunting L, Collins JA *et al.* (2007) International estimates of infertility prevalence and treatment-seeking: potential need and demand for infertility medical care. *Hum Reprod* **22**, 1506–1512.
2. United Nations, Department of Economic and Social Affairs (2022) World Population Prospects 2022: Summary of Results, New York, 2022.
3. Agarwal A, Mulgund A, Hamada A *et al.* (2015) A unique view on male infertility around the globe. *Reprod Biol Endocrinol* **13**, 1–9.
4. Hirsh A (2003) Male subfertility. *Br Med J* **327**, 669–672.
5. Xavier MJ, Salas-Huetos A, Oud MS *et al.* (2021) Disease gene discovery in male infertility: past, present and future. *Hum Genet* **140**, 7–19.
6. Dohle GR, Colpi GM, Hargreave TB *et al.* (2019) EAU Guidelines on male infertility. *Eur Urol* **48**, 703–711.
7. Krausz C (2011) Male infertility: pathogenesis and clinical diagnosis. *Best Pract Res Clin Endocrinol Metab* **25**, 271–285.
8. Jensen MB (2014) Vitamin D and male reproduction. *Nat Rev Endocrinol* **10**, 175–186.
9. Blomberg Jensen M (2012) Vitamin D metabolism, sex hormones, and male reproductive function. *Reproduction* **144**, 647.
10. Blomberg Jensen M, Gerner Lawaetz J, Andersson AM *et al.* (2016) Vitamin D deficiency and low ionized calcium are linked with semen quality and sex steroid levels in infertile men. *Hum Reprod* **31**, 1875–1885.

11. Mendonca V (2023) Modern food habits and its impact on human health. *Int J Innov Res Eng Manag* **10**, 182–185.
12. Lieben L, Carmeliet G & Masuyama R (2011) Calcemic actions of vitamin D: effects on the intestine, kidney and bone. *Best Pract Res Clin Endocrinol Metab* **25**, 561–572.
13. Zhang Y, Fang F, Tang J *et al.* (2019) Association between vitamin D supplementation and mortality: systematic review and meta-analysis. *BMJ* **366**, 14673.
14. Grant WB, Lahore H, McDonnell SL *et al.* (2020) Evidence that vitamin D supplementation could reduce risk of influenza and COVID-19 infections and deaths. *Nutrients* **12**, 988.
15. Keum N, Lee DH, Greenwood DC *et al.* (2019) Vitamin D supplementation and total cancer incidence and mortality: a meta-analysis of randomized controlled trials. *Ann Oncol* **30**, 733.
16. Holt R, Juel Mortensen L, Harpelunde Poulsen K *et al.* (2020) Vitamin D and sex steroid production in men with normal or impaired Leydig cell function. *J Steroid Biochem Mol Biol* **199**, 105589.
17. Blomberg Jensen M, Lawaetz JG, Petersen JH *et al.* (2018) Effects of vitamin D supplementation on semen quality, reproductive hormones, and live birth rate: a randomized clinical trial. *J Clin Endocrinol Metab* **103**, 870–881.
18. Blomberg Jensen M, Nielsen JE, Jørgensen A *et al.* (2010) Vitamin D receptor and vitamin D metabolizing enzymes are expressed in the human male reproductive tract. *Hum Reprod* **25**, 1303–1311.
19. Boisen IM, Nielsen JE, Verlinden L *et al.* (2021) Calcium transport in male reproduction is possibly influenced by vitamin D and CaSR. *J Endocrinol* **251**, 207–222.
20. Heller CG & Clermont Y (1963) Spermatogenesis in man: an estimate of its duration. *Science (1979)* **140**, 184–186.
21. Sharpe RM, McKinnell C, Kivlin C *et al.* (2003) Proliferation and functional maturation of Sertoli cells, and their relevance to disorders of testis function in adulthood. *Reproduction* **125**, 769–784.
22. O'Shaughnessy PJ, Monteiro A, Verhoeven G *et al.* (2010) Effect of FSH on testicular morphology and spermatogenesis in gonadotrophin-deficient hypogonadal mice lacking androgen receptors. *Reproduction* **139**, 177–184.
23. Jarow JP, Wright WW, Brown TR *et al.* (2005) Bioactivity of androgens within the testes and serum of normal men. *J Androl* **26**, 343–348.
24. Skakkebaek NE, Rajpert-De Meyts E, Buck Louis GM *et al.* (2016) Male reproductive disorders and fertility trends: influences of environment and genetic susceptibility. *Physiol Rev* **96**, 55–97.
25. Cheng JB, Levine MA, Bell NH *et al.* (2004) Genetic evidence that the human CYP2R1 enzyme is a key vitamin D 25-hydroxylase. *Proc Natl Acad Sci USA* **101**, 7711–7715.
26. Prosser DE & Jones G (2004) Enzymes involved in the activation and inactivation of vitamin D. *Trends Biochem Sci* **29**, 664–673.
27. Ohyama Y (2004) Eight cytochrome P450s catalyze vitamin D metabolism. *Front Biosci* **9**, 3007.
28. Jones G, Prosser DE & Kaufmann M (2012) 25-Hydroxyvitamin D-24-hydroxylase (CYP24A1): its important role in the degradation of vitamin D. *Arch Biochem Biophys* **523**, 9–18.
29. Gray RW, Omdahl JL, Ghazarian JG *et al.* (1972) 25-Hydroxycholecalciferol-1-hydroxylase. *J Biol Chem* **247**, 7528–7532.
30. Bouillon R, Carmeliet G, Verlinden L *et al.* (2008) Vitamin D and human health: lessons from vitamin D receptor null mice. *Endocr Rev* **29**, 726–776.
31. Yuan Q, Sitara D, Sato T *et al.* (2011) PTH ablation ameliorates the anomalies of Fgf23-deficient mice by suppressing the elevated vitamin D and calcium levels. *Endocrinology* **152**, 4053–4061.
32. Lips P (2006) Vitamin D physiology. *Prog Biophys Mol Biol* **92**, 4–8.
33. Holick MF (2006) High prevalence of vitamin D inadequacy and implications for health. *Mayo Clin Proc* **81**, 353–373.
34. Nagakura K, Abe E, Suda T *et al.* (1986) Inhibitory effect of 1 α ,25-dihydroxyvitamin D₃ on the growth of the renal carcinoma cell line. *Kidney Int* **29**, 834–840.
35. Choudhary D, Jansson I, Stoilov I *et al.* (2005) Expression patterns of mouse and human CYP orthologs (families 1–4) during development and in different adult tissues. *Arch Biochem Biophys* **436**, 50–61.
36. Johnson JA, Grande JP, Roche PC *et al.* (1996) Immunohistochemical detection and distribution of the 1,25-dihydroxyvitamin D₃ receptor in rat reproductive tissues. *Histochem Cell Biol* **105**, 7–15.
37. Mahmoudi AR, Zarnani AH, Jeddi-Tehrani M *et al.* (2013) Distribution of vitamin D receptor and 1 α -hydroxylase in male mouse reproductive tract. *Reprod Sci* **20**, 426–436.
38. Oliveira AG, Dornas RAP, Kalapothakis E *et al.* (2008) Vitamin D₃ and androgen receptors in testis and epididymal region of roosters (*Gallus domesticus*) as affected by epididymal lithiasis. *Anim Reprod Sci* **109**, 343–355.
39. Schleicher G, Privette TH & Stumpf WE (1989) Distribution of soltrilol [1,25(OH)₂-vitamin D₃] binding sites in male sex organs of the mouse: an autoradiographic study. *J Histochem Cytochem* **37**, 1083–1086.
40. Corbett ST, Hill O & Nangia AK (2006) Vitamin D receptor found in human sperm. *Urology* **68**, 1345–1349.
41. Aquila S, Guido C, Perrotta I *et al.* (2008) Human sperm anatomy: ultrastructural localization of 1 α ,25-dihydroxyvitamin D₃ receptor and its possible role in the human male gamete. *J Anat* **213**, 555–564.
42. Merke J, Kreuzer W, Bier B *et al.* (1983) Demonstration and characterisation of a testicular receptor for 1,25-dihydroxycholecalciferol in the rat. *Eur J Biochem* **130**, 303–308.
43. Zanatta L, Zamoner A, Zanatta AP *et al.* (2011) Nongenomic and genomic effects of 1 α ,25(OH)₂ vitamin D₃ in rat testis. *Life Sci* **89**, 515–523.
44. Zanatta L, Bouraïma-Lelong H, Delalande C *et al.* (2011) Regulation of aromatase expression by 1 α ,25(OH)₂ vitamin D₃ in rat testicular cells. *Reprod Fertil Dev* **23**, 725–735.
45. Akerstrom VL & Walters MR (1992) Physiological effects of 1,25-dihydroxyvitamin D₃ in TM4 Sertoli cell line. *Am J Physiol Endocrinol Metab* **262**, E884–E890.
46. Majumdar SS, Bartke A & Stumpf WE (1994) Vitamin D modulates the effects of follicle-stimulating hormone of Sertoli cell function and testicular growth in Siberian hamsters. *Life Sci* **55**, 1479–1486.
47. Menegaz D, Mizwicki MT, Barrientos-Duran A *et al.* (2011) Vitamin D receptor (VDR) regulation of voltage-gated chloride channels by ligands preferring a VDR-alternative pocket (VDR-AP). *Mol Endocrinol* **25**, 1289–1300.
48. Zanatta L, Zamoner A, Gonçalves R *et al.* (2011) 1 α ,25-Dihydroxyvitamin D₃ signaling pathways on calcium uptake in 30-day-old rat Sertoli cells. *Biochemistry* **50**, 10284–10292.

49. Rosso A, Pansera M, Zamoner A *et al.* (2012) $1\alpha,25(\text{OH})_2$ -vitamin D3 stimulates rapid plasma membrane calcium influx via MAPK activation in immature rat Sertoli cells. *Biochimie* **94**, 146–154.
50. Menegaz D, Barrientos-Duran A, Kline A *et al.* (2010) $1\alpha,25(\text{OH})_2$ -Vitamin D3 stimulation of secretion via chloride channel activation in Sertoli cells. *J Steroid Biochem Mol Biol* **119**, 127–134.
51. Foresta C, Strapazzon G, De Toni L *et al.* (2011) Bone mineral density and testicular failure: evidence for a role of vitamin D 25-hydroxylase in human testis. *Endocrinology* **152**, 1195.
52. Aquila S, Guido C, Middea E *et al.* (2009) Human male gamete endocrinology: $1\alpha,25$ -dihydroxyvitamin D3 ($1,25(\text{OH})_2\text{D}_3$) regulates different aspects of human sperm biology and metabolism. *Reprod Biol Endocrinol* **7**, 140.
53. Blomberg Jensen M, Jørgensen A, Nielsen JE *et al.* (2012) Expression of the vitamin D metabolizing enzyme CYP24A1 at the annulus of human spermatozoa may serve as a novel marker of semen quality. *Int J Androl* **35**, 499–510.
54. Blomberg Jensen M, Jørgensen A, Nielsen JE *et al.* (2012) Vitamin D metabolism and effects on pluripotency genes and cell differentiation in testicular germ cell tumors in vitro and in vivo. *Neoplasia* **14**, 952–963.
55. Hirai T, Tsujimura A, Ueda T *et al.* (2009) Effect of $1,25$ -dihydroxyvitamin D on testicular morphology and gene expression in experimental cryptorchid mouse: testis specific cDNA microarray analysis and potential implication in male infertility. *J Urol* **181**, 1487–1492.
56. Kwiecinski GG, Petrie GI & DeLuca HF (1989) Vitamin D is necessary for reproductive functions of the male rat. *J Nutr* **119**, 741–744.
57. Uhland AM, Kwiecinski GG & DeLuca HF (1992) Normalization of serum calcium restores fertility in vitamin D-deficient male rats. *J Nutr* **122**, 1338–1344.
58. Audet I, Laforest J, Martineau GP *et al.* (2004) Effect of vitamin supplements on some aspects of performance, vitamin status, and semen quality in boars. The online version of this article, along with updated information and services, is located at: Effect of vitamin supplement, pp. 626–633.
59. Kinuta K, Tanaka H, Moriwake T *et al.* (2000) Vitamin D is an important factor in estrogen biosynthesis of both female and male gonads. *Endocrinology* **141**, 1317–1324.
60. Fu L, Chen YH, Xu S *et al.* (2017) Vitamin D deficiency impairs testicular development and spermatogenesis in mice. *Reprod Toxicol* **73**, 241–249.
61. Zamani A, Saki F, Hatami N *et al.* (2020) Stereological assessment of the effects of vitamin D deficiency on the rat testis. *BMC Endocr Disord* **20**, 162.
62. Shahreza FD, Hajian M, Gharagozloo P *et al.* (2020) Impact of vitamin D deficiency on mouse sperm structure and function. *Andrology* **8**, 1442–1455.
63. Sun W, Chen L, Zhang W *et al.* (2015) Active vitamin D deficiency mediated by extracellular calcium and phosphorus results in male infertility in young mice. *Am J Physiol Endocrinol Metab* **308**, E51–E62.
64. Merino O, Sánchez R, Gregorio MB *et al.* (2019) Effect of high-fat and vitamin D deficient diet on rat sperm quality and fertility. *Theriogenology* **125**, 6–11.
65. Lorenzen M, Boisen IM, Mortensen LJ *et al.* (2017) Reproductive endocrinology of vitamin D. *Mol Cell Endocrinol* **453**, 103–112.
66. Deng XL, Li YM, Yang XY *et al.* (2015) Efficacy and safety of vitamin D in the treatment of idiopathic oligoasthenozoospermia. *Nat J Androl* **20**, 1082–1085.
67. Padmapriya D, Archana R, Sharma N *et al.* (2021) Effects of vitamin D3 supplementation in male reproductive function: a double-blinded randomized controlled trial. *Curr Top Nutraceutical Res* **20**, 334–339.
68. Gheflati A, Mirjalili SAM, Kaviani M *et al.* (2021) Effects of vitamin D supplementation on semen quality and reproductive hormones in patients with asthenozoospermia: a randomized double-blind placebo-controlled clinical trial. *J Nutr Food Secur* **6**, 31–42.
69. Maghsoumi-Norouzabad L, Zare Javid A, Mansoori A *et al.* (2021) The effects of vitamin D3 supplementation on spermatogram and endocrine factors in asthenozoospermia infertile men: a randomized, triple blind, placebo-controlled clinical trial. *Reprod Biol Endocrinol* **19**, 102.
70. Amini L, Mohammadbeigi R, Vafa M *et al.* (2020) Evaluation of the effect of vitamin D3 supplementation on quantitative and qualitative parameters of spermograms and hormones in infertile men: a randomized controlled trial. *Complement Ther Med* **53**, 102529.
71. Prosser DE & Jones G (2004) Enzymes involved in the activation and inactivation of vitamin D. *Trends Biochem Sci* **29**, 664–673.
72. Akhavizadegan H & Karbakhsh M (2017) Comparison of serum vitamin D between fertile and infertile men in a vitamin D deficient endemic area: a case-control study. *Urol J* **84**, 218–220.
73. Shi X, Chan CPS, Man GKY *et al.* (2021) Associations between blood metal/metalloid concentration and human semen quality and sperm function: a cross-sectional study in Hong Kong. *J Trace Elem Med Biol* **65**, 126735.
74. Tirabassi G, Cutini M, Muscogiuri G *et al.* (2017) Association between vitamin D and sperm parameters: clinical evidence. *Endocrine* **58**, 194–198.
75. Tartagni M, Matteo M, Baldini D *et al.* (2015) Males with low serum levels of vitamin D have lower pregnancy rates when ovulation induction and timed intercourse are used as a treatment for infertile couples: results from a pilot study. *Reprod Biol Endocrinol* **13**, 127.
76. Kidroni G, Har-Nir R, Menczel J *et al.* (1983) Vitamin D3 metabolites in rat epididymis: high $24,25$ -dihydroxy vitamin D3 levels in the cauda region. *Biochem Biophys Res Commun* **113**, 982–989.
77. Clapham DE (2007) Calcium signaling. *Cell* **131**, 1047–1058.
78. Baird GS (2011) Ionized calcium. *Clin Chim Acta* **412**, 696–701.
79. Brown EM, Gamba G, Riccardi D *et al.* (1993) Cloning and characterization of an extracellular Ca^{2+} -sensing receptor from bovine parathyroid. *Nature* **366**, 575–580.
80. Ho C, Conner DA, Pollak MR *et al.* (1995) A mouse model of human familial hypocalciuric hypercalcemia and neonatal severe hyperparathyroidism. *Nat Genet* **11**, 389–394.
81. Boisen IM, Mos I, Lerche-Black EM *et al.* (2020) Heterozygous mutation (Q459R) in the calcium-sensing receptor gene causes familial hypocalciuric hypercalcemia 1 (FHH1). *J Clin Endocrinol Metab* **105**, e1322–e1330.
82. Hoenderop JGJ, Nilius B & Bindels RJM (2005) Calcium absorption across epithelia. *Physiol Rev* **85**, 373–422.
83. VanHouten JN & Wysolmerski JJ (2007) Transcellular calcium transport in mammary epithelial cells. *J Mammary Gland Biol Neoplasia* **12**, 223–235.
84. Loupy A, Ramakrishnan SK, Wootla B *et al.* (2012) PTH-independent regulation of blood calcium

- concentration by the calcium-sensing receptor. *J Clin Invest* **122**, 3355–3367.
85. Dimke H, Desai P, Borovac J *et al.* (2013) Activation of the Ca^{2+} -sensing receptor increases renal claudin-14 expression and urinary Ca^{2+} excretion. *Am J Physiol Renal Physiol* **304**, F761–F769.
 86. Sato T, Courbebaisse M, Ide N *et al.* (2017) Parathyroid hormone controls paracellular Ca^{2+} transport in the thick ascending limb by regulating the tight-junction protein claudin14. *Proc Natl Acad Sci USA* **114**, E3344–E3353.
 87. Clulow J, Jones R & Hansen L (1994) Micropuncture and cannulation studies of fluid composition and transport in the ductuli efferentes testis of the rat: comparisons with the homologous metanephric proximal tubule. *Exp Physiol* **79**, 915–928.
 88. Blomberg Jensen M, Bjerrum PJ, Jessen TE *et al.* (2011) Vitamin D is positively associated with sperm motility and increases intracellular calcium in human spermatozoa. *Hum Reprod* **26**, 1307–1317.
 89. Owen DH & Katz DF (2005) A review of the physical and chemical properties of human semen and the formulation of a semen simulant. *J Androl* **26**, 459–469.
 90. Ford WCL & Harrison A (1984) The role of citrate in determining the activity of calcium ions in human semen. *Int J Androl* **7**, 198–202.
 91. Magnus Ø, Åbyholm T, Kofstad J *et al.* (1990) Ionized calcium in human male and female reproductive fluids: relationships to sperm motility. *Hum Reprod* **5**, 94–98.
 92. Adamopoulos DA & Deliyannis V (1983) Seminal plasma magnesium, calcium and inorganic phosphate concentration in normozoospermic and subfertile men. *Andrologia* **15**, 648–654.
 93. N'Guessan MF, Yaye YG, Coulibaly FA *et al.* (2016) Evaluation of minerals in the seminal plasma of azoospermic semen. *Int J Biomed Res* **7**, 7–11.
 94. Sørensen MB, Bergdahl IA, Hjøllund NHI *et al.* (1999) Zinc, magnesium and calcium in human seminal fluid: relations to other semen parameters and fertility. *Mol Hum Reprod* **5**, 331–337.
 95. Prien SD, Lox CD, Messer RH *et al.* (1990) Seminal concentrations of total and ionized calcium from men with normal and decreased motility. *Fertil Steril* **54**, 171–172.
 96. Salsabili N, Mehrsai AR & Jalaie S (2009) Concentration of blood and seminal plasma elements and their relationships with semen parameters in men with spinal cord injury. *Andrologia* **41**, 24–28.
 97. Kılıç S, Sarıca K, Yaman Ö *et al.* (1996) Effect of total and ionized calcium levels of seminal fluid on sperm motility. *Urol Int* **56**, 215–218.
 98. Umeyama T, Ishikawa H, Takeshima H *et al.* (1986) A comparative study of seminal trace elements in fertile and infertile men. *Fertil Steril* **46**, 494–499.
 99. Abou-Shakra FR, Ward NI & Everard DM (1989) The role of trace elements in male infertility. *Fertil Steril* **52**, 307–310.
 100. Arver S & Sjöberg HE (1982) Calcium fractions in seminal plasma and functional properties of human spermatozoa. *Acta Physiol Scand* **116**, 159–165.
 101. Jenkins AD, Lechene CP & Howards SS (1980) Concentrations of seven elements in the intraluminal fluids of the rat seminiferous tubules, rete testis, and epididymis. *Biol Reprod* **23**, 981–987.
 102. Jenkins AD, Lechene CP & Howards SS (1983) The effect of estrogen administration in vivo on the elemental composition of the intraluminal fluids of the seminiferous tubules, rete testis, and epididymis of the rat. *J Androl* **4**, 272–275.
 103. Boisen IM, Hansen LB, Mortensen LJ *et al.* (2017) *Vitamin D, Reproductive Biology, and Dysfunction in Men*, vol. 1, 4th ed. Vitamin D.
 104. Boisen IM, Rehfeld A, Mos I *et al.* (2021) The calcium-sensing receptor is essential for calcium and bicarbonate sensitivity in human spermatozoa. *J Clin Endocrinol Metab* **106**, e1775–e1792.
 105. Miller D & Ostermeier GC (2006) Towards a better understanding of RNA carriage by ejaculate spermatozoa. *Hum Reprod Update* **12**, 757–767.
 106. Bellastella G, Cooper TG, Battaglia M *et al.* (2010) Dimensions of human ejaculated spermatozoa in papanicolaou-stained seminal and swim-up smears obtained from the integrated semen analysis system (ISAS[®]). *Asian J Androl* **12**, 871–879.
 107. Publicover S, Harper C V & Barratt C (2007) $[\text{Ca}^{2+}]_i$ signalling in sperm – making the most of what you've got. *Nat Cell Biol* **9**, 235–242.
 108. Kelly MC, Brown SG, Costello SM *et al.* (2018) Single-cell analysis of $[\text{Ca}^{2+}]_i$ signalling in sub-fertile men: characteristics and relation to fertilization outcome. *Hum Reprod* **33**, 1023–1033.
 109. Williams HL, Mansell S, Alasmari W *et al.* (2015) Specific loss of CatSper function is sufficient to compromise fertilizing capacity of human spermatozoa. *Hum Reprod* **30**, 2737–2746.
 110. Alasmari W, Barratt CLR, Publicover SJ *et al.* (2018) The clinical significance of calcium–signalling pathways mediating human sperm hyperactivation. *Hum Reprod* **28**, 866–876.
 111. Mendoza FJ, Perez-Marin CC, Garcia-Marin L *et al.* (2012) Localization, distribution, and function of the calcium-sensing receptor in sperm. *J Androl* **33**, 96–104.
 112. Kong W-Y, Tong L-Q, Zhang H-J *et al.* (2015) The calcium-sensing receptor participates in testicular damage in streptozotocin-induced diabetic rats. *Asian J Androl* **17**, 1–6.
 113. Macías-García B, Rocha A & González-Fernández L (2016) Extracellular calcium regulates protein tyrosine phosphorylation through calcium-sensing receptor (CaSR) in stallion sperm. *Mol Reprod Dev* **83**, 236–245.
 114. Wang YJ, Yan J, Yin F *et al.* (2017) Role of autophagy in cadmium-induced testicular injury. *Hum Exp Toxicol* **36**, 1039–1048.
 115. Liu C, Liu Y, Larsen K *et al.* (2018) Calcium-sensing receptor (CASR) is involved in porcine in vitro fertilisation and early embryo development. *Reprod Fertil Dev* **30**, 391.
 116. Macías-García B, García-Marin LJ, Bragado MJ *et al.* (2019) The calcium-sensing receptor regulates protein tyrosine phosphorylation through PDK1 in boar spermatozoa. *Mol Reprod Dev* **86**, 751–761.
 117. Ellinger I (2016) The calcium-sensing receptor and the reproductive system. *Front Physiol* **7**, 371.
 118. Singh R, Sengar G, Singh U *et al.* (2018) Functional proteomic analysis of crossbred (Holstein Friesian × Sahiwal) bull spermatozoa. *Reprod Domest Anim* **53**, 588–608.
 119. Wen N, Yu MF, Liu J *et al.* (2018) Deficiency of MTMR14 impairs male fertility in *Mus musculus*. *PLoS ONE* **13**, e0206224.
 120. Shambharkar PB, Bittinger M, Latario B *et al.* (2015) TMEM203 is a novel regulator of intracellular calcium homeostasis and is required for spermatogenesis. *PLoS ONE* **10**, 1–25.



121. Niemeyer J, Mentrup T, Heidasch R *et al.* (2019) The intramembrane protease SPPL2c promotes male germ cell development by cleaving phospholamban. *EMBO Rep* **20**, 1–20.
122. Okunade GW, Miller ML, Pyne GJ *et al.* (2004) Targeted ablation of plasma membrane Ca²⁺-ATPase (PMCA) 1 and 4 indicates a major housekeeping function for PMCA1 and a critical role in hyperactivated sperm motility and male fertility for PMCA4. *J Biol Chem* **279**, 33742–33750.
123. Prasad V, Okunade GW, Miller ML *et al.* (2004) Phenotypes of SERCA and PMCA knockout mice. *Biochem Biophys Res Commun* **322**, 1192–1203.
124. Weissgerber P, Kriebs U, Tsvilovskyy V *et al.* (2011) Male fertility depends on Ca²⁺ absorption by TRPV6 in epididymal epithelia. *Sci Signal* **4**, ra27.
125. Ma H, Zhang BL, Liu BY *et al.* (2019) Vitamin K2-dependent GGCX and MGP are required for homeostatic calcium regulation of sperm maturation. *iScience* **14**, 210–225.
126. Oury F, Ferron M, Huizhen W *et al.* (2013) Osteocalcin regulates murine and human fertility through a pancreas–bone–testis axis. *J Clin Invest* **123**, 2421–2433.
127. Oury F, Sumara G, Sumara O *et al.* (2011) Endocrine regulation of male fertility by the skeleton. *Cell* **144**, 796–809.
128. Blomberg Jensen M, Andreassen CH, Jørgensen A *et al.* (2021) RANKL regulates male reproductive function. *Nat Commun* **12**, 1–15.
129. Bøllehuus Hansen L, Kaludjerovic J, Nielsen JE *et al.* (2020) Influence of FGF23 and Klotho on male reproduction: systemic vs direct effects. *FASEB J* **34**, 12436–12449.
130. Kaupp UB & Strünker T (2017) Signaling in sperm: more different than similar. *Trends Cell Biol* **27**, 101–109.
131. Huynh T (2002) Selected genetic factors associated with male infertility. *Hum Reprod Update* **8**, 183–198.
132. Juel Mortensen L, Lorenzen M, Jørgensen N *et al.* (2019) Possible link between FSH and RANKL release from adipocytes in men with impaired gonadal function including Klinefelter syndrome. *Bone* **123**, 103–114.
133. Valsa J, Skandhan KP, Sumangala B *et al.* (2016) Time bound changes (in 24 h) in human sperm motility and level of calcium and magnesium in seminal plasma. *Alexandria J Med* **52**, 235–241.
134. Liang H, Miao M, Chen J *et al.* (2016) The association between calcium, magnesium, and ratio of calcium/magnesium in seminal plasma and sperm quality. *Biol Trace Elem Res* **174**, 1–7.
135. Hirsch IH, Jeyendran RS, Sedor J *et al.* (1991) Biochemical analysis of electroejaculates in spinal cord injured Men: comparison to normal ejaculates. *J Urol* **145**, 73–76.
136. Kavanagh JP (1985) Sodium, potassium, calcium, magnesium, zinc, citrate and chloride content of human prostatic and seminal fluid. *Reproduction* **75**, 35–41.
137. Schuh K, Cartwright EJ, Jankevics E *et al.* (2004) Plasma membrane Ca²⁺ ATPase 4 is required for sperm motility and male fertility. *J Biol Chem* **279**, 28220–6.
138. Hansen LB, Rehfeld A, De Neergaard R *et al.* (2017) Selection of high-quality spermatozoa may be promoted by activated vitamin D in the woman. *J Clin Endocrinol Metab* **102**, 950–961.