

DRSP along with the aforementioned mood questionnaires during both their PM and follicular phases.

Results: In the first research phase, the sample was divided into women with probable PMDD diagnosis (PMDD group, $n=68$) and women without probable PMDD diagnosis (nonPMDD group, $n=45$) based on the DSM-5-Based Screening Tool. The PMDD group reported significantly more severe depressive ($F(1; 56.2) = 19.394, p \leq 0.001$) and anxiety ($F(1; 35.6) = 17.714, p \leq 0.001$) symptoms and lower well-being ($F(1; 44.3) = 4.288, p = 0.04$) compared to the non-PMDD group, irrespective of the menstrual phase they experienced.

In the second and third research phases based on the DRSP, the sample was divided into women with probable PMDD diagnosis (PMDD group, $n=3$) and those without probable PMDD diagnosis (nonPMDD group, $n=6$). A statistically significant association was observed between the classifications according to the DSM-5 Based Screening Tool and the DRSP ($p=0.048$; Cramer's $V=0.79$). The PMDD group showed a tendency of lower well-being and more severe anxio-depressive symptoms than the nonPMDD group (Well-being: between phases $p=0.93$, between groups $p=0.06$; BDI-II: between phases $p=0.79$, between groups $p=0.07$; STAI-S: between phases $p=0.87$, between groups $p=0.17$).

Conclusions: The prevalence of PMDD was high in our sample. Women with probable PMDD retrospectively reported substantial affective difficulties and a decline in subjective well-being, regardless of their menstrual cycle. Prospective preliminary findings suggest a trend toward differentiation associated with probable PMDD. These results highlight the need for prospective clinical studies addressing the psychological symptoms of women with PM issues and the importance of appropriate treatment of the clinical appearance of PMDD.

Disclosure of Interest: None Declared

EPP0055

The Mediating Role of Maladaptive Metacognitive Beliefs between Adverse Childhood Experiences and Trait Anxiety

D. Horváth¹, B. Kovács-Tóth¹, B. Oláh¹ and Z. Fekete^{1*}

¹Department of Behavioural Sciences, University of Debrecen, Debrecen, Hungary

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.288

Introduction: Adverse childhood experiences (ACE) have a significant negative impact on health. ACEs lead to more pronounced trait anxiety, among others, which serves as a basis for various mental and somatic symptoms. Recent findings suggest that the fact that individuals with more ACEs also have more maladaptive metacognitive beliefs may contribute to the development of these symptoms.

Objectives: We aim to study the possible mediating role of maladaptive metacognitive beliefs, resulting from adverse childhood experiences, on trait anxiety.

Methods: Data was collected online, anonymously, in a non-clinical population of adults over 18 years of age. The sample consisted of 304 subjects (84.21 % women, 15.79 % men). The applied questionnaires included a demographic questionnaire, the

Adverse Childhood Experiences Questionnaire 10 item version, the Meta-Cognitions Questionnaire, and the Spielberger Trait Anxiety Questionnaire. The system of correlations between the examined variables was explored using structural equation modeling (SEM). The study was carried out with ethical approval and in accordance with the Declaration of Helsinki.

Results: Our results confirm that ACEs have a significant impact on all the measured dimensions of maladaptive metacognitive beliefs. The direct effect of ACEs on adult trait anxiety is also significant. The results of the study on indirect effects support the joint mediating role of the five metacognitive dimensions. The strongest significant mediating effect was found for the uncontrollability and dangerousness of negative beliefs about worry. Both the direct and indirect effects of cognitive self-consciousness on adult trait anxiety are negative, which means that the more the cognitive self-consciousness is characteristic of someone, the lower the degree of trait anxiety in adulthood is.

Conclusions: Our results confirm the mediating effect of metacognitive beliefs on trait anxiety in the context of adverse childhood experiences, which points to the importance of further research on metacognition among the population that suffered early adversities. One of the limitations of the study roots in online data collection: the examined sample is not representative. Moreover, to extend the results, it is recommended to repeat the study on a clinical population. This would enable us to compare our results with those of the clinical population, which could provide further important results in the field of metacognition and adverse childhood experiences.

Disclosure of Interest: None Declared

Psychosurgery and Stimulation Methods (ECT, TMS, VNS, DBS)

EPP0056

Effect of intermittent theta-burst stimulation on the thyroid and adrenal systems in resistant depressed patients

F. Duval^{1*}, V. Danila¹, T. Weiss¹, F. Gonzalez Lopera¹ and M. Tomsa¹

¹Psychiatry, Centre Hospitalier APF2R, Rouffach, France

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.289

Introduction: Disturbances in the hypothalamic-pituitary-thyroid (HPT) and hypothalamic-pituitary-adrenal (HPA) axes have been frequently reported in treatment resistant depressed patients (TRDs). So far, the effects of intermittent theta-burst stimulation (iTBS) treatment—a form of repetitive transcranial magnetic stimulation (rTMS) technique—on the activity of the HPT and HPA axes are poorly understood.

Objectives: The present study aimed to evaluate the effects of iTBS sessions, applied to the left dorsolateral prefrontal cortex, in TRDs with abnormal chronobiological HPT functioning at baseline (BL) possibly associated with hypercortisolemia.

Methods: The $\Delta\Delta\text{TSH}$ test (i.e., the difference between the thyrotropin response to protirelin tests [ΔTSH] performed at 8 AM and 11 PM on the same day) and the dexamethasone suppression test (DST) were performed in 12 TRDs and 14 healthy hospitalized