

On the making of a PN: the interaction of a multiple stellar wind with the ISM

Arturo Manchado^{1,2,3} , Eva Villaver⁴, G. García-Segura⁵ and Luciana Bianchi⁶

¹Instituto de Astrofísica de Canarias, Vía Láctea S/N, E-38200 La Laguna, Tenerife, Spain

²Departamento de Astrofísica, Universidad de La Laguna (ULL), E-38206 La Laguna, Tenerife, Spain

³Consejo Superior de Investigaciones Científicas, Spain

⁴Centro de Astrobiología (CAB, CSIC-INTA), ESAC Campus Camino Bajo del Castillo, s/n, Villanueva de la Cañada, 28692, Madrid, Spain

⁵Instituto de Astronomía-UNAM, Apartado postal 877, Ensenada, 22800 Baja California, México

⁶Department of Physics and Astronomy, Johns Hopkins University, Baltimore, MD 21218, USA

Abstract. NGC 7293, the Helix nebula, represents one of the rare instances in which theoretical predictions of stellar evolution can be accurately tested against observations since the precise parallax distance and the velocity and proper motion of the star are well known. We present numerical simulations of the formation of the Helix PN that are fully constrained by the progenitor stellar mass, stellar evolution history, and star-interstellar medium (ISM) interaction. In the simulations, multiple bow-shock structures are formed by fragmentation of the shock front where the direct interaction of the stellar wind with the ISM takes place.

Keywords. hydrodynamics–ISM: planetary nebula: general; planetary nebula: individual (NGC 7293) – stars: AGB and post-AGB.

1. Introduction

NGC 7293 (a.k.a The Helix, PN G036.1 – 57.1) is one of the planetary nebula (PN) with a very good determination of its distance; 202 pc from Gaia DR3 (González-Santamaría et al. 2021). The central star has a mass of $0.60 \pm 0.02 M_{\odot}$ so (Benedict et al. 2009) and a high temperature $T_{eff} = 104\,000$ K (Guerrero & De Marco 2013) implying that the star+nebula system is in a rather evolved stage. The *Galaxy Evolution Explorer* (GALEX; Martin et al. 2005) wide field-of-view (26 arcminutes) revealed for the first time striking morphological features in the form of extended bow-shocks beyond the nebular halo (Bianchi et al. 2012).

2. Numerical simulations

The numerical simulations have been performed with the fluid solver ZEUS-3D (Stone, Mihalas, & Norman 1992), developed by M. L. Norman and the Laboratory for Computational Astrophysics. The computations have been carried out on a 2D spherical polar grid with the angular coordinate ranging from 0° to 180° and a physical radial extension of 3 pc. They have been run at a resolution of 1600×1440 zones in the radial and angular coordinates of the grid respectively (or equivalently ~ 388 au \times 0.125°).

© The Author(s), 2023. Published by Cambridge University Press on behalf of International Astronomical Union. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1. Left, *GALEX* FUV image of NGC 7293. At a distance of 202 pc, the FOV is 4.23 pc. Right, density map from our simulations, the snapshot corresponds to a ~ 1000 yr old PN. The morphology of the UV features is reproduced by the simulations; more important, the physical scale of the predicted structures matches the observations.

Our boundary conditions are the AGB stellar wind of a $1.5 M_{\odot}$ star and a ISM density of $n_{\text{ISM}} = 0.06 \text{ cm}^{-3}$, and a relative velocity respect to its ambient medium of 40 km s^{-1} . For the post-AGB stage and PN evolution we follow the stellar wind according to the prescription given in Villaver et al. (2002a) by using the post-AGB evolutionary sequence given by Vassiliadis & Wood (1994) for a hydrogen burner with solar metallicity for the assumed stellar mass.

3. RESULTS

In Fig.1, (right panel) we show the result of the numerical simulation after 819000 yr in the AGB, and 1000 yr, after the onset of the photoionization. In the left panel we show the *GALEX* FUV filter image, which at a distance of 202 pc has a FOV of 4.23 pc. It is remarkable that both figures show bow shock structures in the direction of the movement, and that the actual size in parsec is quite similar. We conclude that the morphology of the Helix can be explained by the evolution of a $1.5 M_{\odot}$ star interacting with an ISM with a relative velocity of 40 km s^{-1} . apparent multiple bow-shocks.

Acknowledgments

A. M. acknowledge support from the ACIISI, Gobierno de Canarias and the European Regional Development Fund (ERDF) under grant with reference PROID2020010051 as well as from the State Research Agency (AEI) of the Spanish Ministry of Science and Innovation (MICINN) under grant PID2020-115758GB-I00.

References

- Benedict, G. F., McArthur, B. E., Napiwotzki, R., et al. 2009, *AJ*, 138, 1969
 Bianchi, L., Manchado, A., Forster, K. 2012, *IAU Symp.* 283, p. 308
 González-Santamaría, I., Manteiga, M., Manchado, A., et al. 2021, *A&A*, 656, A51.
 doi:10.1051/0004-6361/202141916
 Guerrero, M. A., & De Marco, O. 2013, *A&A*, 553, A126
 Martin, D. C., Fanson, J., Schiminovich, D., et al. 2005, *ApJ Letters*, 619, L1

- Martin, D. C., et al. 2007, *Nature*, 448, 780
Stone, J. M., Mihalas, D., & Norman, M. L. 1992, *ApJS*, 80, 819
Villaver, E., García-Segura, & Manchado, A. 2002a, *ApJ*, 571, 880
Villaver, E., García-Segura, G., & Manchado, A. 2003, *ApJ Letters*, 585, L49
Vassiliadis, E., & Wood, P. R. 1994, *ApJS*, 92, 125