
Contents—continued from outside back cover]

S. RAMNATH and S. UTHAMASAMY	207	Effect of genotype and cultural practices on <i>Helicoverpa armigera</i> damage to cotton
E. M. MINJA and J. K. AMPOFO	217	Effect of sorghum and cowpea intercropping pattern on the spotted stemborer, <i>Chilo partellus</i> (Swinhoe) larval establishment in southwestern Kenya
J. ALLOTEY and J. A. DANKWAH	223	Some aspects of the biology and control of the cowpea weevil, <i>Callosobruchus maculatus</i> (F.) on bambara groundnut and cowpea
<i>Short communications</i>		
J. V. MUDEGU and T. K. MUKIAMA	229	Electrophoretic variation in wild <i>Anopheles pharoensis</i> Theobald in Mwea Irrigation Scheme, Kenya
A. K. CHANDRA, R. K. SAHA, J. BHATTACHARYA, N. KRISHNAN, S. K. SEN and B. SARATCHANDRA	233	Efficacy of carbendazim as an anti-microsporidial agent and its influence on the growth and cocoon characters of silkworm, <i>Bombyx mori</i> L.
<i>Editor's Note</i>	i	
<i>Announcements</i>	ii	
<i>Instructions to Authors</i>	iii–iv	
<i>Author Checklist</i>	v	
<i>Erratum</i>	vi	

Insect Science and its Application

The International Journal of Tropical Insect Science

VOLUME 16 NUMBER 2

1995

CONTENTS

- S. OGHIAKHE 123 *Mini Review*
Progress and future prospects in utilisation of host plant resistance in the control of insect pests of cowpea, *Vigna unguiculata* in Africa
- D. P. GIGA 145 *Research Articles*
Selection of oviposition sites by the cowpea weevils *Callosobruchus rhodesianus* (Pic.) and *Callosobruchus maculatus* (F.)
- K. TATA-HANGY and J. DEN HOLLANDER 151 Key to the common mealybugs (Homoptera; Coccoidea; Pseudococcidae) occurring in the Central Africa region
- L. W. IRUNGU and R. W. MWANGI 159 Effects of a biologically active fraction from *Melia volkensii* on *Culex quinquefasciatus*
- J. O. MIDIWO, R. W. MWANGI and Y. GHEBREMESKEL 163 Insect antifeedant, growth-inhibiting and larvycidal compounds from *Rapanea melanphloes* (Myrsinaceae)
- P. B. TANZUBIL 167 Effects of neem (*Azadirachta indica* A. Juss) extracts on food intake and utilisation in the African armyworm, *Spodoptera exempta* (Walker)
- L. B. S. ANDRADE, V. M. Q. FLORES, M. P. SALES, R. A. MORAES, A. BLANCO-LABRA and J. XAVIER-FILHO 171 Effect of protein fractions isolated from cereal grains on the development and survival of the cowpea weevil, *Callosobruchus maculatus*
- S. JANARTHANAN, K. M. SUBBU RATHINAM and M. KRISHNAN 177 The effect of photoperiod on the haemolymph protein profile in the silkworm, *Bombyx mori* L.
- E. N. MAGIRI, V. N. KONJI, D. W. MAKAWITI and J. MIDIWO 183 Effect of plant quinones on insect flight muscle mitochondria
- L. F. FORY, F. A. TENJO, C. E. POSSO, C. CARDONA and J. E. MAYER 191 Growth-inhibitory effects of proteinaceous fractions from resistant wild bean accessions on the bean weevil, *Acanthoscelides obtectus*
- V. KUMAR, L. S. YADAV, A. N. VERMA, R. K. KASHYAP and R. KUMAR 199 Development and reproductive potential of *Helicoverpa armigera* (Hubner) on various plant parts of cotton cultivars/hybrid

[continued on inside back cover

Indexed/Abstracted in
Current Contents, CABS, C. A. B. International, BIOSIS, The African
Book Publishing Record

ISSN 0191-9040
ISIADL 16(2) 123-236 (1995)

Typeset by ICIPE Science Press, Nairobi, Kenya
Printed by Regal Press, Nairobi, Kenya