

Inulin supplementation reduces the negative effect of a high-fat diet rich in SFA on bone health of growing pigs

Monika Sobol*, Stanisława Raj and Grzegorz Skiba

Department of Animal Nutrition, The Kielanowski Institute of Animal Physiology and Nutrition, Polish Academy of Sciences, 05-110 Jabłonna, Poland

(Submitted 31 October 2017 – Final revision received 26 January 2018 – Accepted 27 February 2018)

Abstract

Consumption of a high-fat diet, rich in SFA, causes deterioration of bone properties. Some studies suggest that feeding inulin to animals may increase mineral absorption and positively affect bone quality; however, these studies have been carried out only on rodents fed a standard diet. The primary objective of this study was to determine the effect of inulin on bone health of pigs (using it as an animal model for humans) fed a high-fat diet rich in SFA, having an unbalanced ratio of lysine:metabolisable energy. It was hypothesised that inulin reduces the negative effects of such a diet on bone health. At 50 d of age, twenty-one pigs were randomly allotted to three groups: the control (C) group fed a standard diet, and two experimental (T and TI) groups fed a high-fat diet rich in SFA. Moreover, TI pigs consumed an extra inulin supply (7% of daily feed intake). After 10 weeks, whole-body bone mineral content ($P=0.0054$) and bone mineral density ($P=0.0322$) were higher in pigs of groups TI and C compared with those of group T. Femur bone mineral density was highest in pigs in group C, lower in group TI and lowest in group T ($P=0.001$). Femurs of pigs in groups TI and C had similar, but higher, maximum strength compared with femurs of pigs in group T ($P=0.0082$). In conclusion, consumption of a high-fat diet rich in SFA adversely affected bone health, but inulin supplementation in such a diet diminishes this negative effect.

Key words: Bone mineralisation: High-fat diets: Inulin: Pigs: X-ray absorptiometry

Nutrition affects human growth and development from an early age. However, dietary habits of many children and adolescents are characterised by low consumption of fruit, vegetables, whole grains and fibre, and high intake of fast food, soft drinks and salty snacks^(1,2). Over-consumption of energy-rich foods, along with low physical activity, can contribute to an increased risk of civilisational diseases, such as obesity⁽³⁾. In recent years, researchers^(1,4) have reported an alarming increase in the prevalence of childhood overweight and obesity. Globally, childhood overweightness and obesity affect many low- and middle-income countries, particularly in urban settings⁽⁵⁾. Ogden *et al.*⁽⁶⁾ reported that, in the USA, approximately one-third of early adolescents (10–14 years) are overweight or obese. Overweightness and obesity also are a serious public health problem in Europe, where approximately 20% of the early adolescents are overweight or obese⁽⁴⁾. In addition, it is highly probable that overweight and obese children will stay obese during adulthood. It is also more likely that these individuals will develop non-communicable diseases, such as type 2 diabetes and cardiovascular diseases. In addition, it is well known that consumption of a high amount of fat, specifically fat characterised by high

content of unhealthy fatty acids (SFA), causes deterioration of bone properties and consequently can cause osteoporosis, even at a young age⁽⁵⁾.

Because eating habits are difficult to change within a population, one of the ways to improve bone health may be probiotic supplementation. It is well known that the intestinal microbiota influences not only the health of the intestine but also affects the skin, lungs, arteries and bones^(7–9). Inulin is one of the probiotics affecting the composition of gut microbiota and stimulates the growth of bifidobacteria and/or lactobacilli, which increase SCFA formation⁽¹⁰⁾. SCFA directly contribute to increased absorption of Ca, Mg and Fe in the intestines⁽¹¹⁾ and indirectly have an immunostimulating effect⁽¹²⁾. Some studies suggest that feeding inulin to animals may affect bone quality⁽¹³⁾ owing to increased mineral absorption. However, such conclusions were based on the concentration of minerals in the blood plasma and organs such as the liver, kidneys and heart⁽¹⁴⁾, or were based on total mineral balance⁽¹⁵⁾. Thus, it is obvious that high consumption of unhealthy fat (rich in SFA) deteriorates bone quality, whereas inulin improves bones quality.

Abbreviations: BMC, bone mineral content; BMD, bone mineral density; C, group of pigs fed a standard diet; DXA, dual-energy X-ray absorptiometry; ME, metabolisable energy; T, group of pigs fed a high-fat diet rich in SFA; TI, group of pigs fed a high-fat diet rich in SFA and supplemented with inulin.

* **Corresponding author:** Dr M. Sobol, email m.sobol@ifzz.pl

Most studies on the influence of direct effects of inulin on bone properties have been conducted on rodents fed a standard diet. In turn, the influence of high consumption of fat on bone quality usually has been compared with animals fed a standard diet. Recent studies have demonstrated that, for humans, pigs are a better model than rodents owing to several similarities in anatomy, physiology, metabolism and pathology^(16–18). Obviously, there are interspecies differences in the skeletal structure and load, but factors of research interest affect the properties of bones of both pigs and humans in a similar manner.

The aim of this study was to determine the effect of inulin on whole-body mineral content, bone density and femur properties in growing pigs fed a high-fat diet rich in SFA, having an unbalanced ratio of lysine:metabolisable energy (ME). It was hypothesised that inulin would reduce the negative effects of such a diet on bone health.

Methods

According to the 3R (Replacement, Reduction and Refinement) ethical principle, the design of the study and experimental techniques used through the analysis allowed to minimise the number of animals while maintaining high statistical precision. There is no way to completely replace live animals with another research model. Thus, pigs have been chosen as a model for humans because of several similarities in anatomy, physiology, metabolism and pathology.

Experimental procedures used throughout this study were performed in accordance with national/local ethical guidelines and approved by the III Local Ethics Committee on Animal Experimentation of the Warsaw University of Life Sciences – SGGW, Poland.

Animals, diets and housing

This study was carried out at The Kielanowski Institute of Animal Physiology and Nutrition, Polish Academy of Sciences, on twenty-one cross-bred pigs (gilts) (male Duroc × female Polish Large White × Danish Landrace) growing from 50 to 120 d of age. At 50 d of age, animals were randomly allotted to three groups (*n* 7 per group): the control (C) group fed a standard diet (10 g of standardised ileal digestible lysine, 13.5 MJ ME per kg diet, and 0.72 of lysine:ME ratio) balanced according to nutrient requirements of swine⁽¹⁹⁾, two experimental (T and TI) groups fed a high-fat diet rich in SFA (141 g of diethyl ether extract, 7 g of standardised ileal digestible lysine, 15.4 MJ ME per kg diet, and 0.44 of lysine:ME ratio). Pigs fed diet TI received an extra supplement of inulin extracted from chicory root (100% inulin content, 0% sweetness level), with an average degree of polymerisation of at least 23 (Inulin Orafit[®]HPX; BENEIO GmbH). A review of the literature indicated that supplementation of inulin in diets of pigs in previous studies ranged from 0.5 to 9%^(20,21). However, results from some studies^(14,21) indicated that the effect of inulin on the tested parameters depended on its supply. Generally, higher doses of inulin had a greater impact on tested parameters. On the basis of these findings, it was decided that pigs fed diet TI will receive an extra supply of

inulin in an amount of 7% of daily feed intake. This amount of inulin could be successfully used in food products for humans, among others as a component in baked goods^(22,23).

Mineral content in standard and high-fat diets was the same and balanced according to nutrient requirements of swine guidelines⁽¹⁹⁾. Amino acid content in the high-fat diet was not equalised to that in the standardised diet because the aim of the study was to compare the influence of an unbalanced ratio of lysine to ME and elevated consumption of SFA (similar to the Western-type diet in human nutrition) on bone health. Beef tallow, rich in SFA, was used at 13% as fat supplement. Composition and nutritive value of diets are given in Table 1. The fatty acid content (%) of feeds is shown in Table 2 (data analysed). Pigs were fed semi *ad libitum* (95% *ad libitum* intake) twice daily (08.00 and 14.00 hours). This facilitated full control of nutrient intake by pigs.

Piglets were randomly selected from a population and kept individually on a concrete floor without straw in pens (3.3 m²) equipped with nipple drinkers. The environmental conditions in the piggery were as follows: air temperature of 18–20°C, humidity of 60–70% and air flow of 0.2–0.4 m/s, which was regulated by a Fancom ventilation system (Fancom BV) and in accordance with the European Union law⁽²⁴⁾. The pigs were

Table 1. Ingredients, chemical composition and nutritive value of diets

Indices	Standard diet	High-fat diet
Ingredients (g/kg)		
Barley	200	200
Wheat	370	120
Maize	225	360
Rapeseed meal	–	80
Soyabean meal	150	80
Beef tallow	–	130
Rapeseed oil	20	–
Premix A*	35	–
Premix B*	–	35
Chemical composition (g/kg)		
DM	880	884
Ash	25.5	24.1
Organic matter	855	860
Crude protein	152	135
Diethyl ether extract	43.7	141.0
Crude fibre	35.6	34.5
Starch	478	410
Sugar	34	29
Nutritive value (determined) (g/kg)		
Digestible crude protein	124.0	110.0
Apparent ileal digestible		
Lys	10.0	7.0
Met	3.0	2.1
Thr	6.2	4.5
Trp	2.0	1.4
P, digestible	3.0	3.0
Ca	7.8	7.8
Na	1.5	1.5
Metabolisable energy (MJ/kg)	13.5	15.4
Lys:metabolisable energy (g/MJ)	0.74	0.45

* Provided per kg diet: µg: vitamin A, 4500; vitamin D₃, 50; vitamin E, 100; mg: Fe, 240; Zn, 150; Cu, 25; Mn, 20; I, 0.30; Se, 0.30; vitamin K₃, 3.5; vitamin B₁, 3.5; vitamin B₂, 7.0; vitamin B₆, 5.0; vitamin B₁₂, 0.40; biotin, 0.20; folic acid, 3.0; nicotinic acid, 30; Ca -D pantothenate, 15; choline chloride, 400; g: Ca, 6.8 (premix A) and 6.6 (premix B); P, 2.9 (premix A) and 2.8 (premix B); essential amino acids: g: lysine, 4.51 (premix A) and 1.94 (premix B); methionine, 0.90 (premix A); threonine, 2.10 (premix A) and 0.25 (premix B).

Table 2. Fatty acid content of the diets (%)

Fatty acids	Standard diet	High-fat diet
Total SFA	23.67	31.30
C12:0	ND	0.12
C14:0	0.23	0.98
C16:0	20.48	24.13
C18:0	2.96	6.07
Total MUFA	31.17	30.16
C16:1 <i>n</i> -7	0.32	0.82
C18:1 <i>n</i> -7	1.87	1.77
C18:1 <i>n</i> -9	28.98	27.52
Total PUFA	45.16	38.54
C18:2 <i>n</i> -6	41.35	35.69
C18:3 <i>n</i> -3	3.37	2.46
C20:4 <i>n</i> -6	0.39	0.35
C20:5 <i>n</i> -3	ND	ND
C22:5 <i>n</i> -3	0.05	0.04
C22:6 <i>n</i> -3	ND	ND

ND, value below 0.01 was classified as not determined.

maintained on a 12 h day–12 h night cycle, with lights on from 06.00 to 18.00 hours.

Measurements and calculations

At 120 d of age, after 16 h of fasting, each pig was scanned using dual-energy X-ray absorptiometry (DXA) (Norland XR-800™ densitometer scanner with a ‘Whole-Body Scan’ type, Norland, A Cooper Surgical Company) to determine whole-body mineralisation – that is bone mineral content (BMC, g) and bone mineral density (BMD, g/cm²). Before scanning, animals were subjected to short-lasting sedation by injection using a mixture of ketamine hydrochloride (2 mg/kg body weight) and xylazine (0.2 mg/kg body weight)⁽²⁵⁾. Next, pigs were placed in a ventral position with all limbs extended. The DXA scans were obtained according to the standard procedures described by the manufacturer for scanning and analysis. A quality assurance test to verify the stability of the system calibration (control scans) was performed on a daily basis (as indicated by the manufacturers) to ensure precision. The daily calibration procedure was performed using QC Phantom and QA Calibration standard (Norland; A Cooper Surgical Company). The analysis was performed with the whole-body application. Two scans were performed on each pig.

After scanning, animals were slaughtered by exsanguination at the Institute’s experimental slaughter house. After slaughter, the femur was manually dissected from the right side of each carcass. After excision, the femur was cleaned of any remaining flesh, weighed and frozen (–20°C) for subsequent DXA scan (Norland XR-800™ densitometer scanner with a ‘Research Scan’ type, Norland). Specimens for scanning were thawed at room temperature (23°C) for 12 h before use. During scanning, the femur was positioned horizontally with the femoral head facing upwards and the condyles downwards and was scanned from the distal to the proximal end. All scans were performed in duplicate in order to avoid any rotation of the bone, because inconsistencies in their orientation adversely affect test precision. All scans were performed by the same operator. Femur BMC and BMD were recorded. After scanning, the

three-point bending test was applied to determine the mechanical bone characteristics using a TA-HDi Texture Analyser (Godalming) with head speed of 1 kN, 10 mm/min. The relationship between loading force perpendicular to the longitudinal axis of the bone and the resulting displacement was determined graphically. The values of maximum strength (N) and maximum elastic strength (N) were determined. The distance between supports of the bone was set at 40% of the femur length and the measuring head loaded bone samples at the midshaft with a constant speed of 50 mm/min. The geometrical properties of each femur were determined on the basis of the measurements of horizontal and vertical diameters (both external and internal) recorded after cutting the bone. The measurements were collected using an electronic ruler. The values of cross-sectional area (mm²), cortical wall thickness (mm) and cortical index (%) were determined in accordance with the following mathematical formulas:

$$\text{Cross-section area (mm}^2\text{)} = [\pi(HV - bv)] / 4$$

$$\text{Cortical wall thickness (mm)} = [(V + H) - (v + b)] / 4$$

$$\text{Cortical index (\%)} = \{[(H - b / H) + (V - v / V)] / 2\} 100,$$

where *V* is the vertical external diameter (mm), *H* the horizontal external diameter (mm), *v* the vertical internal diameter (mm) and *b* the horizontal internal diameter (mm)⁽²⁶⁾.

Chemical analysis

DM, ash, crude protein, crude fibre, diethyl ether extract, simple sugar and starch contents in the diets were determined as detailed by Horwitz & Latimer⁽²⁷⁾ (procedures: 934.01, 942.05, 984.13, 978.10, 920.39, 974.06 and 920.40, respectively). Content of ME and standardised ileal digestible amino acids in the diets were calculated as detailed by the nutrient requirements of swine⁽¹⁹⁾.

Lipids from the diets were saponified with potassium hydroxide according to the method described by Folch *et al.*⁽²⁸⁾. Next, methyl esters were prepared by esterification with thionyl chloride (4% in methanol) and extracted with *n*-heptane. Fatty acid methyl esters were analysed using a GC-2010AF Shimadzu Gas Chromatograph (Shimadzu Europa GmbH) equipped with a flame ionisation detector. The derivatives were separated on a capillary column (BPX70, 60-m length, 0.25-mm internal diameter and 0.25-µm film thickness; Phenomenex). The operating conditions were as follows: carrier gas, helium; split ratio, 1:100; injector and detector temperature, 260°C; the initial column temperature of 110°C was held for 5 min, then increased to 200°C at a rate of 3.5°C/min, and was held for 2.5 min, then increased to 205°C at a rate of 0.3°C/min, then increased to 215°C at a rate of 1.5°C/min and was held for 3 min. Individual fatty acid peaks were identified in comparison with the commercial standard, Supelco 37 Component FAME Mix (SUPELCO).

Statistical analysis

Statistical analyses were performed using Statgraphics Centurion (version 16.1.18, 2011) software (StatPoint Technologies Inc.). With an α level of 0.05, power established at 80% and an effect size of 0.90, the required total sample size



was 24 (i.e. n 8/group). The hypothesised effect size of 0.90 was calculated from the descriptive statistics of a previous study⁽²⁹⁾. *Post hoc* calculations using the hypothesised effect size and the total sample size of twenty-one (i.e. n 7/group) indicated that the actual power achieved in this study was 79.7%. Data are presented as means and standard deviations. The effect of diets on performance, whole-body mineralisation and femur properties of pigs was analysed using a one-way ANOVA. When the F ratio was significant, Tukey's HSD *post hoc* analysis was performed. Statistical significance was set at $P < 0.05$. A borderline significant trend was set at $P < 0.09$.

Results

Final body weight did not differ among groups C and T; however, body weights of pigs in groups C and T were lower than pigs in group TI ($P = 0.0211$) (Table 3). Daily feed intake ranged from 1.58 kg for pigs in group T to 1.69 kg for pigs in group C ($P = 0.0128$). Pigs in groups T and TI had similar, but higher, daily ME intake from the feed than pigs in group C ($P = 0.0037$). In addition, pigs in group TI received daily approximately 116 g of extra inulin supply, which is 0.72 MJ/d. Pigs in groups T and TI had similar, but lower, total feed intake

($P = 0.0115$) and total lysine intake ($P = 0.0000$) than pigs in group C. Total energy intake from the feed did not differ among groups. Average daily gain was higher ($P = 0.0184$) for pigs in group TI than those in groups C and T; however, the feed conversion ratio ranged from 1.88 kg of feed/kg gain for pigs in group T to 2.03 kg of feed/kg gain for pigs in group C ($P = 0.0434$). Whole-body BMC ($P = 0.0054$) and BMD ($P = 0.0322$) were higher for pigs in groups TI and C compared with pigs in group T (Table 4).

Mass and length of femurs did not differ between treatments (Table 5). Pigs of groups TI and C had similar, but higher, femur BMC compared with group T ($P = 0.0009$). Femur BMD was highest for pigs in group C, followed by pigs in group TI, and lowest for pigs in group T ($P = 0.0001$). Femurs of pigs in groups TI and C had similar, but higher, maximum strength compared with group T ($P = 0.0082$), whereas maximum elastic strength was highest for pigs in group C, followed by pigs in group TI, and lowest in group T ($P = 0.0001$). Femur cross-sectional area tended to be lower for pigs in group TI than for pigs in group C ($P = 0.0753$). Femurs of pigs in groups TI and T had similar, but lower, cortical wall thickness than femurs of pigs in group C ($P = 0.0100$). However, the cortical index ranged from 42.3% for pigs in group T to 45.8% for pigs in group C ($P = 0.0366$).

Table 3. Performance of pigs during experimental period (Mean values and standard deviations)

Items	Group (n 7)						P (ANOVA)
	C		T		TI		
	Mean	SD	Mean	SD	Mean	SD	
Initial BW (kg)	12.7	0.43	12.3	0.31	13.2	0.24	0.2233
Final BW (kg)	71.3 ^a	1.31	71.1 ^a	0.91	74.7 ^b	1.17	0.0211
Daily feed intake (kg)	1.69 ^b	0.03	1.58 ^a	0.05	1.63 ^{a,b}	0.05	0.0128
Daily inulin supply (kg)	–	–	–	–	0.116	0.003	–
Daily metabolisable energy intake from the feed (MJ)	23.2 ^A	0.48	24.4 ^B	0.61	25.1 ^B	0.70	0.0037
Daily metabolisable energy intake from inulin (MJ)	–	–	–	–	0.72	0.020	–
Total feed intake (kg)	124.0 ^b	1.717	115.3 ^a	3.43	114.3 ^a	4.24	0.0115
Total inulin supply (kg)	–	–	–	–	8.0	0.114	–
Total lysine intake (kg)	1.22 ^A	0.03	0.78 ^B	0.02	0.78 ^B	0.04	0.0000
Total energy intake from the feed (MJ ME)	1706.7	23.47	1775.5	52.78	1761.5	57.40	0.2521
Total energy from inulin supply (MJ ME)	–	–	–	–	50.4	2.13	–
Average daily gain (g)	837 ^a	3.30	844 ^a	2.83	896 ^b	2.92	0.0184
Feed conversion ratio (kg feed/kg gain)	2.03 ^b	0.08	1.88 ^a	0.07	1.95 ^{a,b}	0.09	0.0434

C, pigs fed a standard diet; T, pigs fed a high-fat diet rich in SFA; TI, pigs fed a high-fat diet rich in SFA and supplemented with inulin; BW, body weight; ME, metabolisable energy. ^{A,B} Mean values within a row with unlike superscript letters were significantly different ($P < 0.01$). ^{a,b} Mean values within a row with unlike superscript letters were significantly different ($P < 0.05$).

Table 4. Whole-body mineralisation performed at the end of the study (at 120 d of age) (Mean values and standard deviations)

Items	Group (n 7)						P (ANOVA)
	C		T		TI		
	Mean	SD	Mean	SD	Mean	SD	
BMC (g)	1809.8 ^B	21.02	1641.1 ^A	36.51	1752.8 ^B	27.06	0.0054
BMD (g/cm ²)	0.993 ^b	0.02	0.936 ^a	0.05	0.987 ^b	0.02	0.0322

C, pigs fed a standard diet; T, pigs fed a high-fat diet rich in SFA; TI, pigs fed a high-fat diet rich in SFA and supplemented with inulin; BMC, bone mineral content; BMD, bone mineral density. ^{A,B} Mean values within a row with unlike superscript letters were significantly different ($P < 0.01$). ^{a,b} Mean values within a row with unlike superscript letters were significantly different ($P < 0.05$).

Table 5. Femur morphometric, geometric, densitometric and mechanical properties at the end of the study (at 120 d of age) (Mean values and standard deviations)

Items	Group (n 7)						P (ANOVA)
	C		T		TI		
	Mean	SD	Mean	SD	Mean	SD	
Mass (g)	255.3	2.30	245.9	2.20	251.5	4.21	0.5599
Length (cm)	17.54	0.38	17.64	0.23	17.68	0.48	0.8374
BMC (g)	67.98 ^B	2.41	62.02 ^A	1.30	65.48 ^B	1.58	0.0009
BMD (g/cm ²)	1.053 ^C	0.013	0.976 ^A	0.020	1.022 ^B	0.020	0.0001
Maximum strength (N)	430.3 ^B	28.44	368.1 ^A	32.24	420.6 ^B	20.80	0.0082
Maximum elastic strength (N)	349.0 ^C	15.90	276.5 ^A	20.31	320.9 ^B	15.00	0.0001
Cross-sectional area (mm ²)	347.9	34.19	318.5	21.11	297.8	23.10	0.0753
Cortical wall thickness (mm)	5.73 ^B	0.42	5.22 ^A	0.14	5.13 ^A	0.16	0.0100
Cortical index (%)	45.8 ^b	1.78	42.3 ^a	1.64	43.5 ^{a,b}	2.18	0.0366

C, pigs fed a standard diet; T, pigs fed a high-fat diet rich in SFA; TI, pigs fed a high-fat diet rich in SFA and supplemented with inulin; BMC, bone mineral content; BMD, bone mineral density.
^{A,B,C} Mean values within a row with unlike superscript letters were significantly different ($P < 0.01$). ^{a,b} Mean values within a row with unlike superscript letters were significantly different ($P < 0.05$).

Discussion

Composition of pig diets

Studies carried out on animals^(30–33) and humans⁽³⁴⁾ indicated that the amount and source of dietary fat may affect bone metabolism by changing Ca absorption, osteoblast formation, lipid oxidation and synthesis of pro- and anti-inflammatory eicosanoids. However, the aim of this study was to compare the influence of an unbalanced ratio of lysine:ME and elevated consumption of SFA (similar to the Western-type diet in human nutrition) on bone properties.

Performance of pigs

The results of the present study show that pigs fed a high-fat diet supplemented with inulin grew faster than those fed either a high-fat diet that was not supplemented with inulin or a standard diet. However, the feed conversion ratio differed only between pigs fed a standard diet and a high-fat diet not supplemented with inulin. Some authors^(20,29) previously found that feed supplementation with inulin increased mass gain in pigs. However, these authors used only a standard diet supplemented with inulin and suggested that this gain resulted from a favourable impact of inulin on the population of beneficial micro-organisms in the alimentary tract, which in turn protect the host against development of pathogenic strains. Consequently, this protection leads to higher growth rates and better feed utilisation^(10,20). On the other hand, studies carried out on rodents by Duan *et al.*⁽³⁵⁾, Kumar *et al.*⁽³⁶⁾ and Pouyamanesh *et al.*⁽¹⁰⁾ showed that inulin supplementation to a high-fat diet resulted in weight gain attenuation in rodents compared with rodents fed a high-fat diet that was not supplemented with inulin. Discrepancies between our results and results of previous studies may be owing to differences in the kind and amount of fat supplementation, as well as the level of inulin addition. Other reasons for differences between the results of our study and previous studies could be because of differences in tolerance of species for high amounts of dietary fat. Comparing data in the literature concerning the proportion

of energy from fat to total energy in the diet, it seems that rodents are able to tolerate feed with a higher ratio of fat (even 60%) compared with pigs (approximately 28% in the present study).

Whole-body mineralisation

On performing DXA analysis, we were able to determine the BMC and BMD of each pig and relate it to the dietary treatment that was of interest in our study. Recent studies have demonstrated that, for humans, pigs are a better model than rodents owing to several similarities in anatomy, physiology, metabolism and pathology⁽¹⁶⁾. Moreover, pigs have similar lamellar bone structure and similar rates of bone regeneration, cortical bone mineralisation and bone remodelling rates as humans^(17,18). Obviously, there are interspecies differences in the skeletal structure and load, but factors of research interest affect the properties of bones of both pigs and humans in a similar manner. We found that inulin supplementation to a high-fat diet increased the whole-body BMC and BMD compared with animals fed a high-fat diet that was not supplemented with inulin. However, whole-body mineralisation did not differ among pigs fed a standard diet and a high-fat diet supplemented with inulin. Few data have been published on the influence of inulin (and other oligosaccharides) supplementation on the whole-body bone parameters measured by DXA. Studies dealing with this issue have been carried out mainly on rodents^(9,13,37,38); however, a major part of these studies only examined the use of a standard diet. Some authors found that the addition of inulin improved whole-body bone parameters⁽¹³⁾. A similar finding was reported by Devareddy *et al.*⁽³⁷⁾ using a standard diet supplemented with fructo-oligosaccharides. Furthermore, inulin-type fructan supplementation increased whole-body BMC and BMD compared with a placebo group during a long-term study on human adolescents⁽³⁹⁾. In contrast, some authors^(9,38) did not find any effect of inulin supplementation to a standard diet on the whole-body bone parameters of rats.

Numerous studies performed using animal and human models showed that beneficial effects of fructose-based

oligosaccharide supplementation (among other types of inulin) resulted from stimulation of mineral absorption, mainly Ca, but also Mg, Zn and Fe^(40,41). Furthermore, by increasing the absorption of minerals, BMC and BMD improved. Simultaneously, peak bone mass also increased, which reduces the risk of developing osteopaenia and osteoporosis^(13,42). Mechanisms of this response could be associated with lower luminal pH, resulting from SCFA produced by *Bifidobacterium* after prebiotic fermentation⁽⁷⁾. Reduced pH also changed speciation and solubility of minerals and increased their bioavailability⁽⁴³⁾. Few minerals are essential co-factors for enzymes involved in the synthesis of collagen and other bone matrix constituents that are required to build up the organic bone matrix, which is the precondition for mineral accretion (e.g. Mg, Zn, B and Mn)⁽⁷⁾.

On the basis of the literature data, several hypotheses have been suggested to explain how inulin enhances mineral absorption, mainly Ca as the main bone component. First, increased Ca solubility in the colon resulted from pH reduction as a consequence of inulin fermentation^(7,43). Second, osmotic effects increase fluid transfer in the colonic lumen and, as a consequence, increase the permeability between intracellular enterocyte junctions that facilitates diffusion⁽⁴⁴⁾. Third, Ca/H exchange in the distal part of the colon is activated by absorption of short-chain carboxylic acids^(11,45).

Our results showed that feeding pigs a high-fat diet rich in SFA and not supplemented with inulin resulted in strong deterioration of whole-body bone parameters compared with a standard diet. Fehrendt *et al.*⁽³²⁾ found similar results in mice.

Data in the literature show that fatty acid composition of the fat source in the diet also had a significant influence on bone health. In a study by Wang *et al.*⁽⁴¹⁾ on mice, a high-fat diet rich in SFA resulted in lower values of bone mineralisation compared with mice fed a standard diet. These authors did not find such negative effects when a high-fat diet was rich in MUFA. Other authors also found that SFA may have effects that could impair bone health^(46,47), whereas MUFA and PUFA, in particular *n*-3 PUFA, have a beneficial effect on bone health⁽³⁴⁾.

Properties of femur

Densitometry measurements of whole-body bone parameters cannot be compared with data on selected bones because they represent an average value that can mask differences between rapid- and slow-growing bones⁽¹³⁾. Therefore, despite whole-body mineralisation, femur properties also were determined.

A review of the available literature did not provide information on the effects of inulin supplementation to a high-fat diet on the bone health of growing pigs. The present study showed that neither a high-fat diet nor inulin supplementation affected mass and length of femur bones. Inulin supplementation to a high-fat diet, however, improved femur parameters, resulting in values similar to those obtained in pigs fed a standard diet. However, cortical wall thickness and cross-sectional area were similar to those of pigs fed a high-fat diet that was not supplemented with inulin. The results of this study were consistent with those of Fehrendt *et al.*⁽³²⁾, who showed negative impacts of a high-fat diet on bone microstructure in mice, including a

missarrangement of cell-cell and cell-matrix contacts, osteoblast and trabecular structure, and collagen-1 and osteoid synthesis. Bielohuby *et al.*⁽³⁰⁾ and Patsch *et al.*⁽⁴⁸⁾ also showed an adverse effect of high-fat diet on BMD, bone microarchitecture (i.e. trabecular number and connectivity density) and bone strength (i.e. maximum fracture load, bone stiffness and energy to failure) compared with a standard diet. Bielohuby *et al.*⁽³⁰⁾ suggested that lower BMD and mechanical strength of femurs of rats fed a high-fat diet resulted from lower serum concentration of markers of bone formation and, simultaneously, unchanged serum concentration of markers of bone resorption. Moreover, lower rates of mesenchymal cells differentiating into osteoblasts might be explained by reductions of approximately 70–80% in the expression of transcription factors influencing osteoblastogenesis in bone marrow. Furthermore, in studies carried out on mice, some authors found that a high-fat diet reduced femoral trabecular bone mass owing to an increased trabecular separation and a reduced trabecular number and connectivity density⁽³¹⁾. However, a considerable amount of dietary fat was lard, which is characterised by high content of SFA⁽¹⁹⁾. It seems that the amount of fat consumed, as well as its fatty acid profile, could play an important role in organismal response. Results of the present study clearly showed that pigs fed a high-fat diet (rich in SFA) had worse parameters of femurs compared with pigs fed a standard diet. Wang *et al.*⁽⁴¹⁾ also found that rats fed a high-fat diet rich in SFA had lower values of femur volumetric BMD than those fed a control or a high-fat diet rich in MUFA. The negative effect of SFA on bone metabolism can be achieved by several mechanisms. It has been suggested that SFA may interfere with osteoblastogenesis and bone formation⁽⁴⁹⁾. Other authors showed that high content of SFA could increase bone resorption by elevating the expression of inflammatory cytokines and the receptor activator of NF- κ B ligand⁽⁴⁶⁾ or that SFA can participate in osteoclastogenesis resulting in decreased osteoclast apoptosis⁽⁵⁰⁾.

In conclusion, inulin supplementation to a high-fat diet rich in SFA reduced the negative effect of such a diet on whole-body mineralisation, as well as on femur mineral status and strength.

Acknowledgements

This research received no specific grant from any funding agency or from commercial or not-for-profit sectors.

M. S., S. R. and G. S. conceived and designed the study, interpreted the data and drafted the manuscript. M. S. conducted the research and acquired the data and performed the statistical analysis.

None of the authors has any conflicts of interest to declare.

The amount of inulin used in animal studies could reasonably be expected to be achieved in the human population, for example, in food products, among others as a component in baked goods or dairy products.

References

1. Kafeshani O, Sarrafzadegan N, Nouri F, *et al.* (2015) Major dietary patterns in Iranian adolescents: Isfahan Healthy Heart Program, Iran. *ARYA Atheroscler* **11**, 61–68.



2. Mesirow MS & Welsh JA (2015) Changing beverage consumption patterns have resulted in fewer liquid calories in the diets of US children: National Health and Nutrition Examination Survey 2001–2010. *J Acad Nutr Diet* **115**, 559–566.
3. Kar S & Khandelwal B (2015) Fast foods and physical inactivity are risk factors for obesity and hypertension among adolescent school children in east district of Sikkim, India. *J Nat Sci Biol Med* **6**, 356–359.
4. Pigeot I, Barba G, Chadigeorgiou C, *et al.* (2009) Prevalence and determinants childhood overweight and obesity in European countries: pooled analysis of the existing surveys within the IDEFICS Consortium. *Int J Obes (Lond)* **33**, 1103–1110.
5. World Health Organization (2012) *Population-Based Approaches To Childhood Obesity Prevention*. Geneva: WHO Library Cataloguing-in-Publication Data, WHO Document Production Services.
6. Ogden CL, Carroll MD, Kit BK, *et al.* (2014) Prevalence of childhood and adult obesity in the United States, 2011–2012. *J Am Med Assoc* **311**, 806–814.
7. Scholz-Ahrens KE, Ade P, Marten B, *et al.* (2007) Prebiotics, probiotics, and synbiotics affect mineral absorption, bone mineral content, and bone structure. *J Nutr* **137**, 838S–846S.
8. Ortiz LT, Rodriguez ML, Alzueta C, *et al.* (2009) Effect of inulin on growth performance, intestinal tract sizes, mineral retention and tibial bone mineralisation in broiler chickens. *Br Poult Sci* **50**, 325–332.
9. Weaver CM, Martin BR, Story JA, *et al.* (2010) Novel fibers increase bone calcium content and strength beyond efficiency of large intestine fermentation. *J Agric Food Chem* **58**, 8952–8957.
10. Pouyamanesh Z, Amoli MM, Yaghmaei P, *et al.* (2016) Effect of inulin supplementation in male mice fed with high fat diet on biochemical profile and α -amylase gene expression. *Trop J Pharm Res* **15**, 1197–1203.
11. Yasuda K, Roneker KR, Miller DD, *et al.* (2006) Supplemental dietary inulin affects the bioavailability of iron in corn and soybean meal to young pigs. *J Nutr* **136**, 3033–3038.
12. Watzl B, Girrbach S & Roller M (2005) Inulin, oligofructose and immunomodulation. *Br J Nutr* **93**, 49S–55S.
13. Roberfruid MB, Cumps J & Devogelaer JP (2002) Dietary chicory inulin increases whole-body bone mineral density in growing male rats. *J Nutr* **132**, 3599–3602.
14. Jolliff JS & Mahan DC (2012) Effect of dietary inulin and phytase on mineral digestibility and tissue retention in weanling and growing swine. *J Anim Sci* **90**, 3012–3022.
15. Demigné C, Jacobs H, Moundras C, *et al.* (2008) Comparison of native or reformulated chicory fructans, or non-purified chicory, on rat cecal fermentation and mineral metabolism. *Eur J Nutr* **47**, 366–374.
16. Aigner B, Renner S, Kessler B, *et al.* (2010) Transgenic pigs as models for translational biomedical research. *J Mol Med* **88**, 653–664.
17. Pearce AI, Richards RG, Milz S, *et al.* (2007) Animal models for implant biomaterial research in bone: a review. *Eur Cell Mater* **13**, 1–10.
18. Reinwald S & Burr D (2008) Review of nonprimate, large animal models for osteoporosis research. *J Bone Miner Res* **23**, 1353–1368.
19. Nutrient Requirements of Swine (2012) *Committee on Nutrient Requirements of Swine, Board on Agriculture and Natural Resources, Division on Earth and Life Studies*, 11th rev. ed. Washington, DC: National Academies Press.
20. Estrada A, Drew MD & Van Kessel A (2001) Effect of the dietary supplementation of fructooligosaccharides and *Bifidobacterium longum* to early-weaned pigs on performance and fecal bacterial populations. *Can J Anim Sci* **81**, 141–148.
21. Kjos NP, Øverland M, Fauske AK, *et al.* (2010) Feeding chicory inulin to entire male pigs during the last period before slaughter reduces skatole in digesta and backfat. *Livest Sci* **134**, 143–145.
22. Salinas MV, Hamet MF, Binaghi J, *et al.* (2017) Calcium–inulin wheat bread: prebiotic effect and bone mineralisation in growing rats. *Int J Food Sci Tech* **52**, 2463–2470.
23. Peressini D & Sensidoni A (2009) Effect of soluble dietary fibre addition on rheological and breadmaking properties of wheat doughs. *J Cereal Sci* **49**, 190–201.
24. Commission Regulation (EU) (2010) Regulation (EC) No 1924/2006 of the European Parliament and of the Council with regard to the list of nutrition claims. No. 116/2010, Commission Regulation (EU), Brussels, 9 February.
25. Skiba G, Weremko D, Sobol M, *et al.* (2015) Bone mineralisation of weaned piglets fed a diet free of inorganic phosphorus and supplemented with phytase, as assessed by dual-energy X-ray absorptiometry. *Arch Anim Nutr* **69**, 267–275.
26. Ferretti JL, Capozza RF, Mondelo N, *et al.* (1993) Interrelationships between densitometric, geometric and mechanical properties of rat femora: inferences concerning mechanical regulation of bone modelling. *J Bone Miner Res* **8**, 1389–1396.
27. Horwitz W & Latimer GW (editors) (2011) *Official Methods of Analysis of AOAC International, Current through Revision 4*, 18th ed. Gaithersburg, MD: AOAC International.
28. Folch J, Lees M & Sloane Stanley GH (1957) A simple method for the isolation and purification of total lipides from animal tissues. *J Biol Chem* **226**, 497–509.
29. Grela ER, Pietrzak K, Sobolewska S, *et al.* (2013) Effect of inulin and garlic supplementation in pig diets. *Ann Anim Sci* **13**, 63–71.
30. Bielohuby M, Matsuura M, Herbach N, *et al.* (2010) Short-term exposure to low-carbohydrate, high-fat diets induces low bone mineral density and reduces bone formation in rats. *J Bone Miner Res* **25**, 275–284.
31. Cao JJ, Sun L & Gao H (2010) Diet-induced obesity alters bone remodeling leading to decreased femoral trabecular bone mass in mice. *Ann NY Acad Sci* **1192**, 292–297.
32. Fehrendt H, Linn T, Hartmann S, *et al.* (2014) Negative influence of a long-term high-fat diet on murine bone architecture. *Int J Endocrinol* **2014**, 318924.
33. Yanagihara GR, Shimano RC, Tida JA, *et al.* (2017) Influence of high-fat diet on bone tissue: An experimental study in growing rats. *J Nutr Health Aging* **21**, 1337–1343.
34. Griel AE, Kris-Etherton PM, Hilpert KF, *et al.* (2007) An increase in dietary *n*-3 fatty acids decreases a marker of bone resorption in humans. *Nutr J* **6**, 2.
35. Duan Y, An Y, Li N, *et al.* (2013) Multiple univariate data analysis reveals the inulin effects on the high-fat-diet induced metabolic alterations in rat myocardium and testicles in the preobesity state. *J Proteome Res* **12**, 3480–3495.
36. Kumar SA, Ward LC & Brown L (2016) Inulin oligofructose attenuates metabolic syndrome in high-carbohydrate, high-fat diet-fed rats. *Br J Nutr* **116**, 1502–1511.
37. Devareddy L, Khalil DA, Korlagunta K, *et al.* (2006) The effects of fructo-oligosaccharides in combination with soy protein on bone in osteopenic ovariectomized rats. *Menopause* **13**, 692–699.
38. Jamieson JA, Ryz NR, Taylor CG, *et al.* (2008) Dietary long-chain inulin reduces abdominal fat but has no effect on bone density in growing female rats. *Br J Nutr* **100**, 451–459.
39. Abrams SA, Griffin IJ, Hawthorne KM, *et al.* (2005) A combination of prebiotic short- and long-chain inulin-type fructans enhances calcium absorption and bone mineralization in young adolescents. *Am J Clin Nutr* **82**, 471–476.

40. Abrams SA, Hawthorne KM, Aliu O, *et al.* (2007) An inulin-type fructan enhances calcium absorption primarily via an effect on colonic absorption in humans. *J Nutr* **137**, 2208–2212.
41. Wang Y, Dellatore P, Douard V, *et al.* (2016) High fat diet enriched with saturated, but not monounsaturated fatty acids adversely affects femur, and both diets increase calcium absorption in older female mice. *Nutr Res* **36**, 742–750.
42. Kruger MC, Brown KE, Collett G, *et al.* (2003) The effect of fructooligosaccharides with various degrees of polymerization on calcium bioavailability in the growing rat. *Exp Biol Med (Maywood)* **228**, 683–688.
43. Kaur N & Gupta K (2002) Applications of inulin and oligofructose in health and nutrition. *J Biosci* **27**, 703–714.
44. Roberfroid MB & Delzenne N (1998) Dietary fructans. *Annu Rev Nutr* **18**, 117–143.
45. Scharrer E & Lutz T (1991) Effects of short fatty acids on calcium absorption by rat colon. *Exp Physiol* **76**, 615–618.
46. Oh SR, Sul OJ, Kim YY, *et al.* (2010) Saturated fatty acids enhance osteoclast survival. *J Lipid Res* **51**, 892–899.
47. Estadella D, da Penha Oller do Nascimento CM, Oyama LM, *et al.* (2013) Lipotoxicity: effects of dietary saturated and transfatty acids. *Mediators Inflamm* **2013**, 137579.
48. Patsch JM, Kiefer FW, Varga P, *et al.* (2011) Increased bone resorption and impaired bone microarchitecture in short-term and extended high-fat diet-induced obesity. *Metabolism* **60**, 243–249.
49. Parhami F, Tintut Y, Beamer WG, *et al.* (2001) Atherogenic high-fat diet reduces bone mineralization in mice. *J Bone Miner Res* **16**, 182–188.
50. Zhong X, Xiu L, Wei G, *et al.* (2011) Bezafibrate prevents palmitate-induced apoptosis in osteoblastic MC3T3-E1 cells through the NF- κ B signaling pathway. *Int J Mol Med* **28**, 535–542.