

³ CHTMAD – E.P.E., Psychiatry Department, Vila Real, Portugal

* Corresponding author.

The association of antidepressant therapy with mania in people being treated for unipolar depression reinforces the importance of further clarification of this effect, since it has a huge implication in treatment approach and outcome. With this main objective, authors propose to make retrospective sociodemographic and clinical characterization of a group of patients with antidepressant induced mania in diagnosed unipolar depression, admitted in inpatient unit of Psychiatry and Mental Health Department of Centro Hospitalar de Trás-os-Montes e Alto Douro. Authors also aim to identify features that may increase the risk for mania or hypomania in people who present with an episode of depression, which can guide clinical orientation and improve outcome.

Disclosure of interest The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.01.402>

EV0074

Clinical differences between unipolar and bipolar depression

S. Smaoui*, N. Charfi, M. Maâlej Bouali, L. Zouari, N. Zouari, J. Ben Thabeb, M. Maâlej

Hedi Chaker University Hospital, Psychiatry, Sfax, Tunisia

* Corresponding author.

Introduction Epidemiological studies indicate that the majority of patients with bipolar disorder are diagnosed many years later. Unipolar depression represents the most frequent misdiagnosis.

Objectives This study aimed to examine the symptom profiles of depressed patients in order to identify clinical specificities of bipolar depression.

Methods A total of 31 depressed patients were recruited from psychiatry outpatient department of Hedi Chaker university hospital in Sfax (Tunisia), during October and November 2016. Unipolar and bipolar patients were compared on a broad range of parameters, including sociodemographic and clinical characteristics. Depressive symptoms were rated using the Montgomery Asberg Depression Rating Scale (MADRS) and Bipolar Depression Rating Scale (BDRS).

Results The total sample comprised 31 patients with 16 men and 15 women. It involved 20 with unipolar depression and 11 with bipolar depression. Patients with bipolar depression had more family history of bipolar disorder ($P=0.037$) and a triggering factor had been identified less often ($P=0.03$). MADRS scores were similar in bipolar and unipolar patient (median score 28.22 versus 28.36; $P=0.964$). BDRS scores were significantly higher in bipolar depressed patients (median score 33 versus 25; $P=0.01$). The mixed subscale (item 16 to 20) scores were particularly higher (median 6 vs. 1.2; $P\leq 0.01$) especially concerning irritability ($P=0.001$). Increased motor drive ($P=0.004$) and agitation ($P=0.008$).

Conclusion Our findings suggest that the presence of mixed symptoms is very important to recognize depressed patients as having a bipolar disorder. We also recommend routine use of the BDRS for patients presenting for treatment of depression.

Disclosure of interest The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.01.403>

EV0075

A case report of patient who had two manic episodes with psychotic features induced by nasal decongestant

A. Kandeger, R. Tekdemir*, B. Sen, Y. Selvi
Selcuk University, Department of Psychiatry, Konya, Turkey

* Corresponding author.

Objective Phenylephrine, pseudoephedrine and ephedrine are the sympathomimetic drugs that have been used most commonly in oral preparations for the relief of nasal congestion. These drugs stimulate the central nervous system that is affected by the alpha and beta adrenergic agonism. Sympathomimetic agents used in the treatment of flu and common cold with ephedrine and pseudoephedrine are case reports. That the manic and psychotic episodes are triggered. In this article, we would like to present a bipolar manic disorder with two manic episodes and both of them triggered by influenza drugs.

Case A 25-year-old man patient was admitted to psychiatric outpatient clinic with increasing complaints such as increasing energy, speaking much, decreasing sleep, increasing the libido after using the flu drug that prescribed to him containing phenylephrine. Also, 2 years ago, he has manic attack triggered after the flu drug.

Discussion In recent years the study of epilepsy and bipolar disorder in common suggests that bipolar disorder may affect the kindling phenomenon. In our case, two of reasons in the neurobiology of developing manic attacks the kindling phenomenon is likely to be effective. First, the possibility of using cold medicine containing ephedrine or pseudoephedrine in the first manic episode, in the second manic episode having spent the attack with FAQ stimulant effect of lower phenylephrine. Second, in the first episode after using the 5–6 tablets developing manic attacks. In the second attack to be triggered with just 2 doses may indicate the effect of kindling.

Disclosure of interest The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.01.404>

e-Poster viewing: child and adolescent psychiatry

EV0076

Posttraumatic stress disorder symptoms and related factors after circumcision operation performed with general or local anesthesia

B. Kozanhan¹, C. Kocaoğlu², M. Gunduz³, O.F. Akca^{4,*}

¹ Meram Training and Research Hospital, Anesthesia and Reanimation, Konya, Turkey

² Meram Training and Research Hospital, Pediatric Surgery, Konya, Turkey

³ Selcuk University, Pediatric Surgery, Konya, Turkey

⁴ Necmettin Erbakan University Meram School of Medicine, Child and Adolescent Psychiatry, Konya, Turkey

* Corresponding author.

Objective Elective circumcision operation for male children can affect their psychological status.

Aim We aimed to investigate the traumatic effect of the circumcision procedure and the predictive factors that could contribute to this effect.

Method One hundred and thirty-four children who admitted to urology and child surgery clinics to perform a circumcision procedure were included in the study. The whole group was divided into two groups including general anesthesia (GA) ($n=71$) and local anesthesia (LA) ($n=63$) groups. The procedure was performed under general and local anesthesia in the relevant groups. All of the participants filled the Child Depression Inventory (CDI), State (SA), and Trait (TA) Anxiety and Childhood Anxiety Sensitivity index (CASI) before the operation, and the Child Posttraumatic Stress Reaction index (CPSRI) 1 month after the operation.

Results CDI, CASI and TA scores of the children were similar in both groups, however, SA scores of the LA group were higher than GA group ($P<0.001$), and CPSRI scores of the GA group were