

Table Talk: development of an observational tool to assess verbal feeding communications in early care and education settings

Taren Swindle^{1,*}, Julie M Rutledge², Belynda Dix¹ and Leanne Whiteside-Mansell¹

¹Department of Family and Preventive Medicine, University of Arkansas for Medical Sciences, 4301 W. Markham Street, #530, Little Rock, AR 72205-7199, USA: ²School of Human Ecology, Louisiana Tech University, Ruston, LA, USA

Submitted 27 February 2017: Final revision received 19 June 2017: Accepted 21 June 2017: First published online 14 August 2017

Abstract

Objective: Children's dietary intake impacts weight status and a range of short- and long-term health outcomes. Accurate measurement of factors that influence children's diet is critical to the development and evaluation of interventions designed to improve children's diets. The purpose of the current paper is to present the development of the Table Talk observational tool to measure early care and education teachers' (ECET) verbal feeding communications.

Design: An observational tool to assess ECET verbal communication at mealtimes was designed based on the extant literature. Trained observers conducted observations using the tool during lunch for both lead and assistant ECET. Descriptive statistics, test–retest for a subgroup, interclass correlations for each item, and comparisons between leads and assistants were conducted.

Setting: Head Start centres, Southern USA.

Subjects: Seventy-five Head Start educators.

Results: On average, 17.2 total verbal feeding communications (SD 8.9) were observed per ECET. For lead ECET, the most prevalent Supportive Comment was Exploring Foods whereas for assistants Making Positive Comments was the most prevalent. Overall, lead ECET enacted more Supportive Comments than assistant ECET ($F(2,72) = 4.8, P = 0.03$). The most common Unsupportive Comment was Pressuring to Eat, with a mean of 3.8 (SD 4.3) and a maximum of 25. There was no difference in Unsupportive Comments between lead and assistant ECET.

Conclusions: Table Talk may be a useful tool to assess verbal feeding communications of ECET, with potential applications such as informing ECET training and assessing intervention efforts.

Keywords
Early care and education
Childcare
Pre-school
Feeding
Mealtime
Nutrition

The impact of parent feeding practices on children has been studied widely. Restrictive practices are related to increased eating and weight among children⁽¹⁾, while pressure to eat and emotional feeding are related to overeating and emotional eating in children⁽²⁾. Positive effects also are observed, including increased fruit and vegetable consumption within homes where authoritative, supportive feeding practices are used (e.g. modelling intake, making fruits and vegetables available)⁽³⁾. Parents, however, are not the only influence on children's development of food-related behaviours and preferences. Outside the home environment, childcare is often the first influence on children's eating.

The impact of early care and education teachers' (ECET) feeding practices and feeding communications on child outcomes has been researched to a lesser extent than that of parents. While descriptive studies have begun to document ECET feeding interactions^(4–6), the direct link

between these interactions and child outcomes is emerging. A review by Wolfenden *et al.*⁽⁷⁾ of available trials designed to increase fruit and vegetable intake in early childhood found that a pre-school-based intervention resulted in significant increases in fruit consumption for children, whereas home visiting interventions did not have observable impacts. This demonstrates the potential impact of the ECET context and suggests a need to understand the mechanisms that contribute to change in child nutritional outcomes (e.g. dietary variety, neophobia, willingness to try) through early intervention and the interactions with the adults in early care and education settings.

Considering the gap between the potential for ECET to impact children and what is known about the influence of ECET feeding practices on children, additional information on ECET feeding interactions is needed. ECET may eat up to three meals or snacks with children per day, which is

*Corresponding author: Email tswindle@uams.edu

more than 500 meals in a school year. For the most at-risk children, meals in the early education setting may represent most, if not all, of their daily dietary intake⁽⁸⁾. As early childhood is a sensitive time for establishing self-regulation and dietary preferences⁽⁹⁾, ECET are in an important position to create a responsive feeding environment that will support healthy habits for children.

Recognizing the importance of ECET in child feeding, Mita *et al.*⁽¹⁰⁾ presented a conceptual framework of characteristics of a positive meal environment (PME). A PME is conceptualized to consist of opportunities for learning, socializing and eating. The focus in a PME is on the people, emotional tone and rules/routines involved in meals. Mita's model, consistent with the existing literature and recommendations from the American Academy of Nutrition and Dietetics⁽¹¹⁾, provides a strong basis for observational study to quantify PME characteristics such as verbal feeding communications. Extending upon Mita's model, the absence of PME characteristics and the addition of undesirable interactions would relate to a negative mealtime experience (NME), a feeding environment that is not nurturing for healthy habits. Measuring PME and NME characteristics would provide important information on the childcare feeding environment.

Currently, self-report measures are frequently used to assess ECET knowledge, attitudes and self-efficacy about child nutrition and feeding^(12–15). Validated feeding assessment measures developed for use with parents (e.g. Child Feeding Questionnaire (CFQ); Caregiver Feeding Style Questionnaire (CFSQ))^(16,17) have been used in some educational settings, but efforts to understand the psychometric properties in this group have been limited⁽¹⁸⁾. One recently developed tool, the sixty-three-item Child-care Food and Activity Practices Questionnaire (CFAPQ)⁽¹⁹⁾, was developed specifically for use with childcare staff. However, validation efforts of the CFAPQ to date have been limited to internal consistency and item correlation estimates. Self-assessment tools of nutrition policy and procedure are available to support programmes to evaluate and improve the mealtime environment but do not provide tools to quantify distinct verbal feeding communications initiated by ECET (e.g. Nutrition and Physical activity Self-Assessment for Child Care (NAPSACC); Building Mealtime Environments and Relationships (BMER); Environment and Policy Assessment and Observation (EPAO))^(20–22).

Despite the progress of observational studies^(23–25) to describe the pre-school mealtime environment, studies have not used standard instruments to capture the quantity of specific verbal ECET feeding communications in real time. This is a critical gap given that emerging research suggests a lack of correspondence between educator-reported and observed feeding behaviours⁽²⁶⁾. To address this gap, the current study presents the development of a new observational tool to quantify verbal ECET feeding communications (i.e. supportive and unsupportive

comments) consistent with the PME framework, as well as extant literature and recommendations.

Methods

Research design

Head Start programmes serve low-income families (100% of the federal poverty level or below) with children from birth to age 5 years through federal support from the US Administration for Children and Families Early Childhood Learning & Knowledge Center. To represent the childcare setting, Head Start agencies in two Southern states with a high prevalence of overweight and obesity were enrolled to participate in the implementation and evaluation of a nutrition education and food experience curriculum. All Head Start programmes in this sample were full-day programmes. The observations detailed in the present study represent baseline observations of the classrooms before ECET training and implementation of a nutrition education curriculum. Assessment of ECET feeding communications at mealtime was collected as a potential moderating variable of the impact of the curriculum; however, the Table Talk tool was not designed to be specific to the curriculum or training the ECET received and has wider applicability.

Observations were completed in three cohorts across a total of thirty-seven classrooms: (i) ten classrooms in autumn 2013; (ii) twelve classrooms in spring 2014; and (iii) fifteen classrooms in autumn 2015. All classrooms in the study served children between the ages of 3 and 5 years. Of the thirty-seven classrooms, six classrooms in Cohort 1 were in an urban area serving families who were a majority African American (72.4%); sixteen classrooms in Cohorts 1 and 2 were in rural areas serving families who were a majority White (68.1%); and fifteen classrooms (Cohort 3) were in a suburban area serving families who were a majority African American (83.6%). A sub-sample of ECET (lead ECET, *n* 7) was observed twice in spring 2014 within 3 weeks (mean 1.26 weeks) to assess test-retest reliability. The study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving human subjects were approved by the University of Arkansas for Medical Sciences Institutional Review Board. Written informed consent was obtained from all ECET.

Table Talk development

The Table Talk tool was developed based on the empirical literature and preliminary mealtime observations. A review of the extant literature was conducted to compile evidence-based practices and recommendations in child feeding and mealtime interactions. Selected targets for the Table Talk observation tool were based on: (i) the benefits of vocal, positive adult role^(6,27,28); (ii) the counterproductive impact of pressuring children to eat^(29–32); (iii) the value of guiding children to attune to cues of hunger and satiety and valuing

children's choice to eat^(11,25,30,32,33); (iv) the benefits of appropriate food exploration and encouraging children's interest in foods^(11,30,34-36); (v) the undesirable impacts (i.e. overjustification effect) of coercive feeding practices such as promise of a food to increase intake of another food⁽³⁷⁻⁴⁰⁾; and (vi) the advantage of a positive social context at mealtimes^(10,11,41,42). After the first cohort of observations, comments that focused on behavioural control of children (e.g. 'turn around', 'sit up straight') were added to Table Talk to provide additional measurement of the mealtime climate. Consistent with a PME framework⁽¹⁰⁾ and previous research on punitive and authoritarian interactions in childcare settings⁽⁴³⁾, firm, harsh directives and inappropriate transitions or redirections were coded in this category and included in the count of unsupportive behaviours.

Two authors (T.S. and B.D.) led the development of the tool. T.S. is trained as a mixed-methods researcher with graduate education in child development, educational psychology and nutrition. B.D. has graduate training in Applied Communication with a focus on adult learning. Our literature review provided sensitizing concepts (i.e. a start list)⁽⁴⁴⁾, which we observed during our preliminary observations prior to the study. After each preliminary observation, we would meet immediately to discuss application of the list during the observations. That is, we made notes of how we coded communications we heard and compared counts for each category. During this process, we engaged in consensus building to define examples of each category, revise our categories and outline criteria for differentiation between categories. For example, we documented examples of the difference between pressuring children to eat (e.g. 'I need to see you take a bite') and encouraging trying in a positive way (e.g. 'What did you notice about the carrots?'). We developed overarching rules to guide distinctions (e.g. 'Comments that do not respect a child's choice about how much to eat are pressure'). We continued this process until we demonstrated reliability (>85% agreement) on three sequential occasions. We used our notes and examples from this process to develop the training materials.

Table Talk was designed to facilitate quick identification of supportive or unsupportive verbal ECET comments (see first column of Table 1 for content of the observational tool). Observations recorded on the top portion of the tool are positive (*n* 4); observations recorded on the bottom portion of the tool are negative (*n* 8). This promotes efficient categorization of verbal ECET comments. Tally marks are used during the observation and total scores are created for each item by totalling comments after the observation for each ECET.

Observation training

Observers completed a standardized training consisting of an in-person session with instruction on: (i) the intent of each item with provision of examples; (ii) distinguishing between categorizations; and (iii) discreet integration into

Table 1 Mean observed number of communications of early care and education teachers (ECET), by teacher type, in thirty-seven classrooms in eight Head Start centres of two Southern US states, 2013–2015

	Lead ECET			Assistant ECET			Total ECET		
	Mean	sd	Range	Mean	sd	Range	Mean	sd	Range
Supportive Communications									
Positive comments about food served ^(6,26,27) ('I am really enjoying the peas today. Yummy.')	2.8	2.4	0-10	2.3	1.8	0-7	2.5	2.1	0-10
Hunger cues ^(11,25,30,32,33) ('Are you full? How does your belly feel?')	0.4	0.7	0-3	0.1	0.3	0-1	0.3	0.6	0-3
Encourage trying in a positive way ^(6,11,24,26,27,29,31,32) ('Would you like to try the peas?')	3.4	2.8	0-11	2.0	2.5	0-11	2.7	2.7	0-11
Exploring foods ^(11,29,33-35) ('What does it smell like? How does it feel in your mouth?')	4.1	3.7	0-13	2.3	2.5	0-9	3.2	3.3	0-13
Total Supportive Communications	10.7	6.7	0-24	6.7	4.4	0-16	8.7	6.0	0-24
Unsupportive Communications									
Negative comments about the food served ⁽³⁶⁻³⁹⁾ ('I can't believe we're having this again. I don't like peas.')	0.1	0.4	0-2	0.1	0.25	0-1	0.1	0.3	0-2
Pressure to eat ⁽²⁸⁻³¹⁾ ('Eat your food. Take a bite. Clean your plate. Finish.')	3.8	3.5	0-14	3.8	5.0	0-25	3.8	4.3	0-25
Threats (to encourage eating) ^(10,11,28-31,36-39,40,41) ('If you don't eat, you'll be over here by yourself.')	0.0	-	-	0.1	0.2	0-1	0.03	0.1	0-1
Discourage manipulating food ^(11,29,33-35) ('Eat; don't play. That's sticky and nasty.')	0.8	1.3	0-7	0.8	1.2	0-5	0.8	1.2	0-7
Indicate preference for unhealthy food ^(6,26,27) ('I wish we were having French fries today.')	0.3	0.6	0-2	0.1	0.3	0-2	0.2	0.5	0-2
Food as a reward ⁽³⁶⁻³⁹⁾ ('If you eat your vegetable, you can have dessert.')	0.1	0.2	0-1	0.01	0.1	0-1	0.03	0.2	0-1
Hurries to finish eating ^(10,11,28-31,40,41) ('We're waiting on you. Let's hurry so we can go to recess.')	3.7	2.6	0-12	4.6	3.5	0-15	4.2	3.1	0-15
Firm behavioural control ^(10,11,40,41) ('Turn around. Sit up straight. Hands in your lap.')	8.2	5.9	1-31	8.7	8.6	0-47	8.5	7.3	1-47
Total Unsupportive Communications	18.9	8.0	4-42	15.4	9.6	1-49	17.2	8.9	1-49
Total Communications									

the classroom setting. This training included coding a video example with the guidance of a gold-standard observer and then independently. Thereafter, observers completed pilot field observations with the lead investigators to establish inter-rater reliability. Inter-rater reliability was calculated by determining the percentage of items on which observers rated within a narrow margin of error (± 1 for counts ≤ 4 , ± 2 for counts > 4) relative to the gold-standard observer. Before observing classrooms independently, each observer was required to exhibit inter-rater reliability of 85% with one of three gold-standard observers on two occasions. Gold-standard observers exhibited greater than 90% agreement with one another. This level of reliability was obtained within three live observations for all observers after video training. Observers ($n = 10$) included undergraduate students of sociology and child development, graduate-level students in nutrition and psychology, and professionals from education and public health.

Data collection

In total, thirty-seven classrooms across eight centres from three Head Start agencies were observed at lunch. These classrooms included a total of seventy-five ECET – thirty-seven lead ECET, thirty-seven assistant ECET and one family-service coordinator – who regularly ate meals with children. Table 2 presents the demographics of the ECET observed in the study. Observers arrived 10 min prior to the mealtime to select an unobtrusive observation position and to allow children and staff time to adjust to their presence.

Table 2 Demographic characteristics of early care and education teachers (ECET), by teacher type, in thirty-seven classrooms in eight Head Start centres of two Southern US states, 2013–2015

Characteristic	Total ECET (<i>n</i> 75)	Assistant ECET (<i>n</i> 38)	Lead ECET (<i>n</i> 37)
	%	%	%
Age (years)			
19–24	2.0	4.2	0.0
25–34	19.6	33.3	5.4
35–40	21.6	16.7	16.2
≥ 41	56.9	45.8	48.6
Race			
White	28.0	26.3	29.7
African American	64.0	60.5	64.9
American Indian	1.3	2.6	0.0
Asian/Pacific Islander	4.0	2.6	5.3
Other	2.7	7.8	0.0
Hispanic			
No	94.7	94.7	94.3
Yes	5.3	5.3	5.4
Education level			
High school/GED	7.2	14.7	0.0
Some college	15.9	32.4	0.0
Associates degree	34.8	32.4	38.2
Bachelor's degree	34.9	17.6	50.0
Master's degree or above	7.2	2.9	11.8

GED, General Educational Development.

Observations lasted from the time the food was served to the first child to the time the food was removed from the last child. On average, observations lasted 27.8 (SD 6.9) min (range 13–45 min). Observers focused on coding the unique verbal expressions with children as they related to the categories presented in Table 1. That is, if an ECET repeated the same phrase to the same child twice in a row (e.g. 'Eat your green beans', 'Eat your green beans'), a single comment was recorded. If the ECET interacted with another child (i.e. initiated a different interaction) and returned to the previous child to provide the directive again, another comment was recorded. Directing the same phrase to a different child was considered a unique expression. Expressions were considered pressuring rather than encouraging when the child's choice of intake was not recognized (e.g. 'Try the carrots' *v.* 'Would you like to try the carrots with me? I think they're yummy'). When meals were served in the classroom ($n = 32$), one observer recorded communications of both lead and assistant ECET. When meals were served in a cafeteria setting ($n = 5$), two observers were assigned to a class; one observer recorded communications of the lead ECET, and the other observer recorded the communications of the assistant ECET. This allowed for more accurate capturing of verbal ECET communications in the noisier cafeteria environment.

Analyses

Analyses were conducted with using the statistical software package IBM SPSS Statistics version 22.0. Test–retest reliability was determined by correlating items of the first and second observations. Summary scores were created for Supportive and Unsupportive Comments by totalling the number of observed verbal ECET communications within these areas. Means were compared on Table Talk summary scores between lead and assistant ECET using one-way ANOVA, controlling for demographic information. Intraclass correlation coefficients (ICC) were calculated to indicate the variance in Table Talk items and summary scores shared between lead and assistant ECET in the same classroom⁽⁴⁵⁾. Thus, the ICC reflect how ECET communications in the same classroom co-vary. ICC in the present study do not indicate reliability.

Results

Table Talk observations

Table 1 presents a summary of the mean number of recorded verbal feeding communications by lead and assistant ECET. For lead ECET, the most prevalent Supportive Comment was Exploring Foods, whereas for assistants Making Positive Comments was the most prevalent. These were observed 4.1 and 2.3 times per meal, respectively. Lead ECET consistently had higher Supportive Comment scores than assistant ECET, which was reflected by the overall supportive summary scores (leads,

Table 3 One-way ANOVA comparing supportive and unsupportive communications of early care and education teachers (ECET), by teacher type, in thirty-seven classrooms in eight Head Start centres of two Southern US states, 2013–2015

	Lead ECET		Assistant ECET		<i>F</i>	<i>P</i> value
	Mean	SD	Mean	SD		
Supportive Comments	10.7	6.7	6.7	4.4	4.83	0.03
Unsupportive Comments	8.2	5.9	8.7	8.6	0.30	0.59

mean = 10.7; assistants, mean = 6.7). The least used Supportive Comment by both lead and assistant ECET was Hunger Cues, which was used less than 1 time per meal on average. A significant difference was found between lead and assistant ECET on Supportive Comments ($F(2,72) = 4.8$, $P = 0.03$), with lead ECET providing more supportive comments than assistants (see Table 3).

The most common Unsupportive Comment was Pressure to Eat. This was the case for both lead and assistant ECET, with observed means of 3.8 times per meal. The maximum number of Pressure to Eat Comments was 25 times in a meal. For a given classroom, children could hear up to 39 directives to eat in one meal from their lead and assistant ECET combined. This averages to hearing Pressure to Eat comments more frequently than once per minute. The most uncommon Unsupportive Comments were Negative Comments, Threats and Food as a Reward. Notably, up to 47 Unsupportive Comments were observed per meal in a given classroom. Combined, ECET were similar in their use of Unsupportive Comments (mean = 8.5) and Supportive Comments (mean = 8.7). No differences on Unsupportive Comments were found between lead and assistant ECET (Table 3).

Interclass correlation coefficients indicating shared classroom variance

ICC for Supportive Comments examining the shared variance between ECET in the same classroom ranged from a minimum of 0.1 (Hunger Cues) to a maximum of 0.3 (Positive Comments). ICC indicating shared variance between lead and assistant ECET for Unsupportive Comments were a minimum of 0.01 and a maximum of 0.6 (Discourage Manipulation and Hurries to Finish, respectively). ICC could not be estimated for two items (Hunger Cues and Threats) due to restricted variability in their occurrence. The ICC indicating shared variance within classrooms for overall Supportive Comments was 0.1; the ICC for overall Unsupportive Comments was 0.5.

Test–retest reliability

Item-level correlations indicating test–retest reliability ranged from 0.4 (Behaviour Control) to 0.9 (Pressure to Eat). The next lowest observed correlation was 0.5 (Positive Comments). All other correlations were greater

than 0.6; several were greater than 0.8 (Hurries, Discourage Manipulation and Exploring Foods). Correlations could not be estimated for five items (Hunger Cues, Negative Comments, Threats, Preference for Unhealthy Foods, Food as a Reward) that were infrequent and constant across time in this sample.

Discussion

The purpose of the present study was to develop and assess a new tool for quantifying verbal ECET feeding communications. To that end, we proposed a systematic way to measure distinct verbal ECET communications consistent with a PME (i.e. supportive comments) and NME (i.e. unsupportive comments)⁽¹⁰⁾. Results suggest that Table Talk has the potential to capture variability in verbal ECET communications. The observed numbers of Supportive and Unsupportive Comments at a mealtime captured by Table Talk were comparable on average.

Previous reviews on mealtime interactions among ECET have suggested room for improvement^(4,46). To move towards greater adoption of evidence-based practices, the field may benefit from pragmatic tools to measure aspects of the meal environment such as verbal feeding communications in real time. The results of the current study are consistent with previous studies which highlighted areas for improvement in mealtimes in early care and education settings. The study also offers a simple measure that may be the type of pragmatic measure needed to document and address these deficits. In the current study, Table Talk documented that ECET pressured children to eat approximately every 7 min, with a maximum rate of once every 1.2 min. This is greater than the rate reported by Gable and Lutz⁽²³⁾, which was approximately every 15 min, and is consistent with reports from Ramsay *et al.*⁽²⁵⁾ who found that ECET were ten times more likely to direct children to eat than to provide cues to hunger or satiety. Using Table Talk, the current study also documents the frequency with which ECET engage in other unsupportive verbal communications that have not been included in previous studies (e.g. Discourage Manipulation).

Use of the Table Talk tool documented differences between ECET of different roles, with lead ECET providing more supportive comments than assistant ECET. The ability of Table Talk to capture communications of both ECET roles is important as both are typically present and interacting with children during mealtimes. Further, analysis of data collected using the Table Talk tool illustrated that a large amount of variance in communications was shared between ECET in the same classroom (i.e. ICC), especially for Unsupportive Comments. Potential reasons for this could be shared organizational culture or similar training opportunities. Further, previous studies have documented that physical activity and eating behaviours are predicted by those in individuals' immediate

surroundings^(47–49). A similar peer influence could be at work between ECET in the early childhood classroom, and the Table Talk tool may be a useful way to capture this phenomenon. The cause of the shared variance between lead and assistant ECET in the present study is unknown and deserves further exploration.

Some verbal ECET feeding communications included in the Table Talk tool occurred infrequently in this sample. In fact, seven communications occurred less than once per meal on average. The infrequency of these verbal ECET feeding communications in this sample does not necessarily mean that these communications will be infrequent in other settings, such as state-funded childcare, private childcare or family childcare homes. Other studies have shown differences in self-reported feeding practices by programme type, with Head Start teachers faring better than those in other programmes^(50,51). Thus, it is likely that higher rates of these verbal feeding communication behaviours would be seen elsewhere, and these communications should be retained as part of the tool until further work in additional settings is complete.

Table Talk may be a useful for several reasons. First, interrater reliability levels are consistent with NAPSACC⁽²²⁾, a standard measure in the field, and were obtained with a diverse group of observers. High rates were achieved due to the variety of training methods that each observer completed before reliability was measured, including video and field training. Our training methods, topics and standards were comparable to those used to train NAPSACC observers (e.g. observation techniques, mock observations, 85% reliability with gold standard)⁽²²⁾. This provides an indication of the potential feasibility of the instrument for wider use. Second, Table Talk captures observed rather than reported verbal feeding communication behaviour, which eliminates self-report bias that may be an issue with measures such as the CFAPQ⁽¹⁹⁾, BMER⁽²¹⁾ or NAPSACC⁽²⁰⁾ when a self-assessment is completed. Although observation may impact verbal ECET feeding communications, it is reasonable to expect that it would increase the 'correct' communications because of social desirability. Given that unsupportive verbal ECET feeding communications are still observed, there may be other barriers (e.g. lack of knowledge⁽¹⁵⁾, contextual factors such as training and regulations^(50,51), personal dietary factors^(15,50,52)) to ECET being able to consistently use evidence-based feeding communications. Finally, Table Talk may contribute to understanding the predictive power of PME and NME for children. By using a tool such as this in connection with collection of health outcomes data for children, researchers may be able to determine how the mealtime environment in early childhood settings can impact health outcomes in children.

There are also limitations to the Table Talk tool. First, Table Talk was not designed to take the place of other assessment tools that capture environmental characteristics (e.g. NAPSACC, BMER, EPAO^(20–22)) or broader aspects of ECET practice related to nutrition (e.g. CFAPQ⁽¹⁹⁾).

It should be used and further developed (e.g. convergent validity) in combination with other nutrition assessment tools. Further, Table Talk includes more unsupportive communication than supportive communication categories. This reflected the available literature and our pilot observations. We added a category to capture general statements that would capture a harsh mealtime climate (i.e. Behavioural Control) but not a similar category to reflect general statements unrelated to food that would reflect a positive mealtime climate (e.g. 'What did you do last night at home?'). This should be considered for further development of the tool. However, Table Talk was designed as a live observation tool that has the potential for use as an immediate intervention and that would require fewer resource demands than coding recorded mealtimes (e.g. staff time, cost of coding software/equipment). This limits the amount of information that can be collected in the real-time observation, and it is expected the tool will have to remain focused to have validity and reliability. As such, the Table Talk tool will never capture the full complexity of teacher–child interactions at mealtime but is proposed to capture an important aspect.

Implications for research and practice

The current study was restricted to Head Start classrooms in Southern US states. Head Start has specific mealtime regulations which encourage family-style dining, emphasize supportive teacher–child interactions, and require compliance with US Department of Agriculture meal patterns⁽⁵³⁾. Head Start classrooms may be more accustomed to observation as well. Thus, further efforts are needed to assess generalizability to other settings such as state-funded programmes, private childcare and family childcare homes. Pilot observations by our team suggest utility of the Table Talk tool to capture communications at mealtime in these additional settings but suggest different patterns of ECET communication than in Head Start. Comparisons of ECET verbal communications between centre types is a promising area for research which could inform intervention. Additionally, future efforts need to explore the predictive validity of this tool for predicting child outcomes (e.g. intake of healthy foods, willingness to try new foods). A tool shown to predict these outcomes concurrently and/or across time could be valuable to the field.

There are several additional opportunities for future research. First, studies should explore differences in Table Talk scores on ECET characteristics (e.g. level of nutrition training, food security status). Further analyses (e.g. item-level demographic comparisons, correlational analyses) were not conducted in the present study due to the restricted range and limited number of observations of some verbal ECET feeding communications. Collection of Table Talk observations across a wider range of settings and demographic groups could allow for these types of analyses. Cultural differences may be reflected in the communications of ECET, as suggested by recent work

documenting the influence of family history on ECET mealtime practices with children in their classrooms⁽⁵⁴⁾. As no standard, self-report measures of feeding communications or behaviours have been validated in ECET, validation against the CFQ and CFSQ may be useful as well.

Although not documented in the present study, Table Talk is designed to be sensitive to change and to capture wide variability in verbal ECET feeding communications. This is an important feature of the measure because it has the potential to eliminate ceiling effects, which may impact other measures in the field^(19–21), particularly those designed to evaluate the impact of interventions. Future studies should assess if the tool is in fact sensitive to change across time. Table Talk can be used to identify individual verbal ECET feeding communications that naturally occur, offering observable, concrete information that can help shape effective interventions. This measure also has potential for local use at early childhood education agencies or on a broader scale to inform intervention. A trained observer could provide a baseline assessment of how ECET in a given setting relate to supportive and unsupportive verbal feeding communications. Information gathered may inform tailored training at the agency or individual level. Additional observational assessments throughout the school year may be useful to provide ECET with feedback on how their interactions are improving relative to evidence-based practice.

At a broader level and in line with recommendations from Story *et al.*⁽⁵⁵⁾, the current study provides a descriptive environmental exploration of verbal feeding communication as part of the classroom food environment. Apart from the measure development aspect of the study, the observed verbal ECET feeding communications highlight areas for improvement in the use of recommended mealtime interactions in early care and education settings. Due to the shared variance found between lead and assistant ECET in the same classroom, future work can explore the potential mechanisms that lead to these similarities, and verbal feeding communication trainings can be conducted accordingly. Further, policy makers should consider how well current training requirements address supporting ECET to enact supportive verbal feeding communications. A tool such as Table Talk can play a role in identifying parts of the feeding environment that can be improved. Additionally, to individualize feeding communication training, future work in larger samples can explore differences based on ECET characteristics such as education level and ethnicity, as well as diversity among eating settings (e.g. classroom, cafeteria).

Acknowledgements

Acknowledgements: The authors would like to acknowledge and thank the participating Head Start agencies and educators involved in this work. They also thank the research

assistants who aided in data collection and processing. *Financial support:* This project was supported by an Agriculture and Food Research Initiative Competitive Grant (T.S. and L.W.-M., grant number 2011-68001-30014) from the US Department of Agriculture, National Institute of Food and Agriculture. The project was also supported, in part, by the Translational Research Institute (TRI) (T.S., grant numbers UL1TR000039 and KL2TR000063) through the National Institutes of Health, National Center for Research Resources and the National Center for Advancing Translational Sciences; as well as the Lincoln Health Foundation of Lincoln Parish, Louisiana (J.M.R. and T.S.), the Arkansas Biosciences Institute (the major research component of the Arkansas Tobacco Settlement Proceeds Act of 2000), and the National Institutes of Health (T.S., grant number K01 DK110141-01). The content is solely the responsibility of the authors and does not necessarily represent the official views of the funding agencies. *Conflict of interest:* The authors have no financial or other interests in this content to disclose. *Authorship:* T.S. was responsible for the conception of the instrument as well as analysis and interpretation of data and leading the writing of this article; J.M.R. contributed to the acquisition of data, interpretation of the data and drafting the article; B.D. contributed to the conception of the content and the protocol for data acquisition and drafting of the article; and L.W.-M. contributed to the conception of the content, data acquisition and drafting of the article. All authors have provided approval of the final version and agree to be accountable for all aspects of the work. *Ethics of human subject participation:* This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving human subjects were approved by the University of Arkansas for Medical Sciences Institutional Review Board. Written informed consent was obtained from all ECET.

References

1. Faith M, Scanlon K & Birch L (2004) Parent-child feeding strategies and their relationships to child eating and weight status. *Obes Res* **12**, 1711–1722.
2. Rodgers RF, Paxton SJ, Massey R *et al.* (2013) Maternal feeding practices predict weight gain and obesogenic eating behaviors in young children: a prospective study. *Int J Behav Nutr Phys Act* **10**, 24.
3. Blissett J (2011) Relationships between parenting style, feeding style and feeding practices and fruit and vegetable consumption in early childhood. *Appetite* **57**, 826–831.
4. Ward S, Bélanger M, Donovan D *et al.* (2015) Systematic review of the relationship between childcare educators' practices and preschoolers' physical activity and eating behaviours. *Obes Rev* **16**, 1055–1070.
5. Larson N, Ward D, Neelon S *et al.* (2011) What role can child-care settings play in obesity prevention? A review of the evidence and call for research efforts. *J Am Diet Assoc* **111**, 1343–1362.
6. Gibson EL, Kreichauf S, Wildgruber A *et al.* (2012) A narrative review of psychological and educational strategies

- applied to young children's eating behaviours aimed at reducing obesity risk. *Obes Rev* **13**, Suppl. 1, 85–95.
7. Wolfenden L, Wyse RJ, Britton BI *et al.* (2012) Interventions for increasing fruit and vegetable consumption in children aged 5 years and under. *Cochrane Database Syst Rev* **11**, CD008552.
 8. Frisvold DE & Lumeng JC (2011) Expanding exposure. *J Hum Resour* **46**, 373–402.
 9. Fox MK, Devaney B, Reidy K *et al.* (2006) Relationship between portion size and energy intake among infants and toddlers: evidence of self-regulation. *J Am Diet Assoc* **106**, 1 Suppl. 1, S77–S83.
 10. Mita SC, Gray SA & Goodell LS (2015) An explanatory framework of teachers' perceptions of a positive mealtime environment in a preschool setting. *Appetite* **90**, 37–44.
 11. Benjamin Neelon SE & Briley ME (2011) Position of the American Dietetic Association: benchmarks for nutrition in child care. *J Am Diet Assoc* **111**, 607–615.
 12. Derscheid LE, Umoren J, Kim S-Y *et al.* (2010) Early childhood teachers' and staff members' perceptions of nutrition and physical activity practices for preschoolers. *J Res Child Educ* **24**, 248–265.
 13. Freedman MR & Alvarez KP (2010) Early childhood feeding: assessing knowledge, attitude, and practices of multi-ethnic child-care providers. *J Am Diet Assoc* **110**, 447–451.
 14. Nahikian-Nelms M (1997) Influential factors of caregiver behavior at mealtime: a study of 24 child-care programs. *J Am Diet Assoc* **97**, 505–509.
 15. Sharma S, Dortch KS, Byrd-Williams C *et al.* (2013) Nutrition-related knowledge, attitudes, and dietary behaviors among head start teachers in Texas: a cross-sectional study. *J Acad Nutr Diet* **113**, 558–562.
 16. Birch LL, Fisher JO, Grimm-Thomas K *et al.* (2001) Confirmatory factor analysis of the Child Feeding Questionnaire: a measure of parental attitudes, beliefs and practices about child feeding and obesity proneness. *Appetite* **36**, 201–210.
 17. Hughes SO, Cross MB, Hennessy E *et al.* (2012) Caregiver's Feeding Styles Questionnaire. Establishing cutoff points. *Appetite* **58**, 393–395.
 18. Hughes SO, Patrick H, Power TG *et al.* (2007) The impact of child care providers' feeding on children's food consumption. *J Dev Behav Pediatr* **28**, 100–107.
 19. Gubbels JS, Sleddens EF, Raaijmakers LC, Gies JM *et al.* (2016) The Child-care Food and Activity Practices Questionnaire (CFAPQ): development and first validation steps. *Public Health Nutr* **19**, 1964–1975.
 20. Benjamin SE, Ammerman A, Sommers J *et al.* (2007) Nutrition and physical activity self-assessment for child care (NAP SACC): results from a pilot intervention. *J Nutr Educ Behav* **39**, 142–149.
 21. Fletcher J, Branan L, Price B *et al.* (2005) *Building Mealtime Environments and Relationships. An Inventory of Mealtime Practices for Feeding Young Children in Group Settings*. Moscow, ID: University of Idaho.
 22. Ward D, Hales D, Haverly K *et al.* (2008) An instrument to assess the obesogenic environment of child care centers. *Am J Health Behav* **32**, 380–386.
 23. Gable S & Lutz S (2001) Nutrition socialization experiences of children in the Head Start program. *J Am Diet Assoc* **101**, 572–577.
 24. Lanigan JD (2012) The relationship between practices and child care providers' beliefs related to child feeding and obesity prevention. *J Nutr Educ Behav* **44**, 521–529.
 25. Ramsay SA, Branan LJ, Fletcher J *et al.* (2010) 'Are you done?' Child care providers' verbal communication at mealtimes that reinforce or hinder children's internal cues of hunger and satiation. *J Nutr Educ Behav* **42**, 265–270.
 26. Fallon M (2016) Exploring self-reported and observed feeding practices of Rhode Island Head Start teachers. Master's Thesis, University of Rhode Island; available at <http://digitalcommons.uri.edu/theses/853>
 27. Brown KA, Ogden J, Vögele C *et al.* (2008) The role of parental control practices in explaining children's diet and BMI. *Appetite* **50**, 252–259.
 28. Hendy HM & Raudenbush B (2000) Effectiveness of teacher modeling to encourage food acceptance in preschool children. *Appetite* **34**, 61–76.
 29. Batsell WR, Brown AS, Ansfield ME *et al.* (2002) 'You will eat all of that!': a retrospective analysis of forced consumption episodes. *Appetite* **38**, 211–219.
 30. Birch LL, McPhee L, Shoba B *et al.* (1987) 'Clean up your plate': effects of child feeding practices on the conditioning of meal size. *Learn Motiv* **18**, 301–317.
 31. Galloway AT, Fiorito LM, Francis LA *et al.* (2006) 'Finish your soup': counterproductive effects of pressuring children to eat on intake and affect. *Appetite* **46**, 318–323.
 32. Hendy HM (1999) Comparison of five teacher actions to encourage children's new food acceptance. *Ann Behav Med* **21**, 20–26.
 33. Orrell-Valente JK, Hill LG, Brechwald WA *et al.* (2007) 'Just three more bites': an observational analysis of parents' socialization of children's eating at mealtime. *Appetite* **48**, 37–45.
 34. Dazeley P & Houston-Price C (2015) Exposure to foods' non-taste sensory properties. A nursery intervention to increase children's willingness to try fruit and vegetables. *Appetite* **84**, 1–6.
 35. Perry LK, Samuelson LK & Burdnie JB (2014) Highchair philosophers: the impact of seating context-dependent exploration on children's naming biases. *Dev Sci* **17**, 757–765.
 36. Stark LJ, Collins FL, Osnes PG *et al.* (1986) Using reinforcement and cueing to increase healthy snack food choices in preschoolers. *J Appl Behav Anal* **19**, 367–379.
 37. Birch LL, Marlin DW & Rotter J (1984) Eating as the 'means' activity in a contingency: effects on young children's food preference. *Child Dev* **55**, 431–439.
 38. Newman J & Taylor A (1992) Effect of a means–end contingency on young children's food preferences. *J Exp Child Psychol* **53**, 200–216.
 39. Sanders MR, Patel RK, le Grice B *et al.* (1993) Children with persistent feeding difficulties: an observational analysis of the feeding interactions of problem and non-problem eaters. *Health Psychol* **12**, 64–73.
 40. Wardle J, Herrera M-L, Cooke L *et al.* (2003) Modifying children's food preferences: the effects of exposure and reward on acceptance of an unfamiliar vegetable. *Eur J Clin Nutr* **57**, 341–348.
 41. Birch LL, Zimmerman SI & Hind H (1980) The influence of social-affective context on the formation of children's food preferences. *Child Dev* **51**, 856–861.
 42. Koivisto UK, Fellenius J & Sjöden PO (1994) Relations between parental mealtime practices and children's food intake. *Appetite* **22**, 245–257.
 43. Arnett J (1989) Caregivers in day-care centers: does training matter? *J Appl Dev Psychol* **10**, 541–552.
 44. Crabtree BF & Miller WL (1999) *Doing Qualitative Research*. Thousand Oaks, CA: SAGE Publications, Inc.
 45. Snijders TAB & Bosker RJ (1999) *Multilevel Analysis: An Introduction to Basic and Advanced Multilevel Modeling*. Thousand Oaks, CA: SAGE Publications, Inc.
 46. Larson N, Ward DS, Neelon SB *et al.* (2011) What role can child-care settings play in obesity prevention? A review of the evidence and call for research efforts. *J Am Diet Assoc* **111**, 1343–1362.
 47. Ball K, Jeffery RW, Abbott G *et al.* (2010) Is healthy behavior contagious: associations of social norms with physical activity and healthy eating. *Int J Behav Nutr Phys Act* **7**, 86.

48. Wouters EJ, Larsen JK, Kremers SP *et al.* (2010) Peer influence on snacking behavior in adolescence. *Appetite* **55**, 11–17.
49. Pelletier JE, Graham DJ & Laska MN (2014) Social norms and dietary behaviors among young adults. *Am J Health Behav* **38**, 144–152.
50. Dev DA & McBride BA (2013) Academy of Nutrition and Dietetics benchmarks for nutrition in child care 2011: are child-care providers across contexts meeting recommendations? *J Acad Nutr Diet* **113**, 1346–1353.
51. Dev DA, McBride BA, Speirs KE *et al.* (2014) Predictors of Head Start and child-care providers' healthful and controlling feeding practices with children aged 2 to 5 years. *J Acad Nutr Diet* **114**, 1396–1403.
52. Swindle TM, Ward WL, Bokony P *et al.* (2016) A cross-sectional study of early childhood educators' childhood and current food insecurity and dietary intake. *J Hunger Environ Nutr* (Epublication ahead of print version).
53. US Department of Health and Human Services, Administration for Children and Families, Office of Head Start (2016) 1302.31 Teaching and the learning environment. IECLKC. Legislation and regulations: Head Start Program Performance Standards (45 CFR part 1304.23 Child Nutrition). <http://eclkc.ohs.acf.hhs.gov/hslc/standards/>. <http://eclkc.ohs.acf.hhs.gov/policy/45-cfr-chap-xiii/1302-31-teaching-and-learning-environment> (accessed December 2016).
54. Swindle TM, Patterson Z & Boden CJ (2017) A qualitative application of the Belsky model to explore early care and education teachers' mealtime history, beliefs, and interactions. *J Nutr Educ Behav* **49**, 568–578.
55. Story M, Kaphingst KM & French S (2006) The role of child care settings in obesity prevention. *Future Child* **16**, 143–168.