

Connecting the dots: Tracing the evolutionary pathway of Polar Ring Galaxies in the cases of NGC 3718, NGC 2685, and NGC 4262

Krishna R. Akhil, Sreeja S Kartha, Ujjwal Krishnan, Blesson Mathew, Robin Thomas, Shankar Ray, and Ashish Devaraj

Department of Physics and Electronics, CHRIST (Deemed to be University), Bangalore, India

Abstract

Polar Ring Galaxies (PRGs) are a unique class of galaxies characterised by a ring of gas and stars orbiting nearly orthogonal to the main body. This study delves into the evolutionary trajectory of PRGs using the exemplary trio of NGC 3718, NGC 2685, and NGC 4262. We investigate the distinct features of PRGs by analysing their ring and host components to reveal their unique characteristics through Spectral Energy Distribution (SED) fitting. Using CIGALE, we performed SED fitting to independently analyse the ring and host spatially resolved regions, marking the first decomposed SED analysis for PRGs, which examines stellar populations using high-resolution observations from *AstroSat* UVIT at a resolved scale. The UV-optical surface profiles provide an initial idea that distinct patterns in the galaxies, with differences in FUV and NUV, suggest three distinct stages of ring evolution in the selected galaxies. The study of resolved-scale stellar regions reveals that the ring regions are generally younger than their host galaxies, with the age disparity progressively decreasing along the evolutionary sequence from NGC 3718 to NGC 4262. Star formation rates (SFR) also exhibit a consistent pattern, with higher SFR in the ring of NGC 3718 compared to the others, and a progressive decrease through NGC 2685 and NGC 4262. Finally, the representation of the galaxies in the HI gas fraction versus the NUV–r plane supports the idea that they are in three different evolutionary stages of PRG evolution, with NGC 3718 in the initial stage, NGC 2685 in the intermediate stage, and NGC 4262 representing the final stage. This study concludes that PRGs undergo various evolutionary stages, as evidenced by the observed features in the ring and host components. NGC 3718, NGC 2685, and NGC 4262 represent different stages of this evolution, highlighting the dynamic nature of PRGs and emphasising the importance of studying their evolutionary processes to gain insights into galactic formation and evolution.

Keywords: galaxies: evolution –galaxies: peculiar –galaxies: star clusters: general –galaxies: structure –galaxies: photometry –galaxies: star formation –ultraviolet: galaxies

1. Introduction

Whitmore et al. (1990) defined Polar Ring Galaxies (PRGs) as unique galaxies belonging to the lenticular type (S0), characterised by a ring or disc orbiting in a nearly polar plane relative to the main galaxy. Further studies mentioned that the host galaxies are often gas-poor S0 or elliptical (E) types, while the ring consists of gas, dust, and stars (Reshetnikov and Sotnikova 1997; Finkelman et al. 2012). This definition, however, has evolved with recent observations indicating that both red and blue star-forming host galaxies can exhibit orthogonal rings (Reshetnikov and Combes 2015) and that these rings are not always star-forming (Deg et al. 2023). The major scenarios proposed for the formation of PRGs are major galaxy mergers (Bekki 1998; Bournaud and Combes 2003), tidal accretion of mass from a nearby companion (Reshetnikov

and Sotnikova 1997; Bournaud and Combes 2003), and cold gas accretion from the intergalactic medium (Brook et al. 2008).

According to Bekki (1998), PRGs have the capability to transform from a narrow ring structure to peculiar double rings or transient annular rings. Bournaud and Combes (2003) highlight that the ring structure in PRGs can remain stable even after 8 Gyrs in both merging and accretion scenarios. Also, it is seen in simulations that the majority of polar rings exhibit a warp, which typically emerges after 1 to 3 Gyrs of formation. Studies on the stability of self-gravitating polar rings suggest the likelihood of observing warps (Arnaboldi and Sparke 1994; Sparke et al. 2009). Warp signatures are also discernible in the shape profiles of simulated rings, which is in line with the anticipated behaviour of self-gravitating polar rings (Smirnov et al. 2024). Also, the properties of PRGs are more similar to those of early-type galaxies than late-type galaxies (Whitmore et al. 1990; Akhil et al. 2024a). Therefore, exploring the captivating features of the evolution of PRGs will greatly enhance our understanding of the total galaxy evolution framework.

The ring structure of PRGs is typically much bluer than the host component, indicating recent star formation activity (Mosenkov et al. 2022; Akhil et al. 2024b). In this study, we obtain the spectral

Author for correspondence: Sreeja S. Kartha: sreeja.kartha@christuniversity.in,
Akhil Krishna R.: akhil.r@res.christuniversity.in

energy distribution (SED) of the galaxies and their subcomponents to gain further insight into various parameters, such as star formation rate, stellar mass, and age. SED modelling techniques play a crucial role in exploring these galaxies, enabling us to extract valuable insights about their distinct emission sources (Cardelli et al. 1989; Burgarella et al. 2005). In this paper, we employ the SED fitting code-named 'Code Investigating GALaxy Emission' (CIGALE; Burgarella et al. 2005; Noll et al. 2009; Boquien et al. 2019) to analyse a multi-wavelength dataset of a selected sample of PRGs. In this study, we aim to investigate the resolved-scale stellar populations in the host and ring components of PRGs separately across UV, optical, and mid-IR wavelengths. We analyse their SEDs separately for regions belonging to the ring and the host. This separation is challenging, as overlapping components, particularly at certain viewing angles, can affect the photometric data of host galaxies. To assess how these variations and flux measurements influence the SED, we analyse spatially resolved regions within the galaxy, distinguishing between the host, ring, and their overlapping areas. This approach enables us to determine the physical properties and star formation histories of the ring and host components separately, offering key revelations into the evolutionary pathways of PRGs.

To comprehensively investigate and gain a better understanding of the star-forming activities within the ring components, as well as the evolutionary scenarios of PRGs, we use UV data from the Ultra-Violet Imaging Telescope (UVIT). The UVIT onboard *AstroSat* possesses the capacity to provide detailed insights into the star-forming properties of galaxies, as evidenced by the recent studies (George et al. 2018; Ujjwal et al. 2022; Devaraj et al. 2023; Mayya et al. 2024; Sundar Samantaray et al. 2024; Leahy et al. 2024; Ujjwal et al. 2024; Robin et al. 2024a,b; Shashank et al. 2025). Using GALEX data, previous investigations (Marino et al. 2009; Ordenes-Briceño et al. 2016) have explored the star-forming regions and their characteristics in PRGs. However, these studies were constrained by the lack of deep and multi-band photometric data in the UV regime. The multiple filters in the UVIT instrument are useful for better fitting galaxy SEDs. Also, the utilisation of UV images from UVIT to identify and study star-forming regions in the ring is anticipated to provide a more detailed understanding of the evolution of PRGs.

Details regarding the selected sample of galaxies are presented in Section. 2. In Section 3, the observations and data reduction procedures are outlined. Subsequently, in Section 4, the analysis is conducted, and the corresponding results are presented. Finally, we discuss and summarise the results in Sections 5 and 6, respectively.

2. A brief synopsis of the sample of PRGs used for this study.

This paper aims to study the evolution of PRGs through observational effects in the ring structure. From observational data, galaxies such as NGC 3808 (Reshetnikov et al. 1996; Ordenes-Briceño et al. 2016) and NGC 6286 (Shalyapina et al. 2004) are identified as being in the ring formation process through interactions. To find the observational features in the ring evolution of PRGs, we selected a sample of PRGs located within 20 Mpc. The availability of UVIT multiple filter observations will be used to explore the energy distributions of the ring and host separately for galaxies. Moreover, the availability of UVIT observations is crucial for investigating the star-forming regions in the PRGs on

a resolved scale. Furthermore, we require nearby well-studied PRGs; hence, we obtained three highly suitable galaxies for this proposed study: NGC 3718, NGC 2685, and NGC 4262. The physical properties of the three selected galaxies are listed in Table 1.

The discussion on the formation mechanism of the ring structure of NGC 3718 began with Schwarz (1985), and it remains unclear whether it is due to merging or accretion. Markakis et al. (2015) have explored the possibility of a merger. Recently, Watts et al. (2024) proposed that the asymmetric arm structures in NGC 3718 could be due to the passage of a low-mass companion galaxy. These studies indicate that subsequent merging, after the possible accretion of gas into a polar orbit, could also play a role in forming NGC 3718. However, the symmetry of NGC 3718 and the weak bridge between NGC 3718 and NGC 3729 suggest the possibility of accretion events. Pott et al. (2004), Józsa et al. (2004), and Krips et al. (2005) also discussed the gas distributions and symmetry of galaxies in this context.

In the case of NGC 2685, the formation scenario of the ring is confirmed to involve past accretion events based on the amount of molecular and atomic gas content, as discussed by Sil'chenko (1998), Sil'chenko and Moiseev (2014) and Schinnerer and Scoville (2002). Similarly, in the case of NGC 4262, Akhil et al. (2024a) discussed the possibility of recent accretion events based on the properties of the globular cluster systems.

Table 1. The fundamental physical characteristics of the three galaxies. The references for each property are given as footnotes for the table. The parameters listed include right ascension and declination, distance, apparent diameter in kiloparsecs, position angle, total V-band apparent magnitude, NUV magnitude, logarithmic HI total mass, logarithmic stellar mass, and logarithmic star formation rate.

Parameters	NGC 3718	NGC 2685	NGC 4262
RA ¹ (deg)	173.145	133.894	184.877
Dec ¹ (deg)	53.067	58.734	14.877
Distance ² (Mpc)	18.67	16.78	15.5
Apparent diameter ³ (kpc)	46.93	18.98	10.94
PA ⁶ (deg)	11.4	38	145
V _T ⁶ (mag)	10.69	11.35	11.46
NUV ⁷ (mag)	14.15	15.08	16.12
Log HI total mass ⁴ (M _⊙)	10.5	9.2	8.6
Log stellar mass ⁵ (M _⊙)	10.48	10.13	10.06
Log SFR ² (M _⊙ yr ⁻¹)	-0.66	-0.82	-1.97

References 1. Alam et al. (2015), 2. Dállya et al. (2018), 3. de Vaucouleurs et al. (1991) 4. De Vis et al. (2019), 5. Leroy et al. (2019), 6. Paturel et al. (2003), 7. Pak et al. (2014), Zaritsky et al. (2015), Cattorini et al. (2023).

2.1. NGC 3718

NGC 3718, also known as UGC 6524, Arp 214, and PRC D-18, is a member of the loose Ursa Major group, located at a distance of 18.7 Mpc (Tully 1988; Tully et al. 1996). Its morphology is highly peculiar, characterised by a prominent dark dust lane that spans almost perpendicular and straight across the central bulge (Krips et al. 2005). Previous studies, such as Allsopp (1979) and Schwarz (1985), have revealed a significant twist in the gas disc, which warps into an edge-on configuration where the straight dust lane is observed. The complex structure of the twisted HI disc of NGC 3718 is discussed in the studies of Pott et al. (2004), Krips et al.




Figure 1. The UV and Optical colour composites of the selected three PRGs are shown here. The upper panel showcases the UV colour composite, where the blue and yellow colours correspond to the images in UVIT FUV and NUV filters, respectively. The physical scale bar of 5 kpc is shown in the bottom left corner. The lower panel exhibits the optical colour composite, with blue, green, and red colours representing the galaxy images in SDSS g , r , and i bands, respectively.

Table 2. UVIT observation details

Object	RA	Dec	Observation ID	Observation date	P.I.	Available filters	Exposure times (s)
NGC 3718	173.1452	53.067	G05 234T02 9000000	2016-05-29	Ashok Kumar Pati	FUV (CaF2, BaF2)	1102, 4857
						NUV (NUVB13, B4, N2)	2175, 2895, 4959
NGC 2685	133.894	58.734	A04 176T01 9000001	2017-12-21	Kshama S K	FUV (CaF2)	1442
						NUV (NUVB4, Silica15)	2049, 6556
NGC 4262	184.877	14.877	A02 058T02 9000001	2017-02-19	Omkar	FUV (BaF2)	1879
						NUV (Silica15)	1890

(2005) and Sparke et al. (2009). Whitmore et al. (1990) compiled the first PRG catalogue, NGC 3718 was included in the “possibly related object” category (named as PRC D-18). Subsequent research by Sparke et al. (2009) confirmed that the innermost gas that orbits the stellar disc of the galaxy is nearly polar, whereas the outer gas rings are tilted at an angle of 30 to 40 degrees.

2.2. NGC 2685

NGC 2685, also known as the Helix galaxy or the Spindle galaxy (Sandage 1961), is located at a distance of 16.8 Mpc. Whitmore et al. (1990) classified NGC 2685 as a “kinematically confirmed” PRG. The host galaxy exhibits a nearly perpendicular ring structure, with gas and stars orbiting close to the central body (Eskridge and Pogge 1997; Silchenko 1998). The HI observations strongly suggest that the gaseous structure of NGC 2685 forms a coherent, extremely warped disc, with the appearance of two rings being attributed to projection effects. Also, at smaller radii, this disc is kinematically decoupled from the central stellar body (Józsa et al. 2009). NGC 2685 exhibits an outer component of neutral hydrogen. This outer component is positioned perpendicular to the polar

ring and appears to align with the central body of the S0 galaxy (Hagen-Thorn et al. 2005; Józsa et al. 2009).

2.3. NGC 4262

NGC 4262 is a peculiar S0 galaxy in the Virgo cluster region and a classic example of a PRG, which is at a distance of ~ 15 Mpc. Whitmore et al. (1990) have first presented NGC 4262 as a kinematically related possible polar ring candidate. In the ACS VCS survey of 100 ETGs, Ferrarese et al. (2006) explored the morphology, isophotal parameters, and surface brightness profiles of NGC 4262. They investigated and explained that NGC 4262 has a thin dust filament structure, a small regular dusty disc, and a prominent stellar bar. Using GALEX observations of NGC 4262, Buson et al. (2011) detected an extended outer ring studded with UV-bright knots surrounding the galaxy body. This outer ring has an orthogonally inclined host galaxy. From these observations, Moiseev et al. (2011) added NGC 4262 in the new PRG catalogue as SPRC 33. Recently, Akhil et al. (2024a) studied the galaxy using NGVS data and showed an optically faint ring structure that is orthogonal to the host galaxy.




Figure 2. The UV and optical surface brightness profiles of NGC 3718, NGC 2685 and NGC 4262. It is observed that the FUV profile dominates (~ 1 mag/arcsecond² than NUV) in NGC 3718, there is an overlap of FUV and NUV profiles in NGC 2685, and the NUV profile dominates in NGC 4262. The separation radius of the host galaxy obtained is indicated by a green vertical dotted line.




Figure 3. Upper and lower panels show the variations of the optical isophotal parameters: position angle and ellipticity in SDSS *g* band-filter as a function of radius. The separation radius of the host galaxy is indicated by a green vertical dotted line.

3. Data

The observations of three galaxies presented in this paper include UVIT data from the AstroSat ISSDC data archive, optical data from Sloan Digital Sky Survey (York et al. 2000, SDSS), Near-infrared data from Two Micron AllSky Survey (Skrutskie et al. 2006, 2MASS) and Mid-IR data from The Infrared Array Camera (Fazio et al. 2004; Werner et al. 2004, IRAC - Spitzer). UVIT, with a 28' field of view, observes simultaneously in FUV (130–180 nm), NUV (200–300 nm), and VIS (320–550 nm) filters. The spatial resolution of UVIT detectors is $\sim 1.5''$ with a field of view of 0.5 degrees and a pixel size of $0.416''$ per pixel (Agrawal 2006; Kumar et al. 2012; Tandon et al. 2020). The data reduction and calibration processes were conducted using the CCD Lab software (Postma and Leahy 2017). The photometric zero points were taken from Tandon et al. (2020). We followed the procedure outlined in Ujjwal et al. (2022) for the reduction and calibration of UVIT images. The observation log is listed in Table 2. The SDSS, 2MASS, and IRAC data were acquired from their respective public data archives (SDSS Science Archive and NASA/IPAC Infrared Science Archive). The Optical and UV colour composite images of the three PRGs are shown in Figure 1.

4. Analysis and Results

In order to comprehend the evolutionary trajectory of the ring component within PRGs and their broader implications in the

global context, we conducted surface photometry in both UV and optical images for three PRGs. Subsequently, SED analysis was performed on the spatially resolved regions of the galaxy. The exploration extended to identifying and examining the star-forming regions within the ring structure of the galaxies. Ultimately, the study sought to determine the position of these three galaxies within the overarching global evolutionary pathway.

4.1. UV and Optical surface photometry

We have carried out the surface photometry using Python elliptical isophotal analysis^a (Jedrzejewski 1987) in the SDSS *g* band, UVIT FUV, and NUV filters. During the fit, we allowed the isophotes free variation of the centre, ellipticity, and position angle (measured counterclockwise from the west). For NGC 3718, FUV is represented by the UVIT CaF2 filter, and the NUVB13 filter represents NUV. In the case of NGC 2685, FUV corresponds to the UVIT CaF2 filter, while NUV corresponds to the NUVB4 filter. Finally, for NGC 4262, FUV corresponds to the BaF2 filter, and NUV corresponds to the silica filter.

Based on the analysis and observations of optical and UV images, we have determined the size of the host galaxy component for three galaxies. The radial extent of the host galaxy component in these galaxies is determined based on the isophotal parameters' position angle and ellipticity. The abrupt changes in position angle and ellipticity are used to ascertain the radial extent of the host component in each galaxy (Erwin et al. 2005; Li et al. 2011; Akhil et al. 2024b). Figure 3 illustrates the ellipticity and position angle distribution for each galaxy. The host components of NGC 3718, NGC 2685, and NGC 4262 were determined to have radial extents of 50, 40, and 50 arcseconds, and the ring components extended up to 290, 200, and 130 arcseconds, respectively. Note that the radial extent values are derived directly from SDSS *g*-band observations, implying that a slight variation is possible from other studies (e.g. Sparke et al. 2009; Schinnerer and Scoville 2002; Akhil et al. 2024a). Hence, the ring and host components in the images have been distinguished. Subsequently, these specified radii are used to separate the ring and host components of the galaxies in this study from now on.

The UV and optical surface brightness profiles for the sample of three PRGs are illustrated in Figure 2. From the profile of

^a<https://photutils.readthedocs.io/en/stable/isophote.html>

Table 3. Filters and their wavelengths used in the SED fitting

Telescope	Filter	λ (Å)	Telescope	Filter	λ (Å)
UVIT	CaF2	1481	SDSS	u	3572
	BaF2	1541		g	4750
	B15	2196		r	6204
	Silica	2418		i	7519
	B13	2477		z	8992
	B4	2632	2MASS	J	12350
	N2	2792		H	16620
			IRAC	1	36000
				2	45000

NGC 3718, it is revealed that there is ~ 1 mag/arcsecond² dominance of FUV over NUV. It should be noted that in NGC 3718, the host component overlapped with a part of the ring, thereby slightly affecting the brightness distribution of the host component. In NGC 2685, we observe an overlap of FUV and NUV beyond $20''$. In the case of NGC 4262, NUV becomes more dominant (~ 0.9 mag/arcsecond²) than FUV. The brightness profiles of NGC2685 and NGC 4262 given in Figure 2 match with the UV observations reported in Rampazzo et al. (2017) and Bouquin et al. (2018). When FUV emissions in a galaxy decrease, it indicates that OB stars are no longer forming. Also, FUV emission decreases more rapidly than NUV emission, suggesting an older population (compared with FUV) of stars contributing to the NUV. When we compare the results above, these variations in the UV profile are particularly observed beyond the radius of the host galaxy. This provides valuable information about the star formation history and stellar population of the ring component of the galaxies.

4.2. Spectral energy distribution of PRGs

From the section 4.1, we successfully distinguished the ring and host components within the PRGs. In this section, we separated the ring and host components of these galaxies across various spectral regions, including UV, optical, and mid-IR wavelengths. This comprehensive approach allows us to thoroughly examine and understand the SED of ring and host components within these galaxies.

4.2.1. Pre-processing and PSF degradation

To obtain spatially resolved regions in the PRGs and perform SED fitting to study their physical properties, such as SFR, age, and mass, we processed the data in several steps. We removed foreground stars, estimated and subtracted the background, and matched the PSF of each image to the lowest resolution by degrading every band to the PSF of the 2MASS image. We used the proper motions of objects from Gaia Data Release 3 (Gaia DR3; Gaia Collaboration et al. 2023) to remove definite stars from the data (Buzzo et al. 2022; Akhil et al. 2024a). In each photometric band, we used the photutils “MeanBackground” method (Bradley et al. 2023) to estimate and subtract a local background for each star-masked image. For UVIT filters, given the minimal UV background, we manually estimated and subtracted the background. In order to standardise the resolution across datasets, we employed a Gaussian 2D kernel to convolve all images to the resolution ($3.5''$) of the 2MASS. Figure A1 shows all the images of the

Table 4. Models and parameter values used to build the SED. The values are derived from the SED fit and compared with the best available literature values. We use the default values for the parameters that are not listed here. References are: Boquien et al. (2019); Hunt et al. (2019); Turner et al. (2021)

Module	Parameter	Value
Star formation history (sfhdelaydbq)	tau_main	0.1 - 13 Gyr
	age_main	0.1 - 13 Gyr
	age_bq	0.01,0.1,0.5,1 Gyr
	r_sfr	0.01, 0.05,0.5,0.7,0.9
Stellar population (bc03)	IMF	0 (Salpeter)
	metallicity	0.02,0.05
nebular	logU	-2
Dust attenuation (dustatt_modified_starburst)	E(B-V) lines	0.01 - 1
	E(B-V) factor	0.44,0.25
	powerlaw_slope	-1, -0.1, -0.5, 0, 0.5, 1
Dust emission (dl2014)	qpah	0.47,1.77
	umin	0.1,1,5
	alpha	1,2,3
	gamma	0.1,0.5,0.9

three galaxies that were processed. The units and their error conversion procedures are executed in accordance with the guidelines stipulated in the Abdurrouf et al. (2021).

4.2.2. Spatially resolved regions

In our SED fitting analysis, we incorporated UVIT, SDSS, 2MASS, and IRAC photometric data. The specific filters utilised, and their corresponding wavelengths are given in Table 3. In the case of spatially resolved regions in PRGs, we visually categorised them into three components: regions within the ring, the host galaxy, and regions where the host and the ring overlap. We used the ProFound package to identify the brightest regions in the NUV images, where NUV emission remains consistent across all these regions. ProFound (Robotham et al. 2018; Robotham 2018), an astronomical data processing tool in the R programming language, identifies peak flux regions within an image and delineates source segments using watershed deblending. We set a criterion that requires each detected region to cover a minimum of 10 pixels based on the resolution used during convolution. Then, these segments are iteratively expanded to encompass the full photometric profile (Robotham et al. 2018). The segmentation maps derived from the NUV images were overlaid on all other available filters to estimate the flux for each region.

To identify the regions corresponding to the host, ring, and overlap areas, we used IR (IRAC 1) and UV (UVIT FUV) luminosity distributions as overlaid contours. Figure A2 visually represents the selected regions within the host, ring, and overlap areas in the case of NGC 3718 (for NGC 2685, see Figure A3). For region selection, we applied criteria that require at least one filter from the UV, optical, and IR range to show a flux greater than zero. Additionally, the central regions of NGC 3718 and NGC 2685, which might be affected by possible AGN activity, were excluded as they fall within the overlap areas. This exclusion effectively




Figure 4. SED of all the spatially resolved regions identified in galaxies NGC 3718, NGC 2685, and NGC 4262 from left to right. The blue and red shaded regions are based on the 16th and 84th percentiles of SEDs identified in the ring and host components, respectively.

mitigates AGN contamination in the selected regions. Finally, we obtained 407, 70, and 34 spatially resolved regions without any overlap in the central regions of the galaxies NGC 3718, NGC 2685, and NGC 4262, respectively.

4.2.3. SED modelling using CIGALE tool

We used CIGALE^b, a Python-based tool that handles SED modelling and fitting. The SED modelling function allows us to construct comprehensive SEDs for the ring and host regions. This is achieved through the use of a combination of specialised modules designed to understand various characteristics.

We employed the flux of each subcomponent across the wavelength range from FUV to Mid-IR (Table 3) and implemented models to calculate the optimal fit. The images of all the filters of all three galaxies are shown in Figure A1. For the star formation history, we utilised *sfhdelayedbg* model (Ciesla et al. 2017; Boquien et al. 2019), incorporating an instantaneous recent variation in the SFR, either increase or decrease. Additionally, we utilised *BC03* for stellar population modelling (Bruzual and Charlot 2003), *Nebular* for nebular emission contribution (Inoue 2011) and *dustattpowerlaw* for dust attenuation (Boquien et al. 2019). We also utilised the dust emission model by Draine et al. (2014) to model the dust emissions within the galaxies, which builds on earlier work by Draine and Li (2007). This model provides empirical templates for dust emission, dividing the emission into components associated with different radiation field intensities. Table 4 shows the models selected and the parameters chosen for computing every model in detail (Boquien et al. 2019; Boquien 2020).

Using the models and parameters, we derived the SED for spatially resolved regions in the rings and host galaxies of all three galaxies, as shown in Figure 4. Notable differences emerge between the SEDs of the ring and host regions, particularly in the UV region. In NGC 3718 and NGC 2685, the UV flux in the ring regions is higher compared to the host regions. However, in the case of NGC 4262, compared to NGC 3718 and NGC 2685, the flux variation between the host and ring regions of NGC 4262 is less pronounced.

Due to the limited number of regions and to study the ring and host separately for all three galaxies together, we combined all the

ring regions and host regions from the three galaxies. From the total regions identified across all three galaxies, we obtained approximately 70% ring regions and 30% host regions combined. The SED fits for our galaxies are deemed reliable, as their overall chi-square distributions have been thoroughly analysed. A small number of outliers ($\sim 2\%$) with chi-square values exceeding 10 were excluded from the analysis to ensure robustness. The histogram in the top-right corner of Figure 5 illustrates the distribution of reduced chi-square values for the SED fits, combining both host and ring regions across the three PRGs. To examine the host and ring components together, we analysed key physical properties derived from the CIGALE SED fitting, including stellar mass, dust mass, SFR, e-folding time (τ_{main}), and age. Figure 5 presents corner plots for these parameters, highlighting the distinct physical properties of the host and ring regions. The distinct separation observed in these properties highlights significant differences in the stellar populations across these regions in all three galaxies. To quantify these differences, we performed both the Kolmogorov-Smirnov test and the permutation test. The p-values for stellar mass, SFR, τ_{main} , and age are extremely small (essentially zero), indicating significant differences between the ring and host components. In contrast, the p-value for dust mass (~ 0.4) suggests no significant difference. These findings emphasise the presence of distinct stellar populations in the ring and host components overall. The disparity is particularly evident when comparing recent (SFR over 10 Myrs) and older (SFR over 100 Myrs) star formation rates, as shown in the top-right corner of Figure 5. For the logarithmic SFR over 10 Myrs, reflecting the most recent star formation, the mean values for the ring and host regions are -4.02 ± 0.04 and -4.91 ± 0.06 , respectively. In contrast, the logarithmic SFR over 100 Myrs, corresponding to earlier star formation, shows less variation in the ring regions (-4.09 ± 0.03) but higher values for the host (-4.48 ± 0.06). These observations suggest distinct stellar populations in the ring and host components. Moreover, the SFR in the ring regions remains relatively constant, declining rapidly in the host galaxies.

Returning to our initial interpretation, we proposed that the ring components of PRGs evolve progressively in the sequence NGC 3718, NGC 2685 and NGC 4262. To further investigate, we analysed the physical properties of the host and ring regions separately for these galaxies. Since this study focuses on the resolved stellar population regions in the PRGs, we carefully selected regions to

^b<https://cigale.lam.fr/>




Figure 5. The output parameters from SED fitting for the two-by-two combinations of parameters and histograms are presented for all four parameters: stellar mass, dust mass, SFR, e-folding time and age. The black contour regions represent the total stellar population distribution in all three PRGs. The mean values of each parameter are listed at the top of each histogram. Additionally, in each corresponding histogram, a list of three fractional quantiles [0.16, 0.5, 0.84] indicates the upper and lower errors. In the top right corner, the total best-fit chi-squared values, the SFR at 10 and 100 Myr are also shown. In all the plots, the red and blue colours represent regions belonging to the host and ring in the selected sample of PRGs.

minimise contamination between the host and ring components, as discussed in section 4.2.2. Consequently, it is important to note that this study does not provide results representing the entire ring population or the complete properties of the host galaxies. Instead, the analysis reflects the characteristics of the spatially resolved regions that were specifically identified to ensure accurate SED fitting and comparative evaluation. Figure 6 presents histograms of key parameters, including stellar mass, dust mass, SFR, τ_{main} , and age. The mean values obtained for each parameter from the SED analysis of resolved regions in three galaxies host and ring are listed in the table 5. The results reveal distinct stellar populations in the host and ring regions across all three galaxies. Moreover, we observe that the difference in mean SFR values between the host and ring regions decreases from NGC 3718 to NGC 2685, with NGC 4262 showing the smallest difference ($\Delta\text{SFR}_{\text{ring-host}} \sim 1.4 > \sim 0.7 > \sim 0.2$, respectively). This supports the hypothesis of an evolutionary sequence among these galaxies, transitioning from

NGC 3718 to NGC 4262. In terms of stellar mass, e-folding time, and age, the stellar populations exhibit significant differences between the host and ring regions in each galaxy. Interestingly, NGC 3718 shows no significant separation in dust mass between the ring and host components, suggesting that both regions contain similar amounts of dust. This might indicate heightened star formation activity in the ring and host regions.

The τ_{main} is the timescale over which the SFR declines exponentially, reducing to a factor of $1/e$ of its initial value. A larger τ_{main} indicates a slower decay of star formation (Boquien et al. 2019). This suggests that, in all three galaxies, the host is an older component with a star formation history that has gradually declined, while the ring is younger, likely more dynamically evolving, with star formation that has been sustained for longer and is now declining at a slower rate. The hosts consistently exhibit shorter τ_{main} values (NGC 3718: 3.09 ± 0.02 , NGC 2685: 2.93 ± 0.07 , NGC 4262: 3.01 ± 0.06), indicating a rapid decline in star




Figure 6. The histograms for the parameters—stellar mass, dust mass, SFR, e-folding time, and age—are presented separately for the host and ring regions, shown in blue and red, respectively. The first, second, and third rows correspond to the galaxies NGC 3718, NGC 2685, and NGC 4262. In each histogram, the mean values of the density distributions are labelled for both populations. These mean values represent the spatially resolved regions in the galaxies, not the total properties of the entire host and ring components.

Table 5. Mean values of the physical parameters for the spatially resolved ring and host regions of the three galaxies, derived from CIGALE SED fitting. These values represent only the spatially resolved regions and do not reflect the total properties of the host or ring components.

Galaxy	Component	$\log(\text{Age}) [\text{Myr}]$	$\log(M_{\text{dust}}) [M_\odot]$	$\log(M_*) [M_\odot]$	$\log(\text{SFR}) [M_\odot/\text{yr}]$	$\log(\tau_{\text{main}}) [\text{Myr}]$
NGC 3718	Host	3.74 ± 0.02	3.20 ± 0.06	6.66 ± 0.06	-5.19 ± 0.06	3.09 ± 0.02
	Ring	3.19 ± 0.03	3.21 ± 0.06	5.40 ± 0.05	-3.89 ± 0.04	3.74 ± 0.02
NGC 2685	Host	3.84 ± 0.06	3.77 ± 0.14	7.33 ± 0.14	-4.74 ± 0.15	2.93 ± 0.07
	Ring	3.74 ± 0.03	2.94 ± 0.09	6.36 ± 0.09	-4.04 ± 0.08	3.36 ± 0.04
NGC 4262	Host	3.99 ± 0.02	3.44 ± 0.27	7.50 ± 0.27	-4.49 ± 0.40	3.01 ± 0.06
	Ring	3.80 ± 0.06	2.82 ± 0.27	6.58 ± 0.09	-4.17 ± 0.32	3.42 ± 0.10

formation. Their older stellar populations ($\log(\text{Age})$: 3.74–3.99) reflect earlier, more evolved star formation histories. In contrast, the rings show longer τ_{main} values (NGC 3718: 3.74 ± 0.02 , NGC 2685: 3.36 ± 0.04 , NGC 4262: 3.42 ± 0.10), implying sustained star formation over longer periods. The younger stellar populations in the rings ($\log(\text{Age})$: 3.19–3.80) suggest more recent formation, likely triggered by interactions or accretion events. This trend highlights a progression in which NGC 3718 represents an earlier

stage, featuring a dynamically young ring and a relatively evolved host. NGC 2685 shows an intermediate state, where both components have similar ages, while NGC 4262 represents a later stage, with a fully evolved host and a gradually declining ring.




Figure 7. The left panel shows the regions identified in the PRGs overlaid on the UVIT FUV images. Blue regions represent the regions in the ring, while red regions represent those within the host. The right panels illustrate the Radial distance versus SFR of the regions. We observe higher SFR in the ring component for NGC 3718 and NGC 2685 compared to NGC 4262.

4.3. Spatial and radial distribution

Our comprehensive analysis shows that the trio of galaxies NGC 3718, NGC 2685, and NGC 4262 may trace distinct evolutionary phases within the PRGs. NGC 3718 suggests an initial stage, NGC 2685 occupies an intermediary position, and NGC 4262 represents the final stage of the evolution of the ring. To further investigate this progression, we conducted a detailed examination of the star-forming regions and UV properties at resolved scales within both the ring and host components of these galaxies.

Studying star-forming regions within a galaxy is crucial in enhancing our comprehension of its evolution. These regions possess distinctive characteristics, including size, SFR, radial distance, and distribution across morphological features, all of which contribute to our insights into the underlying mechanisms shaping the galaxy (Ujjwal et al. 2022; Robin et al. 2024b). A significant aspect of this analysis involves the utilisation of the intense FUV emissions primarily emitted by massive young hotter O-type and B-type stars to identify and track young massive star-forming regions in a galaxy (Kennicutt 1998).

This study leveraged the advanced capabilities of multiple UVIT filters, which provide substantially higher spatial resolution compared to GALEX, enabling a more detailed examination of PRGs. From the results from the CIGALE SED fitting analysis, we investigated the resolved stellar population regions in PRGs at an unprecedented scale. The left side of Figure 7 depicts the spatial distribution of identified regions within the three PRGs. The right panels of Figure 7 illustrate the SFR of identified star-forming regions in the three PRGs as a function of radius. Our results reveal that NGC 3718 displays a higher number of star-forming regions in the ring structure than the host galaxy ($\sim 72\%$). Furthermore,

we observe that the SFR increases with distance from the centre in this galaxy. Similarly, in NGC 3718 and NGC 2685, star formation is predominantly higher in the ring regions than in the host regions. In contrast, NGC 4262 shows comparatively lower star formation overall, with the host galaxy exhibiting a higher SFR than the ring structure. These observations highlight the distinct star-forming characteristics across the evolutionary stages of these PRGs, providing valuable insights into their star-formation processes.

4.4. The H I gas fraction versus NUV - r plane.

This section consolidates the preceding findings, offering a comprehensive overview of the evolutionary scenario of PRGs. The properties of HI gas and NUV-r colour in galaxies reveal their responsiveness to various physical conditions, providing valuable insights into the processes that drive the transition from blue, star-forming spirals to red, passively evolving ellipticals (Cortese and Hughes 2009; Catinella et al. 2010, 2018). Leveraging literature data, we have examined the HI gas mass fraction in relation to the NUV-r colour plane. In Figure 8, we present the positions of NGC 3718, NGC 2685, and NGC 4262 in the global evolutionary path of galaxies. For comparison, we present the sample of galaxies from the GALEX Arcibo SDSS Survey (Catinella et al. 2010, 2018) in the background, represented by a black-shaded region. To enhance comprehension, we have incorporated the Green Valley region from Salim (2014) (Green dotted lines). The results align with our earlier observations, reinforcing the notion that NGC 3718 resides in the star-forming region, NGC 2685 is transitioning into the green valley stage, and NGC 4262 stands at the conclusion of the green valley phase, poised to evolve into a red sequence galaxy. These outcomes strongly support the claim that these three galaxies stand out as exemplary subjects for future investigations, providing crucial perspectives into the diverse evolutionary trajectories of PRGs. Additionally, we have selected a subset of 11 PRGs from the literature survey, all of which have HI observations, enabling us to observe their evolutionary phases and are represented in Figure 8. The details of the literature-selected galaxies and their properties are listed in Table A1. Furthermore, for comparative analysis of morphological structures, we have included three optical images from the DECaLS survey on the right side of the plot, showcasing well-studied galaxies NGC 4560A, NGC 660, NGC 3656 and NGC 4324. Through these figures, it is evident that NGC 4650A exhibits a well-defined polar structure. According to this study, NGC 4650A may represent an earlier stage of PRG compared to NGC 3718. Subsequently, similar to NGC 3718, we observe NGC 660, and finally, akin to NGC 4262, we find NGC 3656 and NGC 4324.

5. Discussion

PRGs offer crucial insights into galaxy structure and evolution, ranging from understanding gas interaction processes to probing the shapes and distributions of their dark matter halos (Reshetnikov et al. 1996; Reshetnikov and Sotnikova 1997; Bekki 1998; Bournaud and Combes 2003; Macciò et al. 2006; Sparke et al. 2009; Khoperskov et al. 2014; Quiroga et al. 2020; Khoperskov et al. 2021; Smirnov et al. 2024). Since the initial discovery of potential stellar polar structures around nearby galaxies (Sandage 1961), there has been a growing interest in studying these peculiar galaxies in detail. Previous studies, such as




Figure 8. The HI mass fraction as a function of NUV - r colour. The region within the green dashed line corresponds to the green valley from (Salim 2014), and the region on the left and right correspond to the star-forming and quenched region, respectively. The black-shaded regions are the sample of galaxies from the GALEX Arecibo SDSS Survey (Catinella et al. 2010, 2018). NGC 3718 lies within the star-forming region, NGC 2685 is transitioning into the green valley, and NGC 4262 is situated inside the green valley. The blue and orange squares represent the selected 11 sub-sample of PRGs from the literature. To facilitate a comparison of the ring structures, we included well-studied PRGs, specifically NGC 4650A, NGC 660, NGC 3656, and NGC 4324 (represented by blue-filled squares). Also, optical colour images from the DECaLS survey of these well-studied PRGs are displayed on the right side of the plot. These images illustrate various evolutionary phases in the morphology of the galaxies.

Bekki (1998); Bournaud and Combes (2003); Brook et al. (2008), have investigated various formation scenarios for PRGs, including merging, accretion, and cold matter accretion. These studies also deliberate on the potential transformation of PRGs into E/S0-type galaxies. Simulations indicate that polar structures are not permanent, raising the possibility that rings could first form in HI, convert to optical structures, and eventually return to the galaxy host (Bournaud and Combes 2003; Khoperskov et al. 2021; Deg et al. 2023). This underscores the importance of studying the evolution of PRGs and their ring components.

This study identifies the possibility of NGC 3718, NGC 2685, and NGC 4262 as representative of three distinct stages in the evolutionary scenario of PRGs. Surface photometry in FUV and NUV from images provides valuable insights into the evolutionary processes of galaxies. The UV emission is highly sensitive to even subtle levels of star formation (Muñoz-Mateos et al. 2011; Rampazzo et al. 2017; Bouquin et al. 2018). The UV surface photometry reveals distinct patterns in the surface brightness profiles of the sample PRGs, indicating variations in the dominance of FUV and NUV components in the ring structures beyond the radius of the host galaxies. The variation in the FUV luminosity in the ring components is directed towards the direction that NGC 3718 has dominant star formation and NGC 4262 has the least star formation among them. Analysing the UV part of the electromagnetic spectrum offers a straightforward approach to measuring the ongoing SFR in galaxies. Examining resolved-scale regions within both the ring and host components adds a spatial dimension to the analysis. The identified regions exhibit varying levels of star formation, with NGC 3718 displaying higher SFR in the ring structures compared to NGC 2685 and NGC 4262. The SED analysis further provides nuanced insights into their unique characteristics and evolutionary pathways (Figure 6). Notably, the SED

results support a consistent pattern across the three galaxies, suggesting a direct correlation between the evolving nature of the ring structure and the distinct evolutionary stages of PRGs. This observation supports the notion of distinct evolutionary phases within the PRGs, with NGC 3718 representing an initial stage, NGC 2685 occupying an intermediary position, and NGC 4262 indicating a more advanced stage of ring evolution.

NGC 3718 is a well-defined example of a galaxy with a central warp as mentioned by Schwarz (1985), Krips et al. (2005) and Sparke et al. (2009). According to Arnaboldi and Sparke (1994) and Bournaud and Combes (2003), factors like inclination and mass distribution can influence the stability of PRGs and the likelihood of a warp occurring in the ring. In the case of NGC 3718, it is inferred that instability led to the initiation of a warp in the central region. This scenario provides a potential explanation for the second phase, as observed in NGC 2685. Józsa et al. (2009) found that NGC 2685 exhibits two rings, one orthogonal to the host and another situated farther towards the centre, aligning with the host. Their findings suggest that this configuration could be an aftermath of a warp. However, it's crucial to acknowledge that the possibility of other scenarios influencing the formation of the ring structure cannot be dismissed. The integration of HI gas fraction and NUV-r colour plane (Figure 8) analysis in this study aligns with the observed evolutionary stages, reinforcing the classification of NGC 3718 as a star-forming galaxy, NGC 2685 in transition to the green valley, and NGC 4262 positioned close to the termination of the green valley phase. In summary, the study provides a comprehensive narrative of the evolutionary progression within PRGs, combining observational data and modelling techniques to offer enlightening observations into the intricate processes shaping these unique galactic structures. The findings contribute to the broader understanding of galaxy evolution and highlight the potential of NGC 3718, NGC 2685, and NGC 4262 as exemplary subjects for future investigations in the field.

In addition to the three galaxies analysed in this study, a broader perspective on the evolution of PRGs requires examining a larger sample. In future work, we aim to expand our investigation to include a larger sample of galaxies, leveraging dedicated UVIT observations. The analysis of the three PRGs in this study serves as an initial exploration of the evolutionary scenario, while an extended study will enable a deeper investigation into the evolutionary stages of ring structures in PRGs. Such efforts will provide a more comprehensive view of the role of ring structures in galaxy evolution and contribute to the development of robust theoretical models for PRGs.

6. Summary

In this study, we examined the evolutionary trajectory of PRGs by utilising the most exemplary samples of galaxies: NGC 3718, NGC 2685, and NGC 4262. Our comprehensive investigation significantly enhanced our understanding of the star formation and star-forming activities within the ring components, along with shedding light on the evolutionary scenarios of PRGs. Employing SED fitting through CIGALE, we thoroughly explored the SED of the ring and host independently, marking the first instance of generating decomposed SED for PRGs.

- The UV optical surface profiles show that the FUV profile dominates in NGC 3718, FUV overlaps NUV in NGC 2685, and NUV dominate in NGC 4262. This provides an initial insight

into the star formation in the ring component for these galaxies, suggesting the existence of three distinct stages of ring evolution.

- Using SED analysis, we studied stellar populations in the ring and host regions of the three PRGs. We observed notable differences in their physical parameters derived from the SED analysis, with the ring components generally being much younger than the host galaxies. The disparity decreases along the evolutionary sequence proposed from NGC 3718 to NGC 4262, indicating progressive galaxy evolution.
- A consistent pattern emerges when considering the SFR of the components. Our findings indicate that the star-forming regions within the ring component of NGC 3718 exhibit a higher SFR than the other galaxies. The SFR decreases progressively from NGC 3718 to NGC 2685 and then to NGC 4262. Additionally, the difference between the SFR of the host and the ring components approaches zero in the order of NGC 3718 to NGC 4262.
- In the HI gas fraction versus the NUV - r plane, we observed that NGC 3718 lies within the star-forming region, NGC 2685 is transitioning into the green valley, and NGC 4262 is situated inside the green valley.

We suggest that the three galaxies are excellent examples for understanding the evolutionary pathway of PRGs: NGC 3718 is in the initial stage, followed by NGC 2685 (intermediate stage), and finally, NGC 4262 will be in the final stage of the evolution of PRGs. While these results give an initial insight regarding the class of PRGs and their evolution, they emphasise the need for a larger study. As a future endeavour, this study paves the way for modelling these evolutionary stages to facilitate further exploration.

Acknowledgement. We would like to express our sincere gratitude to the referee for their insightful comments and constructive feedback, which significantly improved the quality and clarity of this article. We thank the Center for Research, CHRIST (Deemed to be University), for all their support during this work. We thank Dr Smitha Subramanian, Dr Savithri H. Ezhikode, Dr Arun Roy, and Prajwel Joseph for their valuable comments on this work and wonderful discussions. We acknowledge the facility support from the DST FIST program (SR/FST/PS-I/2022/208). SSK and RT acknowledge the financial support from the Indian Space Research Organisation (ISRO) under the AstroSat archival data utilisation program (No. DS-2B- 3013(2)/6/2019). UK acknowledges the Department of Science and Technology (DST) for the INSPIRE FELLOWSHIP (IF180855). This publication uses UVIT, GALEX, SDSS, and 2MASS data. We gratefully thank all the individuals involved in the various teams for supporting the project from the early stages of the design and observations. This research has used the NASA/IPAC Extragalactic Database (NED), funded by the National Aeronautics and Space Administration and operated by the California Institute of Technology. This work made use of CCDLAB (Postma and Leahy 2017), SAOImageDS9 (Joye and Mandel 2003), TOPCAT (Taylor 2005), Astropy (Astropy Collaboration et al. 2013, 2018), Photutils (Bradley et al. 2023).

Data Availability Statement. All the UVIT data used in this paper are publicly available at https://astrobrowse.issdc.gov.in/astro_archive/archive/Home.jsp. The data underlying this article will be shared with the corresponding author at a reasonable request.

References

- Abdurroúf, Lin, Y.-T., Wu, P.-F., and Akiyama, M. (2021). Introducing piXed-fit: A Spectral Energy Distribution Fitting Code Designed for Resolved Sources. *ApJS*, 254(1):15.
- Agrawal, P. C. (2006). A broad spectral band Indian Astronomy satellite ‘Astrosat’. *Advances in Space Research*, 38(12):2989–2994.
- Akhil, K. R., Kartha, S. S., and Mathew, B. (2024a). Beyond the Rings: Polar Ring Galaxy NGC 4262 and its Globular Cluster System. *mnras*.
- Akhil, K. R., Kartha, S. S., Mathew, B., Ujjwal, K., Ezhikode, S. H., and Robin, T. (2024b). DES J024008.08-551047.5: A new member of the polar ring galaxy family. *A&A*, 681:A35.
- Alam, S., Albareti, F. D., and Allende Prieto, . m. (2015). The Eleventh and Twelfth Data Releases of the Sloan Digital Sky Survey: Final Data from SDSS-III. *ApJS*, 219(1):12.
- Allsopp, N. J. (1979). A neutral hydrogen survey of the peculiar early-type galaxy NGC 3718. *MNRAS*, 186:343–355.
- Arnaboldi, M. and Sparke, L. S. (1994). Self-Gravitation Polar Rings in Axisymmetric and Triaxial Galaxy Potentials. *AJ*, 107:958.
- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., Günther, H. M., Lim, P. L., Crawford, S. M., Conseil, S., Shupe, D. L., Craig, M. W., Dencheva, N., Ginsburg, A., VanderPlas, J. T., Bradley, L. D., Pérez-Suárez, D., de Val-Borro, M., Aldcroft, T. L., Cruz, K. L., Robitaille, T. P., Tollerud, E. J., Ardelean, C., Babej, T., Bach, Y. P., Bachetti, M., Bakanov, A. V., Bamford, S. P., Barentsen, G., Barmby, P., Baumbach, A., Berry, K. L., Biscani, F., Boquien, M., Bostroem, K. A., Bouma, L. G., Brammer, G. B., Bray, E. M., Breytenbach, H., Buddelmeijer, H., Burke, D. J., Calderone, G., Cano Rodríguez, J. L., Cara, M., Cardoso, J. V. M., Cheedella, S., Copin, Y., Corrales, L., Crichton, D., D’Avella, D., Deil, C., Depagne, É., Dietrich, J. P., Donath, A., Droettboom, M., Earl, N., Erben, T., Fabbro, S., Ferreira, L. A., Finethy, T., Fox, R. T., Garrison, L. H., Gibbons, S. L. J., Goldstein, D. A., Gommers, R., Greco, J. P., Greenfield, P., Groener, A. M., Grollier, F., Hagen, A., Hirst, P., Homeier, D., Horton, A. J., Hosseinzadeh, G., Hu, L., Hunkeler, J. S., Ivezić, Ž., Jain, A., Jenness, T., Kanarek, G., Kendrew, S., Kern, N. S., Kerzendorf, W. E., Khvalko, A., King, J., Kirkby, D., Kulkarni, A. M., Kumar, A., Lee, A., Lenz, D., Littlefair, S. P., Ma, Z., Macleod, D. M., Mastroiello, M., McCully, C., Montagnac, S., Morris, B. M., Mueller, M., Mumford, S. J., Muna, D., Murphy, N. A., Nelson, S., Nguyen, G. H., Ninan, J. P., Nöthe, M., Ogaz, S., Oh, S., Parejko, J. K., Parley, N., Pascual, S., Patil, R., Patil, A. A., Plunkett, A. L., Prochaska, J. X., Rastogi, T., Reddy Janga, V., Sabater, J., Sakurikar, P., Seifert, M., Sherbert, L. E., Sherwood-Taylor, H., Shih, A. Y., Sick, J., Silbiger, M. T., Singanamalla, S., Singer, L. P., Sladen, P. H., Sooley, K. A., Sornarajah, S., Streicher, O., Teuben, P., Thomas, S. W., Tremblay, G. R., Turner, J. E. H., Terrón, V., van Kerkwijk, M. H., de la Vega, A., Watkins, L. L., Weaver, B. A., Whitmore, J. B., Woillez, J., Zabalza, V., and Astropy Contributors (2018). The Astropy Project: Building an Open-science Project and Status of the v2.0 Core Package. *AJ*, 156(3):123.
- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., Greenfield, P., Droettboom, M., Bray, E., Aldcroft, T., Davis, M., Ginsburg, A., Price-Whelan, A. M., Kerzendorf, W. E., Conley, A., Crighton, N., Barbary, K., Muna, D., Ferguson, H., Grollier, F., Parikh, M. M., Nair, P. H., Unther, H. M., Deil, C., Woillez, J., Conseil, S., Kramer, R., Turner, J. E. H., Singer, L., Fox, R., Weaver, B. A., Zabalza, V., Edwards, Z. I., Azalee Bostroem, K., Burke, D. J., Casey, A. R., Crawford, S. M., Dencheva, N., Ely, J., Jenness, T., Labrie, K., Lim, P. L., Pierfederici, F., Pontzen, A., Ptak, A., Refsdal, B., Servillat, M., and Streicher, O. (2013). Astropy: A community Python package for astronomy. *A&A*, 558:A33.
- Bekki, K. (1998). Formation of a Polar Ring Galaxy in a Galaxy Merger. *ApJ*, 499(2):635–649.
- Bianchi, L., Herald, J., Efremova, B., Girardi, L., Zabot, A., Marigo, P., Conti, A., and Shiao, B. (2011). GALEX catalogs of UV sources: statistical properties and sample science applications: hot white dwarfs in the Milky Way. *Ap&SS*, 335(1):161–169.
- Boquien, M. (2020). CIGALE: a python Code Investigating GALaxy Emission. In *American Astronomical Society Meeting Abstracts #235*, volume 235 of *American Astronomical Society Meeting Abstracts*, page 228.01.

- Boquien, M., Burgarella, D., Roehlly, Y., Buat, V., Ciesla, L., Corre, D., Inoue, A. K., and Salas, H. (2019). CIGALE: a python Code Investigating GALaxy Emission. *A&A*, 622:A103.
- Bouquin, A. Y. K., Gil de Paz, A., Muñoz-Mateos, J. C., Boissier, S., Sheth, K., Zaritsky, D., Peletier, R. F., Knapen, J. H., and Gallego, J. (2018). The GALEX/S⁴G Surface Brightness and Color Profiles Catalog. I. Surface Photometry and Color Gradients of Galaxies. *ApJS*, 234(2):18.
- Bournaud, F. and Combes, F. (2003). Formation of polar ring galaxies. *A&A*, 401:817–833.
- Bradley, L., Sipőcz, B., Robitaille, T., Tollerud, E., Vinícius, Z., Deil, C., Barbary, K., Wilson, T. J., Busko, I., Donath, A., and et al (2023). *astropy/photutils*: 1.8.0.
- Brook, C. B., Governato, F., Quinn, T., Wadsley, J., Brooks, A. M., Willman, B., Stilp, A., and Jonsson, P. (2008). The Formation of Polar Disk Galaxies. *ApJ*, 689(2):678–686.
- Brown, M. J. I., Moustakas, J., Smith, J. D. T., da Cunha, E., Jarrett, T. H., Imanishi, M., Armus, L., Brandl, B. R., and Peek, J. E. G. (2014). An Atlas of Galaxy Spectral Energy Distributions from the Ultraviolet to the Mid-infrared. *ApJS*, 212(2):18.
- Bruzual, G. and Charlot, S. (2003). Stellar population synthesis at the resolution of 2003. *MNRAS*, 344(4):1000–1028.
- Burgarella, D., Buat, V., and Iglesias-Páramo, J. (2005). Star formation and dust attenuation properties in galaxies from a statistical ultraviolet-to-far-infrared analysis. *MNRAS*, 360(4):1413–1425.
- Buson, L. M., Bettoni, D., and Galletta, G. (2011). An extended ultraviolet ring around the SB0 galaxy NGC 4262. *Ap&SS*, 335(1):231–236.
- Buzzo, M. L., Cortesi, A., Forbes, D. A., Brodie, J. P., Couch, W. J., Barbosa, C. E., de Brito Silva, D., Coelho, P., Chies-Santos, A. L., Escudero, C., Sesto, L., Menéndez-Delmestre, K., Gonçalves, T. S., Bom, C. R., Alvarez-Candal, A., Smith Castelli, A. V., Schoenell, W., Kanaan, A., Ribeiro, T., and Mendes de Oliveira, C. (2022). A new method to detect globular clusters with the S-PLUS survey. *MNRAS*, 510(1):1383–1392.
- Cardelli, J. A., Clayton, G. C., and Mathis, J. S. (1989). The Relationship between Infrared, Optical, and Ultraviolet Extinction. *ApJ*, 345:245.
- Catinella, B., Saintonge, A., Janowiecki, S., Cortese, L., Davé, R., Lemonias, J. J., Cooper, A. P., Schiminovich, D., Hummels, C. B., Fabello, S., Geréb, K., Kilborn, V., and Wang, J. (2018). xGASS: total cold gas scaling relations and molecular-to-atomic gas ratios of galaxies in the local Universe. *MNRAS*, 476(1):875–895.
- Catinella, B., Schiminovich, D., Kauffmann, G., and Fabello, e. a. (2010). The GALEX Arcibo SDSS Survey - I. Gas fraction scaling relations of massive galaxies and first data release. *MNRAS*, 403(2):683–708.
- Cattorini, F., Gavazzi, G., Boselli, A., and Fossati, M. (2023). A complete spectroscopic catalogue of local galaxies in the northern spring sky: Gas properties and nuclear activity in different environments. *A&A*, 671:A118.
- Ciesla, L., Elbaz, D., and Fensch, J. (2017). The SFR-M_{*} main sequence archetypal star-formation history and analytical models. *A&A*, 608:A41.
- Cortese, L. and Hughes, T. M. (2009). Evolutionary paths to and from the red sequence: star formation and HI properties of transition galaxies at $z \sim 0$. *MNRAS*, 400(3):1225–1240.
- Dálya, G., Galgóczi, G., Dobos, L., Frei, Z., Heng, I. S., Macas, R., Messenger, C., Raffai, P., and de Souza, R. S. (2018). GLADE: A galaxy catalogue for multimessenger searches in the advanced gravitational-wave detector era. *MNRAS*, 479(2):2374–2381.
- de Vaucouleurs, G., de Vaucouleurs, A., Corwin, Herold G., J., Buta, R. J., Paturel, G., and Fouque, P. (1991). *Third Reference Catalogue of Bright Galaxies*.
- De Vis, P., Jones, A., Viaene, S., Casasola, V., Clark, C. J. R., Baes, M., Bianchi, S., Cassara, L. P., Davies, J. I., De Looze, I., Galametz, M., Galliano, F., Lianou, S., Madden, S., Manilla-Robles, A., Mosenkov, A. V., Nersesian, A., Roychowdhury, S., Xilouris, E. M., and Ysard, N. (2019). A systematic metallicity study of DustPedia galaxies reveals evolution in the dust-to-metal ratios. *A&A*, 623:A5.
- Deg, N., Palleske, R., Spekkens, K., Wang, J., Jarrett, T., English, J., Lin, X., Yeung, J., and Mould, J. R. e. a. (2023). WALLABY pilot survey: the potential polar ring galaxies NGC 4632 and NGC 6156. *MNRAS*, 525(3):4663–4684.
- Devaraj, A., Joseph, P., Stalin, C. S., Tandon, S. N., and Ghosh, S. K. (2023). UVIT Observations of the Small Magellanic Cloud: Point-source Catalog. *ApJ*, 946(2):65.
- Draine, B. T., Aniano, G., Krause, O., Groves, B., Sandstrom, K., Braun, R., Leroy, A., Klaas, U., Linz, H., Rix, H.-W., Schinnerer, E., Schmiedeke, A., and Walter, F. (2014). Andromeda's Dust. *ApJ*, 780(2):172.
- Draine, B. T. and Li, A. (2007). Infrared Emission from Interstellar Dust. IV. The Silicate-Graphite-PAH Model in the Post-Spitzer Era. *ApJ*, 657(2):810–837.
- Erwin, P., Beckman, J. E., and Pohlen, M. (2005). Antitruncation of Disks in Early-Type Barred Galaxies. *ApJ*, 626(2):L81–L84.
- Eskridge, P. B. and Pogge, R. W. (1997). H II Region Abundances in the Polar Ring of NGC 2685. *ApJ*, 486(1):259–267.
- Fazio, G. G., Hora, J. L., Allen, L. E., Ashby, M. L. N., Barmby, P., Deutsch, L. K., Huang, J. S., Kleiner, S., Marengo, M., Megeath, S. T., Melnick, G. J., Pahre, M. A., Patten, B. M., Polizotti, J., Smith, H. A., Taylor, R. S., Wang, Z., Willner, S. P., Hoffmann, W. F., Pipher, J. L., Forrest, W. J., McMurty, C. W., McCreight, C. R., McKelvey, M. E., McMurray, R. E., Koch, D. G., Moseley, S. H., Arendt, R. G., Mentzell, J. E., Marx, C. T., Losch, P., Mayman, P., Eichhorn, W., Krebs, D., Jhabvala, M., Gezari, D. Y., Fixsen, D. J., Flores, J., Shakoordadeh, K., Jungo, R., Hakun, C., Workman, L., Karpati, G., Kichak, R., Whitley, R., Mann, S., Tollestrup, E. V., Eisenhardt, P., Stern, D., Gorjian, V., Bhattacharya, B., Carey, S., Nelson, B. O., Glaccum, W. J., Lacy, M., Lowrance, P. J., Laine, S., Reach, W. T., Stauffer, J. A., Surace, J. A., Wilson, G., Wright, E. L., Hoffman, A., Domingo, G., and Cohen, M. (2004). The Infrared Array Camera (IRAC) for the Spitzer Space Telescope. *ApJS*, 154(1):10–17.
- Ferrarese, L., Côté, P., Jordán, A., Peng, E. W., Blakeslee, J. P., Piatek, S., Mei, S., Merritt, D., Milosavljević, M., Tonry, J. L., and West, M. J. (2006). The ACS Virgo Cluster Survey. VI. Isophotal Analysis and the Structure of Early-Type Galaxies. *ApJS*, 164(2):334–434.
- Finkelman, I., Funes, J. G., and Brosch, N. (2012). Polar ring galaxies in the Galaxy Zoo. *MNRAS*, 422(3):2386–2398.
- Gaia Collaboration, Vallenari, A., Brown, A. G. A., Prusti, T., de Bruijne, J. H. J., Arenou, F., Babusiaux, C., Biermann, M., Creevey, O. L., and et al... (2023). Gaia Data Release 3. Summary of the content and survey properties. *A&A*, 674:A1.
- Gavazzi, G., Consolandi, G., Dotti, M., Fossati, M., Savorgnan, G., Gualandri, R., and Bruni, I. (2013). Red-channel (6000–8000 Å) nuclear spectra of 376 local galaxies. *A&A*, 558:A68.
- George, K., Joseph, P., Mondal, C., Devaraj, A., Subramaniam, A., Stalin, C. S., Côté, P., Ghosh, S. K., Hutchings, J. B., Mohan, R., Postma, J., Sankarasubramanian, K., Sreekumar, P., and Tandon, S. N. (2018). UVIT observations of the star-forming ring in NGC 7252: Evidence of possible AGN feedback suppressing central star formation. *A&A*, 613:L9.
- Gil de Paz, A., Boissier, S., Madore, B. F., Seibert, M., Joe, Y. H., Boselli, A., Wyder, T. K., Thilker, D., Bianchi, L., Rey, S.-C., Rich, R. M., Barlow, T. A., Conrow, T., Forster, K., Friedman, P. G., Martin, D. C., Morrissey, P., Neff, S. G., Schiminovich, D., Small, T., Donas, J., Heckman, T. M., Lee, Y.-W., Milliard, B., Szalay, A. S., and Yi, S. (2007). The GALEX Ultraviolet Atlas of Nearby Galaxies. *ApJS*, 173(2):185–255.
- Hagen-Thorn, V. A., Shalyapina, L. V., Karataeva, G. M., Yakovleva, V. A., Moiseev, A. V., and Burenkov, A. N. (2005). A Spectroscopic and Photometric Study of NGC 2685, the Prototype Polar-Ring Galaxy. *Astronomy Reports*, 49(12):958–966.
- Hall, M., Courteau, S., Dutton, A. A., McDonald, M., and Zhu, Y. (2012). An investigation of Sloan Digital Sky Survey imaging data and multiband scaling relations of spiral galaxies. *MNRAS*, 425(4):2741–2765.
- Hunt, L. K., De Looze, I., Boquien, M., Nikutta, R., Rossi, A., Bianchi, S., Dale, D. A., Granato, G. L., Kennicutt, R. C., Silva, L., and Ciesla, e. a. (2019). Comprehensive comparison of models for spectral energy distributions from 0.1 μm to 1 mm of nearby star-forming galaxies. *A&A*, 621:A51.
- Inoue, A. K. (2011). Rest-frame ultraviolet-to-optical spectral characteristics of extremely metal-poor and metal-free galaxies. *MNRAS*, 415(3):2920–2931.
- Jedrzejewski, R. I. (1987). CCD surface photometry of elliptical galaxies - I. Observations, reduction and results. *MNRAS*, 226:747–768.

- Joye, W. A. and Mandel, E. (2003). New Features of SAOImage DS9. In Payne, H. E., Jedrzejewski, R. I., and Hook, R. N., editors, *Astronomical Data Analysis Software and Systems XII*, volume 295 of *Astronomical Society of the Pacific Conference Series*, page 489.
- Józsa, G., Oosterloo, T., Klein, U., and Kenn, F. (2004). Kinematics and morphology of warped disk galaxies. In Dettmar, R., Klein, U., and Salucci, P., editors, *Baryons in Dark Matter Halos*, page 109.
- Józsa, G. I. G., Oosterloo, T. A., Morganti, R., Klein, U., and Erben, T. (2009). Kinematic modeling of disk galaxies. III. The warped Spindle NGC 2685. *A&A*, 494(2):489–508.
- Kennicutt, Robert C., J. (1998). Star Formation in Galaxies Along the Hubble Sequence. *ARA&A*, 36:189–232.
- Khoperskov, S., Zinchenko, I., Avramov, B., Khrapov, S., Berczik, P., Saburova, A., Ishchenko, M., Khoperskov, A., Pulsoni, C., Venichenko, Y., Bizyaev, D., and Moiseev, A. (2021). Extreme kinematic misalignment in IllustrisTNG galaxies: the origin, structure, and internal dynamics of galaxies with a large-scale counterrotation. *MNRAS*, 500(3):3870–3888.
- Khoperskov, S. A., Moiseev, A. V., Khoperskov, A. V., and Saburova, A. S. (2014). The shape of dark matter halo in PRG NGC 4262. *arXiv e-prints*, page arXiv:1401.2562.
- Krips, M., Eckart, A., Neri, R., Pott, J. U., Leon, S., Combes, F., García-Burillo, S., Hunt, L. K., Baker, A. J., Tacconi, L. J., Englmaier, P., Schinnerer, E., and Boone, F. (2005). Molecular gas in Nuclei of Galaxies (NUGA). III. The warped LINER NGC 3718. *A&A*, 442(2):479–493.
- Kumar, A., Ghosh, S. K., Hutchings, J., Kamath, P. U., Kathiravan, S., Mahesh, P. K., Murthy, J., Nagbhushana, S., Pati, A. K., Rao, M. N., Rao, N. K., Sriram, S., and Tandon, S. N. (2012). Ultra Violet Imaging Telescope (UVIT) on ASTROSAT. In Takahashi, T., Murray, S. S., and den Herder, J.-W. A., editors, *Space Telescopes and Instrumentation 2012: Ultraviolet to Gamma Ray*, volume 8443 of *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, page 84431N.
- Leahy, D. A., Hansen, J., and Hopkins, A. M. (2024). Far-ultraviolet to Far-infrared Spectral Energy Distribution Modeling of the Star Formation History across M31. *AJ*, 167(5):211.
- Leroy, A. K., Sandstrom, K. M., Lang, D., Lewis, A., Salim, S., Behrens, E. A., Chastenot, J., Chiang, I.-D., Gallagher, M. J., Kessler, S., and Utomo, D. (2019). A $z = 0$ Multiwavelength Galaxy Synthesis. I. A WISE and GALEX Atlas of Local Galaxies. *ApJS*, 244(2):24.
- Li, Z.-Y., Ho, L. C., Barth, A. J., and Peng, C. Y. (2011). The Carnegie-Irvine Galaxy Survey. II. Isophotal Analysis. *ApJS*, 197(2):22.
- Macciò, A. V., Moore, B., and Stadel, J. (2006). The Origin of Polar Ring Galaxies: Evidence for Galaxy Formation by Cold Accretion. *ApJ*, 636(1):L25–L28.
- Marino, A., Iodice, E., Tantalò, R., Piovàn, L., Bettoni, D., Buson, L. M., Chiosi, C., Galletta, G., Rampazzo, R., and Rich, R. M. (2009). GALEX UV properties of the polar ring galaxy MCG-05-07-001 and the shell galaxies NGC 1210 and NGC 5329. *AAP*, 508(3):1235–1252.
- Markakis, K., Dierkes, J., Eckart, A., Nishiyama, S., Britzen, S., García-Marín, M., Horrobin, M., Muxlow, T., and Zensus, J. A. (2015). Subaru and e-Merlin observations of NGC 3718. Diaries of a supermassive black hole recoil? *A&A*, 580:A11.
- Mayya, Y. D., Barway, S., Gómez-González, V. M. A., and Zaragoza-Cardiel, J. (2024). Star formation history of the post-collisional Cartwheel galaxy using AstroSat/UVIT FUV images. *MNRAS*, 527(2):2816–2830.
- Moiseev, A. V., Smirnova, K. I., Smirnova, A. A., and Reshetnikov, V. P. (2011). A new catalogue of polar-ring galaxies selected from the Sloan Digital Sky Survey. *MNRAS*, 418(1):244–257.
- Mosenkov, A. V., Reshetnikov, V. P., Skryabina, M. N., and Shakespear, Z. (2022). Unveiling the Nature of Polar-ring Galaxies from Deep Imaging. *Research in Astronomy and Astrophysics*, 22(11):115003.
- Muñoz-Mateos, J. C., Boissier, S., Gil de Paz, A., Zamorano, J., Kennicutt, R. C., J., Moustakas, J., Prantzos, N., and Gallego, J. (2011). Radial Distribution of Stars, Gas, and Dust in SINGS Galaxies. III. Modeling the Evolution of the Stellar Component in Galaxy Disks. *ApJ*, 731(1):10.
- Noll, S., Burgarella, D., Giovannoli, E., Buat, V., Marcellac, D., and Muñoz-Mateos, J. C. (2009). Analysis of galaxy spectral energy distributions from far-UV to far-IR with CIGALE: studying a SINGS test sample. *A&A*, 507(3):1793–1813.
- Ordenes-Briceño, Y., Georgiev, I. Y., Puzia, T. H., Goudfrooij, P., and Arnaboldi, M. (2016). Compact stellar systems in the polar ring galaxies NGC 4650A and NGC 3808B: Clues to polar disk formation. *AAP*, 585:A156.
- Pak, M., Rey, S.-C., Lisker, T., Lee, Y., Kim, S., Sung, E.-C., Jerjen, H., and Chung, J. (2014). The properties of early-type galaxies in the Ursa Major cluster. *MNRAS*, 445(1):630–647.
- Parkash, V., Brown, M. J. I., Jarrett, T. H., and Bonne, N. J. (2018). Relationships between HI Gas Mass, Stellar Mass, and the Star Formation Rate of HICAT+WISE (H I-WISE) Galaxies. *ApJ*, 864(1):40.
- Paturel, G., Petit, C., Prugniel, P., Theureau, G., Rousseau, J., Brouty, M., Dubois, P., and Cambrésy, L. (2003). HYPERLEDA. I. Identification and designation of galaxies. *A&A*, 412:45–55.
- Postma, J. E. and Leahy, D. (2017). CCDLAB: A Graphical User Interface FITS Image Data Reducer, Viewer, and Canadian UVIT Data Pipeline. *PASP*, 129(981):115002.
- Pott, J. U., Hartwich, M., Eckart, A., Leon, S., Krips, M., and Straubmeier, C. (2004). Warped molecular gas disk in NGC 3718. *A&A*, 415:27–38.
- Quiroga, L. F., Muñoz-Cuartas, J. C., Rodríguez, I., and Libeskind, N. I. (2020). Simulating the galactic system in interaction AM 2229-735 and the formation of its polar structure. *MNRAS*, 491(2):1887–1900.
- Rampazzo, R., Mazzei, P., Marino, A., Uslenghi, M., Trinchieri, G., and Wolter, A. (2017). Investigating early-type galaxy evolution with a multi-wavelength approach. II. The UV structure of 11 galaxies with Swift-UVOT. *A&A*, 602:A97.
- Reshetnikov, V. and Combes, F. (2015). Polar-ring galaxies: the SDSS view on the symbiotic galaxies. *MNRAS*, 447(3):2287–2294.
- Reshetnikov, V. and Sotnikova, N. (1997). Global structure and formation of polar-ring galaxies. *AAP*, 325:933–942.
- Reshetnikov, V. P., Hagen-Thorn, V. A., and Yakovleva, V. A. (1996). Photometric study of polar-ring galaxies. III. Forming rings. *A&A*, 314:729–737.
- Robin, T., Kartha, S. S., Akhil Krishna, R., Krishnan, U., Mathew, B., Cysil, T. B., Patra, N. N., and Shridharan, B. (2024a). The Interaction Jigsaw: investigating star formation in interacting galaxies. *MNRAS*, 534(3):1902–1912.
- Robin, T., Kartha, S. S., Ujjwal, K., Saha, K., Parekh, V., George, K., and Mathew, B. (2024b). What drives the wheels of evolution in NGC 1512? A UVIT study. *A&A*, 681:A7.
- Robotham, A. S. G. (2018). ProFound: Source Extraction and Application to Modern Survey Data. Astrophysics Source Code Library, record ascl:1804.006.
- Robotham, A. S. G., Davies, L. J. M., Driver, S. P., Koushan, S., Taranu, D. S., Casura, S., and Liske, J. (2018). ProFound: Source Extraction and Application to Modern Survey Data. *MNRAS*, 476(3):3137–3159.
- Salim, S. (2014). Green Valley Galaxies. *Serbian Astronomical Journal*, 189:1–14.
- Sandage, A. (1961). *The Hubble Atlas of Galaxies*.
- Schinnerer, E. and Scoville, N. (2002). First Interferometric Observations of Molecular Gas in a Polar Ring: The Helix Galaxy NGC 2685. *ApJ*, 577(2):L103–L106.
- Schwarz, U. J. (1985). HI observations of the peculiar galaxy NGC 3718 and its companion NGC3729. *A&A*, 142:273–288.
- Seibert, M., Wyder, T., Neill, J., Madore, B., Bianchi, L., Smith, M., Shiao, B., Schiminovich, D., Rich, R. M., Conrow, T., Martin, D. C., and GALEX Catalog Team (2012). The Galaxy Evolution Explorer (GALEX) Source Catalogs. In *American Astronomical Society Meeting Abstracts #219*, volume 219 of *American Astronomical Society Meeting Abstracts*, page 340.01.
- Shalyapina, L. V., Moiseev, A. V., Yakovleva, V. A., Hagen-Thorn, V. A., and Burenkov, A. N. (2004). The Phenomenon of the Galaxy NGC 6286: A Forming Polar Ring or a Superwind? *Astronomy Letters*, 30:1–13.
- Shashank, G., Subramanian, S., Muraleedharan, S., Menon, S. H., Mondal, C., Krishna, S., Das, M., and Subramaniam, A. (2025). Tracing hierarchical star formation out to kiloparsec scales in nearby spiral galaxies with UVIT. *A&A*, 693:A188.
- Sil'chenko, O. K. (1998). Central regions of the polar-ring galaxies NGC 2685 and IC 1689. *A&A*, 330:412–418.

- Silchenko, O. K. (1998). Central regions of the polar-ring galaxies NGC 2685 and IC 1689. *A&A*, 330:412–418.
- Sil'chenko, O. K. and Moiseev, A. V. (2014). Inner Polar Gaseous Disks: Incidence, Ages, and Possible Origin. In Iodice, E. and Corsini, E. M., editors, *Multi-Spin Galaxies*, volume 486 of *Astronomical Society of the Pacific Conference Series*, page 27.
- Skrutskie, M. F., Cutri, R. M., Stiening, R., Weinberg, M. D., Schneider, S., Carpenter, J. M., Beichman, C., Capps, R., Chester, T., Elias, J., Huchra, J., Liebert, J., Lonsdale, C., Monet, D. G., Price, S., Seitzer, P., Jarrett, T., Kirkpatrick, J. D., Gizis, J. E., Howard, E., Evans, T., Fowler, J., Fullmer, L., Hurt, R., Light, R., Kopan, E. L., Marsh, K. A., McCallon, H. L., Tam, R., Van Dyk, S., and Wheelock, S. (2006). The Two Micron All Sky Survey (2MASS). *AJ*, 131(2):1163–1183.
- Smirnov, D. V., Mosenkov, A. V., and Reshetnikov, V. P. (2024). Polar-ring galaxies in the Illustris TNG50 simulation. *MNRAS*, 527(2):4112–4128.
- Sparke, L. S., van Moorsel, G., Schwarz, U. J., and Vogelaar, M. (2009). The Remarkable Warped and Twisted Gas Disk in NGC 3718. *AJ*, 137(4):3976–3992.
- Sundar Samantaray, A., Jassal, H. K., Singh, K. P., and Dewangan, G. C. (2024). AstroSat observations of interacting galaxies NGC 7469 and IC 5283. *arXiv e-prints*, page arXiv:2404.12527.
- Tandon, S. N., Postma, J., Joseph, P., Devaraj, A., Subramanian, A., Barve, I. V., George, K., Ghosh, S. K., Girish, V., Hutchings, J. B., Kamath, P. U., Kathiravan, S., Kumar, A., Lancelot, J. P., Leahy, D., Mahesh, P. K., Mohan, R., Nagabhushana, S., Pati, A. K., Rao, N. K., Sankarasubramanian, K., Sriram, S., and Stalin, C. S. (2020). Additional Calibration of the Ultraviolet Imaging Telescope on Board AstroSat. *AJ*, 159(4):158.
- Taylor, M. B. (2005). TOPCAT & STIL: Starlink Table/VOTable Processing Software. In Shopbell, P., Britton, M., and Ebert, R., editors, *Astronomical Data Analysis Software and Systems XIV*, volume 347 of *Astronomical Society of the Pacific Conference Series*, page 29.
- Tempel, E., Tuvikene, T., Kipper, R., and Libeskind, N. I. (2017). Merging groups and clusters of galaxies from the SDSS data. The catalogue of groups and potentially merging systems. *A&A*, 602:A100.
- Tully, R. B. (1988). *Nearby galaxies catalog*.
- Tully, R. B., Verheijen, M. A. W., Pierce, M. J., Huang, J.-S., and Wainscoat, R. J. (1996). The Ursa Major Cluster of Galaxies. I. Cluster Definition and Photometric Data. *AJ*, 112:2471.
- Turner, J. A., Dale, D. A., Lee, J. C., Boquien, M., Chandar, R., Deger, S., and Larson, E. A. (2021). PHANGS-HST: star cluster spectral energy distribution fitting with CIGALE. *MNRAS*, 502(1):1366–1385.
- Ujjwal, K., Kartha, S. S., Akhil, K. R., Mathew, B., Subramanian, S., Sudheesh, T. P., and Thomas, R. (2024). Disentangling the association of PAH molecules with star formation. Insights from the James Webb Space Telescope and from the UltraViolet Imaging Telescope. *A&A*, 684:A71.
- Ujjwal, K., Kartha, S. S., Subramanian, S., George, K., Thomas, R., and Mathew, B. (2022). Understanding the secular evolution of NGC 628 using UVIT. *MNRAS*.
- Watts, C., Das, M., and Barway, S. (2024). Anatomy of the Star Formation in a Tidally Disturbed Disk Galaxy: NGC 3718. *ApJ*, 974(2):206.
- Werner, M. W., Roellig, T. L., Low, F. J., Rieke, G. H., Rieke, M., Hoffmann, W. F., Young, E., Houck, J. R., Brandl, B., Fazio, G. G., Hora, J. L., Gehr, R. D., Helou, G., Soifer, B. T., Stauffer, J., Keene, J., Eisenhardt, P., Gallagher, D., Gautier, T. N., Irace, W., Lawrence, C. R., Simmons, L., Van Cleve, J. E., Jura, M., Wright, E. L., and Cruikshank, D. P. (2004). The Spitzer Space Telescope Mission. *ApJS*, 154(1):1–9.
- Whitmore, B. C., Lucas, R. A., McElroy, D. B., Steiman-Cameron, T. Y., Sackett, P. D., and Olling, R. P. (1990). New Observations and a Photographic Atlas of Polar-Ring Galaxies. *AJ*, 100:1489.
- York, D. G., Adelman, J., Anderson, John E., J., Anderson, S. F., Annis, J., Bahcall, N. A., Bakken, J. A., Barkhouser, R., Bastian, S., Berman, E., Boroski, W. N., Bracker, S., Briegel, C., Briggs, J. W., Brinkmann, J., Brunner, R., Burles, S., Carey, L., Carr, M. A., Castander, F. J., Chen, B., Colestock, P. L., Connolly, A. J., Crocker, J. H., Csabai, I., Czarapata, P. C., Davis, J. E., Doi, M., Dombeck, T., Eisenstein, D., Ellman, N., Elms, B. R., Evans, M. L., Fan, X., Federwitz, G. R., Fiscelli, L., Friedman, S., Frieman, J. A., Fukugita, M., Gillespie, B., Gunn, J. E., Gurbani, V. K., de Haas, E., Haldeman, M., Harris, F. H., Hayes, J., Heckman, T. M., Hennessy, G. S., Hindsley, R. B., Holm, S., Holmgren, D. J., Huang, C.-h., Hull, C., Husby, D., Ichikawa, S.-I., Ichikawa, T., Ivezić, Ž., Kent, S., Kim, R. S. J., Kinney, E., Klaene, M., Kleinman, A. N., Kleinman, S., Knapp, G. R., Korienek, J., Kron, R. G., Kunszt, P. Z., Lamb, D. Q., Lee, B., Leger, R. F., Limmongkol, S., Lindenmeyer, C., Long, D. C., Loomis, C., Loveday, J., Lucinio, R., Lupton, R. H., MacKinnon, B., Mannery, E. J., Mantsch, P. M., Margon, B., McGehee, P., McKay, T. A., Meiksin, A., Merelli, A., Monet, D. G., Munn, J. A., Narayanan, V. K., Nash, T., Neilsen, E., Neswold, R., Newberg, H. J., Nichol, R. C., Nicinski, T., Nonino, M., Okada, N., Okamura, S., Ostriker, J. P., Owen, R., Pauls, A. G., Peoples, J., Peterson, R. L., Petravick, D., Pier, J. R., Pope, A., Pordes, R., Prosapio, A., Rechenmacher, R., Quinn, T. R., Richards, G. T., Richmond, M. W., Rivetta, C. H., Rockosi, C. M., Ruthmansdorfer, K., Sandford, D., Schlegel, D. J., Schneider, D. P., Sekiguchi, M., Sergey, G., Shimasaku, K., Siegmund, W. A., Smee, S., Smith, J. A., Snedden, S., Stone, R., Stoughton, C., Strauss, M. A., Stubbs, C., SubbaRao, M., Szalay, A. S., Szapudi, I., Szokoly, G. P., Thakar, A. R., Tremonti, C., Tucker, D. L., Uomoto, A., Vanden Berk, D., Vogeley, M. S., Waddell, P., Wang, S.-i., Watanabe, M., Weinberg, D. H., Yanny, B., Yasuda, N., and SDSS Collaboration (2000). The Sloan Digital Sky Survey: Technical Summary. *AJ*, 120(3):1579–1587.
- Zacharias, N., Finch, C. T., Girard, T. M., Henden, A., Bartlett, J. L., Monet, D. G., and Zacharias, M. I. (2012). VizieR Online Data Catalog: UCAC4 Catalogue (Zacharias+, 2012). VizieR On-line Data Catalog: I/322A. Originally published in: 2012yCat.1322....0Z; 2013AJ....145...44Z.
- Zaritsky, D., Gil de Paz, A., and Bouquin, A. Y. K. (2015). The connection between the UV colour of early-type galaxies and the stellar initial mass function revisited. *MNRAS*, 446(2):2030–2037.

Appendix A. Additional Information

As discussed in Section 4.4, we have used a set of sample galaxies from the literature to observe their evolutionary phases in the HI gas fraction vs. NUV-r plane relation.




Figure A1. All available images of the three galaxies used in this study are shown here.




Figure A2. The UV (blue) and IR (red) contours obtained from UVIT and IRAC images are overlaid on the optical SDSS image of PRG NGC 3718. On the right side of the image, the top panel shows a zoomed-in view of a ring component with identified regions marked by blue ellipses. Below this, a zoomed-in view of the centre of the galaxy displays identified regions in green (overlap) and red (host).




Figure A3. The UV (blue) and IR (red) contours obtained from UVIT and IRAC images are overlaid on the optical SDSS image of PRG NGC 2685. The identified regions in green, blue and red represent overlap, ring and host regions, respectively.

Table A1. Properties of the selected subset of PRGs from the literature, including galaxy name, right ascension, declination, logarithmic neutral hydrogen mass, stellar mass, NUV, and r magnitudes. References are 1.Hall et al. (2012), 2.Parkash et al. (2018) 3.Gil de Paz et al. (2007), 4.Brown et al. (2014), 5.Seibert et al. (2012), 6.Zacharias et al. (2012), 7.Bouquin et al. (2018), 8.Tempel et al. (2017), 9.Bianchi et al. (2011), 10.Alam et al. (2015), 11.Gavazzi et al. (2013)

name	RA (deg)	Dec (deg)	logMHI (M_{\odot})	logM* (M_{\odot})	NUV (mag)	r (mag)
NGC660	25.760	13.645	9.41 ¹	9.82 ¹	14.80 ³	11.24 ⁴
NGC4650A	91.204	-40.714	10.05 ²	10.02 ²	16.48 ⁵	14.2 ⁶
NGC4324	185.775	5.250	9.28 ¹	10.75 ¹	15.78 ⁷	11.55 ⁸
UGC05791	159.865	47.947	8.43 ¹	8.56 ¹	16.00 ⁷	14.13 ⁸
NGC3656	170.911	53.842	9.35 ¹	10.75 ¹	16.50 ⁹	12.44 ¹⁰
UGC09763	228.009	21.298	9.96 ¹	10.54 ¹	18.80 ⁹	14.21 ¹⁰
UGC09002	211.223	12.721	9.75 ¹	9.58 ¹	17.05 ⁹	14.33 ¹⁰
UGC04385	125.966	14.751	9.22 ¹	9.43 ¹	15.71 ⁵	13.78 ¹⁰
PRCD-51	229.310	21.585	9.88 ¹	10.74 ¹	20.52 ⁹	15.38 ¹⁰
UGC4332	124.907	21.114	9.65 ¹	10.96 ¹	18.39 ⁹	12.96 ¹¹
IC51	11.600	-13.442	8.84 ²	9.44 ²	15.80 ⁹	12.84 ¹⁰