

Aims. The aim of this study was to identify pre-discharge risk factors associated with early inpatient readmission in general adult service users, with a particular focus on modifiable factors. We hypothesised that stability prior to discharge would reduce readmission to inpatient services within 90 days.

Background. Early readmission to inpatient psychiatric services is a poor outcome for service users, staff and the healthcare system. A variety of clinical, demographic and system factors, mostly non-modifiable, have been investigated previously. The identification of pre-discharge and particularly modifiable factors associated with readmission would give an opportunity for intervention and changes in policy.

Method. 272 medical records of all admissions within an 8 month period to a NHS inner city psychiatric inpatient service were reviewed to identify factors associated with readmission within 90 days of discharge. The data were analysed by simple comparison, calculation of odds ratios and logistic regression.

Result. 26% of service users were readmitted to the mental health trust within 90 days of discharge. Incidents (OR = 3.86; 95% CI 1.39–10.75) and psychotropic medication change in the week before discharge (OR = 2.94; 95% CI 1.43–6.03) were significantly associated with readmission, as were the number of previous admissions, and comorbid substance misuse. Successful overnight leave was found to be significantly protective against readmission (OR = 0.29; 95% CI 0.11–0.72).

Conclusion. The ability to predict those at high risk for readmission means they can be targeted for interventions and it can also help develop best practice around inpatient care and the discharge process. The novel findings in this study of pre-discharge modifiable risk factors such as stability and successful overnight leave could have significant implications in discharge planning policy.

Anticonvulsants and antipsychotics for treating anxiety and depressive symptoms in people with alcohol dependence: a systematic review

Ka Yee Chow*, Katy Jones and Musa Sami

University of Nottingham

*Corresponding author.

doi: 10.1192/bjo.2021.801

Aims. Anxiety and depressive symptoms are frequent in people with alcohol use disorders (AUDs) (approximately 55%) and are associated with worse outcomes. Current interventions to treat mood symptoms in alcohol-dependent adults, including psychosocial therapies, anxiolytics and antidepressants have shown limited benefits to date. The efficacy of anticonvulsants and antipsychotics have been studied but never previously systematically reviewed. We aimed to assess the efficacy of anticonvulsants and antipsychotics in treating anxiety and depressive symptoms in alcohol-dependent adults.

Method. A literature search of MEDLINE, EMBASE and PsycINFO was performed through October 2020 to identify all English-language articles of double-blinded randomised controlled trials that included adults with AUDs who were treated with an anticonvulsant or antipsychotic for at least four weeks. A combination of search terms was used to describe the AUDs, study medications and the primary outcome. Participants with other psychiatric disorders were excluded. Mean changes in anxiety and depressive scores were evaluated in addition to the adverse events and withdrawal rates. The risk of bias of each study was also assessed.

Result. Of 393 citations identified, 23 studies (2823 participants) met the inclusion criteria. Eighteen studies examined ten different anticonvulsants, while five studies examined two antipsychotics.

The heterogeneity between the trials' methodology led to conflicting results; however, as the low-quality trials were excluded, the majority agreed that anticonvulsants and antipsychotics were not superior in moderating anxiety and depressive symptoms of alcohol-dependent adults. All antipsychotics were safe and well-tolerated, but adverse events were associated with several anticonvulsants (high dose baclofen, gabapentin, topiramate and valproate).

Conclusion. Current information on anticonvulsants and antipsychotics is insufficient to extrapolate the benefits of treating anxiety and depressive symptoms in alcohol-dependent adults. Further research must be conducted into improving the quality of reporting and understanding the comorbidity's underlying mechanism, and alternative treatment approaches.

Attitudes of medical students towards psychiatry and mental illnesses: a cross-sectional study from Pakistan

Mishal Fatima³, Raja Adnan Ahmed², Muhammad Hamza¹, Syed Muhammad Jawad Zaidi^{1*}, Hassan Nadeem⁴ and Mehwish Kaneez¹

¹Rawalpindi Medical University; ²Consultant Psychiatrist, Cmw Taf Morhannwg University Health board; ³Final Year MBBS student, Rawalpindi Medical University and ⁴Jinnah Sindh Medical University, Dudley group NHS Foundation Trust

*Corresponding author.

doi: 10.1192/bjo.2021.802

Aims. The increasing burden of mental disorders coupled with the social stigmatization in Pakistan is an immense barrier in combating the emerging mental health crisis. The low number of qualified psychiatrists and poor intake in post-graduate psychiatry training programs in the region further complicates the problem. Thus, our study aims to assess the attitudes of Pakistani medical students towards psychiatry. Furthermore, we also aim to evaluate how experience and different levels of exposure to psychiatry among students affect their attitudes towards psychiatry as a career choice.

Method. This cross-sectional study was conducted via an online survey made on Google Forms. A total of 831 medical students studying across various private and public medical institutions of Pakistan responded to the survey. The questionnaire comprised of demographic details (gender, age, institution, and academic year) exposure to psychiatry, duration of psychiatry rotation, and personal experience with mental illness. The attitudes of medical students towards psychiatry were evaluated using the English version of the 30-item Attitudes Towards Psychiatry (ATP-30) scale. Chi-square test and multiple regression with backward method were used to analyze the data.

Result. The Cronbach's alpha value of the ATP-30 scale was 0.830. The participants in our study had a mean score of 107.6 ± 12 on ATP-30. Overall, most participants had a positive attitude towards psychiatry. Multiple regression analysis revealed a significant model pertaining to predictors of attitude toward psychiatry (F (df) = 11.28 (830), P < 0.001). However, the predictors included in the model accounted for only 5.8% of the variation in ATP-30 scores. According to it, those students had a more positive attitude toward psychiatry who identified as female, older and having any sort of exposure toward psychiatric specialty, direct involvement in psychiatric patient care, and reporting personal experience of mental illnesses.

Conclusion. Our study showed that medical students had a positive attitude towards psychiatry but female medical students, students with previous exposure to psychiatry, and students with longer psychiatry rotations tend to view psychiatry more positively. The generally positive trend towards psychiatry in Pakistan indicates the need to sustain improvements through

proactive measures. We recommend longer placements for medical students in mental health settings for at least 4 weeks or longer. Medical schools should also promote research, discussions, and seminars on different psychiatric illnesses in order to enhance awareness among the students.

Psychiatric liaison referrals and the 4PM Rush

Kaj Svedberg¹

¹Rawalpindi Medical University

doi: 10.1192/bjo.2021.956

Aims. Referrals to the psychiatric Liaison team in A&E seem to come in the afternoon in kismet as the day shift is ending. This study looked at the timing distribution of referrals to try improve amount of jobs being handed over to the evening shifts.

Method. Referrals made to Homerton University Hospital (HUH) psychiatric liaison was parsed into 1 hour bins and plotted as a histogram (data between August 2016–October 2019. N = 14182). The data were compared to diurnal human body temperature variation, as well as data published on Hospital Accident & Emergency Activity 2019–20 (digital.nhs.uk) for Ambulance attendances.

Result. Referrals to HUH liaison team appear to closely follow the average human body temperature variations per hour (Pearson Correlation coefficient = 0.90). A peak appears to occur around 4 PM, and a low at 7AM. The referrals data also mirrored timings of official Hospital Episode Statistics (HES) reports 2019–2020 for ambulance attendance in England (Pearson Correlation coefficient = 0.94).

Conclusion. Attendance to A&E and referrals to psychiatric liaison appear correlated to a circadian bound rhythm. “The 4PM referrals rush” appears to be a genuine phenomenon replicated in not only HUH mental health referrals, but general ambulance attendance throughout all of England.

The body temperature analogue for circadian rhythm may be humorously applied here to correlate with the increased referral rates to A&E; the emergency department could be said to be truly heating up in the afternoon. Indeed temperature and activity has already been shown to link strongly via the Arrhenius equation in cricket activity such as chirps per minute.

The conclusions drawn here are that acute mental health attendances, like general health attendances as a whole follow underlying but powerful patterns, and provisions might best be allocated to address this rather than thinking of fixed 9–5 working schedules.

Service Evaluation

Do core psychiatry trainee cognitive behaviour therapy cases meet training needs? An evaluation of core psychiatry trainee delivered CBT cases in Sheffield: implications for training and services

Dasal Abayaratne^{1*}, Russell Birkett² and John Davies¹

¹Sheffield Health and Social Care NHS Foundation Trust and

²University of Sheffield, Sheffield Health and Social Care NHS Foundation Trust

*Corresponding author.

doi: 10.1192/bjo.2021.804

Aims. This evaluation aims to understand if Cognitive Behaviour Therapy (CBT) cases for Core Psychiatry Trainees (CPTs) in Sheffield provide good training in therapy skills and if these can be integrated into general psychiatric practice.

Background. Completion of psychotherapy cases part of the curriculum for CPTs, with cognitive behavior therapy being one of the common modalities used. Whilst there is evidence that trainees often provide competent therapy it is unclear what cases are appropriate and how these contribute to wider CPT learning objectives. **Method.** CPTs who had completed a clinical case in CBT at a tertiary psychotherapy service were identified. All were surveyed and patient demographics and outcomes also collated.

Result. The results showed a significant impact on trainees understanding of CBT, applying theory to clinical context, and changed future practice. Despite being complex, 64% of patients needed no further therapy and 42% were discharged from mental health services.

Conclusion. The evaluation demonstrates the positive outcomes for patients, trainees, future clinical practice, and a move towards collaboration as laid out in the Five-year forward view for mental health. This suggests that medical trainees have a valuable contribution, and role despite minimal experience in CBT.

Are we meeting local and national guidelines for physical health assessment following admission to the meadows?

Yasmin Abbasi*, Ruairidh Morgan and Alice O'Docherty

Surrey and Borders Partnership NHS Foundation Trust

*Corresponding author.

doi: 10.1192/bjo.2021.805

Aims. We audited practice at the Meadows Inpatient Unit regarding physical health assessment, against standards set by Surrey and Borders Partnership and NICE.

Background. SABP policy states that within 24 hours of admission to inpatient services, physical health assessment should be offered. It should be completed within 72 hours. Refusal should be documented.

These guidelines state that within 2 weeks of admission blood tests should be completed, and for specific individuals an ECG should be performed.

NICE guidelines state that “physical healthcare needs” should be discussed with newly admitted patients. NICE guidelines regarding physical health monitoring for individuals with psychosis or schizophrenia recommend that assessment includes “full physical examination to identify physical illness”.

NICE suggests use of antipsychotics for individuals with dementia who have severe distress, or are at risk of harming themselves or others. Those with behavioural and psychological symptoms of dementia (BPSD) should therefore be physically assessed to ensure safe use of antipsychotics may be implemented.

Method. All admissions to The Meadows over seven months were audited retrospectively. The clinical notes were accessed from Systm1.

Consensus medical opinion was reached that full examination should include: GCS/level of consciousness, cardiorespiratory, abdominal and neurological examinations.

Age, gender, diagnosis and prescriptions of psychotropic medication at time of admission were recorded.

The sample included 35 patients.

Result. 55% of patients had a diagnosis of dementia.

63.8% of patients were prescribed antipsychotics on admission, more than other psychotropic medication. This may reflect that the most common diagnosis was dementia, commonly with associated BPSD.

97% of patients had a physical examination completed within 24 hours; most excluded neurological examination. 91% of patients had blood tests completed in two weeks, with the most commonly excluded tests being lipids and glucose. 86% of