## INSTRUCTIONS TO CONTRIBUTORS

The Journal of Helminthology publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance.

Manuscripts, which must be in English or French (with an English summary), should be addressed to:

The Editor, Journal of Helminthology London School of Hygiene and Tropical Medicine Keppel Street, London WC1E 7HT England

Two copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief introduction (unheaded); Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration may be printed is  $13.5 \times 18$  cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space should be indicated. Some indication of scale should normally be given on the figure. Photocopies of illustrations should also be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints of the same size as they are to appear in the Journal (maximum size  $13.5 \times 18$  cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

- DUKE, B. O. L. (1971) The ecology of onchocerciasis in man and animals. In *Ecology* and physiology of parasites (ed A. M. Fallis) pp. 213-222. Adam Hilger Ltd.: London.
- JAMES, C. and WEBBE, G. (1973) A comparison of Egyptian and East African strains of Schistosoma haematobium. Journal of Helminthology, 47, 49-59.

50 offprints are provided free of charge; additional copies may be ordered at the proof stage.

## **Contents**

	Pages
HOLMES, P. H., JAMES, E., MACLEAN, J. M., NELSON, G. S. and TAYLOR, M. G. Erythrokinetic studies on baboons ( <i>Papio anubis</i> ) with acute experimental schistosomiasis mansoni	95—104
CHRISTENSEN, N. O., FRANDSEN, F. and NANSEN, P. Host-finding capacity of schistosome cercariae: comparative efficiency of methods of mouse infection and a radioisotope assay system	105—113
BYLUND, G., BANG, B. and WIKGREN, K. Tests with a new compound (Praziquantel) against <i>Diphyloboth-rium latum</i>	115—119
PALMIERI, J. R. and SULLIVAN, J. T. Stunkardia minuta sp. n. (Trematoda: Paramphistomidae) from the Malayan box-tortoise Cuora amboinensis	121—124
FUJINO, T., HAMAJIMA, F., ISHI, Y. and MORI, R. Development of <i>Microphalloides japonicus</i> (Osborn, 1919) metacercariae <i>in vitro</i> (Trematoda: Microphallidae)	125—129
GOULART, E. G., JOURDAN, M. C., BRAZIL, R. P., BRAZIL, B. G., COSENDEY, A. E., BAR, M., DO CARMO, E. C. and GILBERT, B. Research Note. Ecological control of hookworm and strongyloidiasis	131—132
SUSWILLO, R. R., NELSON, G. S., MULLER, R., McGREEVY, P. B., DUKE, B. O. L. and DENHAM, D. A. Research Note. Attempts to infect jirds ( <i>Meriones unguiculatus</i> ) with <i>Wuchereria bancrofti, Onchocerca volvulus, Loa loa loa and Mansonella ozzardi</i>	132—134
VARMA, S., MALIK, P. D. and LAL, S. S. Research Note. Some new intermediate hosts of <i>Ascarops strongylina</i>	134—135
CHERNIN, J. Aspects of the humoral response of laboratory white mice infected with the metacestodes of Taenia crassiceps	137—142
ENAYAT, M. S. and PEZESHKI, M. The comparison of counterimmuno-electrophoresis with indirect haemagglutination test for detection of antibodies in experimentally infected guinea pigs with <i>Toxocara canis</i>	143—148
STEVENSON, P. and JACOBS, D. E. <i>Toxocara</i> infection in pigs. The use of indirect fluorescent antibody tests and an <i>in vitro</i> larval precipitate test for detecting specific antibodies.	149—154
BLAIR, D. A key to cercariae of British strigeoids (Digenea) for which the life-cycles are known, and notes on the characters used	155—166