

CORRIGENDUM

Towards a finite-time singularity of the Navier–Stokes equations. Part 2. Vortex reconnection and singularity evasion – CORRIGENDUM

H. K. Moffatt and Yoshifumi Kimura

doi:10.1017/jfm.2019.263, Published online by Cambridge University Press,
7 May 2019

Results of numerical integration (using Mathematica) of the following dynamical system were presented in Moffatt & Kimura (2019*b*, hereafter MK19*b*):

$$\frac{ds}{d\tau} = -\frac{\gamma \kappa \cos \alpha}{4\pi} \left[\log \left(\frac{s}{\delta} \right) + 0.4417 \right], \quad (1)$$

$$\frac{d\kappa}{d\tau} = \frac{\gamma \kappa \cos \alpha \sin \alpha}{4\pi s^2}, \quad (2)$$

$$\frac{d\delta^2}{d\tau} = \epsilon - \frac{\gamma \kappa \cos \alpha}{4\pi s} \delta^2, \quad (3)$$

$$\frac{d\gamma}{d\tau} = -\epsilon \frac{s \gamma}{2\sqrt{\pi} \delta^3} \exp[-s^2/4\delta^2], \quad (4)$$

with $\alpha = \pi/4$. Unfortunately, the exponential factor $\exp[-s^2/4\delta^2]$ in (4) was consistently misrepresented as $\exp[-s^2\delta^2/4]$ in the Mathematica code, leading to errors in the numerical results and in figures 4–8, although the conclusions concerning the approach to a ‘near singularity’ of the system (1)–(4) remain unchanged.

Figure 4 of MK19*b* should be replaced by figure 1 below, and figure 5 by figure 2, which exhibit the same qualitative behaviour in each case. The difference, however, is that the spike of vorticity now occurs for significantly larger ϵ (1/45 in figure 1, and 1/200 in figure 2), and at smaller ‘critical time’ t_c (0.7773 in figure 1, and 0.3574 in figure 2). (When $\epsilon = 0$, with the same initial conditions, the singularity occurs at time $t_c \approx 0.2434$; when $\epsilon = 1/4000$, it occurs at the slightly greater time $t_c \approx 0.2522$. As ϵ increases, t_c continues to increase.)

Figures 6–8 and the detailed figures of tables 1 and 2 of MK19*b* are similarly incorrect. Indeed, when $\epsilon = 1/4000$, the minimum value of δ^2 ($\lesssim 10^{-42}$) and the maximum value of $\omega(\tau)/\omega(0)$ can no longer be accurately determined.

The slanted vortex-ring configuration considered by MK19*b* has been investigated by direct numerical simulation (DNS) by Yao & Hussain (2020), and it has become apparent through this work that the model of MK19*b* fails to represent the true nature of the reconnection process. This model assumes that the vortex cores remain compact during reconnection, and that the reconnected vortex strands have negligible




FIGURE 1. (Replacing figure 4 of MK19b) Evolution governed by (1.1)–(1.4), with $\epsilon = 1/45$ and initial conditions $s(0) = 0.1$, $\delta(0) = 0.01$, $\kappa(0) = \gamma(0) = 1$.




FIGURE 2. (Replacing figure 5 of MK19b) Same as figure 1, but with $\epsilon = 1/200$.

effect on the intensification of vorticity in the incident vortices. The DNS indicates that these assumptions are untenable. Nevertheless the MK19b model captures key elements of a possible approach to a singularity, and it is for this reason that the dynamical system (1)–(4) merits critical investigation.

Acknowledgements

We thank F. Hussain and J. Yao, who drew our attention to the incompatibility of our model with the results of their DNS during the reconnection stage, thus leading to identification of the above error.

REFERENCES

- MOFFATT, H. K. & KIMURA, Y. 2019*b* Towards a finite-time singularity of the Navier–Stokes equations. Part 2. Vortex reconnection and singularity evasion. *J. Fluid Mech.* **870**, R1.
 YAO, J. & HUSSAIN, F. 2020 On singularity formation via viscous vortex reconnection. *J. Fluid Mech.* (submitted).