

evidence either for or against the suggestion that ECT may cause permanent cerebral damage.

RAYMOND LEVY

*The Maudsley Hospital,
Denmark Hill, London SE5 8AZ*

ROBIN JACOBY

*Middlesex Hospital,
London W1N 8AA*

References

- CALLOWAY, S. P., DOLAN, R. J., JACOBY, R. J. & LEVY, R. (1982) ECT and cerebral atrophy; A computed tomographic study. *Acta Psychiatrica Scandinavica*, (in press).
- JACOBY, R. J. & LEVY, R. (1980) Computed tomography in the elderly. III. Affective disorder. *British Journal of Psychiatry*, **136**, 270–6.

MAOI FOR OBSESSIVE COMPULSIVE DISORDER

DEAR SIR,

Obsessive compulsive disorder is a relatively uncommon illness, which in severe form is remarkably destructive to the patient and his or her family. The natural history of the disorder is poorly understood.

There are three papers in the literature (Swinson and Thomas, 1970; Annesley, 1969; Jenike, 1981) which present four cases that responded to monoamine oxidase inhibitors. Over the past few years we have had another four cases at our institution where monoamine oxidase inhibitors produced a rapid and sustained remission of symptoms. In all of these cases, numerous prior treatments had been unsuccessful in alleviating the symptoms.

In two of our cases, relapses occurred after the monoamine oxidase inhibitor was stopped and in both cases restarting the drug resulted in loss of symptomatology. All of our patients that responded to monoamine oxidase inhibitors remained free of symptoms at follow-up, which ranged from four months to four years. In the cases previously reported and in our four cases which responded to monoamine oxidase inhibitors, the patients all had phobic anxiety and/or severe anxiety associated with the obsessive compulsive disorder. At our institution, four other cases of severe obsessive compulsive disorder without associated anxiety or panic attacks, did not respond to monoamine oxidase inhibitors.

Our data indicate that at least a subgroup of patients with obsessive compulsive disorder respond to monoamine oxidase inhibitors, sometimes dramatically. Affective illness in the patient or his family was not a good predictor of responsiveness to monoamine oxidase inhibitors in our patients. The presence of severe anxiety or panic attacks, however, was uniformly associated with a good response in our

patients. Sheehan, Ballinger and Jacobson (1980) have shown that phenelzine is effective in treating panic attacks and lowers the obsessive compulsive scores on the SCL-90 scale.

The authors feel that a trial of monoamine oxidase inhibitors is presently indicated in obsessive compulsive disorder, especially when phobic anxiety or panic attacks are part of the clinical presentation.

MICHAEL A. JENIKE

*Massachusetts General Hospital,
Fruit Street, Boston, USA*

References

- JAIN, V. K., SWINSON, R. P. & THOMAS, J. G. (1970) Phenelzine in obsessive neurosis. *British Journal of Psychiatry*, **117**, 237–8.
- ANNESLEY, P. T. (1969) Nardil response in a chronic obsessive compulsive. *British Journal of Psychiatry*, **115**, 748.
- JENIKE, M. A. (1981) Rapid response of severe obsessive compulsive disorder to tranlycypromine. *American Journal of Psychiatry*, **138**, 9.
- SHEEHAN, D. V., BALLINGER, J. & JACOBSON, G. (1980) Treatment of endogenous anxiety with phobic, hysterical, and hypochondriacal symptoms. *Archives of General Psychiatry*, **37**, 51–7.

HYPOALGESIA IN DEPRESSIVE ILLNESS

DEAR SIR,

Dr Hanks (*Journal*, October 1981, 364–5) considers my comments on the analgesic properties of tricyclic antidepressant drugs to be misleading (*Journal*, January 1981, 37–9). They certainly could be but a good deal hinges on the meaning to be attached to the word 'depression' in this context. To be sure, patients experiencing chronic pain can be miserable, unhappy and, on occasions, suicidal and despairing. But such emotions are not necessarily tantamount to a *syndrome* of depression as defined, for example, by Feighner *et al* (1972). In my experience tricyclic antidepressant drugs are rarely successful in relieving 'reactive' depression when experienced as unhappiness in the face of personal and environmental difficulties.

The possibility that some antidepressants have specific analgesic properties apart from their antidepressant ones is suggested by the three following observations:

- (1) Intractable pain is sometimes relieved by quite small doses of anti-depressants, doses far lower than those generally required for the successful treatment of endogenous depression.
- (2) Relief of chronic pain is sometimes achieved after quite short periods of treatment—48 hours in some cases—which are far shorter than the time required for remission of a depressive syndrome (Gade *et al*, 1980; Turkington, 1980).