

Methods: We recorded resting-state EEG in fourteen participants with high Beck Depression Inventory score (24.4 ± 9.7 ; 20.4 ± 1.5 y.o.; 14 females; 1 left-handed) and fourteen participants with a low score (6.8 ± 3.7 ; 21.3 ± 2.0 y.o.; 8 females; 1 left-handed). We applied weighted phase-lag index (wPLI) to construct functional networks at sensors and sources levels and computed characteristic path length (CPL), clustering coefficient (CC), index of modularity (Q), small-world index (SWI) in 4-8, 8-13, 13-30, and 4-30 Hz frequency bands. We used Mann-Whitney U-test ($p < 0.05$) to investigate between-group differences in the graph metrics.

Results: The depressive sample was characterized by increased CC and Q in the 4-30 Hz band networks and decreased CPL in the beta-band network (sensors-level for CPL and CC, and sources-level for Q).

Conclusions: Elevated CC and Q may relate to an increase of intramodular connectivity, and CPL reduction reflects the global connectivity increasing. We hypothesize that intramodular hyperconnectivity could explain the rise of global functional connectivity in participants with depressive symptoms. *Funding:* This research has been supported by the Interdisciplinary Scientific and Educational School of Lomonosov Moscow State University 'Brain, Cognitive Systems, Artificial Intelligence'.

Disclosure: No significant relationships.

Keywords: Depression; Resting-state EEG; Connectivity

EPP0265

Schizotypal Traits are Associated with Decreased Functional Connectivity between Medial Prefrontal Cortex and Cerebellum in a Non-clinical Sample

Y. Panikratova¹, E. Abdullina^{1*}, E. Pechenkova² and I. Lebedeva¹

¹Mental Health Research Center, Laboratory Of Neuroimaging And Multimodal Analysis, Moscow, Russian Federation and ²Higher School of Economics, Laboratory For Cognitive Research, Moscow, Russian Federation

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.558

Introduction: Schizotypy is associated with increased vulnerability to schizophrenia spectrum disorders. Therefore, investigation of its brain correlates seems prominent for better understanding of schizophrenia-spectrum continuum as well as for development of biological treatments for schizotypal personality disorder. Functional alterations of prefrontal cortex (PFC) and their associations with clinical symptoms are well-known to exist in schizophrenia. However, their relevance to schizotypy remains unclear.

Objectives: The aim of the study was to check for associations between schizotypal traits in a non-clinical sample and whole-brain functional connectivity (FC) of lateral as well as medial PFC (IPFC and mPFC, respectively).

Methods: Eighty-two healthy individuals (52 females, mean age 24.8 ± 5.48) filled out the Schizotypal Personality Questionnaire (SPQ-74) and underwent resting-state fMRI (3T). Seeds in IPFC and mPFC were taken from frontoparietal and default mode networks (atlas by Yeo et al., 2011). We analyzed correlations between four schizotypal factors (cognitive/perceptual, paranoid,

negative, and disorganization; Stefanis et al., 2004) and whole-brain FC of the seeds (statistical threshold: $p < .001$ voxelwise; $p [FDR] < .05$ clusterwise).

Results: Cognitive/perceptual factor ('Odd beliefs/magical thinking' and 'Unusual perceptual experiences' SPQ-74 subscales) is negatively correlated to FC of bilateral mPFC with a cluster in the right cerebellum (Crus 1, 2).

Conclusions: Prefrontal-cerebellar dysconnectivity may be one of the neurobiological factors underlying positive-symptoms-like schizotypal traits in non-clinical subjects. To some extent, it coincides with the data on associations between functional features of these brain structures and positive symptoms in schizophrenia (Pinheiro et al., 2021; Goghari et al., 2010).

Disclosure: The study was supported by RFBR Grant 20-013-00748.

Keywords: resting-state fMRI; schizotypy; SPQ-74

EPP0266

Organic lesions and Psychiatry: "A sample on a pendant"

T. Jiménez Aparicio¹, G. Medina Ojeda^{2*}, C. De Andrés Lobo³, C. Vallecillo Adame¹, J. Gonçalves Cerejeira⁴, I. Santos Carrasco⁵, G. Guerra Valera⁴, M. Queipo De Llano De La Viuda⁴, A. Gonzaga Ramírez¹, M. Fernández Lozano¹, B. Rodríguez Rodríguez¹, M.J. Mateos Sexmero¹, N. Navarro Barriga³ and N. De Uribe Viloria⁶

¹Hospital Clínico Universitario, Psiquiatría, Valladolid, Spain; ²Sacyl, Hospital Clínico Universitario Valladolid, Psiquiatría, Valladolid, Spain; ³Hospital Clínico Universitario de Valladolid, Psiquiatría, VALLADOLID, Spain; ⁴Hospital Clínico Universitario de Valladolid, Psiquiatría, Valladolid, Spain; ⁵Clinical Hospital of Valladolid, Psiquiatría, Valladolid, Spain and ⁶Hospital Universitario Fundación de Alcorcón, Psiquiatría, Madrid, Spain

*Corresponding author.


doi: 10.1192/j.eurpsy.2022.559

Introduction: Brain lesions may induce psychiatric symptoms in some cases. Imaging tests are important to make a differential diagnosis, and therefore initiate an appropriate treatment.

Objectives: Presentation of a clinical case about a patient with psychiatric symptoms who presented an organic lesion.

Methods: Bibliographic review including the latest articles in Pubmed about psychiatric symptoms induced by organic lesions.

Results: We present a 51-year-old male patient, with adequate previous functionality, who attended psychiatric consultations due to changes in his character, with delusional mystical and megalomaniac ideation, verbiage, hypoprosexia, memory loss and insomnia (diagnosed with Bipolar Disorder type II, hypomanic episode). Eventually, a brain computed tomography scan was performed, in which meningioma was visualized. The patient underwent surgery, and he asked to keep a sample of his tumor to always carry it with him on a pendant. Psychiatric symptoms induced by organic lesions are highly variable, depending on the location and size of the lesion, and they may be the first and/or only symptom of a meningioma (up to 21% according to various studies), so it is important to perform imaging tests in some cases. At this time, the patient is under follow-up, he has remained euthymic and stable, and he refuses to take psychopharmacological medication.



Conclusions: Psychiatric symptoms may be the first and/or only manifestation of an organic lesion in some cases. Neuroimaging tests (CT and MR) may be useful in the differential diagnosis. It is important to carry out an individualized treatment based on the patient's pathology, which may include surgery and/or drugs.

Disclosure: No significant relationships.

Keywords: Neuroimaging; bipolar disorder; Meningioma; Neurosurgery

EPP0268

Depression among Endometrial Cancer hospitalizations - Preliminary results of a nationwide retrospective study

P. Vieito^{1*}, A.R. Ferreira^{1,2,3}, M. Gonçalves-Pinho^{2,4,5}, F. Costa⁶, M. Coelho⁶, A. Freitas^{2,5} and L. Fernandes^{2,3,7}

¹University of Porto, Faculty Of Medicine, Porto, Portugal; ²CINTESIS - Center for Health Technology and Services, Faculty of Medicine, University of Porto, Department Of Community Medicine, Information And Health Decisions Sciences, Porto, Portugal; ³Faculty of Medicine, University of Porto, Department Of Clinical Neurosciences And Mental Health, Porto, Portugal; ⁴Centro Hospitalar do Tâmega e Sousa, Departamento De Psiquiatria E Saúde Mental, Penafiel, Portugal; ⁵Faculty of Medicine, University of Porto, Department Of Community Medicine, Information And Health Decision Sciences (medcids), Porto, Portugal; ⁶Centro Hospitalar do Tâmega e Sousa, Departamento De Ginecologia E Obstetrícia, Penafiel, Portugal and ⁷Psychiatry Service, Centro Hospitalar Universitário De São João, Porto, Portugal

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.560

Introduction: Uterine cancer is the most common gynecologic malignant neoplasm in developed countries. While depression is up to 3-5 times more common in patients with cancer than in the

general population, literature is still limited regarding the relation between Endometrial Cancer and depression.

Objectives: To analyze Depression among Endometrial Cancer hospitalizations in mainland Portuguese public hospitals (2008-2015).

Methods: A retrospective observational study was conducted using administrative data from all hospitalizations in Portuguese mainland public hospitals between 2008-2015. All women's hospitalizations (≥ 18 years) with a primary diagnosis of Endometrial Cancer (ICD-9-CM 182.x) were selected. Secondary diagnosis of depression was identified with ICD-9-CM 296.2x, 296.3x and 311x codes. Surgical procedures codes 68.4x, 65.6x, 40.3x, 40.5x, 68.6x, 68.9x and 68.8x were used to divide the hospitalizations into surgical vs non-surgical. Groups were compared with Pearson Chi-square test and crude odds ratio(OR) was used to estimate the association between surgery and depression.

Results: From 10227 hospitalizations with a primary diagnosis of Endometrial Cancer, 533 had a registry of depression(5.2%). Annual depression frequency rose from 2.0% (2008) to 8.3% (2015). Among patients with a record of depression, 73.2% had surgery. Women who had surgery were significantly more likely to have registered depression ($p < 0.001$). The OR for depression in surgical vs non-surgical patients was 1.73 (95%IC:1.42-2.10).

Conclusions: Patients hospitalized due to Endometrial Cancer and submitted to surgery had almost two-fold more risk of having a registry of depression. This trend reinforces the importance of early depression screening of these patients, enabling the implementation of timely therapeutic strategies before and after surgery.

Disclosure: No significant relationships.

Keywords: Endometrial Cancer; surgery; Depression; Administrative data

EPP0269

Barriers to access cancer screening and treatment services in Germany

F. Baessler* and A. Zafar

Centre for Psychosocial Medicine, Department Of General Internal And Psychosomatic Medicine, Heidelberg, Germany

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.561

Introduction: individual attitudes and structural inadequacies act as major barriers towards non-utilization of cancer screening and treatment offers in many high-income countries with subsidized public healthcare.

Objectives: Our interdisciplinary research group at Heidelberg University is studying the underlying individual perceptions, attitudes and experiences of age- and gender-specific barriers against cancer-related medical and psychosocial offers available in Germany.

Methods: We designed a mixed-methods, sequential explanatory study using two quantitative instruments to determine the most important age- and gender-specific barriers for non-patients and cancer patients and survivors. In the second phase, semi-structured interviews will be conducted via selective sampling to record participant opinions, experiences and expectations of using cancer-related health services.