


# Detection of echinostomatid trematode eggs at the forest–oil palm interface in Sabah, Malaysia

Liesbeth Frias<sup>1,2</sup> , Benny Obrain Manin<sup>3</sup>, Sergio Guerrero-Sánchez<sup>4</sup>, Symphorosa Sipangku<sup>5</sup> and Tock H. Chua<sup>3</sup>

## Research Article

\*Current affiliation.

**Cite this article:** Frias L, Manin BO, Guerrero-Sánchez S, Sipangku S, Chua TH (2024). Detection of echinostomatid trematode eggs at the forest–oil palm interface in Sabah, Malaysia. *Parasitology* **151**, 181–184. <https://doi.org/10.1017/S0031182023001257>

Received: 17 October 2023  
Revised: 26 November 2023  
Accepted: 27 November 2023  
First published online: 5 December 2023

### Keywords:

flukes; food-borne trematodes; land-use change; tropical ecosystems; wildlife–human interfaces; zoonotic parasites

### Corresponding author:

Liesbeth Frias;  
Email: [lfrias@duke-nus.edu.sg](mailto:lfrias@duke-nus.edu.sg)

<sup>1</sup>Programme in Emerging Infectious Diseases, Duke-NUS Medical School, Singapore; <sup>2</sup>Asian School of the Environment, Nanyang Technological University, Singapore; <sup>3</sup>Faculty of Medicine and Health Sciences, Universiti Malaysia Sabah, Kota Kinabalu, Sabah, Malaysia; <sup>4</sup>Centre for Applied One Health Research and Policy Advice, Jockey Club College of Veterinary Medicine and Life Sciences, City University of Hong Kong, Kowloon, Hong Kong and <sup>5</sup>Sabah Wildlife Department, Kota Kinabalu, Sabah, Malaysia

### Abstract

In this study, we report the occurrence of echinostomatid eggs in feces of wildlife, domestic animals and humans frequenting the forest–oil palm plantation interface in the Kinabatangan (Sabah, Malaysia), and discuss potential implications for public health. Using microscopy, we detected echinostomatid eggs in six host species, including Asian palm civets (*Paradoxurus hermaphroditus* [13/18]), leopard cats (*Prionailurus bengalensis* [3/4]), long-tailed macaques (*Macaca fascicularis* [1/10]), domestic dogs [3/5] and cats [1/1], and humans [7/9]. Molecular analysis revealed a close genetic proximity of civet echinostomatids to *Artyfechinostomum malayanum*, a zoonotic parasite of public health relevance. The intermediate hosts for *A. malayanum* have been reported in at least 3 districts in Sabah, suggesting that all the necessary elements required for the completion of the parasite's life cycle are present. Our findings point at the presence of zoonotic trematodes in an area with high human–wildlife interaction and highlight the potential public and animal health concern of zoonotic trematode infection in the context of Southeast Asia's rapidly changing ecosystems.

## Introduction

Trematodiasis are an important group of neglected tropical diseases caused by digenetic trematodes, which include the intestinal, liver, lung and blood flukes (Saijuntha *et al.*, 2021). Intestinal trematodes and their hosts are highly diverse in Southeast Asia, where rapid and extensive changes in land use and land conversion can intensify the interactions among wildlife, domestic animals and humans, thereby creating new opportunities for parasite transmission (Jones *et al.*, 2013). Over the last 7 decades, the Kinabatangan floodplain, an important biodiversity hotspot in Southeast Asia, has faced significant forest loss due to extensive commercial logging and industrial oil palm (Abram *et al.*, 2014). Intensive human activity in the area (e.g. oil palm agriculture, ecotourism, low-scale fisheries), and high occurrence of wild and of domestic animals, can increase the likelihood of zoonotic pathogen transmission, particularly in areas frequented by infected livestock or where open defecation occurs (Fried *et al.*, 2004). With large-scale conversion of forests to oil palm plantations, the dynamics of zoonotic diseases, including lesser-known parasites like echinostomatids, are inevitably being modified. Consequently, this study was undertaken to evaluate the presence of echinostomatid eggs in the feces of wildlife inhabiting the forest–oil palm plantation interface. The implications for potential public health concern in these human-dominated landscapes are also examined and discussed.

## Materials and methods

Non-invasive surveys were carried out along 2 km linear transects running along the boundaries of each one of the 3 oil palm estates and the natural forest. Each transect was visited 3 consecutive days per month between April and July 2019, where fresh feces from wild and free-roaming domestic species were collected. A modified formalin–ethyl acetate sedimentation protocol was used to inspect samples for the presence of trematode eggs (Frias *et al.*, 2021). Parasite prevalence, or the proportion of the host population infected with at least one trematode egg, was measured and the eggs per gram of feces (EPG) was calculated by taking the average of 5 replicate counts of all observed trematode eggs under a McMaster chamber's grid. Host species identification was conducted by amplifying a small fragment of the *cytochrome b* (*cytb*) gene using the L14724/H15915 primer pair (Irwin *et al.*, 1991). Samples positive for trematodes through microscopy underwent DNA sequencing of both the nuclear ribosomal 18S gene and the mitochondrial cytochrome oxidase I (COI) gene. Amplification and sequencing of these genes were performed using 18S gene specific primers (18S9modF, 18S637modR) and COI specific primers (Dice1F, Dice11R and Dice14R) following established PCR protocols (Moszczyńska *et al.*, 2009; Van Steenkiste *et al.*, 2015).

© The Author(s), 2023. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided that no alterations are made and the original article is properly cited. The written permission of Cambridge University Press must be obtained prior to any commercial use and/or adaptation of the article.

**Table 1.** Prevalence and abundance of trematode eggs found in feces of wildlife, domestic animals and humans, as detected through microscopy

Host species	Prevalence (%)	EPG (mean $\pm$ s.d.)
Asian palm civet ( <i>Paradoxurus hermaphroditus</i> )	72.2 (13/18)	28–616 (123.4 $\pm$ 177.3)
Leopard cat ( <i>Prionailurus bengalensis</i> )	75 (3/4)	28–56 (46.3 $\pm$ 15.9)
Asian otter ( <i>Lutra sumatrana</i> )	0 (0/2)	0
Domestic dog ( <i>Canis lupus familiaris</i> )	60 (3/5)	55–21 252 (7126.7 $\pm$ 12 232.9)
Domestic cat ( <i>Felis catus</i> )	100 (1/1)	280
Long-tailed macaque ( <i>Macaca fascicularis</i> )	10 (1/10)	84.0
Human	77.7 (7/9)	14.7–396 (131 $\pm$ 171)


## Results

A total of 49 feces from 3 different forest–oil palm interface areas were collected. Trematode eggs were detected through microscopy in 28 samples (57.14%) from Asian palm civets (*Paradoxurus hermaphroditus*), leopard cats (*Prionailurus bengalensis*), domestic dogs (*Canis lupus familiaris*), domestic cats (*Felis catus*), long-tailed macaques (*Macaca fascicularis*) and humans (Table 1). Echinostomatid eggs detected in feces were ovoid to ellipsoid with a pronounced operculum, and 135 (129–145)  $\mu\text{m} \times 75$

(70–82)  $\mu\text{m}$  ( $n = 6$ ) in size (Fig. S1), compatible with size and shape descriptions of human-infecting echinostomes (Esteban *et al.*, 2019). Parasite DNA was only successfully amplified for COI in 6 samples from civets (GenBank accession numbers LC773275–LC773280), displaying a close genetic similarity of 97.05%–97.64% with parasites of the genus *Artyfechinostomum* spp. (Echinostomatidae) in the NCBI database, and not previously reported in Sabah (Fig. 1).

## Discussion

Despite being one of the fastest-growing economies in Southeast Asia, parasitic infections are still widespread in Malaysia. Rural and agricultural areas in Sabah are often remote, with limited access to healthcare, drinking water and poor environmental sanitation, and where the presence of intermediate and reservoir hosts can perpetuate parasite transmission (Zin *et al.*, 2015). In this study, leopard cats and Asian palm civets exhibited the highest parasite prevalence. Leopard cats and civets are not only native to most of Asia but are also more likely to frequent human settings, providing an insight into their potential as reservoir hosts for various pathogens (Guerrero-Sánchez *et al.*, 2022). Free-ranging cats are not frequently found in the studied interface areas; however, plantation workers do own domestic cats and dogs that are often allowed to roam freely, as observed during our surveys. Despite their smaller population, dogs are a constant presence in these areas, as they play an important role in accompanying plantation workers. Dogs had the highest parasite egg output in this study, suggesting that infected individuals can end up contributing significantly to parasite environmental pollution, and highlighting the importance of including domestic



**Figure 1.** Phylogenetic tree depicting the relationship between our samples (508 bp) and other related echinostomes inferred from 1000 replicates of partial COI sequences based on the neighbour-joining method and by using *Fasciolopsis buski* as an outgroup. Evolutionary distances were computed by using the Kimura 2-parameter methods and are presented as number of base substitutions per site. Terminal nodes in colour denote specimens collected in the current study and the other sequences were retrieved from GenBank. Tree reconstruction was conducted in Geneious Prime v. 2023.2.

animals in public health strategies aimed at controlling food-borne trematodiasis in humans. Additionally, 9 samples originally presumed to be from carnivores were found to be from humans, with a reported prevalence (77.7%) that is notably higher than previous studies on echinostomes in Southeast Asia (7%–17% and 0.7%–1.8% for *E. ilocanum* [Sohn *et al.*, 2011a; Chai and Jung, 2019], 7.5%–22.4% for *E. revolutum* [Sohn *et al.*, 2011b] and an average of 43% for *E. lindoense* [Carney *et al.*, 1980], but see [Chai, 2009]). Our findings highlight the importance of implementing epidemiological surveys in these human–wildlife interfaces using a more holistic approach. In Sabah, the intermediate hosts for *E. lindoense* and *E. revolutum* have not yet been found, whereas the intermediate freshwater snail hosts for *A. malayanum* have been reported from at least three districts, including the study area (Table S1), suggesting that all the necessary components for the completion of this parasite's life cycle are present.

Despite having its limitations, fecal samples are often the only non-invasive method available to obtain information on gastrointestinal parasites in wildlife (Bechert, 2012). Only 21.4% of the samples detected as positive by microscopy were amplified by PCR. The low amplification rate could be attributed to the presence of DNA polymerase inhibitors in feces (Stensvold *et al.*, 2006; Lamaningao *et al.*, 2017), but it is also possible that more than one trematode species was present in the samples, and that primer affinity was not sufficient to characterize all of them.

The Kinabatangan floodplain's fragmented landscape creates a network of intricate channels and drains interconnected with the river and lakes (Harun *et al.*, 2014), teeming with abundant fish, snails and crustaceans. This ecosystem serves as an abundant food source for generalist species like civets, whose activity ranges coincide with those of humans and domestic animals (Wells *et al.*, 2018). The overlapping presence of definitive and intermediate hosts, combined with the unique characteristics of the floodplain, plays a significant role in facilitating the persistence of multi-host parasites (Anh *et al.*, 2009). Despite the significant public health importance of *A. malayanum* and *A. sufrartyfex*, there is limited information available in the literature. Moreover, since the symptoms of echinostomiasis due to *Artyfechinostomum* spp. tend to appear long after these parasites have established themselves in the host and are similar to those of other diarrhoeal diseases, medical diagnosis is particularly challenging (Ghatani *et al.*, 2018).

Although there are obvious links between extensive/industrial oil palm agriculture, riverine/estuarine ecology, and food safety, little attention has been given to the potential threat that oil palm agriculture poses to public health through changes in parasite ecology and food safety. Swamp aquatic ecosystems and flooded areas, prevalent in oil palm-dominated landscapes, can sustain habitats and reservoirs ideal for the development of invertebrate populations that can act as intermediate hosts for trematodes. Therefore, the negative impact of extensive agriculture on the health of aquatic ecosystems and food safety needs to be considered and addressed, as it may have significant implications for the development of effective control strategies in the region.

**Supplementary material.** The supplementary material for this article can be found at <https://doi.org/10.1017/S0031182023001257>.

**Data availability statement.** Data available within the article or its supplementary materials.

**Acknowledgements.** Permits to access natural resources were reviewed and granted by Sabah Wildlife Department and the Sabah Biodiversity Centre (JKM/MBS.1000-2/2 JLD.8 [110]; JK M/MBS.1000-2/2 JLD.10 [57,59]). We thank plantation managers for granting us access to plantation estates. Graphic abstract was created using BioRender.

**Author contributions.** LF and SGS conceived and designed the study. LF and SGS conducted data gathering. LF, BOM and THC conducted the analyses. SS provided support throughout the application process for permits. LF and SGS wrote the original draft. BOM, THC and SS reviewed and edited the original draft.

**Financial support.** This research was supported by a Nanyang Technological University Presidential Postdoctoral Fellowship (#020617-00001) to LF.

**Competing interests.** None.

**Ethical standards.** Not applicable.

## References

- Abram NK, Xofis P, Tzanopoulos J, MacMillan DC, Ancrenaz M, Chung R, Peter L, Ong R, Lackman I, Goossens B and Ambu L (2014) Synergies for improving oil palm production and forest conservation in floodplain landscapes. *PLoS ONE* **9**, e106391.
- Anh NT, Phuong NT, Murrell KD, Johansen MV, Dalsgaard A, Thu LT, Chi TT and Thamsborg SM (2009) Animal reservoir hosts and fish-borne zoonotic trematode infections on fish farms, Vietnam. *Emerging Infectious Diseases* **15**, 540–546.
- Bechert U (2012) Noninvasive techniques to assess health and ecology of wildlife populations. In Miller RE and Fowler M (eds), *Fowler's Zoo and Wild Animal Medicine*. St. Louis, MO: Elsevier Saunders, pp. 60–70. doi: 10.1016/b978-1-4377-1986-4.00009-3.
- Carney WP, Sudomo M and Purnomo A (1980) Echinostomiasis: a disease that disappeared. *Tropical and Geographical Medicine* **32**, 101–105.
- Chai, JY (2009) Echinostomes in humans. In Fried B and Toledo R (eds), *The Biology of Echinostomes*. New York, USA: Springer, pp. 147–183.
- Chai, JY and Jung, BK (2019) Epidemiology of trematode infections: an update. In Toledo R and Fried B (eds), *Digenetic Trematodes*. Cham: Springer International Publishing, pp. 359–409. doi: 10.1007/978-3-030-18616-6\_12.
- Esteban JG, Muñoz-Antoli C, Toledo R and Ash LR (2019) Diagnosis of human trematode infections. *Advances in Experimental Medicine and Biology* **1154**, 437–471.
- Frias L, Hasegawa H, Chua TH, Sipangkui S, Stark DJ, Salgado-Lynn M, Goossens B, Keuk K, Okamoto M and MacIntosh AJ (2021) Parasite community structure in sympatric Bornean primates. *International Journal of Parasitology* **51**, 925–933.
- Fried B, Graczyk TK and Tamang L (2004) Food-borne intestinal trematodiasis in humans. *Parasitology Research* **93**, 159–170.
- Ghatani S, Arya LK and Tandon V (2018) Emerging trematode zoonoses in India. In Yadav AK, Tandon V and Hoti SL (eds), *Advances in Medico-Veterinary Parasitology: An Indian Perspective*. New Delhi: Panima Publishing Corporation, pp. 340–354.
- Guerrero-Sánchez S, Wilson A, González-Abarzúa M, Kunde M, Goossens B, Sipangkui R and Frias L (2022) Serological evidence of exposure of Bornean wild carnivores to feline-related viruses at the domestic animal–wildlife interface. *Transboundary and Emerging Diseases* **69**, e3250–4.
- Harun S, Dambul R, Abdullah MH and Mohamed M (2014) Spatial and seasonal variations in surface water quality of the Lower Kinabatangan River Catchment, Sabah, Malaysia. *Journal of Tropical Biology and Conservation* **11**, 117–131.
- Irwin DM, Kocher TD and Wilson AC (1991) Evolution of the cytochrome b gene of mammals. *Journal of Molecular Evolution* **32**, 128–144.
- Jones BA, Grace D, Kock R, Alonso S, Rushton J, Said MY, McKeever D, Mutua F, Young J, McDermott J and Pfeiffer DU (2013) Zoonosis emergence linked to agricultural intensification and environmental change. *Proceedings of the National Academy of Sciences* **110**, 8399–8404.
- Lamaningao P, Kanda S, Laimanivong S, Shimono T, Darcy AW, Phyaluanglath A, Mishima N and Nishiyama T (2017) Development of a PCR assay for diagnosing trematode (*Opisthorchis* and *Haplorchis*) infections in human stools. *The American Journal of Tropical Medicine and Hygiene* **96**, 221–228.
- Moszczyńska A, Locke SA, McLaughlin JD, Marcogliese DJ and Crease TJ (2009) Development of primers for the mitochondrial cytochrome c oxidase I gene in digenetic trematodes (Platyhelminthes) illustrates the challenge of barcoding parasitic helminths. *Molecular Ecology Resources* **9**, 75–82.
- Saijuntha W, Andrews RH, Sithithaworn P and Petney TN (2021) Biodiversity of human trematodes and their intermediate hosts in Southeast Asia. In Petney TN, Saijuntha, W and Mehlhorn H (eds),

- Biodiversity of Southeast Asian Parasites and Vectors Causing Human Disease*. Cham: Springer International Publishing, pp. 63–95. doi: 10.1007/978-3-030-71161-0\_4.
- Sohn WM, Kim HJ, Yong TS, Eom KS, Jeong HG, Kim JK, Kang AR, Kim MR, Park JM, Ji SH and Sinuon M** (2011a) *Echinostoma ilocanum* infection in Oddar Meanchey province, Cambodia. *The Korean Journal of Parasitology* **49**, 187.
- Sohn WM, Chai JY, Yong TS, Eom KS, Yoon CH, Sinuon M, Socheat D and Lee SH** (2011b) *Echinostoma revolutum* infection in children, Pursat Province, Cambodia. *Emerging Infectious Diseases* **17**, 117.
- Stensvold CR, Saijuntha W, Sithithaworn P, Wongratanacheewin S, Strandgaard H, Ørnbjerg N and Johansen MV** (2006) Evaluation of PCR based coprodiagnosis of human opisthorchiasis. *Acta Tropica* **97**, 26–30.
- Van Steenkiste N, Locke SA, Castelin M, Marcogliese DJ and Abbott CL** (2015) New primers for DNA barcoding of digeneans and cestodes (Platyhelminthes). *Molecular Ecology Resources* **15**, 945–952.
- Wells K, Gibson DI, Clark NJ, Ribas A, Morand S and McCallum HI** (2018) Global spread of helminth parasites at the human–domestic animal–wildlife interface. *Global Change Biology* **24**, 3254–3265.
- Zin T, SabaiAung T, Saupin S, Myint T, KhinSN D, Aung MS and Shamsu BS** (2015) Influencing factors for cholera and diarrhoea: water sanitation and hygiene in impoverished rural villages of Beluran district. *Sabah Malaysia. Malaysian Journal of Public Health Medicine* **15**, 30–40.