

Spectrophotometry of Selected AGN Seyfert Galaxy AKN 564

L.S. Slavcheva, B.M. Mihov, G.T. Petrov, R.S. Bachev
Institute for Astronomy, Bulgarian Academy of Sciences,
lslav@astro.bas.bg

Akn 564 ($\alpha_{1950} = 22^h40^m18.3^s$, $\delta_{1950} = 29^\circ27'47''$) is a Sy1.5G SBb type galaxy. According to Zwicky (1966) it has a photographic magnitude $m_p = 14.4$ and a redshift of 0.025. The spectra of the galaxy were obtained at the 2.6-m telescope of the Crimean Astrophysical Observatory with a spectrograph having a dispersion of 100 Å mm⁻¹. They were processed with the help of SPEC and LONG packages integrated in MIDAS. As a result of the spectrophotometry we obtain the fluxes at $\lambda\lambda$ 4363, 4959, 5007 Å: I(4363), I(4959), I(5007). The spectrum of the galaxy in $\lambda\lambda$ 4000-7000 is shown in Figure 1. We use the relation of the fluxes of those narrow forbidden emission lines:

$$R = [j(\lambda 4959) + j(\lambda 5007)]/j(\lambda 4363) \quad (1)$$

$$= [8.32 \exp(3.29 \times 10^4/T)] / (1 + 4.5 \times 10^{-4} N_e/T^{1/2}), \quad (2)$$

sensitive at a greater extent to the electron temperature T_e than to the electron density n_e . The value of $R = 74.3$ we got, having a typical value of $n_e = 5 \times 10^5$ cm⁻³ for the NLR (Narrow Line Region), leads to the estimation of a typical temperature of $T_e = 10^4$ K.

We can evaluate the effective volume V_{eff} and respectively the size R_{eff} , the mass M_g and the kinetic energy E_k of the emitting gas in the NLR with $n_e = 5 \times 10^5$ cm⁻³ and $T_e = 10^4$ K assumed and I(5007) measured via the equations (Dibay 1980):

$$L(H\beta) = 4\pi R^2(1+z)^2 I(H\beta); \quad (3)$$

$$V_{\text{eff}} = R^2 I(H\beta) / j(H\beta); \quad (4)$$

$$R = cz/H; \quad (5)$$

$$V_{\text{eff}} = fV; \quad (6)$$

$$R_{\text{eff}} = (3V_{\text{eff}}/4)^{1/3}; \quad (7)$$

$$M_g = n_e m_p V_{\text{eff}} / M_\odot; \quad (8)$$

$$E_k = 1/2 M_g v^2 = 1/4 M_g FWHM; \quad (9)$$

$$M_c = 3v_\zeta^2 R/G, \quad (10)$$

where V is the geometrical volume of the region, $f \approx 10^{-3}$ is the filling factor and j is the emission coefficient.

T_e and n_e in the BRL (Broad Line Region) cannot be estimated directly. We accept representative of the BLR values of $n_e = 5 \times 10^5$ cm⁻³ and $T_e =$

10^4K acquired by comparing photoionizational models with some observational parameters. As a result we evaluate V_{eff} , R_{eff} , M_g , E_k and the mass of the central object M_c , all of them given in the following table:

NLR		BLR	
n_e , [cm^{-3}]	5×10^3	n_e , [cm^{-3}]	10^9
T_e , [K]	10^4	T_e , [K]	10^4
$I([\text{OIII}] \lambda 5007)$, [$\text{erg.cm}^{-2}.\text{s}^{-1}$]	1.04×10^{-12}	$I(\text{H}\beta)$, [$\text{erg.cm}^{-2}.\text{s}^{-1}$]	5.85×10^{-13}
$\text{FWHM}([\text{OIII}] \lambda 5007)$, [cm.s^{-1}]	663×10^3	$\text{FWHM}(\text{H}\beta)$, [cm.s^{-1}]	899×10^3
$L([\text{OIII}] \lambda 5007)$, [erg.s^{-1}]	9.18×10^{41}	$L(\text{H}\beta)$, [erg.s^{-1}]	5.18×10^{41}
$j([\text{OIII}] \lambda 5007)$, [$\text{erg.cm}^{-3}.\text{s}^{-1}$]	1.15×10^{-19}	$j(\text{H}\beta)$, [$\text{erg.cm}^{-3}.\text{s}^{-1}$]	6.63×10^{-9}
V_{eff} , [cm^3]	1.6×10^{55}	V_{eff} , [cm^3]	6.19×10^{48}
R , [pc]	5	R , [pc]	0.037
M_g , [Mo]	6.68×10^3	M_g , [Mo]	5.17
E_k , [erg]	7.34×10^{51}	E_k , [erg]	1.04×10^{49}
		M_c , [Mo]	0.52×10^7

The errors of the fluxes are about $7 \times 10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1}$ and the errors of the other parameters are about 10-30 %.




Figure 1. Energy distribution in $\lambda\lambda$ 4000-7000 Å for Akn564. The data reduction was made by MIDAS 95NOV packages. The strongest forbidden and permitted lines are marked.

References

- Arakelian, M., 1975, *Publ.Bjurak.Obs.*, 47, 3
 Dibay, E., 1980, *Astron. J.*, , 57, 677
 Zwicky, F., 1966, *Ap.J.*, , 143, 192