

Psychiatry, Psychology, And Neuroscience, King's College London, London, United Kingdom and ⁴Department Of Psychiatry, Massachusetts General Hospital, Boston, United States of America

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.341

Introduction: Sedentary behaviour is potentially a modifiable risk factor for depression and anxiety disorders, but findings have been inconsistent.

Objectives: To assess associations of sedentary behavior with depression and anxiety symptoms and estimate the impact of replacing daily time spent in sedentary behaviors with sleep, light, or moderate-to-vigorous physical activity, using novel compositional data analysis methods.

Methods: Prospective cohort study in with 60,235 UK Biobank participants (mean age: 56; 56% female). Exposure was baseline daily movement behaviours (accelerometer-assessed sedentary behaviour, physical activity, and self-reported total sleep). Outcomes were depression and anxiety symptoms (Patient Health Questionnaire-9 and Generalised Anxiety Disorders-7) at follow up.

Results: Replacing 60 minutes of sedentary behaviour with light activity, moderate-to-vigorous activity, and sleep was associated with lower depression symptom scores by 1.3% (95%CI, 0.4%-2.1%), 12.5% (95%CI, 11.4%-13.5%), and 7.6% (95%CI, 6.9%-8.4%), and lower odds of depression by 0.95 (95%CI, 0.94-0.96), 0.75 (95%CI, 0.74-0.76), and 0.90 (95%CI, 0.90-0.91) at follow-up. Replacing 60 minutes of sedentary behaviour with moderate-to-vigorous activity and sleep was associated with lower anxiety symptom scores by 6.6% (95%CI, 5.5%-7.6%) and 4.5% (95%CI, 3.7%-5.2%), and lower odds of meeting the threshold for an anxiety disorder by 0.90 (95%CI, 0.89-0.90) and 0.97 (95%CI, 0.96-0.97) at follow-up. However, replacing 60 minutes of sedentary behaviour with light activity was associated with higher anxiety symptom scores by 4.5% (95%CI, 3.7%-5.3%) and higher odds of an anxiety disorder by 1.07 (95%CI, 1.06-1.08).

Conclusions: Sedentary behaviour is a risk factor for increased depression and anxiety symptoms in adults, but different replacement activities differentially influence mental health.

Disclosure: No significant relationships.

Keywords: sedentary behaviour; Physical Activity; Depression; Anxiety

O148

Psychiatric morbidity among undergraduate students of university of Ilorin, Kwara state, Nigeria

O. Buhari^{1*}, A.J. Ogunmodede², O. Bolarinwa³, O. Adegunloye¹, A. Adegoke⁴ and R. Oguntay⁵

¹Dept Of Behavioural Sciences, UNIVERSITY OF ILORIN & UNIVERSITY OF ILORIN TEACHING HOSPITAL, ILORIN, Nigeria;

²Dept Of Behavioural Sciences, UNIVERSITY OF ILORIN TEACHING HOSPITAL, ILORIN, Nigeria; ³Department Of Epidemiology And Community Health, Faculty Of Clinical Sciences, University of Ilorin/ University of Ilorin Teaching Hospital, ilorin, Nigeria; ⁴Department Of Counseling Education, University of Ilorin, ilorin, Nigeria and ⁵Department Of Psychology, university of Ilorin, Ilorin, Nigeria

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.342

Introduction: The Nigerian tertiary education system admits mostly teenagers and young adults from different ethno-religious and family backgrounds, some of whom may have inherent risks and predisposition to mental illness. They then undergo stressful conditions related to the university life such as long durations of lectures, over-crowding, and lack of social amenities, haphazard lecture schedules as well as incessant industrial strike actions of academic and non-academic staff. In spite of these, there appears to be few studies on the burden of emotional and mental disorders among Nigerian University students, and none was cited suggesting interventions that may be appropriate.

Objectives: The objectives of the study is to determine the prevalence of psychiatric morbidity and its associated factors among undergraduate students of University of Ilorin

Methods: This is a cross-sectional study using multi staged systematic randomization. A self-administered sociodemographic questionnaire and the 12 item general health questionnaire (GHQ -12) was administered on 3,300 students.

Results: Psychiatric morbidity was found to be 23.6% of the 3179 analyzable returned questionnaires. Factors found to be significantly associated with psychiatric morbidity included female gender, relationship with parents, parental employment status and family structure. Students on scholarship were more likely to have mental illness. Other associated factors include whether course of study was the preferred one and relationships with peers and lecturers on campus. About 46.6% of the students were willing to have internet based mental health intervention programmes.

Conclusions: The data obtained from this study is relevant for the formation of mental health promotion and prevention programs on our campus.

Disclosure: this study is part of the first phase of a three phase study. it aims to explore the factors associated with psychiatric morbidity among University student as a precursor for determining appropriate mental health interventions. it was partly funded by the

Keywords: psychiatric morbidity; undergraduate students

O149

Prolonged length of stay in acute psychiatric wards: A descriptive study

L. Lopes^{1*}, M. Gonçalves-Pinho^{2,3,4}, S. Pereira¹, J.P. Ribeiro³ and A. Freitas^{2,4}

¹Department Of Psychiatry And Mental Health, Centro Hospitalar de Vila Nova de Gaia e Espinho, Vila Nova de Gaia, Portugal; ²Cintesis – Center For Health Technology And Services Research, Faculty of Medicine, University of Porto, Porto, Portugal; ³Department Of Psychiatry And Mental Health, Centro Hospitalar do Tâmega e Sousa, Penafiel, Portugal, Penafiel, Portugal and ⁴Department Of Community, Information And Health Decision Sciences (medcids), Faculty of Medicine, University of Porto, Porto, Portugal, Porto, Portugal

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.343

Introduction: The psychiatric care paradigm has shifted towards community-centered models. Yet, prolonged hospitalizations are still a reality, with debated impact at healthcare systems and patients.

Objectives: This work aims to describe prolonged hospitalizations in acute psychiatric wards through patients' sociodemographic and clinical data.