

psychiatric approach or a prison sanction. Sometimes a previous feigned symptom does not mean to have new episode with psychiatric symptomatology that should be treated. In this paper we'll focus in the prison psychiatry (that probably include all the situations that has been describe above.) and to give some clinical tips to deal with this kind of situation in the everyday work and casualty job.

Disclosure: No significant relationships.

Keywords: Faking symptoms; malingering; feigned symptoms and prison

Challenging stigma attached to mental disorders in different european countries: Understanding and doing something

S0070

Stigma towards patients with schizophrenia and other mental disorders: Challenges and interventions in Italy

B. Carpiello

Italian Psychiatric Association; Secretary Of The Epa-council Of Npas, Department of Medical Sciences and Public Health University of Cagliari, Cagliari, Italy
doi: 10.1192/j.eurpsy.2021.91

Stigma toward mental illness is considered a major public health problem, being a significant obstacle for the access to care by people with psychiatric disorders, not only the severe ones but also those improperly called "minor" disorders, as recent research demonstrates. Moreover, stigma per se causes further suffering, undermining the quality of life of those who suffer from mental disorders due to discrimination, social isolation and lack of opportunities. Thus, combating stigma is one of the main goal of mental health policies worldwide. After the 1978 Reform Act, substantial ideological and practical changes were introduced in Italy, such as, among others, the abandonment of custodial care and of the dangerousness criterion for involuntary treatments, along with the development of a nationwide system in mental health care. Notwithstanding there relevant changes and more than forty years of experience in community treatment of mental disorders and the widespread implementation of interventions oriented to social inclusion, no data about significant changes in public stigma toward mentally ill people could be registered in our country. Moreover, a quite limited number of specific anti-stigma programmes and campaigns at a national or local level were developed with a correspondently paucity of research regarding the evaluation of these interventions with specific reference to their quality and effectiveness

Disclosure: No significant relationships.

S0072

Challenging stigma attached to mental disorders: A psychosocial perspective

A. Fiorillo

Department Of Psychiatry, University of Campania "L. Vanvitelli", Naples, Italy
doi: 10.1192/j.eurpsy.2021.92

Stigma attached to mental disorders represent one of the main obstacles to patients' full recovery and empowerment. In the last decades, many anti-stigma programmes have been implemented worldwide, but stigma still represents a major obstacle for people with severe mental disorders, their family members, friends and also healthcare professionals. Stigma is a complex social phenomenon, which entails a lack of knowledge, discriminating attitudes and excluding behaviours in the general population, which deserves a multi-level approach. In particular, anti-stigma strategies combining the three most common approaches, including contact, education, and organization of protest activity, are the most effective. Interventions should contain age-appropriate information and should be provided at an early age (e.g., in schools). Interdisciplinary approaches are recommended. In particular, contact strategies are important to reduce prejudice and change attitudes towards people with mental disorders and may be implemented either by video (interviews/personal testimonies), but ideally in person with affected individuals, reporting their real life experiences. In this workshop, the role of advocacy associations together with all stakeholders of mental health will be discussed in the process of fighting stigma according to a psychosocial perspective.

Disclosure: No significant relationships.

Keywords: discrimination; social inclusion; Stigma; Mental disorders

Behavioral addictions during social-distancing for the COVID-19 pandemic

S0078

The impact of physical distancing on body-image and social media use

O. Corazza

Department Of Clinical And Pharmaceutical Sciences, University of Hertfordshire, Hatfield, United Kingdom
doi: 10.1192/j.eurpsy.2021.93

The current coronavirus pandemic (Covid-19) is posing new critical challenges on mental health due to widespread social alarm as well as long lasting "physical distancing" as a result of public health protection measures or voluntary conduct. In a period of uncertainty, certain rewarding behaviors, such as the use of the Internet, exercise among other coping strategies might have increased considerably. We will share the results of an international cross-sectional investigation on the impact of physical-distancing on such potentially addictive behaviours to mitigate the pandemic effects, while identifying the most risky patterns and vulnerable populations. The studied sample consists of 3161 participants from Italy (41%), Spain (16%), the UK (12%), Lithuania (12%), Portugal (11%), Japan (6%), and Hungary (4%). Results are currently being analysed.

Disclosure: No significant relationships.

Keywords: Problematic use of the internet; self-distancing; Covid-19; self-image