

Reported provision of analgesia to patients with acute abdominal pain in Canadian paediatric emergency departments

Naveen Poonai, MD*[†]; Allyson Cowie, BSc*; Chloe Davidson, MD*; Andréanne Benidir, MD[‡]; Graham C. Thompson, MD^{§¶}; Philippe Boisclair, MD^{||}; Stuart Harman**; Michael Miller, PhD*; Andriana Butter, MD^{††}; Rod Lim, MD*; Samina Ali, MD^{††§§}

ABSTRACT

Objectives: Evidence exists that analgesics are underutilized, delayed, and insufficiently dosed for emergency department (ED) patients with acute abdominal pain. For physicians practicing in a Canadian paediatric ED setting, we (1) explored theoretical practice variation in the provision of analgesia to children with acute abdominal pain; (2) identified reasons for withholding analgesia; and (3) evaluated the relationship between providing analgesia and surgical consultation.

Methods: Physician members of Paediatric Emergency Research Canada (PERC) were prospectively surveyed and presented with three scenarios of undifferentiated acute abdominal pain to assess management. A modified Dillman's Tailored Design method was used to distribute the survey from June to July 2014.

Results: Overall response rate was 74.5% (149/200); 51.7% of respondents were female and mean age was 44 (SD 8.4) years. The reported rates of providing analgesia for case scenarios representative of renal colic, appendicitis, and intussusception, were 100%, 92.1%, and 83.4%, respectively, while rates of providing intravenous opioids were 85.2%, 58.6%, and 12.4%, respectively. In all 60 responses where the respondent indicated they would obtain a surgical consultation, analgesia would be provided. In the 35 responses where analgesia would be withheld, 21 (60%) believed pain was not severe enough, while 5 (14.3%) indicated it would obscure a surgical condition.

Conclusions: Pediatric emergency physicians self-reported rates of providing analgesia for acute abdominal pain scenarios were higher than previously reported, and appeared unrelated to request for surgical consultation. However, an unwillingness to provide opioid analgesia, belief that analgesia can obscure a surgical condition, and failure to

take self-reported pain at face value remain, suggesting that the need exists for further knowledge translation efforts.

RÉSUMÉ

Objectifs: D'après des données, il y a une sous-utilisation, un report de l'administration et un dosage insuffisant de l'analgésie dans les services des urgences (SU) chez les patients souffrant de douleurs abdominales aiguës. En ce qui concerne les médecins qui pratiquent dans les services des urgences pédiatriques (SUP) au Canada, les auteurs ont : 1) examiné les différences de pratique théorique dans l'administration de l'analgésie chez les enfants souffrant de douleurs abdominales aiguës; 2) cerné les motifs à l'appui du report de l'administration de l'analgésie; et 3) évalué le lien entre l'administration de l'analgésie et les consultations en chirurgie.

Méthode: Les médecins membres du Groupe de Recherche en Urgence Pédiatrique du Canada ont répondu de manière prospective à un questionnaire d'enquête, et on leur a soumis trois cas de douleurs abdominales aiguës indifférenciées afin d'en évaluer la prise en charge. La distribution du questionnaire s'est faite selon une version modifiée de la méthode de Dillman, de juin à juillet 2014.

Résultats: Le taux de réponse général s'est élevé à 74,5 % (149/200); 51,7 % des répondants étaient des femmes et l'âge moyen était de 44 ans (écart type : 8,4). Les taux d'administration de l'analgésie dans les scénarios soumis, présentant des cas de colique néphrétique, d'appendicite et d'invagination, étaient de 100 %, de 92,1 % et de 83,4 %, respectivement, tandis que les taux d'administration d'opioïdes par voie intraveineuse atteignaient 85,2 %, 58,6 % et 12,4 %, respectivement. Dans les 60 réponses dans lesquelles on avait

From the *Division of Emergency Medicine, London Health Sciences Centre, Western University, London, ON; [†]Children's Health Research Institute, London Health Sciences Centre, London, ON; [‡]Hospital for Sick Children, Department of Gastroenterology, University of Toronto, Toronto, ON; [§]Department of Pediatrics, University of Calgary, Calgary, AB; [¶]Alberta Children's Hospital Research Institute, Calgary, AB; ^{||}Centre hospitalier de l'Université Laval, CHU de Québec, QC; **Children's Hospital of Eastern Ontario, University of Ottawa, Ottawa, ON; ^{††}Division of Paediatric Surgery, London Health Sciences Centre, Western University, London, ON; ^{‡‡}Department of Pediatrics, University of Alberta, Edmonton, AB; and ^{§§}Women and Children's Health Research Institute, Edmonton, AB.

Correspondence to: Dr. Naveen Poonai, London Health Sciences Centre, London, Ontario, 800 Commissioners Road East, London, ON, N6A 5W9; Email: naveen.poonai@lhsc.on.ca

indiqué demander une consultation en chirurgie, il y aurait eu administration de l'analgésie. Dans les 35 réponses dans lesquelles on avait indiqué différer l'administration de l'analgésie, 21 médecins (60 %) étaient d'avis que la douleur n'était pas assez forte, tandis que 5 autres (14,3 %) ont indiqué que la mesure masquerait la nécessité d'une intervention chirurgicale.

Conclusions: Les taux autodéclarés d'administration de l'analgésie pour des douleurs abdominales aiguës dans les scénarios soumis, par les médecins travaillant aux services des urgences pédiatriques, étaient plus élevés que les taux

antérieurs, et ils ne semblaient pas liés à la demande de consultation en chirurgie. Toutefois, la réticence à prescrire des analgésiques opioïdes, croyance selon laquelle l'analgésie masquerait la nécessité d'une intervention chirurgicale, ainsi que la persistance du refus d'accepter tel quel le degré de douleur décrit par les malades donnent à penser qu'il faudrait poursuivre les efforts d'application des connaissances en la matière.

Keywords: abdominal pain, analgesia, appendicitis, survey, pediatric

INTRODUCTION

Evidence suggests that analgesia is underutilized for acute abdominal pain¹⁻⁵, delayed in its administration,¹⁻³ and dosed insufficiently^{1,3} in the emergency department (ED) setting. Compared to adults, children are at particular risk for suboptimal analgesia and have been found to receive analgesia less often.⁶⁻⁸

Abdominal pain is the most frequent clinical feature of acute appendicitis,^{9,10} which is the most common pediatric condition requiring urgent surgical intervention.¹¹ In 2003, Kim and colleagues found that over one-third of pediatric emergency physicians (PEPs) were unlikely to provide analgesia before establishing a definitive diagnosis in children with acute abdominal pain.¹² Disapproval by surgeons was identified as the main barrier.¹² In the last decade, many studies have disputed the notion that providing analgesia is associated with an increased risk of diagnostic or management errors.¹³⁻¹⁵

The importance of providing optimal pain treatment has been echoed by several national and international policy statements. In addition to the mandate by the World Health Organization (WHO) that adequate pain treatment should be a fundamental human right,¹⁶ the American Academy of Pediatrics (AAP) recently reaffirmed its position that adequate analgesia be provided for children.¹⁷ Moreover, untreated pain in childhood has been reported to lead to long-term negative outcomes such as anxiety, hyperesthesia, and needle phobia.¹⁸

Notwithstanding the above, few EDs have policies guiding pain management in patients with acute abdominal pain,^{1,19} and a 2012 study reported that analgesia is not provided to one-third of children with abdominal pain.²⁰ It is thus imperative to explore reasons behind withholding analgesia, and, more specifically, the relationship of the practice of withholding analgesia

with surgical consultation, in order to inform knowledge translation initiatives to improve care. The objectives of this study were to: (1) explore theoretical practice variation in the provision of analgesia to children with acute abdominal pain; (2) identify reasons for withholding analgesia; and (3) evaluate the relationship between providing analgesia and surgical consultation for physicians practicing in a Canadian pediatric ED setting.

MATERIALS AND METHODS

Study design

A cross-sectional survey of PEPs was designed to test the hypothesis that there remains a reluctance to provide analgesia to children with acute abdominal pain and that this decision is related to surgical consultation.

Protocol

Potential participants were contacted from June to July 2014 through a database of PEPs administrated by Paediatric Emergency Research Canada (PERC). A modified Dillman's Tailored Design Method for mail and internet surveys was used to optimize responses.²¹ A pre-notification email was sent to physicians in the database on day 0, followed by electronic survey dissemination on days 3, 10, 17, 24, and 31. A paper-based survey copy was mailed to non-respondents on day 38. Members of the research team were blinded to the identity of electronic or paper-based participants. Surveys were administered using the SurveyMonkey platform (www.surveymonkey.com). Consent to participate was implied by completion of any portion of the electronic or paper-based survey. This study received approval from Western University's Research Ethics Board.

Participants

The participants included consenting physicians within the PERC database as of March 2014. PERC is a network of health care providers whose primary clinical, administrative, and academic appointments are at EDs within tertiary care paediatric centres across Canada, and it includes physicians who consented to have their email addresses distributed for research purposes.

Instrument

The survey instrument included demographic questions, followed by three scenarios based on actual clinical cases of intussusception, renal colic, and appendicitis (see supplementary material). After each scenario, the participants were asked: (i) whether they would offer analgesia; (ii) whether they would obtain a surgical consultation; (iii) what their analgesic choices might be; and (iv) their reasons behind a decision not to offer analgesia (if applicable). Finally, the survey asked respondents to choose from a list of clinical conditions for which they would routinely provide analgesia and, using a 5-point Likert scale, rate the degree to which they believed that analgesia could mask important physical signs. Responses to all survey questions included multiple choice responses, Likert scale ratings, and free-text. Data were coded in duplicate by two co-investigators (AC,CD), and the survey was available in both English and French.

The survey was developed based on the approach outlined by Burns and colleagues²² using a focus group of four investigators (NP, RL, AC, CD). After a pre-testing phase, the survey was pilot tested among seven emergency physicians and two surgical residents who were asked to rate it for face validity, clarity, length, comprehensiveness, and bias.

Statistical analysis

Response rates, demographic variables, number of participants indicating they would or would not provide analgesia, reasons for not providing analgesia, and types of analgesia were summarized using means, frequencies, and percentages, as appropriate. The relationship between providing analgesia, obtaining a surgical consultation, and demographic variables were summarized using the Fisher exact or chi-square test, as appropriate. The primary outcome variable was the reported frequency of providing analgesia for each scenario.

Secondary outcomes included the reasons for withholding analgesia, frequency of opioid use, and the relationship of opioid provision to surgical consultation for each scenario. Exploratory analyses included the exploration of the relationship between providing any analgesia and the following covariates, defined a priori: years of independent practice (greater than or less than 10), and type of training (pediatric emergency medicine (PEM) or other). Data were analyzed using SPSS (version 19, IBM SPSSTM, New York, NY). A *p*-value of <0.05 was considered statistically significant.

RESULTS

Respondents

The survey was distributed to 200 physicians. One hundred thirty completed the electronic version and 19 completed the paper-based survey, resulting in an overall response rate of 74.5%. Respondents were permitted to skip questions and therefore the response rates were variable for each question. On average, there was a 10% increase in responses with each additional dissemination of the survey. All of the respondents worked at least one clinical shift per month. The demographic characteristics of the participants are provided in Table 1.

Provision of analgesia

The characteristics of participants' answers to questions pertaining to the provision of analgesia based on three scenarios are presented in Table 2. The proportions of any analgesic provision, for undifferentiated abdominal pain arising from intussusception, renal colic, and appendicitis, were 83.4%, 100%, and 92.1%, respectively, while 12.4%, 85.2%, and 58.6% of participants indicated they would provide intravenous opioids, respectively, for each case.

In 35 responses, participants indicated they would not provide analgesia, and the most common reason (21/35, 60%) for this decision was a belief that pain was not sufficiently severe (Figure 1). In all 61 responses where respondents indicated they would obtain surgical consultation, they also indicated they would provide analgesia. There was no significant relationship between the provision of analgesia and type of training or years of practice (up to 10 versus greater than 10 years, Case 1: *p* = 0.27; Case 3: *p* = 0.72).

Table 1. Demographic features of study participants (n = 149)

Characteristic	
Mean age in years (SD)	43.6 (8.4)
Number of females (%)	77 (52)
Highest level of training (%)	
PEM	88 (59.1)
General pediatrics	26 (17.5)
FRCP-emergency medicine	23 (15.4)
CCFP-emergency medicine	9 (6)
Family medicine	2 (1.3)
Other	1 (0.7)
Number (%) by years in practice	
Greater than 20	24 (16.1)
16-20	15 (10.1)
11-15	35 (23.5)
6-10	38 (25.5)
Up to 5 years	35 (23.5)
Currently in fellowship	2 (1.3)
Number (%) by shifts per month	
At least 12	51 (34.2)
6-11	71 (47.7)
Fewer than 6	27 (18.1)
Number (%) with >50% of shifts in a tertiary care centre	142 (95.3)
Number (%) of patients who are under 18 years	
80-100%	123 (82.6)
60-79%	3 (2.0)
40-59%	17 (11.4)
20-39%	5 (3.4)
Less than 20%	1 (0.6)

PEM = paediatric emergency medicine; FRCP = Fellow of the Royal College of Physicians; CCFP = Certification in the College of Family Physicians

Indications for analgesia

Table 3 provides results by etiology for the acute abdominal conditions for which participants indicated they would routinely provide analgesia. The most common was renal colic (138/149, 92.6%), followed by appendicitis (134/149, 89.9%).

Most participants either disagreed (48/139, 34.5%) or strongly disagreed (85/139, 61.1%) that analgesia can mask physical findings enough to miss a diagnosis of appendicitis. Three of 139 participants (2.2%) agreed with this statement.

DISCUSSION

The results of our scenario-based survey of Canadian PEPs' self-reported rates of analgesia provision for acute abdominal pain are higher than rates reported

approximately one decade ago.^{12,23} However, unwillingness to provide opioids for severe pain and concerns regarding analgesia obscuring a surgical diagnosis remain. Our results support the possibility that awareness has increased regarding the importance of providing analgesia to children with acute abdominal pain. Our findings are also consistent with a 2013 Canadian survey of PEPs²⁴ that found only 4% of respondents stated they would withhold analgesia in the case of a child with suspected "surgical abdomen." However, both the 2013 self-reported finding and our findings may be incongruent with directly observed practice. In a 2004 retrospective medical record review of 290 children referred to the surgical service with abdominal pain in a Canadian tertiary care centre, only 14% received analgesia.²⁵ More recently, a 2012 large Canadian multi-centre retrospective medical record review found that two-thirds of children with suspected appendicitis received analgesia.²⁰ Other investigators have found that most survey respondents (64%) supported the concept of providing pre-diagnostic analgesia; however, almost 70% reported that pain treatment was rarely, if ever, given.²⁶ Wolfe and colleagues found that 75% of emergency physicians reported that patients received analgesia, but this contrasted with institutional audits revealing an actual administration rate of only 30%.²³ A plausible explanation for our findings is social desirability bias, a well-described phenomenon in survey research.²⁷ We sought to identify this bias by providing, as one of our three scenarios, a child with abdominal pain rated 4 out of 10. The fact that 23/145 (15.8%) of respondents indicated they would provide intravenous analgesia in a case of relatively mild abdominal pain suggests that social desirability bias may have played a role in participants' responses. Another possible contributing factor for this difference between self-report and practice might be patient refusal of analgesia. Whatever the reasons, this discrepancy highlights the need for knowledge translation initiatives such as the development of evidence-based pain management policies across EDs.

Historically, the reluctance among clinicians to provide analgesia to patients with acute abdominal pain^{12,28,29} was thought to be due to concerns of obscuring the diagnosis of appendicitis,^{30,31} leading to a delay in surgical management.^{32,33} In a number of previous surveys of emergency physicians^{23,28} and surgeons¹⁹, a large proportion of respondents chose not to provide analgesia until after surgical consultation.

Table 2. Provision of analgesia by case

	Case 1 2-year-old male with 4/10 abdominal pain limiting activities (suspected intussusception) (n = 145)	Case 2 16-year-old female with history of renal stones and 10/10 flank pain (suspected renal colic) (n = 142)	Case 3 6-year-old male with vomiting, fever, 8/10 suprapubic pain (suspected appendicitis) (n = 140)
Participants' Analgesia Provision			
Number (%) offering immediate analgesia	121 (83.4)	142 (100)	129 (92.1)
Number (%) arranging immediate surgical consultation	8 (5.5)	13 (9.2)	39 (27.9)
Number (%) offering respective analgesia¹			
Acetaminophen	101 (70)	11 (7.7)	64 (45.7)
Ibuprofen	101 (70)	21 (14.8)	64 (45.7)
Oral ketorolac	0	9 (6.3)	0
IV ketorolac	5 (3.4)	91 (64.1)	24 (17.1)
Oral opioid	8 (5.5)	15 (10.6)	13 (9.3)
IV opioid	18 (12.4)	121 (85.2)	82 (58.6)
Intranasal fentanyl	5 (3.4)	9 (6.3)	9 (6.4)
IV ketamine	0	1 (0.7)	0
Diclofenac	0	1 (0.7)	0
Naproxen	0	2 (1.4)	0
Combination analgesia	93 (64.1)	139 (97.9)	116 (82.9)

¹Participant could choose more than one answer.
IV = intravenous




Figure 1. Self-reported reasons for withholding analgesia among participants who indicated they would not provide analgesia. (n = 35)

This practice has long impeded timely administration of analgesia³⁴ or led to analgesia being withheld altogether.³⁵ In contrast, in all cases in our survey where participants sought surgical consultation, they indicated they would provide analgesia. Among cases where participants indicated they would withhold analgesia, only 5/35 (14%) indicated this was because they believed it would obscure a surgical condition. This shift in self-reported practice may reflect an increased

Table 3. Reported routine analgesic provision by etiology (n = 149)

Etiology	Number (%) of respondents¹ 'routinely' providing analgesia of any type
Renal colic	138 (92.6)
Appendicitis	134 (89.9)
Ovarian torsion	132 (88.6)
Testicular torsion	131 (87.9)
Bowel obstruction	107 (71.8)
Ectopic pregnancy	99 (66.4)
Intussusception	89 (59.7)
Mesenteric adenitis	88 (59.1)
UTI or pyelonephritis	66 (44.3)
Constipation	22 (14.8)
Gastroesophageal reflux	21 (14.1)
Gastroenteritis	5 (3.4)

¹ Respondents were permitted to choose more than one etiology
UTI = urinary tract infection

acceptance that analgesia does not hinder the physical examination. Alternatively, it might suggest greater reliance on diagnostic imaging,^{11,36,37} compared to the physical examination.³⁸ Still, we believe that 14% remains unacceptably high; ample evidence currently supports the pre-diagnostic administration of analgesia.^{13,14,39-46} Furthermore, although the proportion of

respondents withholding analgesia due to the belief it may mask a surgical diagnosis is significantly lower than reported in other studies, it still portends a delayed approach to providing analgesia in children with abdominal pain and emphasizes the need for wider knowledge translation.

Although the majority of respondents indicated they would provide analgesia in an appendicitis scenario (and AAP recommends to provide systemic opioids for severe pain⁴⁰), less than two-thirds of survey respondents reported a willingness to provide intravenous opioids, despite a pain score of 8 out of 10. There are several possible explanations. First, despite ample evidence demonstrating opioids to be effective agents for pain associated with appendicitis,^{13,14,39,41} concerns of adverse effects in children may still exist. Second, uncertainty of the diagnosis presented in the case scenario may have resulted in less willingness to provide opioids. Goldman and colleagues described this phenomenon, whereby morphine was given more commonly to children with a higher probability of appendicitis.⁴ Similarly, a significant number of respondents indicated they would provide immediate oral analgesia to patients who were vomiting or due for surgical consultation. This may reflect a reluctance to provide intravenous opioids as a first-line therapy.

Our findings highlight an important phenomenon regarding the reasons that physicians reportedly choose not to administer analgesia. All the scenarios in our survey depicted children with at least 4/10 abdominal pain. Among cases where respondents withheld analgesia, the most common reason cited was a belief that pain was not severe enough. This finding is incongruent with the WHO recommendations¹⁶ that analgesia be routinely provided for children with pain scores of 4/10 or greater. These recommendations further advise that physicians base their decision to offer analgesia on the patient's self-report of pain, rather than the clinician's opinion of how much pain should exist for a particular clinical situation.¹⁶

Assuming our results are more indicative of opinions rather than actual practice, our findings suggest that the change in reported practice of providing analgesia to children with acute abdominal pain has altered clinical opinion in favor of therapy. This may be due to increased awareness of the importance of appropriate pain management, improved understanding of analgesic effectiveness, or increased use of diagnostic imaging. More importantly, our findings suggest that in contrast to several decades ago, PEPs today may be willing to adopt

such initiatives because, in general, they widely endorse providing analgesia. As a result, we feel future pain management policies should incorporate tools to help clinicians recognize and quantify pain in children and identify appropriate evidence-based therapies so that actual practice patterns can better reflect what is reported in surveys.

In addition to limitations inherent to any survey design, there are several additional limitations specific to our study that should be considered. Evidence suggests that pain score documentation in the ED is associated with increased use of analgesia.^{7,47} Providing participants with a pain score may have thus artificially inflated their decision to provide analgesia. In addition, the PERC database included physicians who practiced primarily in a tertiary care setting. Increased familiarity with pediatric abdominal emergencies, more timely access to diagnostic imaging, and potentially greater awareness of current literature may have resulted in higher rates of reported analgesic provision. For these reasons, our findings may not be generalizable to community settings and general emergency physicians. In addition, our scenarios—based on actual cases of abdominal pain—were chosen because they varied in their diagnostic clarity. It has been shown that analgesia is more likely given in cases with a greater diagnostic certainty⁴ and there remains the possibility that the scenarios were sufficiently clear to the respondent such that this inflated reported rates of analgesic provision. Despite our favorable response rate of over 70%, up to 9/149 (6%) of respondents did not answer questions pertaining to the primary outcome. We do not feel that this constituted a threat to external validity or overall results of the study because the response options were comprehensive and open-ended and the number of non-respondents was relatively low. In keeping with good practice for clinician-led surveys,²² and the requirements of our ethics board, we did not force responses. Finally, the results of this survey did not evaluate PEPs' actual practice regarding analgesic timing or dosing. These are all well-described components of suboptimal analgesia in children¹⁻⁵ and issues that could be explored in future work.

CONCLUSIONS

PEPs' self-reported rates of providing analgesia for acute abdominal pain were higher than previously reported, and appeared unrelated to requests for

surgical consultation. However, an unwillingness to provide opioid analgesia, belief that analgesia can obscure a surgical condition, and failure to take patient self-reported pain at face value remain, suggesting that the need exists for further knowledge translation efforts.

Competing Interests: Funding received from Department of Paediatrics Resident Research Grant, Western University, London, ON. No other competing interests declared.

SUPPLEMENTARY MATERIAL

To view Supplementary material for this article, please visit <http://dx.doi.org/doi:10.1017/cem.2015.112>

REFERENCES

1. Ali S, Chambers AL, Johnson DW, et al. Paediatric pain management practice and policies across Alberta emergency departments. *Paediatr Child Health* 2014;19(4):190-4.
2. Furyk J, Sumner M. Pain score documentation and analgesia: a comparison of children and adults with appendicitis. *Emerg Med Australas* 2008;20(6):482-7.
3. Todd KH, Ducharme J, Choiniere M, et al. Pain in the emergency department: results of the pain and emergency medicine initiative (PEMI) multicenter study. *J Pain* 2007; 8(6):460-6.
4. Goldman RD, Crum D, Bromberg R, et al. Analgesia administration for acute abdominal pain in the pediatric emergency department. *Pediatr Emerg Care* 2006;22(1): 18-21.
5. Herd DW, Babl FE, Gilhotra Y, et al. Pain management practices in paediatric emergency departments in Australia and New Zealand: a clinical and organizational audit by National Health and Medical Research Council's National Institute of Clinical Studies and Paediatric Research in Emergency Departments International Collaborati. *Emerg Med Australas* 2009;21(3):210-21.
6. Bauman BH, McManus JG Jr. Pediatric pain management in the emergency department. *Emerg Med Clin North Am* 2005;23(2):393-414.
7. Drendel AL, Brousseau DC, Gorelick MH. Pain assessment for pediatric patients in the emergency department. *Pediatrics* 2006;117(5):1511-8.
8. Brown JC, Klein EJ, Lewis CW, et al. Emergency department analgesia for fracture pain. *Ann Emerg Med* 2003; 42(2):197-205.
9. Addiss DG, Shaffer N, Fowler BS, et al. The epidemiology of appendicitis and appendectomy in the United States. *Am J Epidemiol* 1990;132(5):910-25.
10. Rasmussen OO, Hoffmann J. Assessment of the reliability of the symptoms and signs of acute appendicitis. *J R Coll Surg Edinb* 1991;36(6):372-7.
11. Sivit CJ. Imaging the child with right lower quadrant pain and suspected appendicitis: current concepts. *Pediatr Radiol* 2004;34(6):447-53.
12. Kim MK, Galustyan S, Sato TT, et al. Analgesia for children with acute abdominal pain: a survey of pediatric emergency physicians and pediatric surgeons. *Pediatrics* 2003;112(5):1122-6.
13. Manterola C, Astudillo P, Losada H, et al. Analgesia in patients with acute abdominal pain. *Cochrane Database Syst Rev* 2011;3:CD005660.
14. Poonai N, Paskar D, Konrad S, et al. Opioid analgesia for acute abdominal pain in children: A systematic review and Meta-analysis. *Acad Emerg Med* 2014;21(11):1183-92.
15. Ranji SR, Goldman LE, Simel DL, et al. Do opiates affect the clinical evaluation of patients with acute abdominal pain? *JAMA* 2006;296(14):1764-74.
16. Brennan F, Carr DB, Cousins M. Pain management: a fundamental human right. *Anesth Analg* 2007;105(1):205-21.
17. Fein JA, Zempsky WT, Cravero JP, et al. Relief of pain and anxiety in pediatric patients in emergency medical systems. *Pediatrics* 2012;130(5):e1391-405.
18. Taddio A. Effects of early pain experience: The human literature. In *Chronic and Recurrent Pain in Children and Adolescents: Progress in Pain Research and Management* (eds. McGrath PJ, Finley GA). Seattle: IASP Press; 1999: 57-74.
19. Villain C, Wyen H, Ganzera S, et al. Early analgesic treatment regimens for patients with acute abdominal pain: a nationwide survey among general surgeons. *Langenbecks Arch Surg* 2013;398(4):557-64.
20. Thompson GC, Schuh S, Gravel J, et al. Variations in the diagnosis and management of appendicitis at Canadian pediatric emergency hospitals. *Acad Emerg Med* 2015; 22(7):811-22.
21. Dillman DA. *Mail and Internet Surveys: The Tailored Design Method: 2007 Update with New Internet, Visual, and Mixed-Mode Guide*. Hoboken, NJ: John Wiley & Sons; 2011.
22. Burns KE, Duffett M, Kho ME, et al. A guide for the design and conduct of self-administered surveys of clinicians. *CMAJ* 2008;179(3):245-52.
23. Wolfe JM, Lein DY, Lenkoski K, et al. Analgesic administration to patients with an acute abdomen: a survey of emergency medicine physicians. *Am J Emerg Med* 2000; 18(3):250-3.
24. Ali S, Chambers A, Johnson DW, et al. Reported practice variation in pediatric pain management: A survey of Canadian pediatric emergency physicians. *CJEM* 2014; 16(5):352-60.
25. Green RS, Kabani A, Dostmohamed H, et al. Analgesic use in children with acute abdominal pain. *Pediatr Emerg Care* 2004;20(11):725-9.
26. Zimmerman O, Halpern P. Opinion survey of analgesia for abdominal pain in Israeli emergency departments. *Isr Med Assoc J* 2004;6(11):681-5.
27. Mindell JS, Coombs N, Stamatakis E. Measuring physical activity in children and adolescents for dietary surveys: practicalities, problems and pitfalls. *Proc Nutr Soc* 2014; 73(2):218-25.
28. Ayoade BA, Tade AO, Salami BA, et al. Administration of analgesics in patients with acute abdominal pain: a survey of the practice of doctors in a developing country. *Int J Emerg Med* 2009;2(4):211-5.

29. Jawaid M, Masood Z, Ayubi TK. Pre-operative analgesia in the accident and emergency department. *J Coll Physicians Surg Pak* 2009;19(6):350-3.
30. Falch C, Vicente D, Häberle H, et al. Treatment of acute abdominal pain in the emergency room: a systematic review of the literature. *Eur J Pain* 2014;18(7):902-13.
31. Nissman SA, Kaplan LJ, Mann BD. Critically reappraising the literature-driven practice of analgesia administration for acute abdominal pain in the emergency room prior to surgical evaluation. *Am J Surg* 2003;185(4):291-6.
32. Armstrong FD. Analgesia for children with acute abdominal pain: a cautious move to improved pain management. *Pediatrics* 2005;116(4):1018-9.
33. Vane DW. Efficacy and concerns regarding early analgesia in children with acute abdominal pain. *Pediatrics* 2005;116(4):1018.
34. Mills AM, Shofer FS, Chen EH, et al. The association between emergency department crowding and analgesia administration in acute abdominal pain patients. *Acad Emerg Med* 2009;16(7):603-8.
35. Grundmann RT, Petersen M, Lippert H, et al. [The acute (surgical) abdomen - epidemiology, diagnosis and general principles of management]. [Article in German]. *Z Gastroenterol* 2010;48(6):696-706.
36. Smink DS, Finkelstein JA, Garcia Peña BM, et al. Diagnosis of acute appendicitis in children using a clinical practice guideline. *J Pediatr Surg* 2004;39(3):458-63.
37. Neighbor ML, Baird CH, Kohn MA. Changing opioid use for right lower quadrant abdominal pain in the emergency department. *Acad Emerg Med* 2005;12(12):1216-20.
38. Hashikawa C, Burke TF, Pallin DJ, et al. Analgesia administration for acute abdominal pain: a survey of emergency physicians. *Ann Emerg Med* 2007;50(1):91-2.
39. Gallagher EJ, Esses D, Lee C, et al. Randomized clinical trial of morphine in acute abdominal pain. *Ann Emerg Med* 2006;48(2):150-60; 160.e1-4.
40. Fein J, Zempsky W, Cravero J, and the Committee on Pediatric Emergency Medicine and Section on Anesthesiology and Pain Medicine. Relief of pain and anxiety in pediatric patients in emergency medical systems; American Academy of Pediatrics. *Pediatrics* 2012;130(5):e1391-405.
41. Sharwood LN, Babl FE. The efficacy and effect of opioid analgesia in undifferentiated abdominal pain in children: a review of four studies. *Paediatr Anaesth* 2009;19(5):445-51.
42. Amoli HA, Golozar A, Keshavarzi S, et al. Morphine analgesia in patients with acute appendicitis: a randomised double-blind clinical trial. *Emerg Med J* 2008;25(9):586-9.
43. LoVecchio F, Oster N, Sturmman K, et al. The use of analgesics in patients with acute abdominal pain. *J Emerg Med* 1997;15(6):775-9.
44. Mahadevan M, Graff L. Prospective randomized study of analgesic use for ED patients with right lower quadrant abdominal pain. *Am J Emerg Med* 2000;18(7):753-6.
45. Pace S, Burke TF. Intravenous morphine for early pain relief in patients with acute abdominal pain. *Acad Emerg Med* 1996;3(12):1086-92.
46. Thomas S, Silen W. Effect on diagnostic efficiency of analgesia for undifferentiated abdominal pain. *Br J Surg* 2003;90(1):5-9.
47. Kellogg KM, Fairbanks RJ, O'Conn AB, et al. Association of pain score documentation and analgesic use in a pediatric emergency department. *Pediatr Emerg Care* 2012;28(12):1287-92.