



Access  
leading  
journals in  
your subject

# Cambridge Core

Explore today at [cambridge.org/core](https://www.cambridge.org/core)

Cambridge **Core**



**CAMBRIDGE**  
UNIVERSITY PRESS

The background of the advertisement is a dark, monochromatic photograph of several jellyfish. The jellyfish are illuminated from below, creating a glowing effect against the dark water. They are scattered across the frame, with some in the foreground and others in the background, creating a sense of depth. The jellyfish have a translucent, bell-shaped body with a pattern of small white spots, and long, flowing tentacles.

# Life Sciences

Books and Journals from  
Cambridge University Press

Cambridge is one of the leading publishers in ecology and conservation biology and publishes high quality texts and research across the breadth of the life sciences, focusing particularly on animal behaviour, biological anthropology, evolutionary biology, computational and systems biology, as well as statistics and professional development titles for biologists.

We also have an extensive portfolio of established journals in agriculture, ecology and conservation, and animal science.

For further details visit:

[cambridge.org/core-life-sciences](http://cambridge.org/core-life-sciences)

Cambridge  
Core



CAMBRIDGE  
UNIVERSITY PRESS

# THE MARINE BIOLOGICAL ASSOCIATION OF THE UNITED KINGDOM

## TERMS OF MEMBERSHIP FOR 2017

MBA MEMBERSHIP (prices per annum)	Price from 1st March 2017
Young Marine Biologist (YMB)	£12.50
Student Member	£25.00
Associate Member	£40.00 (direct debit discount £5)
Professional Member, Mem.MBA	£45.00 (direct debit discount £5)
MBA Fellow, FMBA	£120.00 (direct debit discount £5)
Institutional Member	£340.00

JOURNAL SUBSCRIPTION (MBA price added to the above membership fees, with the exception of YMB as no Journal option for this category)	
Online Journal	£20.00
Print & Online Journal	£80.00

The Association is a learned society founded in 1884 and incorporated by Royal Charter in 2013. The Marine Biological Association (MBA) provides a unified, clear, independent voice on behalf of the marine biological community. In 1888, the MBA opened the Plymouth laboratory at Citadel Hill.

The MBA is also a registered charity with its key aims being to promote scientific research into all aspects of life in the sea, including the environment on which it depends, and to disseminate to the public the knowledge gained. The Patron of the Association is His Royal Highness the Duke of Edinburgh, who is also an Honorary Fellow. The other Honorary Fellows are His Serene Highness Prince Albert of Monaco, Dr Sylvia Earle, Sir Tim Hunt and Professor James Lovelock.

The work of the Association is controlled by the Council elected annually by its subscribing members.

The Marine Biological Association, under the direction of its Council, has undertaken research in all branches of marine science and many of the results have been published in this Journal. An account of the history of the first hundred years of the Association and Journal may be found in Volume 67 (pp. 463 and 465).

Since its foundation the Association has been supported by grants from Research Councils and by subscriptions and donations

from private members, universities, learned societies, the Fishmongers' Company and other public bodies.

The Marine Biological Association is a very active scientific learned society with a large international membership. The MBA also manages an internationally renowned research programme.

Subscriptions can be paid by cash/cheque, direct debit from a UK bank account or credit card. \*Subscriptions are reduced by £5 per annum if paying by direct debit.

Members of the Association have a wide range of rights and privileges including the use of post-nominals (FMBA for MBA Fellows, Mem.MBA for Professional Members) and being entitled to receive The Marine Biologist Magazine. The Commissioners of Inland Revenue have approved the Association for the purposes of Section 16, Finance Act, 1958, and that the whole of the annual subscriptions paid by a member who qualifies for relief under the section will be allowable as a deduction from his emoluments assessable to income tax.

All correspondence should be addressed to: The Membership Secretary, Marine Biological Association, The Laboratory, Citadel Hill, Plymouth, PL1 2PB, UK (Tel. +44 (0) 1752 426493; e-mail: [membership@mba.ac.uk](mailto:membership@mba.ac.uk)).

PUBLISHED BY THE PRESS SYNDICATE OF THE UNIVERSITY OF CAMBRIDGE  
The Pitt Building, Trumpington Street, Cambridge, CB2 1RP, United Kingdom

This journal issue has been printed on FSC-certified paper and cover board. FSC is an independent, non-governmental, not-for-profit organization established to promote the responsible management of the world's forests. Please see [www.fsc.org](http://www.fsc.org) for information.

Printed by Bell and Bain Limited, Glasgow

CAMBRIDGE UNIVERSITY PRESS

Journals Fulfillment Department, UPH, Shaftesbury Road, Cambridge CB2 8BS, United Kingdom  
1 Liberty Plaza, Floor 20, New York, NY 10006, USA  
477 Williamstown Road, Port Melbourne, VIC 3207, Australia  
C/ Orense, 4, planta 13, 28020, Madrid, Spain  
Dock House, The Waterfront, Cape Town 8001, South Africa

## CONTENTS

### ARTICLES

- Jacob, B.G., Von Dassow, P., Salisbury, J.E., Navarro, J.M. and Vargas, C.A. Impact of low pH/high  $p\text{CO}_2$  on the physiological response and fatty acid content in diatom *Skeletonema pseudocostatum* 225
- Duarte, R.C. and Flores, A.A.V. Morph-specific habitat and sex distribution in the caridean shrimp *Hippolyte obliquimanus* 235
- Kassuga, A.D., Marafon-Almeida, A. and Masunari, S. Population structure of *Sinobos cf. stanfordi* (Crustacea, Tanaidacea) from Pinheiros River mangrove, Guaratuba Bay, southern Brazil 243
- Márquez, F. and Averbuj, A. Sexual dimorphism in the shell of a nassariid gastropod. A 3D geometric morphometrics approach 249
- Staaf, D.J., Camarillo-Coop, S., Haddock, S.H.D., Nyack, A.C., Payne, J., Salinas-Zavala, C.A., Seibel, B.A., Trueblood, L., Widmer, C. and Gilly, W.F. Natural egg mass deposition by the Humboldt squid (*Dosidicus gigas*) in the Gulf of California and characteristics of hatchlings and paralarvae – ERRATUM 257
- Zaidi, R., Derbal, F. and Kara, M.H. Temporal and ontogenic variations of diet of the goldblotch grouper *Epinephelus costae* (Serranidae) in the eastern coast of Algeria 259
- Quintão Araújo, T. and Hochberg, R. First description of a gastrotrich with four scale types, including feathered biancres, a novel find for Gastrotricha 269
- Jawad, L.A. and Jig, L. Comparative osteology of the axial skeleton of the genus *Pampus* (Family: Stromateidae, Perciformes) 277
- Gonçalves, G.R.L., Júnior, E.A.B., Negreiros-Fransozo, M.L. and Castilho, A.L. Morphometric and gonad maturity of the spider crab *Libinia ferreirae* Brito Capello, 1871 (Decapoda: Majoidea: Epialtidae) on the south-eastern Brazilian coast 289
- Batista, F.M., Grade, A., Power, D.M., Ruano, F. and Harper, E.M. Occurrence and characterization of pearls from oysters of the genus *Crassostrea* 297
- Valdés, Á., Murillo, F.J., McCarthy, J.B. and Yedinak, N. New deep-water records and species of North Atlantic nudibranchs (Mollusca, Gastropoda: Heterobranchia) with the description of a new species 303
- Morandini, A.C., Stampar, S.N., Maronna, M.M. and Da Silveira, F.L. All non-indigenous species were introduced recently? The case study of *Cassiopea* (Cnidaria: Scyphozoa) in Brazilian waters 321
- Miyamoto, Y. and Iwanaga, C. Effects of sulphide on anoxia-driven mortality and anaerobic metabolism in the ark shell *Anadara kagoshimensis* 329
- Flores-Martínez, I.A., Torres-Rojas, Y.E., Galván-Magaña, F. and Ramos-Miranda, J. Diet comparison between silky sharks (*Carcharhinus falciformis*) and scalloped hammerhead sharks (*Sphyrna lewini*) off the south-west coast of Mexico 337
- Lusher, A.L., Pope, N. and Handy, R.D. Reproductive effects of endocrine disrupting chemicals, bisphenol-A and 17 $\beta$ -oestradiol, on *Cerastoderma edule* from south-west England: field study and laboratory exposure 347
- Wei, J., Zhao, C., Zhang, L., Yang, L., Zuo, R., Hou, S. and Chang, Y. Effects of short-term continuous and intermittent feeding regimes on food consumption, growth, gonad production and quality of sea urchin *Strongylocentrotus intermedius* fed a formulated feed 359
- Maciver, S.K., Evans, J., Borg, J.A., Ramos-Esplá, A.A. and Schembri, P.J. Status of the 'Mangrove tunicate' *Ecteinascidia turbinata* (Ascidacea: Perophoridae) in the Mediterranean Sea 369
- De Souza Silva-Camacho, D., Gomes, R.S., Santos, J.N.S., and Araújo, F.G. Distribution of benthic fauna in sediment grains and prop roots of a mangrove channel in south-eastern Brazil 377
- Kanamori, M., Baba, K., Natsuike, M. and Goshima, S. Life history traits and population dynamics of the invasive ascidian, *Ascidella aspersa*, on cultured scallops in Funka Bay, Hokkaido, northern Japan 387
- Ulusoy, Ş., Özden, Ö. and Pöpke, O. Organochlorine pesticide and polychlorinated biphenyl levels of horse mackerel (*Trachurus* sp.) caught from Marmara Sea coastal sites 401
- Sauty, S., Ingole, B., Jones, D.O.B., Ray, D. and Kameshraj, K.A. First quantitative exploration of benthic megafaunal assemblages on the mid-oceanic ridge system of the Carlsberg Ridge, Indian Ocean 409
- Chakraborty, K., Krishna Raola, V., Joy, M. and Makkar, F. Nutritional attributes in the fillet of skipjack tuna (*Katsuwonus pelamis*) from the Arabian Sea near the south-west coast of India 419
- Le Garrec, V., Grall, J., Chevalier, C., Guyonnet, B., Jourde, J., Lavesque, N., Bonifácio, P. and Blake, J.A. *Chaetozona corona* (Polychaeta, Cirratulidae) in the Bay of Biscay: a new alien species for the North-east Atlantic waters? 433
- Machado, F.M., Morton, B. and Passos, F.D. Functional morphology of *Cardiomya cleryana* (d'Orbigny, 1842) (Bivalvia: Anomalodesmata: Cuspidariidae) from Brazilian waters: new insights into the lifestyle of carnivorous bivalves 447