

Cu²⁺-ADSORPTION CHARACTERISTICS OF ALUMINUM HYDROXIDE AND OXYHYDROXIDES¹

M. B. McBRIDE

Department of Agronomy, Cornell University, Ithaca, New York 14853

Abstract—The nature of Cu²⁺ adsorption by boehmite, gibbsite, and noncrystalline alumina was studied over a range of equilibrium pH (4.5–7.5) and Cu²⁺ concentration (10⁻³–10⁻⁶ M) by electron spin resonance (ESR). Available chemisorption sites at pH 4.5 were the most numerous for noncrystalline alumina (~1 mmole/100 g), less for boehmite, and least for gibbsite as indicated by the relative strength of the rigid-limit ESR signal attributed to Cu²⁺ adsorbed at discrete sites. The chemisorption process involved immobilization of Cu²⁺ by displacement of one or more H₂O ligands by hydroxyl or surface oxygen ions, with the formation of at least one Cu–O–Al bond. As the pH was raised from 4.5 to 6.0, essentially all of the solution Cu²⁺ appeared to be adsorbed by the solids. However, the noncrystalline alumina and boehmite chemisorbed much of the total adsorbed Cu²⁺ (10 mmole/100 g), whereas precipitation or nucleation of Cu(OH)₂ in the gibbsite system was indicated. Precipitated Cu²⁺ was more readily redissolved by exposure to NH₃ vapor than chemisorbed Cu²⁺.

Key words—Adsorption, Alumina, Aluminum, Boehmite, Copper, Electron spin resonance, Gibbsite.

INTRODUCTION

The relative importance of oxides and oxyhydroxides of aluminum and iron in the adsorption of Cu²⁺ by clays and soils is difficult to evaluate. Clay-size phyllosilicates probably contain oxide impurities to various extents, with the result that “specific adsorption” of metal ions by clays may arise from metal-oxide interactions (Jenne, 1968). Strong evidence exists that most of the adsorption sites of pure layer silicate clays maintain a “loose” electrostatic bond with ions such as Cu²⁺ (Clementz *et al.*, 1973; McBride *et al.*, 1975; McBride, 1976) unless the pH is raised to a level which induces surface hydrolysis of the metal. The effect of this hydrolysis is to cause Cu²⁺ to become less exchangeable (i.e., specifically adsorbed) at higher pH. The strong bonding of Cu²⁺ by organic matter in soils provides an additional adsorption mechanism but there is not general agreement regarding the relative importance of organic and mineral forms of Cu²⁺ in soils (Shuman, 1979).

It is well known that pure Fe and Al oxides and oxyhydroxides are capable of adsorbing Cu²⁺ in a nonexchangeable form (Kinniburgh *et al.*, 1976; Forbes *et al.*, 1976). Adsorption by noncrystalline alumina involves a direct Al–O–Cu bond (McBride, 1978). Because truly noncrystalline oxide minerals probably do not exist in soils, the present study was undertaken to compare Cu²⁺ adsorption on noncrystalline alumina, boehmite, and gibbsite. The results should determine whether mechanisms of trace-metal bonding are comparable on different alumina minerals.

MATERIALS AND METHODS

Adsorption of Cu²⁺ on boehmite, gibbsite, and noncrystalline alumina was measured by equilibrating 0.10 g of solid with 20 ml of Cu²⁺ solution in 0.05 M NaCl solution at room temperature. The samples were shaken in capped centrifuge tubes for one day, or for much longer periods of time to assess the importance of slow adsorption reactions. The samples were then centrifuged, and the supernatants were analyzed for Cu²⁺ by atomic absorption spectrophotometry. The amount of adsorbed Cu²⁺ was determined by the measured difference in solution Cu²⁺ concentration before and after reaction with the mineral. In addition, the pH of supernatants was measured and ESR spectra were obtained on the wet unwashed mineral samples using a Varian E-104 (X-band) spectrometer. Selected mineral samples were washed with CaCl₂ solution or water, and the ESR spectra were obtained after the washing procedure.

Copper ions were coprecipitated with Al(OH)₃ by adding 10⁻⁴ moles of Cu²⁺ in solution (as CuCl₂) to 10⁻² moles of Al³⁺ in solution (as Al(NO₃)₃) and rapidly adding enough NaOH to precipitate the Al³⁺ completely (as Al(OH)₃). The product was then dialyzed for one week, and the ESR spectrum of the precipitate was obtained.

The effect of pH on Cu²⁺ adsorption by noncrystalline alumina, boehmite, and gibbsite was determined by placing 0.100 g samples of the solids in centrifuge tubes and adding 20 ml of 5 × 10⁻⁴ M Cu²⁺ in 0.05 M NaCl. The pH values of the mixtures were adjusted with NaOH over a range from 5 to 7, and the samples were shaken overnight. After centrifuging, the pH and Cu²⁺ concentrations of the supernatants were determined. However, at higher pH values, Cu²⁺ concentrations

¹ Agronomy paper No. 1399.




Figure 1. Adsorption of Cu^{2+} by noncrystalline alumina as a function of time.

were generally below the detection limit of flame atomic absorption, and the Cu^{2+} activity was measured with an Orion specific ion electrode. The unwashed, undried minerals were analyzed by ESR at the pH of equilibration.

In the Cu^{2+} -adsorption experiments described above, the availability of the adsorbed Cu^{2+} to ligand-displacement reactions was tested by exposing the minerals to NH_3 vapor (from concentrated NH_4OH) overnight. Changes in the ESR spectrum after this treatment were used to indicate the degree of Cu^{2+} - NH_3 bond formation.

Surface areas of the alumina minerals as determined by B.E.T. isotherm analysis of N_2 adsorption data were $111 \text{ m}^2/\text{g}$ for noncrystalline alumina, $143 \text{ m}^2/\text{g}$ for boehmite, and $5.9 \text{ m}^2/\text{g}$ for gibbsite. Noncrystalline alumina was prepared by rapid addition of NaOH to an AlCl_3 solution followed by washing and freeze drying without aging, whereas boehmite was prepared by aging $\text{Al}(\text{NO}_3)_3$ solutions at 180°C for 12 hr. A surface area of $143 \text{ m}^2/\text{g}$ was determined on a sample of boehmite which was somewhat more crystalline than that used for the adsorption experiments. Therefore, the value of $143 \text{ m}^2/\text{g}$ may be a somewhat low estimate of the surface area. The prepared boehmites were characterized by X-ray powder diffraction and infrared spectroscopy. The gibbsite sample was obtained from the Aluminum Company of America.

RESULTS AND DISCUSSION

Cu^{2+} adsorption on noncrystalline alumina

The initial adsorption of ~ 1.0 mmole $\text{Cu}^{2+}/100$ g on noncrystalline alumina was quite rapid (Figure 1). This




Figure 2. ESR spectra of Cu^{2+} chemisorbed by noncrystalline alumina as affected by reaction time. The position of the isotropic signal of solution $\text{Cu}(\text{H}_2\text{O})_6^{2+}$ is denoted by a vertical line, and the free electron resonance position ($g = 2.0023$) is indicated by the high-field vertical line in this and the following figures.

was followed by a slow adsorption over two weeks, producing a maximum adsorption of 6.0 mmole/100 g. During this period the solution pH remained in the range 4.9 – 5.1 . However, after 60 days the pH dropped to 4.4 , and the total adsorbed Cu^{2+} decreased to 5.1 mmole/100 g (Figure 1). Evidently, aging of the alumina gel released protons which in turn caused a partial desorption of Cu^{2+} .

The ESR spectrum of Cu^{2+} adsorbed on the alumina became more intense with longer equilibration times because of the slow adsorption process, and showed changes in the measured hyperfine splitting (A) and g -values (Figure 2). Close inspection of the spectrum of Cu^{2+} initially adsorbed (Figure 2, 1-hr reaction time) revealed a rigid-limit spectrum as well as an isotropic resonance characteristic of free $\text{Cu}(\text{H}_2\text{O})_6^{2+}$ (indicated by the vertical line in Figure 2). The latter resonance




Figure 3. ESR spectra of Cu²⁺ coprecipitated in dialyzed aluminum hydroxide (Cu/Al mole ratio of 0.01). Shown are (a) air-dry aluminum hydroxide powder, (b) undried aluminum hydroxide gel, and (c) undried gel after a few minutes exposure to NiSO₄ salt. Positions of g_{\parallel} and g_{\perp} hyperfine lines are denoted by vertical lines.

disappeared after washing the alumina with water, indicating that this signal arose from nonadsorbed Cu²⁺ in the aqueous phase. The ESR parameters of the rigid-limit Cu²⁺ were estimated to be $g_{\parallel} = 2.32$, $g_{\perp} = 2.07$, and $A_{\parallel} = 155$ gauss (0.0168 cm⁻¹), $A_{\perp} = 17.7$ gauss (0.0017 cm⁻¹). Resolution of the hyperfine components of g_{\perp} is not commonly achieved because of the small value of A_{\perp} , although this level of spectral detail has been observed in dry Mg²⁺-smectites with small amounts of Cu²⁺ occupying exchange sites (McBride *et al.*, 1975) and also in strongly dehydrated zeolites with only a fraction of the exchange sites occupied (Nicula *et al.*, 1965). Evidently the adsorbed Cu²⁺ ions are well dispersed on the alumina surface; otherwise dipolar broadening effects would prevent observation of the g_{\perp} hyperfine components. An adsorption level of 1 mmole/100 g on noncrystalline alumina with a surface area of 111 m²/g should result in an average Cu²⁺-Cu²⁺ separation distance on the surface of about 43 Å. Even at the highest adsorption level of 6 mmole/100 g, the average separation was 18 Å, still large enough to prevent significant Cu²⁺-Cu²⁺ dipolar interaction.

After longer reaction times, the isotropic resonance

of free Cu²⁺ disappeared as a result of increased adsorption, and a second rigid-limit spectrum progressively increased in intensity. This is most readily seen as a change in shape of the g_{\perp} signal (Figure 2). The parameters of this second signal after 60 days of reaction were $g_{\parallel} = 2.37$, $g_{\perp} = 2.08$, and $A_{\parallel} = 125$ gauss (0.0138 cm⁻¹). No estimate of A_{\perp} could be made, possibly because of a decrease in the magnitude of A_{\perp} which prevented resolution of individual hyperfine lines.

The data suggest that two separate mechanisms are involved in Cu²⁺ adsorption on noncrystalline alumina. The first is characterized by a rapid adsorption process and a relatively low adsorption level (~ 1 mmole Cu²⁺/100 g) and produces an ESR signal with a low g_{\parallel} value and high A_{\parallel} value. The second adsorption process occurs over several weeks; it results in a greater amount of adsorbed Cu²⁺ (~ 5 mmole/100 g) and produces an ESR signal with a high g_{\parallel} and low A_{\parallel} value. Since "specific" Cu²⁺ adsorption on the hydroxide surface involves the displacement of H₂O ligands on Cu²⁺ by OH⁻ or oxygen anions, shifts in the ESR g and A parameters probably reflect changes in the number of surface groups bonded to Cu²⁺. The Cu(H₂O)₆²⁺ ion has $g_{\parallel} = 2.40$ and $A_{\parallel} = 0.0128$ cm⁻¹ (Lewis *et al.*, 1966), whereas the planar Cu(OH)₄²⁻ complex has $g_{\parallel} = 2.26$ and $A_{\parallel} = 0.0186$ cm⁻¹ (Ottaviani and Martini, 1980). Evidently, increased coordination of Cu²⁺ to OH⁻ at equatorial ligand positions causes a decrease in g_{\parallel} and an increase in A_{\parallel} . By comparison, coprecipitated Cu²⁺ in aged aluminum hydroxide produces a rigid-limit ESR spectrum that can be resolved into two sets of resonances with parameters similar to those of adsorbed Cu²⁺ ($g_{\parallel} = 2.31$ - 2.32 and $A_{\parallel} = 156$ gauss (0.0169 cm⁻¹), $g_{\perp} = 2.36$ and $A_{\parallel} = 130$ gauss (0.0143 cm⁻¹)) as shown in Figure 3a. A comparison of these ESR parameters with those of Cu(H₂O)₆²⁺ and Cu(OH)₄²⁻ suggests that the chemisorbed and coprecipitated Cu²⁺ exists in two ligand environments. The spectrum with low g_{\parallel} and high A_{\parallel} may result from Cu²⁺ coordinated to several surface oxygen or hydroxyl ions, while that with high g_{\parallel} and low A_{\parallel} may have a more limited association with the surface. Estimates of degree of covalency of the Cu-O σ bonds from these two sets of ESR parameters produce values of $\alpha^2 = 0.83$ - 0.85 , where α^2 is a function of bond covalency (Kivelson and Neiman, 1961). Similarly, the parameters for Cu(H₂O)₆²⁺ and Cu(OH)₄²⁻ in aqueous solutions are in the same range, indicating that the Cu-O bonds of chemisorbed Cu²⁺ are fairly ionic and similar in degree of covalency to the Cu-O bonds of Cu(H₂O)₆²⁺ and Cu(OH)₄²⁻. Spectral data (unpublished) obtained for vanadyl (VO²⁺) adsorbed on noncrystalline alumina and coprecipitated in Al(OH)₃ generally agree with those obtained for Cu²⁺, indicating metal-OH or metal-O-Al bonding of the rigidly bound metal with little difference observed between the ligand fields of chemisorbed and coprecipitated metal.




Figure 4. Adsorption isotherm of Cu^{2+} on noncrystalline alumina after 90 min of reaction time at a pH of 4.5–4.6.

The predominant ESR spectrum of Cu^{2+} coprecipitated in the undried alumina (Figure 3b) had $g_{\parallel} = 2.32$, $g_{\perp} = 2.05$, $A_{\parallel} = 156$ gauss (0.0169 cm^{-1}), and $A_{\perp} = 21.3$ gauss (0.0020 cm^{-1}), but air-drying or freezing (at liquid N_2 temperature) tended to increase the relative intensity of the $g_{\parallel} = 2.36$ signal. Much of the Cu^{2+} substituted in this alumina was probably at or near the particle surfaces, because rapid treatment with dithionite or Ni^{2+} salt eliminated most of the Cu^{2+} ESR signal, leaving a broad, weak spectrum (Figure 3c). Dithionite chemically reduces Cu^{2+} to nonparamagnetic species, whereas Ni^{2+} ions broaden ESR signals beyond detection by magnetic dipolar interactions if they are able to diffuse near the paramagnetic species. In addition, exposure to NH_3 vapor for one day shifted much of the Cu^{2+} spectrum toward lower g values, indicating the availability of the coprecipitated Cu^{2+} to ligand displacement. The fact that divalent metal substitution increases the surface area of Al and Fe oxides precipitated from solution (McBride, 1978; Nalovic *et al.*, 1975) suggests that these ionic impurities preferentially occupy positions at or near the oxide surfaces to avoid internal charge imbalances.

The ESR parameters of Cu^{2+} initially chemisorbed on hydrated alumina are very similar to those of Cu^{2+} in type X and Y zeolites dehydrated at 400°C (Nicula *et al.*, 1965; Conesa and Soria, 1978). Dehydration of zeolites at successively higher temperatures produced a reduction in the values of g_{\perp} and g_{\parallel} while increasing




Figure 5. ESR spectrum of chemisorbed Cu^{2+} on boehmite at low pH after 48 hr of reaction time (a), and the spectrum of the same sample after one day of exposure to NH_3 (b).

A_{\parallel} and allowing the hyperfine lines of g_{\perp} to be resolved. Temperatures of 200°C and higher caused the partial removal of inner-sphere hydration water from $\text{Cu}(\text{H}_2\text{O})_6^{2+}$ and permitted the remaining hydration water to hydrolyze. Thus, dehydration of Type Y zeolite at temperatures above 100°C converted one type of spectrum ($g_{\parallel} = 2.38$, $g_{\perp} = 2.09$, $A_{\parallel} = 0.0133 \text{ cm}^{-1}$) into another ($g_{\parallel} = 2.33$, $g_{\perp} = 2.07$, $A_{\parallel} = 0.0166 \text{ cm}^{-1}$, $A_{\perp} = 0.0017 \text{ cm}^{-1}$), a result consistent with the thermally induced hydrolysis of $\text{Cu}(\text{H}_2\text{O})_6^{2+}$ to form hydroxy- Cu^{2+} .

The initial rapid chemisorption of Cu^{2+} on noncrystalline alumina depended upon the equilibrium Cu^{2+} concentration, and approached an apparent maximum at ~ 1.0 mmole/100 g (Figure 4). The available sites for this adsorption mechanism were evidently quite limited at the low pH (4.5) used to obtain this isotherm.

Cu^{2+} adsorption on boehmite and gibbsite

The adsorption of Cu^{2+} on boehmite at low pH (4.5) was much less (< 0.5 mmole/100 g) than on noncrystalline alumina, with no significant reduction in the Cu^{2+} concentration of 20 ml of 5×10^{-4} M CuCl_2 after reaction with 0.100 g of boehmite for a week. This low level of adsorption occurred despite the high surface area of the boehmite. However, a small amount of adsorption was detected by ESR, as shown by the appearance of a rigid-limit spectrum in the wet, unwashed boehmite with $g_{\parallel} = 2.36$ – 2.37 , $g_{\perp} = 2.07$, and $A_{\parallel} = 125$ – 130 gauss (Figure 5a). This signal was apparent after 1.5 hr of reaction, but did not appear to increase in intensity after one day of equilibration. No change in g or A parameters occurred over 8 days of equilibration, a result unlike that for the noncrystalline alumina. The higher g values and lower hyperfine splitting of Cu^{2+} on boehmite, when compared with the initially adsorbed Cu^{2+} on alumina, suggests less Cu^{2+} bonding to surface groups on boehmite than on alumina. Exposure of the boehmite sample to NH_3 vapor caused the rigid-limit signal of adsorbed Cu^{2+} to be converted to a four-line isotropic signal with $A_0 = 66$ gauss, $g_0 = 2.12$ (Figure 5b). The g -value, indicative of $\text{Cu}(\text{NH}_3)_4^{2+}$ (Nicula *et al.*, 1965), and the isotropic nature of the signal are evidence that NH_3 readily removed Cu^{2+} from




Figure 6. Effect of pH on Cu²⁺ adsorption by noncrystalline alumina, boehmite, and gibbsite.

adsorption sites on the boehmite surface, causing the ion to rotate freely in solution.

The gibbsite sample adsorbed measurable Cu²⁺ (~0.7 mmole/100 g) with no pH adjustment, probably due to the somewhat higher pH (~4.8) attained in the gibbsite system when no acid or base was added. However, no well-defined ESR signal could be observed for this adsorbed Cu²⁺. Despite this lack of signal, exposure of the gibbsite to NH₃ vapor produced the isotropic Cu(NH₃)₄²⁺ spectrum, proving that Cu²⁺ "adsorbed" by the gibbsite was readily desorbed by ligand-displacement reactions.

Effect of pH on Cu²⁺ adsorption

All three alumina materials demonstrated Cu²⁺ adsorption that was highly dependent on equilibrium pH (Figure 6). The pH-pCu equilibrium data (Figure 7) demonstrate that the Cu-alumina systems were not oversaturated with respect to the solubility of CuO or Cu(OH)₂. The noncrystalline alumina became more undersaturated with increasing time. The boehmite and gibbsite systems were often very little different in Cu²⁺ solubility at a given pH than comparable aqueous solutions without minerals added. It is likely that neither boehmite nor gibbsite possessed enough chemisorption sites to control the solubility of Cu²⁺ in the concentration range used in this study. It is not certain from these solubility data, however, if the boehmite and gibbsite surfaces provided nucleation sites for adsorption of hydroxy-Cu or if Cu²⁺ precipitated as a separate hydroxide phase.

The ESR spectra of Cu²⁺ adsorbed in the boehmite system increased in intensity as the pH was raised (Figure 8a), although the signal intensity did not increase as rapidly as the adsorption level. The existence of the rigid-limit spectrum at adsorption levels as high as 10 mmole/100 g indicates that individual Cu²⁺ ions were




Figure 7. Solubility diagram for Cu²⁺ after one day in aqueous solution (star) noncrystalline alumina (triangle), boehmite (square), and gibbsite (circle) systems. Open symbols represent Cu²⁺ activity measurements made by ion selective electrode while closed symbols represent activities determined from atomic absorption (concentrations corrected to activities by the Debye-Huckel equation). Arrows mark the shift of solubility data after four weeks equilibration time for the Cu²⁺ mineral systems and after three weeks for aqueous Cu²⁺ with no mineral present. Solubility lines are drawn from the known solubility products of Cu(OH)₂ and CuO (Baes and Mesmer, 1976).

adsorbed at discrete surface sites as the pH was raised; however, small amounts of precipitated Cu(OH)₂ would not have been detected by ESR because of the very broad signal produced by Cu-Cu interaction in Cu(OH)₂. The increase in adsorption from 0.7 mmole/100 g at pH = 4.75 to 10 mmole/100 g at pH = 6.64 had little qualitative effect on the observed rigid-limit spectrum (Figure 8a).

In contrast to boehmite, gibbsite showed only slight evidence of rigid-limit Cu²⁺ at low pH and no increase in signal intensity with increasing pH and increasing Cu²⁺ adsorption (Figure 8b). Evidently, the "adsorbed Cu²⁺" in this system did not occupy discrete sites on gibbsite surfaces, but was nucleated or precipitated so that Cu-Cu dipolar interactions prevented observation of an ESR signal. There is no evidence from this study to prove that the gibbsite surfaces had any effect on Cu²⁺ solubility as the pH was raised.

The intensity of the rigid-limit Cu²⁺ signal increased




Figure 8. ESR spectra of alumina minerals as affected by pH: (a) boehmite at pH = 4.75 and 6.64 with 0.7 and 10.0 mmole Cu^{2+} adsorbed per 100 g, respectively; (b) gibbsite at pH = 4.77 and 7.56 with 0.7 and 9.9 mmole Cu^{2+} adsorbed per 100 g, respectively; (c) noncrystalline alumina at pH = 4.95 and 6.07 with 3.0 and 9.6 mmole Cu^{2+} adsorbed per 100 g, respectively.

in the noncrystalline alumina system as the pH was increased, as a result of greater Cu^{2+} adsorption (Figure 8c). Like boehmite, the alumina provided separate surface sites for adsorption of Cu^{2+} ions as the pH increased. The nature of these sites did not appear to change with pH and adsorption level, because the A_{\parallel} value remained at about 140 gauss, and the g_{\parallel} and g_{\perp} values were about 2.34 and 2.06, respectively. Drying the noncrystalline gels at 100°C affected the spectrum, shifting g_{\parallel} to lower values and eliminating evidence of hyperfine splitting in the g_{\perp} component.

NH₃ ligand displacement reactions of adsorbed Cu^{2+}

The exposure of moist alumina materials to NH_3 vapor following Cu^{2+} adsorption permitted the "avail-




Figure 9. ESR spectra of alumina minerals after exposure to NH_3 vapor for one day: (a) noncrystalline alumina at pH = 4.95 (3.0 mmole Cu^{2+} /100 g) and 6.07 (9.6 mmole Cu^{2+} /100 g); (b) boehmite at pH = 4.75 (0.7 mmole Cu^{2+} /100 g) and 6.64 (10.0 mmole Cu^{2+} /100 g); and (c) gibbsite at pH = 4.77 (0.7 mmole Cu^{2+} /100 g) and 7.56 (9.9 mmole Cu^{2+} /100 g). The pH and Cu^{2+} adsorption values represent the equilibrium conditions before addition of NH_3 .


ability" of adsorbed Cu^{2+} to be assessed. Chemisorbed Cu^{2+} , with one or more bonds to surface Al-O groups, would be expected to resist Cu-NH₃ bond formation, while physically adsorbed (surface-nucleated) or precipitated $\text{Cu}(\text{OH})_2$ might be solubilized by NH_3 by the formation of $\text{Cu}(\text{NH}_3)_4^{2+}$ complexes. Generally, exposure of the Cu^{2+} -alumina systems to NH_3 caused the isotropic $\text{Cu}(\text{NH}_3)_4^{2+}$ spectrum (see Figure 5b) to appear. However, the noncrystalline alumina, with the largest quantity of chemisorbed Cu^{2+} , retained a portion of the Cu^{2+} in a bound form. Evidence for this surface-bound Cu^{2+} is apparent as a rigid-limit spectrum with low g_{\parallel} and g_{\perp} values, attributed to adsorbed Cu^{2+} with partial H_2O ligand displacement by NH_3 . This bound form of Cu^{2+} was most evident on the alumina material that had adsorbed the greatest amount of Cu^{2+} (at high pH), as shown in Figure 9a. Like the noncrystalline alumina, NH_3 -treated boehmite also retained some Cu^{2+} in a bound form with partial NH_3 -ligand displacement, although the $\text{Cu}(\text{NH}_3)_4^{2+}$ spectrum dominated in boehmite that had adsorbed large amounts of Cu^{2+} at higher pH (Figure 9b). It is likely that very little of the 10.0 mmole/100 g of Cu^{2+} adsorbed on boehmite at high pH actually formed direct Cu-O-Al bonds; therefore, most of this Cu^{2+} was readily desorbed as

Cu(NH₃)₄²⁺. It is not surprising, then, that the NH₃ caused Cu²⁺ to readily desorb from gibbsite as shown by the predominance of the isotropic 4-line Cu(NH₃)₄²⁺ spectrum (Figure 9c), because the structure of gibbsite is expected to have few Al-OH groups available to chemisorb Cu²⁺.


The NH₃-displacement results are consistent with a model of Cu²⁺ chemisorption which assumes reaction at surface OH-groups which are coordinated to only one Al ion. Parfitt (1978) has noted that Al-OH groups on the (001) face of gibbsite are nonreactive toward chemisorption of anions, while edge Al-OH groups are reactive. The gibbsite sample used in the anion-adsorption study had about 6 mmole/100 g of edge Al-OH groups and a surface area of 45 m²/g (Parfitt *et al.*, 1977). Because the gibbsite used in the present study had a much lower surface area, the quantity of edge Al-OH groups was much less than 6 mmole/100 g. Therefore, if Cu²⁺ chemisorption occurred only at the singly coordinated OH⁻ groups of edges, most of the 10 mmole/100 g of Cu²⁺ that was "adsorbed" on the gibbsite at high pH could not have been bonded by Al-O-Cu linkages. The ease of Cu²⁺ dissolution by NH₃, the weakness of the rigid-limit ESR spectrum of adsorbed Cu²⁺ and the pH-pCu solubility data all support the hypothesis that most of the Cu²⁺ was precipitated or nucleated as Cu(OH)₂ on gibbsite surfaces at high pH. Because the ideal boehmite surface has OH⁻ ions coordinated to one, two, and three Al ions (Parfitt, 1978), whereas the poorly organized surface of noncrystalline alumina is probably dominated by singly coordinated OH⁻, these materials should chemisorb more Cu²⁺ than gibbsite. Thus, the quantity of reactive surface-OH⁻ ions per unit weight of mineral seems to determine the level of chemisorption (noncrystalline alumina > boehmite > gibbsite).

SUMMARY

Investigations of adsorbed Cu²⁺ on aluminum hydroxide and oxyhydroxide indicate that Cu²⁺ is rapidly immobilized by chemisorption at Al-OH sites with the probable insertion of several OH⁻ or surface oxygen ions into coordination positions of the hydrated Cu²⁺ ion. At low pH, this mechanism accounts for about 1 mmole/100 g of Cu²⁺ adsorption on noncrystalline alumina, considerably less than this amount on boehmite, and almost none on gibbsite. The active bonding sites are probably surface hydroxyls coordinated to single Al ions, because these sites are numerous in noncrystalline alumina, less prevalent in boehmite, and present only at edges in gibbsite. Chemisorption may involve the formation of a surface complex with adjacent oxygen atoms as shown below:



with the release of two H⁺ ions for each Cu²⁺ ion adsorbed, a result consistent with published data on proton release (Forbes *et al.*, 1976). Bonding to a single surface oxygen may also occur, as recently suggested by Ottaviani and Martini (1980), with hydrogen bonding holding the Cu²⁺ in a rigid arrangement:



As the pH is raised, "adsorption" of Cu²⁺ increases in all alumina systems, but part of this increase in boehmite and almost all of the increase in gibbsite is a result of precipitation or surface nucleation of Cu(OH)₂ as shown by solubility data, the lack of an observable ESR signal, and the ease of dissolution of the Cu²⁺ by NH₃. However, both boehmite and noncrystalline alumina gave evidence of greater amounts of chemisorbed Cu²⁺ with increasing pH (as high as 10 mmole/100 g for noncrystalline alumina). Unlike precipitated forms of Cu²⁺, chemisorbed Cu²⁺ ions were bound at discrete sites on the surfaces.

Reaction time was an important factor for noncrystalline alumina, with adsorption at low pH increasing over a period of several weeks. However, this slow process involved a separate bonding mechanism in which the Cu²⁺ may have been coordinated to fewer hydroxyl or surface oxygen ions.

ACKNOWLEDGMENTS

Research was supported by the Earth Sciences Section, National Science Foundation Grant EAR 77-10215.

REFERENCES

- Baes, C. F. and Mesmer, R. E. (1976) *The Hydrolysis of Cations*: Wiley, New York, 489 pp.
- Clementz, D. M., Pinnavaia, T. J., and Mortland, M. M. (1973) Stereochemistry of hydrated copper(II) ions on the interlamellar surfaces of layer silicates. An electron spin resonance study: *J. Phys. Chem.* **77**, 196-200.
- Conesa, J. C. and Soria, J. (1978) Electron spin resonance study of dipole-coupled copper(II) pairs in Y zeolites: *J. Phys. Chem.* **82**, 1575-1578.
- Forbes, E. A., Posner, A. M., and Quirk, J. P. (1976) The specific adsorption of divalent Cd, Co, Cu, Pb and Zn on goethite. *J. Soil Sci.* **27**, 154-166.
- Jenne, E. A. (1968) Controls on Mn, Fe, Co, Ni, Cu and Zn concentrations in soils and water: The significant role of hydrous Mn and Fe oxides: *Adv. Chem. Ser.* **73**, 337-387.
- Kinniburgh, D. G., Jackson, M. L., and Syers, J. K. (1976) Adsorption of alkaline earth, transition, and heavy metal cations by hydrous oxide gels of iron and aluminum: *Soil Sci. Soc. Amer. J.* **40**, 796-799.
- Kivelson, D. and Neiman, R. (1961) ESR studies on the bonding in copper complexes: *J. Chem. Phys.* **35**, 149-155.
- Lewis, W. B., Alei, M., and Morgan, L. O. (1966) Magnetic resonance studies on copper(II) complex ions in solution.

- I. Temperature dependences of the ^{17}O NMR and copper(II) EPR linewidths of $\text{Cu}(\text{H}_2\text{O})_6^{2+}$: *J. Chem. Phys.* **44**, 2409–2417.
- McBride, M. B. (1976) Origin and position of exchange sites in kaolinite: An ESR study: *Clays & Clay Minerals* **25**, 6–13.
- McBride, M. B. (1978) Retention of Cu^{2+} , Ca^{2+} , Mg^{2+} , and Mn^{2+} by amorphous alumina: *Soil Sci. Soc. Amer. J.* **42**, 27–31.
- McBride, M. B., Pinnavaia, T. J., and Mortland, M. M. (1975) Electron spin resonance studies of cation orientation in restricted water layers on phyllosilicate (smectite) surfaces: *J. Phys. Chem.* **79**, 2430–2435.
- Nalovic, L., Pedro, G., and Janot, C. (1975) Demonstration by Mössbauer spectroscopy of the role played by transitional trace elements in the crystallogenesis of iron hydroxides (III): in *Proc. Int. Clay Conf., Mexico City, 1975*, S. W. Bailey, ed., Applied Publishing, Wilmette, Illinois, 601–610.
- Nicula, A., Stamires, D., and Turkevich, J. (1965) Paramagnetic resonance absorption of copper ions in porous crystals: *J. Chem. Phys.* **42**, 3684–3692.
- Ottaviani, M. F. and Martini, G. (1980) Adsorption of the $\text{Cu}(\text{OH})_4^{2-}$ complex on aluminas studied by electron spin resonance: *J. Phys. Chem.* **84**, 2310–2315.
- Parfitt, R. L. (1978) Anion adsorption by soils and soil materials: *Adv. Agron.* **30**, 1–50.
- Parfitt, R. L., Fraser, A. R., Russell, J. D., and Farmer, V. C. (1977) Adsorption on hydrous oxides II. Oxalate, benzoate and phosphate on gibbsite: *J. Soil Sci.* **28**, 40–47.
- Shuman, L. M. (1979) Zinc, manganese, and copper in soil fractions: *Soil Sci.* **127**, 10–17.

(Received 20 November 1980; accepted 22 April 1981)

Резюме—Природа адсорбции Cu^{2+} бемитом, гиббситом, и некристаллическим глиноземом исследовалась в пределах равновесия pH (4,5–7,5) и концентрации Cu^{2+} (10^{-3} – 10^{-8} м) посредством электронного спинового резонанса (ЭСР). Места доступной хемосорбции при pH 4,5 были наиболее многочисленными для некристаллических глиноземов (1 ммоль/100 г), менее для бемита и менее всего для гиббсита, как это показала относительная сила жесткого предела сигнала ЭСР, относящегося к Cu^{2+} , адсорбированному в дискретных местах. Процесс хемосорбции включает иммобилизацию Cu^{2+} посредством замещения одного или более лигандов H_2O гидроксилем либо поверхностными ионами кислорода с образованием по крайней мере одной связи $\text{Cu}-\text{O}-\text{Al}$. Когда pH увеличивалось от 4,5 до 6,0, по существу все Cu^{2+} раствора были, по-видимому, адсорбированы твердыми телами. Однако, некристаллический глинозем и бемит хемосорбировали большую часть целого количества адсорбированного Cu^{2+} (10 ммоль/100 г), тогда как в системе гиббсита происходило осаждение и зародышеобразование $\text{Cu}(\text{OH})_2$. Осажденные Cu^{2+} более легко растворялись повторно при доступе пара NH_3 , чем хемосорбированные Cu^{2+} . [E.C.]

Resümee—Es wurde die Art und Weise der Cu^{2+} -Adsorption durch Boehmit, Gibbsite, und nichtkristallisiertem Aluminiumoxid im pH-Gleichgewichtsbereich (4,5–7,5) und bei Cu^{2+} -Konzentrationen von 10^{-3} bis 10^{-8} M durch Elektronenspinresonanz (ESR) untersucht. Die meisten verfügbaren Chemisorptionsstellen bei pH 4,5 waren bei nichtkristallisiertem Aluminiumoxid (1 mMol/100 g), weniger bei Boehmit und am wenigsten bei Gibbsite vorhanden. Dies ging aus der relativen Stärke des ESR-Signals hervor, das von der Menge des gebundenen Kupfers abhängt und von an bestimmten Stellen adsorbiertem Cu^{2+} herrührte. Der Chemisorptionsprozeß umfaßte eine Immobilisierung des Cu^{2+} durch den Ersatz von einem oder mehr H_2O -Liganden durch Hydroxyl- oder Oberflächensauerstoffionen und die Bildung von mindestens einer $\text{Cu}-\text{O}-\text{Al}$ -Bindung. Wenn der pH-Wert von 4,5 auf 6,0 angehoben wurde, schien im wesentlichen alles Cu^{2+} der Lösung durch die Festsubstanzen adsorbiert worden zu sein. Das nichtkristallisierte Aluminiumoxid und der Boehmit chemisorbierten jedoch viel mehr vom gesamten adsorbierten Cu^{2+} (10 mMol/100 g), während im Gibbsite-System eine Ausfällung oder Nukleation von $\text{Cu}(\text{OH})_2$ angezeigt wurde. Ausgefälltes Cu^{2+} wurde durch NH_3 -Dampf leichter wieder aufgelöst als chemisorbiertes Cu^{2+} . [U.W.]

Résumé—On a étudié par résonance de spin d'électrons (ESR) la nature de l'adsorption de Cu^{2+} par la boémite, la gibbsite et l'alumine noncristalline sur une étendue de pH d'équilibre (4,5–7,5) et de concentration de Cu^{2+} (10^{-3} – 10^{-8}). Les sites disponibles pour la chemisorption au pH 4,5 étaient les plus nombreux pour l'alumine noncristalline (1,0 mmole/g), moins pour la boémite, et le moins pour la gibbsite comme indiqué par la force relative du signal ESR de limite-rigide attribuée au Cu^{2+} adsorbé à des sites discrets. Le procédé de chemisorption impliquait l'immobilisation de Cu^{2+} par déplacement d'un ou de plusieurs ligands H_2O par des ions hydroxydes ou des ions oxygènes de surface, avec la formation d'au moins un lien $\text{Cu}-\text{O}-\text{Al}$. A fur et à mesure que le pH était élevé de 4,5 à 6,0, toute la solution Cu^{2+} semblait essentiellement être adsorbée par les solides. L'alumine noncristalline et la boémite, cependant, ont chemisorbé une grande partie du Cu^{2+} total adsorbé (10 mmole/100 g), tandis qu'une précipitation ou une nucléation du $\text{Cu}(\text{OH})_2$ dans le système gibbsite était indiquée. Exposé à une vapeur NH_3 , le Cu^{2+} précipité était plus facilement redissout que le Cu^{2+} chemisorbé. [D.J.]