

Revision of *Eothinoceras* and the status of the Eothinoceratidae (Cyrtocerinida, Multiceratoidea, Cephalopoda)

David H. Evans^{1*}  and Marcela Cichowolski²

¹Natural England, Worcester County Hall, Spetchley Road, Worcester WR5 2NP, UK <devanscephs@gmail.com>

²Instituto de Estudios Andinos 'Don Pablo Groeber,' CONICET and Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Ciudad Universitaria, Pab. 2, AR-C1428EGA, Buenos Aires, Argentina <mcicho@gl.fcen.uba.ar>

Abstract.—The type specimens of *Eothinoceras americanum* Ulrich et al., 1944, type species of the genus *Eothinoceras* Ulrich et al., 1944, are revised based on new photographic material. The resulting interpretation of the conch shape of *Eothinoceras* shows that the type species is an endogastric cyrtocone, necessitating the restriction of *Eothinoceras* to the type species and requiring the revision of the family Eothinoceratidae and the order Cyrtocerinida. A new scheme for classifying the genera of the order Cyrtocerinida into its three families is proposed: (1) Eothinoceratidae Ulrich et al., 1944, containing the genera *Eothinoceras*; *Protothinoceras* Chen and Teichert, 1987; *Mesothinoceras* Chen and Teichert, 1987; and *Conothinoceras* Chen and Teichert, 1987; (2) Cyrtocerinidae Flower, 1946, including *Cyrtocerina* Billings, 1865; *Tangshanoceras* Chen, 1976; and *Centrocyrtocerina* Stait, 1983; (3) Bathmoceratidae Gill, 1871, containing *Bathmoceras* Barrande, 1865; *Saloceras* Evans, 2005; *Sacerdosoceras* Evans, 2005; *Margaritoceras* Cecioni and Flower, 1985; and *Mutveiceras* Cichowolski et al., 2014.

Introduction

While reviewing the order Cyrtocerinida Mutvei, 2015, in preparation for the revision of Part K of the Treatise on Invertebrate Paleontology, it has become clear that a re-evaluation of *Eothinoceras* Ulrich et al., 1944, was required because it appears that the type species, *E. americanum* Ulrich et al., 1944, is markedly different from the species subsequently assigned to the genus. This has several implications for the systematics of the order Cyrtocerinida.

The type specimens of *Eothinoceras americanum* consist of a set of randomly oriented sections representing four individuals preserved on three blocks of limestone collected by W.B. Dwight during the latter half of the 19th century from the Lower Ordovician (Stairsian, late Tremadocian) Rochdale Formation in the vicinity of Rochdale, (Dutchess County, New York). In erecting *E. americanum* and the family Eothinoceratidae, Ulrich et al. (1944, p. 131) drew attention to the similarity of the connecting rings on this material to those of *Cyrtocerina* Billings, 1865, from the Upper Ordovician of North America, but they assigned the specimens to the new genus *Eothinoceras* on the grounds that the conch was regarded as orthoconic and breviconic, and that its resemblance to *Cyrtocerina* was believed to be superficial.

Teichert and Glenister (1954, p. 48, 49) summarized the history of the systematics of the Eothinoceratidae and Cyrtocerinidae Flower, 1946, noting that Flower and Kummel (1950) included *Eothinoceras* and *Cyrtocerina* in the Cyrtocerinidae. Flower and Kummel (1950) regarded evidence that

Eothinoceras possessed an orthoconic conch as inconclusive. Based on the similarity of the connecting rings of the type species, Teichert and Glenister (1954) erected *E. maitlandi* Teichert and Glenister, 1954 from the Lower Ordovician (late Tremadocian–early Floian) Emanuel Limestone Formation of northwestern Australia, employing the characters of this species to supplement and emend the diagnosis of *Eothinoceras*. Since this emendation, two other species of *Eothinoceras* have been erected (*E. marchense* Balashov, 1960 and *E. renatae* Cecioni and Flower, 1985) and four genera (*Margaritoceras* Cecioni and Flower, 1985; *Saloceras* Evans, 2005; *Sacerdosoceras* Evans, 2005; and *Mutveiceras* Cichowolski et al., 2014) have been assigned to the Eothinoceratidae.

The illustrations of the type material of *Eothinoceras americanum* figured by Ulrich et al. (1944) have proved difficult to interpret. A restudy of the type species of *E. americanum*, based on high-resolution images kindly provided by the New York State Museum, supplies the basis for this reassessment of the status of *Eothinoceras*. *Eothinoceras americanum* is here found to be a moderately breviconic endogastric cyrtocone, and thus differs markedly from those taxa subsequently assigned to the genus.

Materials and methods

Repository and institutional abbreviation.—NYSM, New York State Museum, Albany.

Type material of *Eothinoceras americanum*

Details of the holotype and paratypes of *Eothinoceras americanum* are set out in Tables 1 and 2.

*Corresponding author.

Table 1. Relationship of the type series of *Eothinoceras americanum* to illustrations by Ulrich et al. (1944) with observations regarding the condition of each specimen.

Status	Accession No.	Ulrich et al., 1944	Comments
Holotype	NYSM 10376	pl. 67, fig. 5	Block contains holotype only. Longitudinal lateral sections visible on polished top and unpolished underside of block. A partial transverse section showing conch walls is visible at adoral end (Fig. 1.1–1.3).
Paratype	NYSM 10377	pl. 67, figs. 2, 3	On same block as NYSM 10378. Although figured by Ulrich et al. as separate individuals, NYSM 10377 appears to represent a single individual because the septa, although broken, can be traced along length of block (Fig. 1.4, 1.6).
Paratype	NYSM 10378	pl. 67, fig. 1	On same block as NYSM 10377. A discrete portion of phragmocone clearly separate and distinct from NYSM 10377 (Fig. 1.8).
Paratype	NYSM 10379	pl. 67, fig. 4	A siphuncle exposed on three faces, two of which are at high angles to each other (Fig. 1.5, 1.7).

Preservation of type material.—In their study of the Rochdale Formation cephalopod faunas of eastern New York State, Kröger and Landing (2008) were unable to identify the locations of D.W. Dwight's collecting sites in Dutchess County with confidence but noted that their section Wap-R might be along strike from Dwight's locality F (Kröger and Landing, 2008, fig. 6). They observed that the top of the Rochdale Formation in section Wap-R consists of a thrombolitic limestone containing many basslerocerids and endocerids. The clotted textures visible in the block containing the type series of *E. americanum* indicate that they are preserved in a thrombolite, suggesting that they could have originated from the thrombolite forming the top of the Wap-R section. The cephalopods themselves consist of fragments of conchs that were most probably broken prior to final burial, and then underwent a degree of stylolitis during their diagenesis, resulting in truncated septa and missing conch walls. The holotype (NYSM 10376) represents the most intact individual, whereas NYSM 10377 (Ulrich et al., 1944, pl. 67, figs. 2, 3) consists of a single, heavily stylolitized phragmocone that includes the siphuncle.

Description of holotype NYSM 10376.—The illustration of the holotype (Ulrich et al., 1944, pl. 67, fig. 5; Figs. 1.1–1.3, 2.1, 2.2) represents an oblique sagittal section through a phragmocone that is truncated adorally. The visible section is 17.5 mm long and increases in width from ~3 mm (where the phragmocone is truncated by breakage) to 12.9 mm at mid-length and 11.8 mm at the truncated adoral end. The siphuncle is visible at the apical end and shows an elliptical section. The length of siphuncle exposed is 4.67 mm and its maximum diameter is 3.1 mm. Because the conch wall curves in toward the siphuncle at the apical end (Figs. 1.2, 2.1), the siphuncle is interpreted as being located close to the phragmocone wall. The septal necks are achoanitic to loxochaoanitic with a very slight apical deflection of the septa around the septal foramen. The connecting rings are concave and substantially thickened, developing prominent subtriangular projections into the lumen of the siphuncle. Each connecting ring reaches its maximum thickness at approximately three-fifths of its length adorally. Septa are 0.75 mm apart at the apical end of the phragmocone and increase to 1.2 mm apart adorally.

The section visible on the reverse side of the block is 10.9 mm long. Adorally, the width of the section increases rapidly, reaching 9.1 mm over 4 mm, with most of the expansion occurring within the first 1.5 mm. The section reaches

13.2 mm in diameter at the truncated adoral end. Septa are 0.95 mm apart. There is no sign of the siphuncle.

The adoral transverse section shows the block to be 8.44 mm thick but appears to have been scooped out so that the area exposing the underside of the specimen is only 3.85 mm thick. The lateral surfaces of the phragmocone wall are visible in the section and reach a maximum separation of 14.3 mm, undergoing inflection within the section.

Interpretation.—As interpreted here, the proximity of the siphuncle to the conch wall at the apical end of the section (Fig. 1.2) is taken to indicate that the siphuncle is located close to the conch wall, if not in contact with it. The cross section of the siphuncle can be interpreted as a markedly oblique longitudinal section through the siphuncle, but its maximum diameter is less than that seen in any of the paratypes, whereas the ratio of the lateral diameter to the apparent dorsoventral diameter is substantially smaller than those of the paratypes. Because the conchs of two of the paratypes (NYSM 10377, 10378) are similar in diameter to that of the holotype at their apical ends, a difference in the lateral diameter of the siphuncle in the holotype seems unlikely to be the explanation for the differences in apparent diameter with those of the paratypes. An alternative would be that the section of the siphuncle seen in the holotype represents a tangential cut through one side of a siphuncle curved along its axis. Consequently, the plane of section does not penetrate the siphuncle to a sufficient depth to generate values for the siphuncle diameter like those of the paratypes.

The siphuncle is not visible in the section on the underside of the block (Fig. 1.3). Although the surface of the section is roughened and visibility is poor, the septa are visible, and it would be expected that the siphuncle would also be visible if it were present within the section. Because the siphuncle lies close to the conch wall on the upper side of the specimen, it would be expected to appear at the apical end of this underside section were it present. Because the siphuncle is not seen on the underside, it is concluded that it must be close to the conch wall on the opposing side of the phragmocone. The offset between the intersection of the conch wall on the upper- and undersides of the block indicates that the phragmocone expands rapidly, with the conch wall having a convex curvature on the underside of the block. The increase in the apparent distance between septa along the length of the section figured by Ulrich et al. (1944, pl. 67, fig. 5) could also be a consequence of the curvature of the phragmocone. The slight decrease seen in the width of the section adorally (Fig. 1.2) can be interpreted as indicating that the conch is either breviconic or curved. The fact that the adoral



width of the section on the underside of the block is greater than on the topside could be an indication of the curvature of the conch axis and/or an ovoid cross section with the antisiphuncular side of the conch more broadly rounded than the siphuncular side.

Comparison of the combined sections of the holotype (Fig. 2.1, 2.2) with a range of hypothetical dorsoventral sagittal sections through cyrtoconic phragmocones (Fig. 3) indicates that the best fit is with that of a rapidly expanding endogastric cyrtocone.

Description of paratype NYSM 10377.—This specimen (Ulrich et al., 1944, pl. 67, figs. 2, 3; Fig. 1.4, 1.6), is represented by two sections that are in continuity but lie at ~45° to each other. The larger section shows a portion of an incomplete phragmocone 17 mm long, probably close to the lateral plane. The conch wall is not preserved and appears to have been lost to stylolization whereas the other lateral half of the phragmocone is missing and was probably lost through damage to the conch prior to burial. The apparent distance between the septa is 0.89 mm parallel to the conch axis. The apical end of the specimen is connected to the adoral portion by a tenuous but continuous series of septa. The section lies in a lateral plane but is oblique to the conch axis. The apparent distance between the septa is 0.33 mm. At the apical end, the siphuncle is almost in contact with the conch wall and the width of the section increases to an estimated 12 mm over 8 mm. The lateral diameter of the siphuncle is 4.27 mm and the length visible is 5.1 mm. The septal necks are loxochonitic. The connecting rings are concave facing into the camerae, thickened and subtriangular in section. The projections of the connecting rings into the lumen of the siphuncle are elongated and markedly acute, indicating that the section is oblique dorsoventrally relative to the axis of the siphuncle.

Interpretation.—The loss of one lateral side of the phragmocone makes interpretation of this specimen more difficult because there is no evidence on which to base an interpretation of the shape of the adoral portion of the phragmocone. Nevertheless, the apical portion of the conch agrees with the holotype in possessing a marginal or submarginal siphuncle and an apparently rapid rate of expansion of the conch. The section through the siphuncle rises from the apical end adorally toward the anti-siphuncular side of the phragmocone. It is not possible to determine the direction of curvature of the conch because the adoral portion of the specimen is so incomplete that the siphuncle, if it was ever visible on this surface, has not been preserved. Thus, the absence of the siphuncle cannot be used as evidence that the conch had a concave curvature on the siphuncular side of the phragmocone.

Description of paratype NYSM 10378.—This specimen (Ulrich et al., 1944, pl. 67, fig. 1; Fig. 1.8), consists of a short (4.46 mm) length of phragmocone that is truncated adorally and stylolitized apically. It comprises a siphuncle 6.36 mm in diameter and the remnants of the septa and phragmocone wall. The siphuncle lies close to the conch wall. Within the section, the width of the phragmocone reaches at least 9.6 mm and the apparent distance between the septa is 0.7 mm. The septal necks are loxochonitic and the connecting rings are concave facing into the camerae but are thickened so that they project into the lumen of the siphuncle. The difference in the acuteness of the thickenings of the connecting rings on either side of the siphuncle could suggest that the plane of section is oblique to its long axis and lies somewhere between the lateral and dorsoventral planes.

Interpretation.—The truncation of the section of the siphuncle makes interpretation difficult, but as with NYSM 10377, with the rapid increase in the width of the section of the phragmocone, it is likely that the section through the siphuncle is an oblique cut from one side to the other.

Description of paratype NYSM 10379.—The upper surface of the block (Ulrich et al., 1944, pl. 67, fig. 4; Figs. 1.5, 1.7, 4) exposes a rough surface through a section of the siphuncle that at one end is attached to a small area of intact septa and phragmocone wall. What little remains of the septa and phragmocone wall suggests that the width of the phragmocone could increase at a high angle. The apical side of the phragmocone appears to be truncated, or possibly stylolitized, so that it is not clear whether the apparent extended/elongated apical side of the siphuncle is a real feature. Where visible, the apparent distance between the septa is 0.5 mm and the septal necks are loxochonitic. The connecting rings are thickened, and where the cut of the section appears nearly parallel with the siphuncle wall, they are elongate and acute. On the adoral side of the siphuncle there appear to be traces of the siphuncle wall, but as with the opposite side of the siphuncle, it is difficult to discern the connecting rings and this surface could also be stylolitized. The trace of the siphuncle walls continues onto the adjacent face of the block, where it closes and inflects. The region of the inflection also preserves a few septa 0.5 mm distant, as well as a fragment of the conch wall. The connecting rings in this zone are elongate and acute, matching those of the other surface in the angle that they are directed.

Interpretation.—This specimen appears to represent three cuts at different angles that penetrate deep into the siphuncle but do not go right through it (Fig. 4), instead exiting on the same side as the cut entered, so that the two inflections are on the same side of the siphuncle and connected by the wall of

Figure 1. Type material of *Eothinoceras americanum* Ulrich et al., 1944, from the Rochdale Formation, Lower Ordovician (Stairsian), near Rochdale, Dutchess County, New York: (1–3) holotype NYSM 10376: (1) section through adoral end of specimen showing trace of phragmocone wall on right; (2) upper face of holotype (also figured by Ulrich et al., 1944, pl. 67, fig. 5) showing section of siphuncle at apical end and decreasing width of section adorally; (3) roughly prepared section through back of specimen (although septa are intact at apical end in vicinity of phragmocone wall, there is no evidence of presence of siphuncle on this surface); (4, 6) paratype NYSM 10377 (also figured by Ulrich et al., 1944, pl. 67, figs. 2, 3): (4) section through whole specimen showing continuity of septa (surface containing siphuncle is oblique to rest of section); (6) detail of siphuncle and adoral end of phragmocone; (5, 7) paratype NYSM 10379 (also figured by Ulrich et al., 1944, pl. 67, fig. 4): (5) section through siphuncle and surviving septa (wall of siphuncle extends all the way to distant edge of block); (7) section through siphuncle on face as continuing from far edge of that shown in (5); (8) paratype NYSM 10378 (also figured by Ulrich et al., 1944, pl. 67, fig. 1), oblique dorsoventral section through siphuncle and septa with possibly intact phragmocone wall on left. a = siphuncle wall and connecting ring protruding into lumen of siphuncle; b = septa; c = wall of phragmocone. Scale bars = 10 mm (1–5), 5 mm (6–8).

Table 2. Comparison of the type material based in available measurements of sections. AR = ratio SD/LD, providing an indication of the obliquity of the siphuncle within the section; CD = relative cameral depth as ratio of distance between septa to dorsoventral diameter of phragmocone; DS = apparent distance between septa; LD = longest diameter of the siphuncle within the section; MD = maximum diameter of phragmocone visible; SD = shortest diameter of the siphuncle within the section.

Specimen	MD (mm)	SD (mm)	LD (mm)	AR	DS (mm)	CD
10376	12.9	3.1	4.65	0.659	0.75	0.09
10377	-	3.50	4.27	0.82	1.20	-
10378	-	6.36	-	-	0.33	-
10379	-	5.82	-	-	0.89	-
					0.70	-
					0.50	-

the siphuncle (Fig. 4). Because the exposed lateral surfaces of the siphuncle appear to be stylolitized and the rest of the siphuncle wall is only exposed in oblique transverse sections, it is not possible to determine the direction of curvature of the siphuncle axis, or for that matter, whether it is curved at all.

Discussion

The holotype and all three of the paratypes possess siphuncles with loxochoanitic septal necks and thick connecting rings. The surfaces of the connecting rings facing into the camerae are concave, whereas the surfaces facing into the lumen of the siphuncle are strongly convex and possesses triangular cross sections. In all of the specimens, the siphuncle is located close to the conch wall whereas the conch appears to increase in diameter at a rapid rate. Two of the paratypes (NYSM 10377, 10378) provide no additional information on the morphology of *Eothinoceras americanum*, but the holotype (NYSM 10376), based on the section through the siphuncle combined with the absence of the siphuncle on the underside of the specimen, indicates that the conch has a markedly endogastric curvature. As with the other paratypes, NYSM 10379 does not provide any unequivocal evidence with reference to the degree of curvature of the conch but would be likely to do so if sectioned. In conclusion, contrary to previous interpretations of *Eothinoceras*, *E. americanum* is here interpreted as a moderately breviconic endogastric cyrtocone with a compressed cross section.

The state of preservation of the type material is such that it is not possible to constrain the degree of curvature of the conch of *Eothinoceras americanum* with confidence, but it appears most similar to that of *Protothinoceras* Chen and Teichert, 1987, and differs from that of *Tangshanoceras* Chen, 1976 in being less strongly curved although the diameter of the siphuncle increases more slowly. Chen and Teichert (1987, p. 155) regarded the oldest member of the family Cyrtoceriniidae, *Tangshanoceras*, as having been derived from *Protothinoceras*. Here we note that *Tangshanoceras* might as easily have been derived from *Eothinoceras*.

Protothinoceras and *Tangshanoceras* came from the Langchiashan Formation of Hebei Province, North China (Chen, 1976; Chen and Teichert, 1987). The Liangchiashan and Rochdale formations are both Tremadocian in age and broadly coeval with each other (An et al., 1983; Chen and Teichert, 1987; Zhen and Nicoll, 2009). They were deposited at similar latitudes and




Figure 2. Partial reconstruction of the phragmocone of *Eothinoceras americanum* based on the holotype NYSM 10376: (1) sketches of visible sections from images (gray shading marks zone where thickness of block rapidly decreases); (2) interpreted reconstruction of dorsoventral section (pale gray shading = siphuncle; darker gray shading = septate part of phragmocone). i = adoral end; ii = back of specimen (dashed lines); iii = front of specimen (continuous lines) overlain to match positions on either side of block.

in similar environments: stromatolitic reefs (Chen and Teichert, 1987) and thrombolitic facies (Kröger and Landing, 2008, fig. 5), respectively. Although only two cephalopod genera are common to both formations (see Chen and Teichert, 1987,




Figure 3. Hypothetical dorsoventral sections through exogastric (top left) and endogastric (top right) cyrtoconic brevicones showing the traces of lateral sections (lower 1–7). Of the hypothetical lateral sections, (6) appears to match the section of NYSM 10376 (Fig. 1.2), the holotype of *Eothinoceras americanum*, most closely.

p. 151, 152; Kröger and Landing, 2008, p. 497), this increases to six if other coeval Laurentian faunas are included (see Ulrich et al., 1942, 1943, 1944), although many of these genera have long stratigraphical ranges.

Implications for the systematics of the Cyrtocerinida

Because all other species assigned to *Eothinoceras* differ from *E. americanum* in possessing a relatively slowly expanding

orthoconic conch that can have a slight exogastric curvature and an almost circular conch cross section, it seems untenable to retain those species in the genus. Moreover, because the preservation of the type material of *E. americanum* is too poor to permit the genus to be recognized as the senior synonym of *Protothinoceras* or *Tangshanoceras* with any certainty, the genus is best regarded as monospecific, and represented only by the type material of *E. americanum*. For now, a pragmatic solution would be to retain a family Eothinoceratidae,




Figure 4. Interpretation of paratype NYSM 10379. Shaded areas represent the surfaces visible on the actual specimen. The leftmost section is that represented by Ulrich et al. (1944, pl. 67, fig. 4). The specimen represents a short length of the siphuncle where one side has been lost to weathering.

containing *Eothinoceras*, *Protothinoceras*, and *Mesothinoceras* Chen and Teichert, 1987. Those genera possessing a very marked endogastric curvature and high rate of conch expansion, including *Cyrtocerina*, *Conothinoceras* Chen and Teichert, 1987, *Centrocyrtocerina* Stait, 1983, and *Tangshanoceras*, are placed in the Cyrtocerinidae Flower, 1946 (Table 4).

Those species that were previously assigned to *Eothinoceras* are here placed in *Saloceras*. *Saloceras* (Table 3, column 6) is an orthoconic longicone that has a moderately broad marginal siphuncle with achoanitic to weakly orthochoanitic septal necks and connecting rings that are thickened and protrude strongly into the lumen of the siphuncle. Externally, the distance between the sutures ranges from 0.07–0.17 of the conch diameter, whereas the sutures form narrow but prominent saddles over the siphuncle. Other than *Eothinoceras americanum*, all three of the species previously assigned to *Eothinoceras* (Table 3, columns 3–5) are morphologically very similar to *Saloceras* and are here reassigned to that genus.

Saloceras appears to be closely related to *Bathmoceras* Barrande, 1865. The latter could arise from the former through

Table 4. Proposed revised familial composition of the Cyrtocerinida.

Family Eothinoceratidae Ulrich et al., 1944
<i>Eothinoceras</i> Ulrich et al., 1944
<i>Protothinoceras</i> Chen and Teichert, 1987
<i>Mesothinoceras</i> Chen and Teichert, 1987
<i>Conothinoceras</i> Chen and Teichert, 1987
Cyrtocerinidae Flower, 1946
<i>Cyrtocerina</i> Billings, 1865
<i>Centrocyrtocerina</i> Stait, 1983
<i>Tangshanoceras</i> Chen, 1976
Bathmoceratidae Gill, 1871
<i>Bathmoceras</i> Barrande, 1865
<i>Saloceras</i> Evans, 2005
<i>Sacerdosoceras</i> Evans, 2005
<i>Margaritoceras</i> Cecioni and Flower, 1985
<i>Mutveiceras</i> Cichowolski et al., 2014

an increasing height of the ventral saddle. This led to an increase in the circumference of the connecting ring but did not alter the circumference of the siphuncle as illustrated by Mutvei (2015, fig. 1c). This, when combined with the extension of the inner surface of the connecting ring into the lumen of the siphuncle, resulted in Dapingian and Darriwilian forms in which the ratios of the surface area of the connecting ring to the cameral volume are increasingly enhanced. For this reason, *Saloceras* is assigned to the Bathmoceratidae Gill, 1871. The orthoconic *Margaritoceras* and the slightly exogastric *Sacerdosoceras* and *Mutveiceras* all exhibit similar siphuncle morphologies to *Saloceras sericeum* (Salter in Ramsay, 1866) and are here assigned to the Bathmoceratidae (Table 4). The enigmatic genus *Desioceras* Cecioni, 1953 from northwestern Argentina cannot be included in this new scheme until more material is available to permit a revision of its morphology.

Acknowledgments

We are most grateful to L. Amati (NYSM) for providing the high-resolution images of the type material of *Eothinoceras americanum*. Without her help in this matter, it would not have been possible to conduct this investigation. We are grateful to the reviewers M. Aubrechtová and K. Histon and to the editor M. Yacobucci for their helpful suggestions to improve this manuscript.

Declaration of competing interests

The authors declare that they have no competing interests.

Table 3. Comparison of the type species, *Eothinoceras americanum*, with other species subsequently assigned to the genus and with *Saloceras sericeum*. AA = apical angle; CC = conch curvature; CD = relative cameral depth as ratio of distance between septa to dorsoventral diameter of phragmocone; CS = conch shape; CX = conch cross section; RSD = relative diameter of siphuncle as ratio of siphuncle diameter to phragmocone diameter; SP = relative position of the siphuncle; VS = ventral saddle; * = estimated from reconstruction (see Figs. 1, 2).

	<i>Eothinoceras americanum</i>	<i>Eothinoceras maitlandi</i>	<i>Eothinoceras marchense</i>	<i>Eothinoceras renatae</i>	<i>Saloceras sericeum</i>
CC	Endogastric	Weakly exogastric	Weakly exogastric	Unknown	Orthoconic
CS	Breviconic	Longiconic	Longiconic	Longiconic	Longiconic
CX	Compressed	Depressed 0.85	Circular	Cross section unknown	Circular to slightly depressed
AA (°)	~30*	7	9–10	12	6.5
SP	Marginal or submarginal	Marginal	Almost marginal	Marginal	Marginal
SD	~0.34	0.29	0.25	0.19	0.36
CD	0.09	0.07	0.05	0.11	0.14
VS	Unknown; conch surface not visible	Shallow, 0.5–1 camera in height	Unknown; conch surface too poorly preserved to interpret	Shallow, 0.5 camera in height	0.5–1.5 camerae in height

References

- An, T., Zhang, F., Xiang, W., Zhang, Y., Xu, W., Zhang, H., Jiang, D., Yang, C., Lin, L., Cui, Z., and Yang, X., 1983, [The Conodonts of North China and the Adjacent Regions]: Beijing, Science Press, 223 p., 38 pls. [in Chinese with English abstract]
- Balashov, Z.G., 1960, Novye ordovikskie nautiloidei SSSR: Novye Vidy Drevnikh Rasteniy i Bespozvonochnykh SSSR, Ministerstva Geologii i Okhrany Nedr SSSR, Moscow, p. 123–136, pls. 26–32.
- Barrande, J., 1865, Systeme Silurien du Centre de la Bohême, Première Partie, Recherches Paléontologiques, Volume 2, Classe des Mollusques, Ordre des Céphalopodes: Prague, pls. 1–107.
- Billings, E., 1865, Paleozoic Fossils, Volume 1, Containing Descriptions and Figures of New and Little Known Species of Organic Remains from the Silurian Rocks, 1861–1865: Geological Survey Canada, Montreal, p. 169–426.
- Cecioni, G., 1953, Contribución al conocimiento de los nautiloideos eopaleozoicos argentinos: Parte 1, Protocycloceratidae-Cyclostomiceratidae: Boletín del Museo Nacional de Historia Natural de Santiago de Chile, v. 26, no. 2, p. 55–110, pls. 1–3.
- Cecioni, G., and Flower, R.H., 1985, Bathmoceratidae (Nautiloideos, Ordovícico) de Sud América: Neues Jahrbuch für Geologie und Paläontologie, v. 176, no. 3, p. 341–357.
- Chen, J.Y., 1976, [Advances in the Ordovician stratigraphy of North China with a brief description of nautiloid fossils]: Acta Palaeontologica Sinica, v. 15, no. 1, p. 57–74. [in Chinese with English abstract.]
- Chen, J.Y., and Teichert, C., 1987, The Ordovician cephalopod suborder Cyrtocerinina (order Ellesmerocerida): Paleontologia Cathayana, v. 2, p. 145–229.
- Cichowolski, M., Waisfeld, B.G., Vaccari, N.E., and Marengo, L., 2014, The nautiloid family Eothinoceratidae from the Floian of the Central Andean Basin (NW Argentina and South Bolivia): Geological Journal, v. 50, no. 6, p. 764–782, <https://doi.org/10.1002/gj.2595>.
- Evans, D.H., 2005, The Lower and Middle Ordovician cephalopod faunas of England and Wales: Monograph of the Palaeontographical Society, v. 158, no. 623, p. 1–81, <https://doi.org/10.1080/25761900.2022.12131803>.
- Flower, R.H., 1946, Ordovician cephalopods of the Cincinnati region, part 1: Bulletins of American Paleontology, v. 29, no. 116, 556 p., 50 pls.
- Flower, R.H., and Kummel, B., 1950, Classification of the Nautiloidea: Journal of Paleontology, v. 24, p. 604–615.
- Gill, T., 1871, Arrangement of the families of the Mollusca: Smithsonian Miscellaneous Collections, v. 227, 49 p.
- Kröger, B., and Landing, E., 2008, Onset of the Ordovician cephalopod radiation—Evidence from the Rochdale Formation (middle Early Ordovician, Stairsian) in eastern New York: Geological Magazine, v. 145, no. 4, p. 490–520, <https://doi.org/10.1017/S0016756808004585>.
- Mutvei, H., 2015, Characterization of two new superorders Nautilosiphonata and Calciosiphonata and a new order Cyrtocerinida of the subclass Nautiloidea: Siphuncular structure in the Ordovician nautiloid *Bathmoceras* (Cephalopoda): GFF, v. 137, no. 3, p. 164–174, <https://doi.org/10.1080/11035897.2015.1061592>.
- Ramsay, A.C., 1866, The Geology of North Wales, with an Appendix on the Fossils by J.W. Salter: Memoirs of the Geological Survey of Great Britain, v. 3, vii + 381 p., 28 pls.
- Stait, B., 1983, Ordovician nautiloids of Tasmania, Australia: Ellesmerocerida and Tarphycerida: Alcheringa, v. 7, p. 253–261.
- Teichert, C., and Glenister, B.F., 1954, Early Ordovician cephalopod fauna from northwestern Australia: Bulletins of American Paleontology, v. 35, no. 150, p. 7–112.
- Ulrich, E.O., Foerste, A.F., Miller, A.K., and Furnish, W.M., 1942, Ozarkian and Canadian cephalopods, Part I: Nautilicones: Geological Society of America Special Papers, v. 37, p. 1–157.
- Ulrich, E.O., Foerste, A.F., and Miller, A.K., 1943, Ozarkian and Canadian cephalopods, Part 2: Brevicones: Geological Society of America Special Papers, v. 49, p. 1–240.
- Ulrich, E.O., Foerste, A.F., Miller, A.K., and Unklesbay, A.G., 1944, Ozarkian and Canadian cephalopods, Part 3: Longicones and summary: Geological Society of America Special Papers, v. 58, p. 1–226.
- Zhen, Y.Y., and Nicoll, R.S., 2009, Biogeographic and biostratigraphic implications of the *Serratognathus bilobatus* fauna (Conodonta) from the Emanuel Formation (Early Ordovician) of the Canning Basin, Western Australia: Records of the Australian Museum, v. 61, p. 1–30, <https://doi.org/10.3853/j.0067-1975.61.2009.1520>.

Accepted: 7 October 2022