



# Stable reduction of finite covers of curves

Qing Liu

## ABSTRACT

Let  $K$  be the function field of a connected regular scheme  $S$  of dimension 1, and let  $f : X \rightarrow Y$  be a finite cover of projective smooth and geometrically connected curves over  $K$  with  $g(X) \geq 2$ . Suppose that  $f$  can be extended to a finite cover  $\mathcal{X} \rightarrow \mathcal{Y}$  of semi-stable models over  $S$  (it is known that this is always possible up to finite separable extension of  $K$ ). Then there exists a unique minimal such cover. This gives a canonical way to extend  $X \rightarrow Y$  to a finite cover of semi-stable models over  $S$ .

Let  $S$  be a Dedekind scheme (i.e. a connected Noetherian regular scheme of dimension 1), with field of functions  $K := K(S)$ . Let  $f : X \rightarrow Y$  be a finite morphism of smooth geometrically connected projective curves over  $K$ . We can ask how to extend, in some canonical way, the morphism  $f$  to a morphism of models of  $X$  and  $Y$  over  $S$ . It is proved in [LL99, Proposition 4.4] that if  $X$  and  $Y$  have stable models  $\mathcal{X}^{\text{st}}, \mathcal{Y}^{\text{st}}$  over  $S$ , then  $f$  extends uniquely to a morphism  $\mathcal{X}^{\text{st}} \rightarrow \mathcal{Y}^{\text{st}}$ . However, we will in general lose the finiteness of  $f$ . On the other hand, after a finite separable extension of  $K$ ,  $f$  extends to a finite morphism of semi-stable models  $\mathcal{X} \rightarrow \mathcal{Y}$  over  $S$  (see [Col03]; [LL99, Remark 4.6] and Corollary 3.10 below). Following Coleman [Col03], such a pair  $\mathcal{X} \rightarrow \mathcal{Y}$  is called a *semi-stable model* of  $f$ , and it is called *stable* if, moreover, it is minimal among the semi-stable models of  $f$  (cf. Definition 3.1). The stable model of  $f$  (if it exists) is unique up to isomorphism.

**THEOREM 0.1** (Corollary 4.6). *Suppose that either  $g(X) \geq 2$ , or  $g(X) = 1$  and  $X$  has potentially good reduction. Then there exists a finite separable extension  $K'$  of  $K$  such that  $X_{K'} \rightarrow Y_{K'}$  admits a stable model  $\mathcal{X}' \rightarrow \mathcal{Y}'$  over  $S'$ , where  $S'$  is the integral closure of  $S$  in  $K'$ . Moreover, for any Dedekind scheme  $T$  dominating  $S'$ ,  $\mathcal{X}'_T \rightarrow \mathcal{Y}'_T$  is the stable model of  $X_{K(T)} \rightarrow Y_{K(T)}$  over  $T$ .*

This gives a canonical way to extend a finite cover of projective smooth curves over  $K$  to a finite cover of semi-stable models over (some finite cover of)  $S$ . The last part of the theorem states that the stable model of  $f$  commutes with flat base change. The stable model of  $f$  should be seen as an analogue of the stable model of a curve. In [Mau05], this theorem is used to construct a compactification of Hurwitz moduli spaces of finite covers of curves. We also prove the theorem for smooth marked curves  $X, Y$ . Note that, in general,  $\mathcal{X}', \mathcal{Y}'$  are not the respective stable models of the curves  $X_{K'}, Y_{K'}$ . The proof of Theorem 0.1 is based on the following more general result.

**THEOREM 0.2** (Theorem 4.5). *Let  $f : X \rightarrow Y$  be a finite morphism of smooth geometrically connected projective curves over  $K$ . Let  $\mathcal{X}, \mathcal{Y}$  be respective models of  $X$  and  $Y$  over  $S$ . Then there exists a finite separable extension  $K'$  of  $K$ , and models  $\mathcal{X}', \mathcal{Y}'$  of  $X_{K'}$  and  $Y_{K'}$  over  $S'$  (integral closure of  $S$  in  $K'$ ) such that:*

- (1)  $\mathcal{X}', \mathcal{Y}'$  dominate  $\mathcal{X}_{S'}, \mathcal{Y}_{S'}$ , respectively, and are semi-stable over  $S'$ ;
- (2)  $X_{K'} \rightarrow Y_{K'}$  extends to a finite morphism  $\mathcal{X}' \rightarrow \mathcal{Y}'$ ;

---

Received 14 September 2004, accepted in final form 3 January 2005.

2000 Mathematics Subject Classification 14G20, 14H30, 14D10, 11G20.

Keywords: curves, finite covers, stable reduction.

This journal is © Foundation Compositio Mathematica 2006.

- (3) the pair  $(\mathcal{X}', \mathcal{Y}')$  is minimal: for any pair  $(\mathcal{X}'', \mathcal{Y}'')$  satisfying properties (1) and (2),  $\mathcal{X}''$  and  $\mathcal{Y}''$  dominate  $\mathcal{X}'$  and  $\mathcal{Y}'$ , respectively; the cover  $\mathcal{X}' \rightarrow \mathcal{Y}'$  is called the stable hull of  $\mathcal{X}_{S'} \dashrightarrow \mathcal{Y}_{S'}$ ;
- (4) the formation of  $\mathcal{X}', \mathcal{Y}'$  commutes with flat base change  $T \rightarrow S'$ : if  $T$  is a Dedekind scheme dominating  $S'$ , then  $\mathcal{X}'_T \rightarrow \mathcal{Y}'_T$  is the stable hull of  $\mathcal{X}_T \dashrightarrow \mathcal{Y}_T$ .

The paper is organized as follows. In § 1,  $S = \text{Spec } \mathcal{O}_K$  is local. We discuss some sufficient conditions for a model of  $X_{\hat{K}}$  over  $\hat{\mathcal{O}}_K$  (completion of  $\mathcal{O}_K$ ) to be defined over  $\mathcal{O}_K$ . For example, this is true for semi-stable or regular models. This is used to reduce the proof of Proposition 4.4 to the case of a complete local base  $S$ . In § 2, we prove the special case of Theorem 0.2 when  $X = Y, \mathcal{X} = \mathcal{Y}$  and  $f = \text{Id}$ . Parts (1) and (2) of Theorem 0.2 are proved in § 3. The existence of a stable model (part (3)) is proved in § 4.

**Convention**

Through this paper,  $S$  is a Noetherian regular connected scheme of dimension 1 (Dedekind scheme), and, unless otherwise specified,  $X, Y$  are projective, smooth and geometrically connected curves over  $K := K(S)$ . When we state that a property (P) holds for some model  $\mathcal{X}$  of  $X$  after finite separable extension of  $K$ , this means that there exists a finite separable extension of  $K'/K$ , such that (P) is satisfied for  $\mathcal{X}_{S'}$ , where  $S'$  is the normalization of  $S$  in  $K'$ .

Note that the hypothesis  $X, Y$  geometrically connected (instead of connected) is not serious. Actually,  $X$  is geometrically connected over the finite separable extension  $H^0(X, \mathcal{O}_X)$  of  $K$ .

**1. Descent from the completion**

Let  $\mathcal{O}_K$  be a discrete valuation ring, and let  $X$  be a geometrically connected smooth projective curve  $K$ . We give some sufficient conditions for a model of  $X_{\hat{K}}$  over  $\hat{\mathcal{O}}_K$  to be defined over  $\mathcal{O}_K$ .

LEMMA 1.1. *Let  $\mathcal{O}_K$  be a discrete valuation ring, let  $A$  be a flat local  $\mathcal{O}_K$ -algebra, localization of a finitely generated  $\mathcal{O}_K$ -algebra. Suppose that  $\text{Spec}(A \otimes \hat{K})$  is integral, one-dimensional and smooth over  $\hat{K}$ . Let  $\mathcal{W} \rightarrow \text{Spec } \hat{A}$  be a projective birational morphism. Then under any of the following conditions,  $\mathcal{W}$  and the morphism  $\mathcal{W} \rightarrow \text{Spec } \hat{A}$  are defined over  $A$ :*

- (a)  $\text{Pic}(\hat{A} \otimes \hat{K})$  is a torsion group (i.e. every element has finite order);
- (b)  $\hat{A}$  is  $\mathbb{Q}$ -factorial.

*Proof.* The morphism  $\mathcal{W} \rightarrow \text{Spec}(\hat{A})$  is the blowing-up along a closed subscheme  $V(\mathfrak{J})$  of  $\text{Spec}(\hat{A})$ . Let  $t$  be a uniformizing element of  $\mathcal{O}_K$ . Let us show that  $\mathfrak{J}$  can be chosen in such a way that  $t^n \in \mathfrak{J}$  for some  $n \geq 1$ . Let us suppose that condition (a) is satisfied. Then there exists some  $m \geq 1$  such that the restriction of  $\mathfrak{J}^m$  to the generic fiber is principal, generated by a  $f \in \mathfrak{J}^m \otimes \hat{K}$ . Under condition (b), let  $D$  be the scheme theoretical closure of  $V(\mathfrak{J}) \cap \text{Spec}(\hat{A} \otimes \hat{K})$  in  $\text{Spec}(\hat{A})$ , considered as a Weil divisor. Then  $mD$  is principal for some  $m > 0$ , generated by a  $f \in \mathfrak{J}^m$ .

In both cases, there exist  $a, b \in \mathbb{Z}$  such that  $t^a \mathfrak{J}^m \subseteq f \hat{A} \subseteq t^b \mathfrak{J}^m$ . Replacing  $\mathfrak{J}$  by the ideal  $t^a f^{-1} \mathfrak{J}^m$  (which does not change the blowing-up along  $\mathfrak{J}$ ), we can suppose that the restriction of  $\mathfrak{J}$  to the generic fiber is trivial, and  $t^n \in \mathfrak{J}$  for some  $n \geq 0$  ( $n = a - b$ ). Since  $\mathcal{O}_K$  is dense in  $\hat{\mathcal{O}}_K$ ,  $\mathfrak{J}$  is then generated by an ideal  $I$  of  $A$ . Let  $\mathcal{X} \rightarrow \text{Spec } A$  be the blowing-up along  $V(I)$ , then  $\mathcal{W} \rightarrow \text{Spec}(\hat{A})$  is obtained from  $\mathcal{X} \rightarrow \text{Spec } A$  by the base change  $\hat{\mathcal{O}}_K/\mathcal{O}_K$ . □

*Remark 1.2.* Let  $\mathcal{W}$  be a projective regular model of  $X_{\hat{K}}$  over  $\hat{\mathcal{O}}_K$  dominating  $\mathcal{X}_{\hat{\mathcal{O}}_K}$  for some model  $\mathcal{X}$  of  $X$  over  $\mathcal{O}_K$ . Then  $\mathcal{W}$  and  $\mathcal{W} \rightarrow \mathcal{X}_{\hat{\mathcal{O}}_K}$  are defined over  $\mathcal{O}_K$ . Actually, if  $\tilde{\mathcal{X}} \rightarrow \mathcal{X}$  is the minimal desingularization of  $\mathcal{X}$ , then  $\tilde{\mathcal{X}}_{\hat{\mathcal{O}}_K} \rightarrow \mathcal{X}_{\hat{\mathcal{O}}_K}$  is the minimal desingularization of  $\mathcal{X}_{\hat{\mathcal{O}}_K}$  (same proof as in [Liu02, Proposition 9.3.28]). Hence,  $\mathcal{W} \dashrightarrow \tilde{\mathcal{X}}_{\hat{\mathcal{O}}_K}$  is a birational morphism. It is a sequence of




FIGURE 1. A birational morphism over  $\hat{\mathcal{O}}_K$  not defined over  $\mathcal{O}_K$  (see Remark 1.4).

blowing-ups of closed points [Liu02, 9.2.2]. Hence,  $\mathcal{W}$  is defined over  $\mathcal{O}_K$ , and so is the morphism  $\mathcal{W} \rightarrow \mathcal{X}_{\hat{\mathcal{O}}_K}$  by faithfully flat descent of rational maps.

PROPOSITION 1.3. *Let  $S = \text{Spec } \mathcal{O}_K$  be local. Let  $\mathcal{X}$  be a model of  $X$  over  $S$  dominating some semi-stable or regular model. Let  $\varphi : \mathcal{W} \rightarrow \mathcal{X}_{\hat{\mathcal{O}}_K}$  be a projective birational morphism over  $\hat{\mathcal{O}}_K$ . Then  $\mathcal{W}$  and  $\varphi$  are defined over  $S$ .*

*Proof.* It is enough to show that  $\mathcal{W}$  is defined over  $S$ . We can suppose that  $\mathcal{X}$  itself is semi-stable or regular. The morphism  $\varphi$  is an isomorphism outside of a finite set  $F$  of closed points of  $(\mathcal{X}_{\hat{\mathcal{O}}_K})_s = \mathcal{X}_s$ . Let  $x \in F$ , and let  $A_x = \mathcal{O}_{\mathcal{X},x}$ . If  $\mathcal{X}$  is regular, then  $\hat{A}_x$  is regular and thus factorial. If  $\mathcal{X}$  is semi-stable, there exists a finite (étale) extension  $\mathcal{O}_L/\hat{\mathcal{O}}_K$  such that  $\hat{A}_x \otimes_{\hat{\mathcal{O}}_K} \mathcal{O}_L$  is a finite direct sum of rings  $\mathcal{O}_L[[u]]$  or  $\mathcal{O}_L[[u, v]]/(uv - a)$ ,  $a \in \mathcal{O}_L$ . It is well known that Pic of these rings tensored by  $\hat{L}$  are trivial (see, for instance, [Hen99, Corollary 2.2], for  $\mathcal{O}_L[[u, v]] \otimes \hat{L}/(uv - a)$ ). Hence,  $\text{Pic}(\hat{A}_x)$  is torsion [Liu02, Theorem 7.2.18 and Remark 7.2.19]. By Lemma 1.1,  $\mathcal{W} \times_{\mathcal{X}_{\hat{\mathcal{O}}_K}} \text{Spec } \hat{A}_x \rightarrow \text{Spec } \hat{A}_x$  is defined over  $A_x$ . By glueing the morphisms above  $\text{Spec } A_x$ , when  $x$  varies in  $F$ , and the isomorphism above  $\mathcal{X} \setminus F$ , we find a morphism over  $\mathcal{X}$  which is equal to  $\varphi$  when base changed to  $\hat{\mathcal{O}}_K$ .  $\square$

Remark 1.4. In general, not every birational projective morphism  $\mathcal{W} \rightarrow \mathcal{X}_{\hat{\mathcal{O}}_K}$  is defined over  $\mathcal{O}_K$ . In other words, even if  $\mathcal{W} \rightarrow \mathcal{X}_{\hat{\mathcal{O}}_K}$  is an isomorphism outside of the special fiber, it is not necessarily the blowing-up along a closed subscheme with support in the special fiber. To construct such a counterexample, we will imitate the example of a non-contractible component given in [BLR90, Lemma 6.7.6]. Let us consider the smooth elliptic curve  $\mathcal{E}/\mathcal{O}_K$  and the point  $a_k \in \mathcal{E}_k(k)$  as given in [BLR90, p. 171]. The point  $a_k$  satisfies the property that no multiple (in the sense of the group law on  $\mathcal{E}$ )  $na_k$ ,  $n > 0$ , can be lifted to a section in  $\mathcal{E}(\mathcal{O}_K)$ .

Let  $\mathcal{X}' \rightarrow \mathcal{E}$  be the blowing-up of  $\mathcal{E}$  along  $\{a_k, e_k\}$ , where  $e$  is the unit section of  $\mathcal{E}$ . Let  $P_a$  and  $P_e$  be the respective inverse images of  $a_k, e_k$  in  $\mathcal{X}'$ . They are projective lines over  $k$ . Let  $\tilde{\mathcal{E}}_k$  be the strict transform of  $\mathcal{E}_k$  in  $\mathcal{X}'$ . The unit section of  $E_K$  gives rise to a section of  $\mathcal{X}'$  which meets  $\mathcal{X}'_k$  at an interior point of  $P_e$ . Hence, there exists a contraction map  $\mathcal{X}' \rightarrow \mathcal{X}$  of  $P_a \cup \tilde{\mathcal{E}}_k$  (see [BLR90, Corollary 6.7.3]). Now over  $\hat{\mathcal{O}}_K$ , we can contract  $\tilde{\mathcal{E}}_k$  in  $\mathcal{X}'_{\hat{\mathcal{O}}_K}$  (see [BLR90, 6.7.4]), which gives a model  $\mathcal{W}/\hat{\mathcal{O}}_K$  and we have birational projective morphisms  $\mathcal{X}'_{\hat{\mathcal{O}}_K} \rightarrow \mathcal{W} \rightarrow \mathcal{X}_{\hat{\mathcal{O}}_K}$ . (See Figure 1.) We want to show that  $\mathcal{W}$  is not defined over  $\mathcal{O}_K$ , or, equivalently, that  $\tilde{\mathcal{E}}_k$  cannot be contracted over  $\mathcal{O}_K$ . Suppose that  $\mathcal{W}$  exists over  $\mathcal{O}_K$ , then there exists a relative effective Cartier divisor  $D$  on  $\mathcal{X}'$  such that  $\text{Supp } D$  meets  $P_a$  and  $P_e$  but not  $\tilde{\mathcal{E}}_k$ . Taking the Zariski closure of  $D_K$  in  $\mathcal{E}$  gives a relative effective Cartier divisor  $D'$  on  $\mathcal{E}$  such that  $D'_k = na_k + me_k$  for some integers  $n, m > 0$ . However,  $D' - (n + m)e \in \text{Pic}^0_{\mathcal{E}/\mathcal{O}_K}(\mathcal{O}_K) \simeq \mathcal{E}(\mathcal{O}_K)$  then defines a section  $b \in \mathcal{E}(\mathcal{O}_K)$  such that  $D' - (n + m)e \sim b - e$ . In the special fiber we then have  $n(a_k - e_k) \sim b_k - e_k$ ; hence  $na_k = b_k$  in the group  $\mathcal{E}_k(k)$ , contradicting the assumption on  $a_k$ .

LEMMA 1.5. *Suppose that  $X$  has semi-stable reduction over  $S$ . Then every relatively minimal semi-stable model  $\mathcal{W}$  of  $X_{\hat{K}}$  over  $\hat{\mathcal{O}}_K$  is defined over  $\mathcal{O}_K$ .*

*Proof.* If  $g(X) \geq 2$ , then the unique relatively minimal semi-stable model of  $X$  is the stable model of  $X$  over  $S$ . Since the stable model is unique and commutes with flat base change [Liu02, 10.3.36],  $\mathcal{W}$  over  $\hat{\mathcal{O}}_K$  is defined over  $\mathcal{O}_K$ . The same is true if  $g(X) = 1$  and  $X$  has good reduction.

Suppose that  $g(X) = 1$  and  $X$  has multiplicative reduction. Let  $\mathcal{X}$  be a relatively minimal semi-stable model. Let  $\tilde{\mathcal{X}}$  be its minimal desingularization. Let us show that  $\tilde{\mathcal{X}}$  is the minimal regular model of  $X$  over  $S$ . Let  $\Gamma$  be an irreducible component of  $\mathcal{X}_s$  and let  $\tilde{\Gamma}$  be its strict transform in  $\tilde{\mathcal{X}}$ . By Lemma 2.13(a),  $\deg \omega_{\mathcal{X}/S}|_{\Gamma} > 0$  if  $\mathcal{X}_s$  is reducible. If  $\mathcal{X}_s$  is irreducible, then  $\deg \omega_{\mathcal{X}/S}|_{\Gamma} = 2g(X) - 2 = 0$ . So in any case  $\deg \omega_{\tilde{\mathcal{X}}/S}|_{\tilde{\Gamma}} = \deg \omega_{\mathcal{X}/S}|_{\Gamma} \geq 0$ . Hence,  $\tilde{\mathcal{X}}$  has no exceptional divisor. Since any strict subset of the set of irreducible components of  $\tilde{\mathcal{X}}_s$  can be contracted [Liu02, Example 9.4.19] into a semi-stable model (Lemma 2.13(a)),  $\mathcal{X}_s$  is irreducible. So the relatively minimal semi-stable models of  $X$  correspond bijectively to the irreducible components of the minimal regular model of  $X$ . The same is true over  $\hat{\mathcal{O}}_K$ . Since the minimal regular model commutes with the base change  $\hat{\mathcal{O}}_K/\mathcal{O}_K$  (see [Liu02, 9.3.28]), we see that the relatively minimal semi-stable models of  $X_{\hat{K}}$  are exactly those of  $X$  base changed to  $\hat{\mathcal{O}}_K$ .

Suppose that  $g(X) = 0$ . Let  $\mathcal{X}$  be a relatively minimal semi-stable model of  $X$  over  $S$  and let  $\Gamma_1, \dots, \Gamma_n$  be the irreducible components of  $\mathcal{X}_s$ . Then

$$\sum_i \deg \omega_{\mathcal{X}/S}|_{\Gamma_i} = 2g(X) - 2 < 0.$$

So  $\omega_{\mathcal{X}/S}$  has negative degree on at least one component  $\Gamma_i$ . By Lemma 2.13(a),  $n = 1$ , thus  $\mathcal{X}_s$  is irreducible, semi-stable and has arithmetic genus 0. So  $\mathcal{X}_s$  is smooth. The same reasoning shows that  $\mathcal{W}$  is smooth. Let  $\mathcal{L}$  be the dual of the dualizing sheaf on  $\mathcal{W}$ . Let  $s_0, s_1 \in H^0(\mathcal{W}, \mathcal{L})$  be a basis with  $s_i \in H^0(X, \omega_{X/K}^\vee)$ . Then  $s_0, s_1$  define a closed immersion  $\mathcal{W} \rightarrow \mathbb{P}^2$  over  $\hat{\mathcal{O}}_K$ . Its image is a conic defined by a polynomial with coefficients in  $K$ . Hence,  $\mathcal{W}$  is defined over  $\mathcal{O}_K$ . □

The next corollary is weaker than Proposition 1.3, but sufficient for the purpose of Theorem 4.5. It is an immediate consequence of Proposition 1.3 and Lemma 1.5. However, we give a direct proof without using Proposition 1.3.

COROLLARY 1.6. *Let  $S = \text{Spec } \mathcal{O}_K$  be local. Let  $X$  be a smooth geometrically connected projective curve over  $K$ . Then every semi-stable model  $\mathcal{W}$  of  $X_{\hat{K}}$  over  $\hat{\mathcal{O}}_K$  is defined over  $\mathcal{O}_K$ .*

*Proof.* The model  $\mathcal{W}$  dominates a relatively minimal semi-stable model  $\mathcal{Z}$  of  $X_{\hat{K}}$  which is then defined over  $\mathcal{O}_K$  by Lemma 1.5. Let  $\tilde{\mathcal{W}} \rightarrow \mathcal{W}$  be the minimal desingularization of  $\mathcal{W}$ . Then  $\tilde{\mathcal{W}} \rightarrow \mathcal{Z}$  is defined over  $\mathcal{O}_K$  (1.2). The irreducible components of  $\tilde{\mathcal{W}}_s$  in the exceptional locus of  $\tilde{\mathcal{W}} \rightarrow \mathcal{W}$  are  $(-2)$ -curves and can be contracted over  $\mathcal{O}_K$  (Lemma 2.13(a)). Hence,  $\mathcal{W}$  is defined over  $\mathcal{O}_K$ . □

Remark 1.7. If  $X = \mathbb{P}_K^1$ , then every normal model  $\mathcal{W}$  of  $X_{\hat{K}}$  is defined over  $\mathcal{O}_K$ . Indeed, if  $\tilde{\mathcal{W}}$  is the minimal desingularization of  $\mathcal{W}$ , then  $\tilde{\mathcal{W}}$  dominates a relatively minimal regular model. The latter is smooth and defined over  $\mathcal{O}_K$ . Hence,  $\tilde{\mathcal{W}}$  is defined over  $\mathcal{O}_K$ . Now every strict subset of the set of irreducible components of  $\tilde{\mathcal{W}}_s$  is contractible over  $\mathcal{O}_K$  (see [Liu02, Exercise 9.4.5]). So  $\mathcal{W}$  is defined over  $\mathcal{O}_K$ .

## 2. Stable hull of a model

DEFINITION 2.1. Let  $S$  be a connected Noetherian regular scheme of dimension 1 (i.e. a *Dedekind scheme*). Let  $X$  be an integral projective variety over  $K$ . A *model  $\mathcal{X}$  of  $X$  over  $S$*  is an integral

projective scheme over  $S$  whose generic fiber is isomorphic to  $X$ . Recall that  $\mathcal{X}$  is said to be *semi-stable* if its geometric fibers are reduced with only ordinary double points as singularities. A morphism of models is defined in an obvious way.

DEFINITION 2.2. Let  $X$  be a connected projective smooth curve over  $K$ , and let  $\mathcal{X}$  be a model of  $X$  over  $S$ . A *stable hull* of  $\mathcal{X}$  is a semi-stable model  $\mathcal{W}$  of  $X$  dominating  $\mathcal{X}$ , and minimal for these properties (i.e. every semi-stable model dominating  $\mathcal{X}$  dominates  $\mathcal{W}$ ).

The aim of this section is to prove the following result.

THEOREM 2.3. *Let  $X$  be a geometrically connected projective smooth curve over  $K$  and let  $\mathcal{X}$  be a model of  $X$  over  $S$ .*

- (a) *The stable hull of  $\mathcal{X}$  is unique (up to isomorphism) when it exists. In general, there exists a finite separable extension  $K'/K$  such that  $\mathcal{X}_{S'}$  (where  $S'$  is the integral closure of  $S$  in  $K'$ ) has a stable hull over  $S'$ .*
- (b) *The stable hull commutes with flat base change: suppose that  $\mathcal{X}$  admits a stable hull  $\mathcal{W}$  over  $S$  and let  $S' \rightarrow S$  be a flat morphism of Dedekind schemes, then  $\mathcal{W}_{S'}$  is the stable hull of  $\mathcal{X}_{S'}$  over  $S'$ .*

The proof of the theorem is postponed to § 2.17.

LEMMA 2.4. *Let  $G$  be a finite group acting on  $X$ . Let  $\mathcal{X}$  be a model of  $X$  over  $S$ . Then there exists a model  $\mathcal{Z}$  of  $X$  dominating  $\mathcal{X}$ , endowed with an action of  $G$ , and minimal for these properties.*

*Proof.* (See [Dej96, 7.6].) Let  $\sigma \in G$ , then there exists a model  $\mathcal{X}^\sigma$  such that  $\sigma : X \rightarrow X$  extends to an isomorphism  $\sigma : \mathcal{X} \rightarrow \mathcal{X}^\sigma$ . If  $\tau \in G$ , by composing  $(\sigma^{-1}\tau) : \mathcal{X}^{\tau^{-1}\sigma} \rightarrow \mathcal{X}$  with  $\sigma : \mathcal{X} \rightarrow \mathcal{X}^\sigma$ , we obtain an isomorphism  $\mathcal{X}^{\tau^{-1}\sigma} \rightarrow \mathcal{X}^\sigma$  denoted by  $\tau$ . Let  $\mathcal{P}$  be the fiber product  $\prod_{S, \sigma \in G} \mathcal{X}^\sigma$  over  $S$ . Then we can make  $G$  act on  $\mathcal{P}$  by

$$\tau : (x_\sigma)_\sigma \mapsto (\tau(x_{\tau^{-1}\sigma}))_\sigma.$$

Moreover, the diagonal morphism  $\Delta : X \rightarrow \mathcal{P}_K, x \mapsto (x, \dots, x)$  is  $G$ -equivariant. Let  $\mathcal{Z}$  be the Zariski closure of  $\Delta(X)$  in  $\mathcal{P}$ , endowed with the reduced structure. Then  $G$  leaves stable  $\mathcal{Z}$ . Note that  $\mathcal{Z}$  dominates  $\mathcal{X}$  because the projection morphism  $\mathcal{P} \rightarrow \mathcal{X}$  induces a morphism  $\mathcal{Z} \rightarrow \mathcal{X}$  which is an isomorphism on the generic fibers.

Let us prove that  $\mathcal{Z}$  is minimal. Let  $\mathcal{W}$  be a model endowed with an action of  $G$  and a birational morphism  $\mathcal{W} \rightarrow \mathcal{X}$ . Then we have a morphism  $\mathcal{W} \rightarrow \mathcal{X}^\sigma$  for all  $\sigma$  and, hence, a morphism  $h : \mathcal{W} \rightarrow \mathcal{P}$ . Since  $h(\mathcal{W})$  is irreducible and contains  $\mathcal{Z}$ ,  $h$  induces a morphism of models  $\mathcal{W} \rightarrow \mathcal{Z}$ . □

Let us give a corollary of Theorem 2.3.

COROLLARY 2.5. *Let  $G$  be a finite group acting on  $X$ . Let  $\mathcal{X}$  be a model of  $X$  over  $S$ . Then after finite separable extension of  $K$ ,  $\mathcal{X}$  is dominated by a semi-stable (respectively, semi-stable and regular) model  $\mathcal{W}$  such that the action of  $G$  extends to  $\mathcal{W}$ . Moreover, there exists a minimal such a model  $\mathcal{W}$ .*

*Proof.* Let  $\mathcal{Z}$  be the model defined in Lemma 2.4. Let  $\mathcal{W}$  be the stable hull of  $\mathcal{Z}_{S'}$  over some  $S'/S$  (Theorem 2.3(a)). By the uniqueness property, the action of  $G$  on  $\mathcal{Z}_{S'}$  extends to  $\mathcal{W}$ . It is clear that  $\mathcal{W}$  is minimal with respect to the required properties. To have a minimal semi-stable regular model, it is enough to take the minimal desingularization of  $\mathcal{W}$ . □

LEMMA 2.6. *Let  $S$  be local with separably closed residue field. Let  $\mathcal{X}$  be a model of  $X$  with minimal desingularization  $\tilde{\mathcal{X}} \rightarrow \mathcal{X}$ . Suppose that  $\tilde{\mathcal{X}}$  dominates a regular model  $\mathcal{Z}$  and that  $\mathcal{X}_s, \mathcal{Z}_s$  are geometrically reduced. Then  $\tilde{\mathcal{X}}_s$  is geometrically reduced.*

*Proof.* Note that if  $\mathcal{W}$  is a normal model of  $X$ , then  $\mathcal{W}_s$  verifies the property  $(S_1)$ , thus  $\mathcal{W}_s$  is geometrically reduced if and only if every irreducible component of  $\mathcal{W}_s$  has geometric multiplicity [BLR90, Definition 9.1.3] equal to 1 in  $\mathcal{W}_s$ . The latter condition depends only on the generic points of  $\mathcal{W}_s$ . We can decompose  $\tilde{\mathcal{X}} \rightarrow \mathcal{Z}$  into a sequence of blowing-ups

$$\tilde{\mathcal{X}} =: \mathcal{Z}_0 \rightarrow \mathcal{Z}_1 \rightarrow \cdots \rightarrow \mathcal{Z}_n := \mathcal{Z}$$

such that  $\mathcal{Z}_i \rightarrow \mathcal{Z}_{i+1}$  consists of blowing-down an exceptional divisor  $\Theta_i$  contained in  $\mathcal{Z}_i$ . We will show by induction that  $\Theta_i$  has multiplicity 1 in  $(\mathcal{Z}_i)_s$  and  $\Theta_i \simeq \mathbb{P}_{k(s)}^1$ . Since  $(\mathcal{Z}_n)_s$  is geometrically reduced, this will imply that  $\tilde{\mathcal{X}}_s$  is geometrically reduced.

By the minimality of  $\tilde{\mathcal{X}} \rightarrow \mathcal{X}$ ,  $\Theta_0$  is not mapped to a closed point in  $\mathcal{X}$ . Thus,  $\tilde{\mathcal{X}} \rightarrow \mathcal{X}$  is an isomorphism in a neighborhood of the generic point of  $\Theta_0$ . In particular,  $\Theta_0$  has geometric multiplicity 1. Since  $\Theta_0$  is a projective line by the Castelnuovo criterion, it is isomorphic to  $\mathbb{P}_{k(s)}^1$ . Suppose that the same holds for  $\Theta_j$ ,  $0 \leq j \leq i - 1$ . Let  $\Theta_{i,j}$  be the strict transform of  $\Theta_i$  in  $\mathcal{Z}_j$ ,  $0 \leq j \leq i - 1$ . If  $\Theta_{i,j}$  meets  $\Theta_j$  for some  $j$ , then the computation of  $\Theta_j^2 = -1$  shows that  $\Theta_{i,j}$  has multiplicity 1 and cuts  $\Theta_j$  at a rational point. Thus,  $\Theta_i$  has multiplicity 1 and is isomorphic to  $\mathbb{P}_{k(s)}^1$ . Otherwise,  $\mathcal{Z}_0 \rightarrow \mathcal{Z}_i$  is an isomorphism in a neighborhood of  $\Theta_{i,0}$ . In particular,  $\Theta_{i,0}$  is an exceptional divisor. Then we can conclude exactly as for  $\Theta_0$ .  $\square$

PROPOSITION 2.7. *Let  $\mathcal{X}$  be a normal model of  $X$  over  $S$ . Then the following properties are equivalent:*

- (i)  $\mathcal{X}$  is dominated by a semi-stable model over  $S$ ;
- (ii)  $X$  admits a semi-stable model over  $S$  and  $\mathcal{X}_s$  is geometrically reduced for all  $s \in S$ ;
- (iii) the minimal desingularization  $\tilde{\mathcal{X}}$  of  $\mathcal{X}$  is semi-stable over  $S$ .

*Proof.* (i)  $\implies$  (ii) If  $\mathcal{X}$  is dominated by a semi-stable model  $\mathcal{X}'$ , then any irreducible component  $\Gamma$  of  $\mathcal{X}_s$  is birational to an irreducible component of  $\mathcal{X}'_s$ . The latter being geometrically reduced,  $\mathcal{X}_s$  is geometrically reduced.

(ii)  $\implies$  (iii) We know that  $\tilde{\mathcal{X}}$  dominates a relatively minimal regular model  $\mathcal{Z}$ . The semi-stable reduction hypothesis implies that  $\mathcal{Z}$  is semi-stable (see [Liu02, 10.3.34(a)] if  $g(X) \geq 1$ ; if  $g(X) \leq 0$ , then  $\mathcal{Z}$  is smooth). Since the minimal desingularization commutes with étale base change (see the proof of [Liu02, Proposition 9.3.28]), we can suppose that  $S$  is local with separably closed residue field. By Lemma 2.6,  $\tilde{\mathcal{X}}_s$  is geometrically reduced. The map  $\tilde{\mathcal{X}} \rightarrow \mathcal{Z}$  consists of blowing-up successively closed points. The fact that  $\tilde{\mathcal{X}}_s$  is geometrically reduced implies that we only blow-up rational points in the smooth locus. Since  $\mathcal{Z}$  is semi-stable, then so is  $\tilde{\mathcal{X}}$ .  $\square$

COROLLARY 2.8. *There exists a finite separable extension  $K'/K$  such that  $\mathcal{X}_{S'}$ , where  $S'$  is the normalization of  $S$  in  $K'$ , is dominated by a semi-stable model of  $X_{K'}$ .*

*Proof.* We can suppose that  $X$  has semi-stable reduction over  $S$ . Since  $X$  has good reduction over an open dense subset of  $S$ , we can suppose that  $S$  is local. By the finiteness theorem of Grauert–Remmert [GR66] (see also [BLR95, Theorem 1.3]) applied to the formal completion of  $\mathcal{X}$  along its special fiber, there exists a finite Galois extension  $L/\hat{K}$  such that the normalization of  $\mathcal{X}_{\mathcal{O}_L}$  has geometrically reduced special fiber. See [Epp73, p. 247] for how to descend the result to  $K$  (note that [Kuh03] fills a gap in the proof of a main theorem in [Epp73]). We then apply Proposition 2.7.  $\square$

Remark 2.9. The corollary is useful in a recent work of Deninger and Werner on vector bundles and representations of the fundamental group of  $p$ -adic curves [DW04]. In fact, de Jong [Dej97, Theorem 2.4] already proved it in the situation when  $S$  is a Noetherian integral excellent scheme of any dimension, and when  $X$  is an integral curve over  $K(S)$ . The scheme  $S'$  is then proper and generically finite over  $S$ . The proof here for one-dimensional  $S$  is simpler and more effective in some sense.

*Remark 2.10.* When  $S$  is local and complete, the corollary can be reformulated in terms of rigid analytic geometry as follows: let  $\mathcal{U}$  be a formal covering of  $X$ . Then after finite separable extension of  $K$ ,  $\mathcal{U}$  can be refined to a distinguished formal covering  $\mathcal{V}$  with semi-stable reduction. As such, the statement can be easily worked out using Theorem 5.5, and step 2 in the proof of Lemma 7.3 of [BL85]. The non-complete case can then be obtained using Proposition 1.3.

*Remark 2.11* (Effective reduced fiber theorem). In the case of Corollary 2.8, we can give an effective method to eliminate the multiplicities of  $\mathcal{X}_s$ , without using Grauert–Remmert’s theorem. Suppose that  $X$  has semi-stable reduction. Let  $\Gamma$  be an irreducible component of  $\mathcal{X}_s$  of geometric multiplicity  $d > 1$ . If we choose two closed points  $P_1, P_2$  of  $X$  which specialize to two distinct points in the interior of  $\Gamma$ , and if we take  $L = K(P_1, P_2)$ , then the irreducible components of  $(\mathcal{X}_{\mathcal{O}_L})'_s$  (where  $(\mathcal{X}_{\mathcal{O}_L})'$  denotes the normalization of  $\mathcal{X}_{\mathcal{O}_L}$ ) lying above  $\Gamma$  are geometrically reduced. If  $K$  is strictly Henselian, we can bound  $[L : K]$  by  $d^2$ . Note that  $L$  can be chosen to be separable over  $K$ . If  $X$  has not necessarily semi-stable reduction, then we can bound  $[L : K]$  by the max of  $d^2$  and a constant depending only on  $g$ .

**2.12 The stable hull**

Now let us construct a minimal semi-stable model dominating  $\mathcal{X}$ . Let  $\mathcal{Z}$  be a locally complete intersection (e.g. regular or semi-stable) model of  $X$  over  $S$ . Let  $\omega_{\mathcal{Z}/S}$  be the (invertible) dualizing sheaf of  $\mathcal{Z}/S$ . Recall that a  $(-2)$ -curve on  $\mathcal{Z}$  is an irreducible component  $\Gamma$  of a closed fiber  $\mathcal{Z}_s$  such that  $\deg \omega_{\mathcal{Z}/S}|_{\Gamma} = 0$ . If  $\mathcal{Z}$  is semi-stable and  $k(s)$  is algebraically closed, and  $\Gamma$  is not a connected component of  $\mathcal{Z}_s$ , then this is equivalent to  $\Gamma \simeq \mathbb{P}^1_{k(s)}$  and  $\Gamma$  meets the other irreducible components at exactly two points. Recall that the *exceptional locus* of a birational projective morphism  $\pi : \mathcal{Z} \rightarrow \mathcal{X}$  is by definition the complementary of  $\pi^{-1}(U)$ , where  $U$  is the largest open subscheme of  $\mathcal{X}$  such that  $\pi^{-1}(U) \rightarrow U$  is an isomorphism. When  $\mathcal{X}$  is normal, the exceptional locus is equal to the union of the prime divisors of  $\mathcal{Z}$  which map to closed points in  $\mathcal{X}$ . A semi-stable model  $\mathcal{Z}$  dominating  $\mathcal{X}$  will be called *relatively minimal* if there is no semi-stable model between  $\mathcal{X}$  and  $\mathcal{Z}$ , except  $\mathcal{Z}$  itself.

LEMMA 2.13. *Let  $\mathcal{Z}$  be a semi-stable model of  $X$  over  $S$ . Let  $V$  be an effective vertical divisor on  $\mathcal{Z}$  such that for all  $s \in S$ , no connected component of  $\mathcal{Z}_s$  is contained in  $V$ .*

- (a) *If  $\deg \omega_{\mathcal{Z}/S}|_{\Gamma} \leq 0$  for all components  $\Gamma$  of  $V$ , then there exists a contraction map  $\mathcal{Z} \rightarrow \mathcal{W}$  of  $V$  and  $\mathcal{W}$  is semi-stable.*
- (b) *If there exists a contraction map  $\mathcal{Z} \rightarrow \mathcal{W}$  of  $V$  with  $\mathcal{W}$  semi-stable, then  $\deg \omega_{\mathcal{Z}/S}|_{\Gamma} \leq 0$  for at least one irreducible component  $\Gamma$  of  $V$ .*

*Proof.* (a) This is well known but we were not able to find a proper reference. We can suppose that  $S$  is local. Let  $\rho : \tilde{\mathcal{Z}} \rightarrow \mathcal{Z}$  be the minimal desingularization. Then the components  $\Theta$  of the exceptional locus  $E$  of  $\rho$  are  $(-2)$ -curves. Let  $\tilde{V}$  be the strict transform of  $V$  in  $\tilde{\mathcal{Z}}$ . Let us show that  $V' := E + \tilde{V}$  can be contracted. We have  $\omega_{\tilde{\mathcal{Z}}/S} = \rho^* \omega_{\mathcal{Z}/S}$  because  $\mathcal{Z}$  is semi-stable. Hence,  $\deg \omega_{\tilde{\mathcal{Z}}/S}|_{\Gamma'} \leq 0$  for all components  $\Gamma'$  of  $V'$ . If there exists a  $\Gamma'$  such that  $\deg \omega_{\tilde{\mathcal{Z}}/S}|_{\Gamma'} < 0$ , then  $\Gamma'$  is an exceptional divisor. Let  $\pi : \tilde{\mathcal{Z}} \rightarrow \mathcal{Z}'$  be the contraction of  $\Gamma'$ . Then  $\omega_{\tilde{\mathcal{Z}}/S} = \pi^* \omega_{\mathcal{Z}'/S}(\Gamma')$ . We deduce easily that  $\deg \omega_{\mathcal{Z}'/S}|_{\Gamma''} \leq 0$  for all  $\Gamma''$  in  $\pi(V')$ . So by successively blowing-down exceptional divisors, we can suppose that  $V'$  consists only of  $(-2)$ -curves. By Artin’s criterion of contractibility [Liu02, Corollary 9.4.7],  $V'$  can be contracted. Therefore,  $V$  can be contracted.

It remains to see that  $\mathcal{W}$  is semi-stable. Let  $\mathcal{O}_{K'}$  be a discrete valuation ring containing  $\mathcal{O}_S$  and let  $S' = \text{Spec } \mathcal{O}_{K'}$ . It is enough to show that  $\mathcal{W}_{S'}$  is semi-stable. The map  $\mathcal{Z}_{S'} \rightarrow \mathcal{W}_{S'}$  is the contraction of  $V_{k(s')}$ . We have  $\omega_{\mathcal{Z}_{S'}/S'} = \omega_{\mathcal{Z}/S} \otimes \mathcal{O}_{S'}$ . If  $\Gamma'$  is a component of  $\mathcal{Z}_{s'}$  lying over  $\Gamma \subseteq V$ , then

$$[k(s') : k(s)] \deg_{k(s')} \omega_{\mathcal{Z}_{S'}/S'}|_{\Gamma'} = [k(\Gamma') : k(\Gamma)] \deg_{k(s)} \omega_{\mathcal{Z}/S}|_{\Gamma} \leq 0.$$

So we can reduce the lemma to the case  $k(s)$  algebraically closed. Let  $\Gamma \subseteq V$ . Then  $p_a(\Gamma) = 0$ ,  $\Gamma \simeq \mathbb{P}^1_{k(s)}$ , and  $\Gamma$  meets the other components of  $\mathcal{Z}_s$  in at most two points. Now it is well known that  $\mathcal{W}_s$  is semi-stable (see, for instance, [Liu02, Lemma 10.3.31]).

(b) The previous computations show that  $\deg \omega_{\mathcal{Z}/S}|_\Gamma = \deg \omega_{\tilde{\mathcal{Z}}/S}|_{\tilde{\Gamma}}$  for any irreducible component  $\Gamma$  of  $\mathcal{Z}_s$ . Let  $\tilde{\mathcal{W}}$  be the minimal desingularization of  $\mathcal{W}$ . Suppose that  $\tilde{\mathcal{Z}} \rightarrow \tilde{\mathcal{W}}$  is not an isomorphism. Let  $\Gamma$  be a component of  $\mathcal{Z}_s$  whose strict transform in  $\tilde{\mathcal{Z}}$  is an exceptional divisor contracted into a closed point in  $\tilde{\mathcal{W}}$ . Then  $\Gamma \subseteq V$  and  $\deg \omega_{\mathcal{Z}/S}|_\Gamma = \deg \omega_{\tilde{\mathcal{Z}}/S}|_{\tilde{\Gamma}} < 0$ . If  $\tilde{\mathcal{Z}} = \tilde{\mathcal{W}}$ , then  $\deg \omega_{\mathcal{Z}/S}|_\Gamma = \deg \omega_{\tilde{\mathcal{W}}/S}|_{\tilde{\Gamma}} = 0$  for all  $\Gamma$  in  $V$  because  $\tilde{\Gamma}$  (the strict transform of  $\Gamma$ ) is a  $(-2)$ -curve in  $\tilde{\mathcal{W}}$ . □

PROPOSITION 2.14. *Let  $\mathcal{X}$  be a model of  $X$  over  $S$  dominated by a semi-stable model.*

- (a) *A semi-stable model  $\mathcal{Z}$  dominating  $\mathcal{X}$  is relatively minimal if and only if for all irreducible components of the exceptional locus of  $\mathcal{Z} \rightarrow \mathcal{X}$ , we have  $\deg \omega_{\mathcal{Z}/S}|_\Gamma > 0$ .*
- (b) *Let  $\tilde{\mathcal{X}} \rightarrow \mathcal{X}$  be the minimal desingularization of  $\mathcal{X}$ . Let  $\tilde{\mathcal{X}} \rightarrow \mathcal{W}$  be the contraction of the  $(-2)$ -curves contained in the exceptional locus of  $\tilde{\mathcal{X}} \rightarrow \mathcal{X}$ . Then  $\mathcal{W}$  is the stable hull of  $\mathcal{X}$ .*

*Proof.* (a) This is an immediate consequence of Lemma 2.13.

(b) By Lemma 2.13(a),  $\mathcal{W}$  is semi-stable and dominates  $\mathcal{X}$ . Let  $\mathcal{Z}$  be a semi-stable model dominating  $\mathcal{X}$ , and relatively minimal. Let us first show that  $\tilde{\mathcal{X}}$  dominates  $\mathcal{Z}$ . Let  $\tilde{\mathcal{Z}}$  be the minimal desingularization of  $\mathcal{Z}$ . Then  $\tilde{\mathcal{Z}}$  dominates  $\tilde{\mathcal{X}}$ . Suppose that  $\tilde{\mathcal{Z}} \rightarrow \tilde{\mathcal{X}}$  is not an isomorphism. Let  $\Theta$  be an exceptional divisor of  $\tilde{\mathcal{Z}}$  mapped to a closed point of  $\tilde{\mathcal{X}}$ . Since  $\tilde{\mathcal{Z}} \rightarrow \mathcal{Z}$  is minimal,  $\Theta$  maps to an irreducible component  $\Gamma$  of  $\mathcal{Z}_s$  which is then contained in the exceptional locus of  $\mathcal{Z} \rightarrow \mathcal{X}$ . We have  $\deg \omega_{\mathcal{Z}/S}|_\Gamma = \deg \omega_{\tilde{\mathcal{Z}}/S}|_\Theta < 0$  – contradiction. Therefore,  $\tilde{\mathcal{Z}} \simeq \tilde{\mathcal{X}}$  and  $\tilde{\mathcal{X}} \rightarrow \mathcal{Z}$  consists in contracting some  $(-2)$ -curves in  $\mathcal{X}_s$ . Hence,  $\mathcal{Z}$  dominates  $\mathcal{W}$ . □

Remark 2.15. A *Du Val model* of  $X$  over  $S$  is a model  $\mathcal{X}$  such that  $\deg \omega_{\tilde{\mathcal{X}}/S}|_\Gamma = 0$  where  $\tilde{\mathcal{X}} \rightarrow \mathcal{X}$  is the minimal desingularization and where  $\Gamma$  is any irreducible component of the exceptional locus of  $\tilde{\mathcal{X}} \rightarrow \mathcal{X}$ . The above results (Lemma 2.13 and Proposition 2.14) still hold when ‘semi-stable’ is replaced by ‘Du Val’, except the base change property. The point is that Du Val models do not commute with base change.

Remark 2.16. Let  $\mathcal{W}$  be semi-stable and dominating  $\mathcal{X}$ . Then  $\mathcal{W}$  is the stable hull of  $\mathcal{X}$  if and only if  $\text{Aut}_{\mathcal{X}_s}(\mathcal{W}_s)$  is finite for all  $s \in S$ .

### 2.17 Proof of Theorem 2.3

(a) This is contained in Corollary 2.8 and Proposition 2.14.

(b) It is enough to show that  $\mathcal{W}_{S'}$  is relatively minimal. Let  $\Gamma'$  be an irreducible component of  $\mathcal{W}_{S'}$  contained in the exceptional locus of  $\mathcal{W}_{S'} \rightarrow \mathcal{X}_{S'}$ . The image of  $\Gamma'$  in  $\mathcal{W}$  is in the exceptional locus of  $\mathcal{W} \rightarrow \mathcal{X}$ . Similarly to the proof of Lemma 2.13(a), we have  $\deg \omega_{\mathcal{W}_{S'}/S'}|_{\Gamma'} > 0$ . Therefore,  $\mathcal{W}_{S'}$  is relatively minimal by Proposition 2.14(a). □

Remark 2.18. Suppose that  $g(X) \geq 1$  and  $S$  is affine. Let  $\mathcal{X}$  be the minimal regular model of  $X$  over  $S$  and let  $\mathcal{W}$  be the stable hull of  $\mathcal{X}_{S'}$  over some extension  $S'/S$ . Let  $\mathcal{X}'$  be the stable or minimal regular model of  $X_{K'}$  over  $S'$ . Then  $H^0(\mathcal{W}, \omega_{\mathcal{W}/S'}) = H^0(\mathcal{X}', \omega_{\mathcal{X}'/S'})$ . One should be able to recover some arithmetic information on  $\mathcal{X}$  from the sheaf  $\omega_{\mathcal{W}/S'} \otimes ((\omega_{\mathcal{X}/S})^\vee \otimes \mathcal{O}_{S'})$ . Let us consider the ideal

$$\mathcal{O}_{S'}(-\delta) := H^0(\mathcal{W}, \omega_{\mathcal{W}/S'}) \otimes (H^0(\mathcal{X}, \omega_{\mathcal{X}/S})^\vee \otimes \mathcal{O}_{S'}).$$

For example, if  $X$  is an elliptic curve over  $K$ , then for every closed point  $s \in S$ , we can show that

$$12 \text{ord}_s(\delta) = \text{ord}_s(\Delta) + a_s \text{ord}_s(j),$$



where  $\Delta$  is the minimal discriminant divisor of  $X$  over  $S$ , and  $a_s = 0$  if  $X$  has potentially good reduction at  $s$ ,  $a_s = 1$  otherwise.

**2.19 Marked curves**

Recall that a (proper) *marked curve*  $Z \rightarrow T$  over a scheme  $T$  is a proper flat scheme of relative dimension 1 over  $T$  endowed with a finite set  $M \subset Z(T)$  of sections with pairwise disjoint supports contained in the smooth locus of  $Z/T$  (for our purpose, it is not necessary to order these sections). Note that if  $T$  is irreducible with generic point  $\xi$ , then  $M$  is determined by its generic fiber  $M \cap Z_\xi$ . We say that  $(Z, M)$  is *semi-stable* if  $Z \rightarrow T$  is semi-stable. We say that  $(Z, M)$  is *stable* if it is semi-stable and if for any geometric point  $\bar{t}$  of  $T$ ,  $Z_{\bar{t}}$  is connected and for any irreducible component  $\Gamma$  of  $Z_{\bar{t}}$ ,  $\Gamma$  meets the other components in at least  $1 - (2p_a(\Gamma) - 2) - |M \cap \Gamma|$  points. This amounts to saying that  $\omega_{Z/T}(M)$  is ample.

A *morphism* of marked curves  $(Z, M) \rightarrow (Z', M')$  over  $T$  is a  $T$ -morphism  $f : Z \rightarrow Z'$  such that  $f(M) \subseteq M'$ .

Let  $(X, M)$  be a smooth marked curve over  $K = K(S)$ . A *marked model* of  $(X, M)$  over  $S$  is a marked curve  $(\mathcal{X}, \mathcal{M})$  over  $S$  whose generic fiber is isomorphic to  $(X, M)$ . Since  $\mathcal{M}$  is uniquely determined by  $M$  and  $\mathcal{X}$ , we will omit  $\mathcal{M}$  in the notation  $(\mathcal{X}, \mathcal{M})$  and we will simply say that  $\mathcal{X}$  is a marked model of  $(X, M)$ .

Let  $\mathcal{X}$  be a model of  $X$  over  $S$ . The *stable marked hull* of  $\mathcal{X}$  is the minimal semi-stable marked model of  $(X, M)$  dominating  $\mathcal{X}$ . Note that a stable marked hull is not necessarily a stable marked curve.

**COROLLARY 2.20.** *Let  $(X, M)$  be a smooth marked curve over  $K$  and let  $\mathcal{X}$  be a (non-marked) model of  $X$ . Then after finite separable extension of  $K$ ,  $\mathcal{X}$  admits a stable marked hull. More precisely, if  $\mathcal{X}$  has a stable hull over some extension  $S'/S$ , then it has a stable marked hull over  $S'$ . Moreover, the formation of stable marked hull commutes with flat base change of Dedekind schemes.*

*Proof.* We can suppose that  $\mathcal{X}$  has a stable hull  $\mathcal{Z}$  over  $S$ . Let  $\tilde{\mathcal{Z}}$  be a desingularization of  $\mathcal{Z}$ , let  $\overline{M}$  be the Zariski closure of  $M$  in  $\tilde{\mathcal{Z}}$  and let  $\tilde{\mathcal{Z}}_{s_1}, \dots, \tilde{\mathcal{Z}}_{s_n}$  be the fibers such that  $\overline{M} \rightarrow \overline{M} \cap \tilde{\mathcal{Z}}_{s_i}$  is not injective. Let  $\mathcal{Z}' \rightarrow \tilde{\mathcal{Z}}$  be an embedded resolution of  $\overline{M} + \sum_i \tilde{\mathcal{Z}}_{s_i}$  in  $\tilde{\mathcal{Z}}$  so that the Zariski closure  $\mathcal{M}'$  of  $M$  in  $\mathcal{Z}'$  is a disjoint union of sections (contained in the smooth locus because  $\mathcal{Z}'$  is regular). Then  $\mathcal{Z}'$  is a semi-stable marked model dominating  $\mathcal{X}$ .

Let  $\mathcal{W}$  be any semi-stable marked model of  $(X, M)$  over  $S$ . Similarly to the non-marked case, we can show that  $\mathcal{W}$  is relatively minimal if and only if  $\omega_{\mathcal{W}/S}(\mathcal{M})|_\Gamma$ , where  $\mathcal{M}$  is the Zariski closure of  $M$  in  $\mathcal{W}$ , has positive degree for all  $\Gamma$  in the exceptional locus of  $\mathcal{W} \rightarrow \mathcal{X}$ . This then implies that the stable marked hull is obtained by contracting prime divisors  $\Gamma$  in the exceptional locus of  $\mathcal{Z}' \rightarrow \mathcal{Z}$  such that  $\deg(\omega_{\mathcal{Z}'/S}(\mathcal{M}')|_\Gamma) \leq 0$ , and that the stable marked hull commutes with flat base change. □

*Remark 2.21.* If  $(X, M)$  is stable (meaning that  $2g(X) - 2 + |M| \geq 1$ ) and if  $X$  has semi-stable reduction over  $S$ , then there exists a semi-stable marked model  $\mathcal{X}$  of  $(X, M)$  over  $S$  and minimal for this property. This model is the stable marked model of  $(X, M)$ . It is characterized by the property that for all irreducible components  $\Gamma$  of  $\mathcal{X}_s$ , one has  $\deg \omega_{\mathcal{X}/S}(\mathcal{M})|_\Gamma > 0$ , where  $\mathcal{M}$  is the Zariski closure of  $M$  in  $\mathcal{X}$ . As above, this implies that the stable marked model commutes with base change.

**3. Semi-stable models of finite covers**

We (re)prove that any finite morphism of projective smooth curves over  $K$  extends, after finite separable extension of  $K$ , to a finite morphism of semi-stable models.

DEFINITION 3.1. Let  $f : X \rightarrow Y$  be a finite morphism of smooth connected projective curves over  $K(S)$ . A *model (or extension) of  $f$  over  $S$*  consists of a morphism  $\mathcal{X} \rightarrow \mathcal{Y}$  over  $S$  extending  $f$ , where  $\mathcal{X}$  and  $\mathcal{Y}$  are models over  $S$  of  $X$  and  $Y$ , respectively. A model of  $f$  is said to be *finite* if it is a finite morphism, and *semi-stable* (see [Col03]) if it is finite and if  $\mathcal{X}, \mathcal{Y}$  are semi-stable. We say that a model  $\mathcal{X} \rightarrow \mathcal{Y}$  of  $f$  dominates another model  $\mathcal{X}' \rightarrow \mathcal{Y}'$  if there are birational morphisms  $\mathcal{X} \rightarrow \mathcal{X}'$ ,  $\mathcal{Y} \rightarrow \mathcal{Y}'$  making the following diagram commutative.

$$\begin{array}{ccc} \mathcal{X} & \longrightarrow & \mathcal{Y} \\ \downarrow & & \downarrow \\ \mathcal{X}' & \longrightarrow & \mathcal{Y}' \end{array}$$

A model of  $f$  is *stable* if it is semi-stable and minimal (for the domination relation) among semi-stable models of  $f$ . If  $\mathcal{X}, \mathcal{Y}$  are respective models of  $X, Y$ . Then the semi-stable model  $\mathcal{X}' \rightarrow \mathcal{Y}'$  of  $f$  such that  $\mathcal{X}'$  dominates  $\mathcal{X}$ ,  $\mathcal{Y}'$  dominates  $\mathcal{Y}$ , and which is minimal for this property, is called the *stable hull* of (the rational map)  $\mathcal{X} \dashrightarrow \mathcal{Y}$ . We can obviously make similar definitions for marked curves.

Remark 3.2. For a given  $f : X \rightarrow Y$ , the semi-stable models are not unique: let  $\mathcal{X} \rightarrow \mathcal{Y}$  be a semi-stable model of  $f$ , let  $\mathcal{Y}' \rightarrow \mathcal{Y}$  be a blowing-up along a closed point, then the stable hull of  $\mathcal{X} \dashrightarrow \mathcal{Y}'$  (see Theorem 4.5) is a new semi-stable model.

### 3.3 Decomposition of inseparable morphisms

Let us first deal with purely inseparable morphisms  $X \rightarrow Y$ . The next two statements are well known at least over perfect base fields.

LEMMA 3.4. Let  $K$  be a field of characteristic  $p > 0$ . Let  $E/F$  be a finite extension of function fields of one variable over  $K$ , with  $E$  separable over  $K$ . Then there exists a unique purely inseparable sub-extension  $L/F$  of  $E/F$  such that  $E/L$  is separable. Moreover,  $F = KL^{p^r}$  for some  $r \geq 0$ .

Proof. Let  $F_s$  be the separable closure of  $F$  in  $E$ . By [Liu02, Corollary 3.2.27] (here we use the hypothesis  $E$  separable over  $K$ ), there exists  $r \geq 0$  such that  $F_s = KE^{p^r}$ . Let

$$L := \{e \in E \mid e^{p^r} \in F\}.$$

Then  $L/F$  is a purely inseparable extension,  $F = KL^{p^r}$  and  $E/L$  is separable because otherwise  $L \subseteq KE^{p^r}$  and  $F_s \subseteq KE^{p^{r+1}}$ . The uniqueness of  $L$  is obvious because it is necessarily equal to the radical closure of  $F$  in  $E$ . □

PROPOSITION 3.5. Let  $f : X \rightarrow Y$  be a finite morphism of normal connected curves over a field  $K$  of characteristic  $p > 0$ . Suppose that  $X$  is smooth. Then  $f$  can be decomposed into a finite separable morphism  $X \rightarrow Z$  followed by  $Z \rightarrow Y$  which can be identified to a Frobenius map  $Z \rightarrow Z^{(p^r)}$  for some  $r \geq 0$ . Moreover,  $Z$  is smooth.

Proof. Let  $L$  be the radical closure of  $K(Y)$  in  $K(X)$ , and let  $Z$  be the normalization of  $Y$  in  $L$ . Then  $f$  induces a finite separable morphism  $X \rightarrow Z$ . It is flat because  $Z$  is regular of dimension 1. Let  $\bar{K}$  be an algebraic closure of  $K$ , then  $X_{\bar{K}} \rightarrow Z_{\bar{K}}$  is flat, hence  $Z_{\bar{K}}$  is regular. Finally,  $Z \rightarrow Y$  can be identified to  $Z \rightarrow Z^{(p^r)}$  by [Liu02, Proposition 7.4.21]. □

Note that  $f$  can also be decomposed into  $X \rightarrow X^{(p^r)}$  followed by a separable morphism  $X^{(p^r)} \rightarrow Y$ .

**3.6 Semi-stable models**

LEMMA 3.7. *Let  $f_i : X \rightarrow Y_i$ ,  $i = 1, \dots, n$  be finite surjective morphisms of integral projective varieties over  $K$  and let  $\mathcal{Y}_i$  be a model of  $Y_i$  over  $S$ . Then there exists a model  $\mathcal{X}$  of  $X$  over  $S$  such that  $\mathcal{X}$  dominates  $\mathcal{Y}_i$  (that is,  $X \rightarrow Y_i$  extends to  $\mathcal{X} \rightarrow \mathcal{Y}_i$ ) for all  $i$ .*

*Proof.* For  $N$  large enough, we have a closed immersion  $g : X \rightarrow \mathbb{P}_K^N \times_K (\prod_{K,i} Y_i)$  induced by the projective morphism  $(f_1, \dots, f_n) : X \rightarrow \prod_{K,i} Y_i$ . Now take  $\mathcal{X}$  to be the Zariski closure of  $g(X)$  in  $\mathbb{P}_S^N \times_S (\prod_{S,i} \mathcal{Y}_i)$ , endowed with the reduced structure. Note that if  $X$  is geometrically reduced, we can also use Lemma 4.3 with  $\mathcal{X}_i = N(\mathcal{Y}_i, K(X))$ . □

PROPOSITION 3.8. *Let  $S$  be a Dedekind scheme, and let  $f : X \rightarrow Y$  be a finite morphism of smooth geometrically connected projective curves over  $K := K(S)$ . Let  $\mathcal{X}, \mathcal{Y}$  be respective models of  $X$  and  $Y$ . Then there exists a finite separable extension  $K'/K$  such that over the normalization  $S'$  of  $S$  in  $K'$ , the cover  $X_{K'} \rightarrow Y_{K'}$  extends to a finite morphism  $\mathcal{X}' \rightarrow \mathcal{Y}'$ , where  $\mathcal{X}'$  (respectively  $\mathcal{Y}'$ ) is a semi-stable model of  $X$  (respectively  $Y$ ) over  $S$  dominating  $\mathcal{X}_{S'}$  (respectively,  $\mathcal{Y}_{S'}$ ).*

*Proof.* Let  $X \rightarrow Z \rightarrow Y$  be the decomposition given by Proposition 3.5. Let  $\hat{X} \rightarrow Z$  be the Galois closure of  $X \rightarrow Z$ . After a finite separable extension of  $K$ ,  $\hat{X}$  is smooth over  $K$ . Let  $\hat{\mathcal{X}}_0/S$  be a model of  $\hat{X}$  dominating  $\mathcal{X}$  and  $\mathcal{Y}$  (3.7). Let  $G := \text{Gal}(K(\hat{X})/K(Z))$ . By Corollary 2.5, after a finite separable extension of  $K$ , there exists a semi-stable model  $\hat{\mathcal{X}}$  of  $\hat{X}$  endowed with an action of  $G$  and dominating  $\hat{\mathcal{X}}_0$ . Let  $\mathcal{X}' = \hat{\mathcal{X}}/H$  where  $H = \text{Gal}(K(\hat{X})/K(X))$ , and  $\mathcal{Z}' = \hat{\mathcal{X}}/G$ . Then  $\mathcal{X}' \rightarrow \mathcal{Z}'$  is a finite morphism of semi-stable models of  $X$  and  $Z$ , respectively [Ray90, Proposition 5]. Let  $\mathcal{Y}' = \mathcal{Z}'^{(p')}$ . Then the canonical map  $\mathcal{Z}' \rightarrow \mathcal{Y}'$  is finite and  $\mathcal{Y}'$  is semi-stable (or [Liu02, Exercise 10.3.19(a)]). Since  $\hat{\mathcal{X}}$  dominates  $\mathcal{Y}$  and is finite over  $\mathcal{Y}'$ , we see easily that  $\mathcal{Y}'$  dominates  $\mathcal{Y}$  (use, for instance, [LL99, Lemma 4.1]). Hence, the proposition is proved with  $f'$  equal to the composition  $\mathcal{X}' \rightarrow \mathcal{Z}' \rightarrow \mathcal{Y}'$ . □

Remark 3.9. If  $S$  is any Noetherian integral excellent scheme, then using the result of de Jong [Dej97] as quoted in Remark 2.9, we see that the proposition is still true.

The following corollary was known for separable morphisms (see [Col03] when  $K$  is complete; [LL99, Remark 4.6] when  $g(X) \geq 1$ ).

COROLLARY 3.10. *Let  $f : X \rightarrow Y$  be a finite morphism of smooth geometrically connected projective curves over  $K$ . Then after a finite separable extension of  $K$ , there exists a finite morphism  $\varphi : \mathcal{X} \rightarrow \mathcal{Y}$  of semi-stable models of  $X, Y$ , respectively.*

*Proof.* Apply Proposition 3.8 to any pair of models of  $X, Y$ . □

LEMMA 3.11. *Let  $S$  be local. Let  $F$  be a finite closed subset of  $X$ . Let  $\mathcal{Z}$  be a semi-stable model of  $X$  over  $S$ . Then there exists an integer  $d > 0$  such that for any finite extension  $\mathcal{O}_{K'}/\mathcal{O}_S$  of discrete valuation rings with ramification index divisible by  $d$ , if  $\tilde{\mathcal{Z}}'$  denotes a desingularization of  $\mathcal{Z}_{\mathcal{O}_{K'}}$ , then the Zariski closure of  $F_{K'}$  in  $\tilde{\mathcal{Z}}'$  is contained in the smooth locus of  $\tilde{\mathcal{Z}}'$ .*

*Proof.* Let  $\alpha \in F$ , and let  $x \in \mathcal{Z}_s$  be a singular specialization of  $\alpha$ . The local ring of an étale neighborhood of  $x \in \mathcal{Z}$  is isomorphic to  $\mathcal{O}_K[[u, v]]/(uv - a)$ , with  $a$  a power of a uniformizing element of  $\mathcal{O}_K$ . Let  $u(\alpha)$  be the image of  $u$  in  $K(\alpha)$ . Then  $|a| < |u(\alpha)| < 1$ . After an extension of large enough ramification index,  $|u(\alpha)|$  belongs to  $|K|$ . If this condition is satisfied for all  $\alpha \in F$  and for all singular specializations of  $\alpha$ , then it is easy to see that the specializations of  $F$  in  $\tilde{\mathcal{Z}}'$  are smooth points. Indeed, if  $\mathcal{Z}$  is regular, the parameter  $a$  in the above local ring is a uniformizing element, hence  $|a| < |u(\alpha)| < 1$  cannot hold in  $|K|$ , so  $F$  must specialize to smooth points. □

PROPOSITION 3.12. *Let  $f : (X, M) \rightarrow (Y, N)$  be a finite morphism of smooth geometrically connected marked projective curves over  $K$ , let  $\mathcal{X}, \mathcal{Y}$  be respective models of  $X, Y$ . Then after a finite separable extension of  $K$ , there exists a semi-stable marked model  $\mathcal{X}'$  (respectively,  $\mathcal{Y}'$ ) of  $(X, M)$  (respectively,  $(Y, N)$ ) such that  $\mathcal{X}'$  and  $\mathcal{Y}'$  dominate  $\mathcal{X}$  and  $\mathcal{Y}$ , respectively, and  $f$  extends to a finite morphism  $\mathcal{X}' \rightarrow \mathcal{Y}'$ .*

*Proof.* After enlarging  $K$  and replacing  $\mathcal{X}$  and  $\mathcal{Y}$  by their respective stable marked hull, we can suppose that  $\mathcal{X}, \mathcal{Y}$  are semi-stable and that the Zariski closure  $\overline{M}$  of  $M$  in  $\mathcal{X}$  is a disjoint union of sections, and the same for  $N$  in  $\mathcal{Y}$ . Let  $X \rightarrow Z \rightarrow Y$  be the decomposition as given by Proposition 3.5, and let  $\hat{X} \rightarrow Z$  be the Galois closure of  $X \rightarrow Z$ . Let  $f : \hat{X} \rightarrow X$  and  $g : \hat{X} \rightarrow Y$  be the canonical morphisms. By Lemma 3.11, after a finite separable extension, and after replacing  $\hat{X}$  by its minimal desingularization (the group  $G$  still acts), we can suppose that the Zariski closure of  $f^{-1}(M) \cup g^{-1}(N)$  in  $\hat{X}$  is contained in the smooth locus. Then the Zariski closure of  $M$  in  $\mathcal{X}'$  is contained in the smooth locus because  $\hat{\mathcal{X}}_{\text{sm}}/H$  is smooth, and it is a disjoint union of sections because  $\mathcal{X}'$  dominates  $\mathcal{X}$  and  $\overline{M}$  is already a disjoint union of sections. Hence,  $\mathcal{X}'$  is semi-stable marked for  $(X, M)$ . The same arguments hold for  $\mathcal{Y}'$ .  $\square$

#### 4. Stable hull of a morphism

DEFINITION 4.1. Let  $f : X \rightarrow Y$  be a finite morphism of connected smooth projective curves over  $K$ . Let  $\mathcal{X}, \mathcal{Y}$  be respective models of  $X, Y$  over  $S$ . The *finite hull of  $\mathcal{X} \dashrightarrow \mathcal{Y}$*  is a finite model  $\mathcal{X}^f \rightarrow \mathcal{Y}^f$  of  $f$  over  $S$ , such that  $\mathcal{X}^f$  and  $\mathcal{Y}^f$  are normal models of  $X, Y$  dominating  $\mathcal{X}$  and  $\mathcal{Y}$ , respectively, and which is minimal (for the domination relation) with respect to these properties.

LEMMA 4.2. *Let  $\mathcal{Y}$  be an integral scheme locally of finite type over  $S$ , let  $L$  be a finite extension of  $K(\mathcal{Y})$ , separable over  $K = K(S)$ , and let  $\mathcal{X}$  be the normalization of  $\mathcal{Y}$  in  $L$ . Then  $\mathcal{X}$  is finite over  $\mathcal{Y}$ .*

*Proof.* The assertion is, of course, trivial if  $S$  is excellent. Let  $y \in \mathcal{Y}_s$ , let  $A = \mathcal{O}_{\mathcal{Y},y}$  and let  $B$  be the integral closure of  $A$  in  $L$ . We have to show that  $B$  is finite over  $A$ . Let  $\mathcal{O}_K = \mathcal{O}_{S,s}$ . Let  $C$  be the integral closure of  $A \otimes_{\mathcal{O}_K} \hat{\mathcal{O}}_K$  in  $L \otimes_K \hat{K}$ . The latter is reduced (because  $L$  is separable over  $K$ ) and finite over  $K(\mathcal{Y}) \otimes_K \hat{K}$ , the total ring of fractions of  $A \otimes \hat{\mathcal{O}}_K$ . Since  $\hat{\mathcal{O}}_K$  is excellent,  $C$  (and, thus,  $B \otimes \hat{\mathcal{O}}_K$ ) is finitely generated over  $A \otimes \hat{\mathcal{O}}_K$ . Then Nakayama’s lemma implies that  $B \otimes \hat{\mathcal{O}}_K$  is generated over  $A \otimes \hat{\mathcal{O}}_K$  by finitely many elements of  $B$ . These elements generate  $B$  over  $A$  because  $\mathcal{O}_K \rightarrow \hat{\mathcal{O}}_K$  is faithfully flat.

Note that the proof still work if  $S$  is any Noetherian integral scheme such that  $\hat{\mathcal{O}}_{S,s}$  is reduced for all  $s \in S$ .  $\square$

LEMMA 4.3. *Let  $X$  be an integral projective variety over  $K$ , and let  $\mathcal{X}_1, \mathcal{X}_2, \dots, \mathcal{X}_n$  be models of  $X$  over  $S$ .*

- (a) *There exists a smallest model  $\mathcal{X}$  of  $X$  dominating  $\mathcal{X}_i$  for all  $i$ . Let us denote  $\mathcal{X}$  by  $\mathcal{X}_1 \vee \dots \vee \mathcal{X}_n$ .*
- (b) *If  $X$  is geometrically integral, then for any flat morphism of Dedekind schemes  $S' \rightarrow S$ , we have  $(\mathcal{X}_1 \vee \dots \vee \mathcal{X}_n)_{S'} = (\mathcal{X}_1)_{S'} \vee \dots \vee (\mathcal{X}_n)_{S'}$ .*
- (c) *If  $\dim X = 1$ , then every irreducible component of  $(\mathcal{X}_1 \vee \dots \vee \mathcal{X}_n)_s$  dominates an irreducible component of  $(\mathcal{X}_i)_s$  for some  $i$ .*

*Proof.* The proof is similar to that of Lemma 2.4. Let  $\mathcal{P}$  be the fiber product  $\prod_{S,i} \mathcal{X}_i$  over  $S$ . Then the diagonal map makes  $X$  a closed subscheme of  $\mathcal{P}_K$ . Let  $\mathcal{X}$  be the Zariski closure of  $X$  in  $\mathcal{P}$  endowed with the reduced structure. Then  $\mathcal{X}$  is a model of  $X$  over  $S$  dominating the  $\mathcal{X}_i$ . Let  $\mathcal{Z}$  be a model of  $X$  over  $S$  dominating the  $\mathcal{X}_i$ . Then we have a natural morphism  $f : \mathcal{Z} \rightarrow \mathcal{P}$  whose image  $f(\mathcal{Z})$  is irreducible, with generic fiber  $\mathcal{X}_K$ . Hence,  $f(\mathcal{Z}) = \mathcal{X}$  and  $f$  factorizes through  $\mathcal{Z} \rightarrow \mathcal{X} \rightarrow \mathcal{P}$ .

So  $\mathcal{X}$  is minimal. If  $X$  is geometrically integral, then  $\mathcal{X}_{S'}$  is an integral closed subscheme of  $\mathcal{P}' := \prod_{S',i}(\mathcal{X}_i)_{S'}$ , with generic fiber isomorphic to the diagonal of  $\mathcal{P}'_{K'}$ . By construction,  $\mathcal{X}_{S'}$  is equal to  $(\mathcal{X}_1)_{S'} \vee \cdots \vee (\mathcal{X}_n)_{S'}$ .

Let  $s \in S$ . Then  $\mathcal{X}_s$  is a closed subscheme of  $\mathcal{P}_s = \prod_{k(s),i}(\mathcal{X}_i)_s$ , pure of dimension  $\dim X$ . Let  $\Gamma$  be an irreducible component of  $\mathcal{X}_s$ . If  $\dim X > 0$ , then the image of  $\Gamma$  in  $\mathcal{X}_{i_0}$  has positive dimension for some  $i_0$ . If  $\dim X = 1$ , then the image of  $\Gamma$  in  $(\mathcal{X}_{i_0})_s$  is an irreducible component.  $\square$

PROPOSITION 4.4. *Let  $\mathcal{X} \dashrightarrow \mathcal{Y}$  be a rational map. Suppose that either  $S$  is local and Henselian, or  $\mathcal{X}$  and  $\mathcal{Y}$  are semi-stable or regular. Then the finite hull of  $\mathcal{X} \dashrightarrow \mathcal{Y}$  exists. It commutes with base changes in the following sense: let  $S' \rightarrow S$  be a flat morphism of Dedekind schemes. Then the finite hull of  $\mathcal{X}_{S'} \dashrightarrow \mathcal{Y}_{S'}$  exists and is equal to  $((\mathcal{X}^f)_{S'})^\sim \rightarrow ((\mathcal{Y}^f)_{S'})^\sim$ , where  $\sim$  means normalization.*

*Proof.* The rational map  $\mathcal{X} \dashrightarrow \mathcal{Y}$  is defined and finite above an open dense subset of  $S$ . So we can suppose that  $S = \text{Spec } \mathcal{O}_K$  is local with closed point  $s$ . The case when  $\mathcal{X}, \mathcal{Y}$  are semi-stable or regular is easily reduced to the Henselian case by passing to the completion of  $\mathcal{O}_K$  and using Proposition 1.3. Suppose that  $\mathcal{O}_K$  is Henselian. Let  $\mathcal{X}_1$  be the normalization of  $\mathcal{X} \vee N(\mathcal{Y}, K(X))$  (see Lemma 4.3). We have a morphism  $\mathcal{X}_1 \rightarrow \mathcal{Y}$ . By [LL99, Lemma 4.14], there exists a (unique) normal model  $\mathcal{Y}^f$  such that the rational map  $\mathcal{X}_1 \dashrightarrow \mathcal{Y}^f$  is quasi-finite and surjective in codimension 1 (in other words, if  $\mathcal{U}$  is the domain of definition of  $\mathcal{X}_1 \dashrightarrow \mathcal{Y}^f$ , then  $\mathcal{U}_s$  is dense in  $(\mathcal{X}_1)_s$ ,  $\mathcal{U}_s \rightarrow \mathcal{Y}_s$  is quasi-finite and has dense image). Since  $\mathcal{X}_1 \rightarrow \mathcal{Y}$  is a morphism,  $\mathcal{Y}^f$  dominates  $\mathcal{Y}$ . Let  $\mathcal{X}^f$  be the normalization of  $\mathcal{Y}^f$  in  $K(X)$ . Then it is easy to see that  $\mathcal{X}^f \rightarrow \mathcal{Y}^f$  is the finite hull of  $\mathcal{X} \dashrightarrow \mathcal{Y}$  (use [LL99, Lemma 4.1] for instance).

It remains to prove the base change property. The rational map  $(\mathcal{X}_{S'})^\sim \dashrightarrow ((\mathcal{Y}^f)_{S'})^\sim$  is quasi-finite and surjective in codimension 1. By the above construction,  $(\mathcal{Y}_{S'})^f = ((\mathcal{Y}^f)_{S'})^\sim$  and  $(\mathcal{X}_{S'})^f$  is the normalization of  $((\mathcal{Y}^f)_{S'})^\sim$  in  $K(X_{K(S)})$ . Since  $((\mathcal{X}^f)_{S'})^\sim \rightarrow ((\mathcal{Y}^f)_{S'})^\sim$  is finite,  $(\mathcal{X}_{S'})^f$  is isomorphic to  $((\mathcal{X}^f)_{S'})^\sim$ .  $\square$

THEOREM 4.5. *Let  $S$  be a connected Noetherian regular scheme of dimension 1, let  $f : X \rightarrow Y$  be a finite morphism of smooth geometrically connected projective curves over  $K := K(S)$ . Let  $\mathcal{X}, \mathcal{Y}$  be respective models of  $X, Y$  over  $S$ . Then after a finite separable extension of  $K$ ,  $\mathcal{X} \dashrightarrow \mathcal{Y}$  admits a stable hull  $\mathcal{X}' \rightarrow \mathcal{Y}'$ . Moreover, the formation of  $\mathcal{X}' \rightarrow \mathcal{Y}'$  commutes with flat base change.*

*Proof.* After a finite separable extension of  $K$ , we can suppose that there exists a semi-stable model  $\mathcal{X}_\infty \rightarrow \mathcal{Y}_\infty$  of  $f$  dominating  $\mathcal{X} \dashrightarrow \mathcal{Y}$  (Corollary 3.10). Let us show that  $\mathcal{X} \dashrightarrow \mathcal{Y}$  then admits a stable hull over  $S$ . Consider the stable hull  $\mathcal{Y}_1$  of  $\mathcal{Y}$  and the stable hull  $\mathcal{X}_1$  of  $N(\mathcal{Y}_1, K(X))$ . Then  $\mathcal{X}_1$  and  $\mathcal{Y}_1$  are dominated by  $\mathcal{X}_\infty$  and  $\mathcal{Y}_\infty$ , respectively. Let  $\mathcal{X}_2 \rightarrow \mathcal{Y}_2$  be the finite hull of  $\mathcal{X}_1 \rightarrow \mathcal{Y}_1$ . Then it is also dominated by  $\mathcal{X}_\infty \rightarrow \mathcal{Y}_\infty$ . Now restart again the process of taking stable hull and finite hull with  $\mathcal{X}_2 \rightarrow \mathcal{Y}_2$ . We construct in this way an increasing sequence of (normal) models  $\mathcal{X}_n \rightarrow \mathcal{Y}_n$  of  $f$  over  $S$  which are dominated by  $\mathcal{X}_\infty \rightarrow \mathcal{Y}_\infty$ . This sequence is stationary at some rank  $n_0$ . Then  $\mathcal{X}_{n_0} \rightarrow \mathcal{Y}_{n_0}$  is a semi-stable model of  $f$ . Note that the construction of  $\mathcal{X}_n \rightarrow \mathcal{Y}_n$  does not depend on the choice of  $\mathcal{X}_\infty \rightarrow \mathcal{Y}_\infty$ . In particular,  $\mathcal{X}_n \rightarrow \mathcal{Y}_n$  is dominated by any semi-stable model of  $f$  dominating  $\mathcal{X} \dashrightarrow \mathcal{Y}$ . Therefore,  $\mathcal{X}_{n_0} \rightarrow \mathcal{Y}_{n_0}$  is the stable hull of  $\mathcal{X} \dashrightarrow \mathcal{Y}$ . Finally, the formation of the stable hull commutes with flat base change because the stable hull of a model and the finite hull of a morphism commute with flat base change (Theorem 2.3 and Proposition 4.4).  $\square$

COROLLARY 4.6. *Suppose that either  $g(X) \geq 2$ , or  $g(X) = 1$  and  $X$  has potentially good reduction. Then there exists a finite separable extension of  $K'$  of  $K$  such that  $X_{K'} \rightarrow Y_{K'}$  admits a stable model  $\mathcal{X}' \rightarrow \mathcal{Y}'$  over  $S'$ , where  $S'$  is the integral closure of  $S$  in  $K'$ . Moreover, for any Dedekind scheme  $T$  dominating  $S'$ ,  $\mathcal{X}'_T \rightarrow \mathcal{Y}'_T$  is the stable model of  $X_{K(T)} \rightarrow Y_{K(T)}$ .*

*Proof.* We can suppose that  $X$  has semi-stable reduction over  $S$ . The cover  $X \rightarrow Y$  extends to a finite morphism of smooth projective models over a dense open subset of  $S$ . So we can suppose




FIGURE 2. A stable model not coming from the Galois closure (see Remark 4.8).

that  $S = \text{Spec } \mathcal{O}_K$  is local. Let  $\mathcal{X}^{\text{st}}$  be the stable (respectively, smooth projective) model of  $X$  if  $g(X) \geq 2$  (respectively, if  $g(X) = 1$ ). Suppose first that  $\mathcal{O}_K$  is complete (hence, Henselian). Let  $\mathcal{X}^{\text{st}} \dashrightarrow \mathcal{Y}''$  be the rational map extending  $X \rightarrow Y$  and which is quasi-finite and surjective in codimension 1 (see [LL99, Lemma 4.14]). Then the stable hull  $\mathcal{X} \rightarrow \mathcal{Y}$  of  $\mathcal{X}^{\text{st}} \dashrightarrow \mathcal{Y}''$  is clearly the stable model of  $X \rightarrow Y$ . The construction of  $\mathcal{X} \rightarrow \mathcal{Y}$  commutes with flat base change because that of  $\mathcal{Y}''$  and the stable hull commute with flat base change. If  $\mathcal{O}_K$  is non-necessarily complete, we can use Corollary 1.6.  $\square$

*Remark 4.7.* Let  $X \rightarrow Y$  be as above. If  $X \rightarrow Y$  has a semi-stable model over  $S$ , then it has a stable model over  $S$ . This can be seen in the proof of Corollary 4.6. If  $X \rightarrow Y$  is, moreover, Galois of group  $G$ , and if  $\mathcal{X}$  is the stable model (or smooth model if  $g(X) = 1$ ) of  $X$  over  $S$ , then the stable model of  $X \rightarrow Y$  is equal to  $\mathcal{X} \rightarrow \mathcal{X}/G$ .

*Remark 4.8.* Suppose, moreover, that  $X \rightarrow Y$  is separable, and that the Galois closure  $\hat{X}$  of  $X \rightarrow Y$  is smooth and geometrically connected over  $K$  (which is true after a finite separable extension of  $K$ ). Let  $\hat{\mathcal{X}}$  be the stable model of  $\hat{X}$ , and let  $G = \text{Gal}(K(\hat{X})/K(Y))$ ,  $H = \text{Gal}(K(\hat{X})/K(X))$  as in the proof of Proposition 3.8. Then we can ask whether  $\hat{\mathcal{X}}/H \rightarrow \hat{\mathcal{X}}/G$  is the stable model of  $X \rightarrow Y$ . The answer is no in general. Let us give an example with  $X$  and  $Y$  having good reduction.

Let  $S$  be local, complete, with algebraically closed residue field  $k$ . Let  $C_1 \rightarrow D_1$  be a finite separable morphism of degree  $d \geq 3$  with  $C_1, D_1 \simeq \mathbb{P}_k^1$ , totally ramified above some point  $y_1 \in D_1$ , and such that the Galois closure  $E$  of  $C_1 \rightarrow D_1$  is a curve of genus  $g(E) \geq 1$ . Let  $C_2 \rightarrow D_2$  be a finite separable morphism of degree  $d$  of smooth connected projective curves over  $k$ , totally ramified above a point  $y_2 \in D_2$  and such that  $g(C_2) \geq 1$ . Let  $D$  be the semi-stable curve over  $k$  obtained by identifying  $y_1$  and  $y_2$ . Let  $C$  be the semi-stable curve defined in a similar way. Then we have a finite morphism  $\rho : C \rightarrow D$ , which is generically étale, and such that  $C_1 \rightarrow D_1, C_2 \rightarrow D_2$  have the same ramification index at  $y_1$  and  $y_2$ . (See Figure 2.) By [Liu03, Proposition 5.4], the cover  $C \rightarrow D$  lifts to a finite morphism  $\mathcal{C} \rightarrow \mathcal{D}$  over  $S$  with smooth generic fibers  $X, Y$ . Let  $\mathcal{C} \rightarrow \mathcal{C}_2$  (respectively,  $\mathcal{D} \rightarrow \mathcal{D}_2$ ) be the contraction of  $C_1$  (respectively, of  $D_1$ ). Then  $\mathcal{C}_2, \mathcal{D}_2$  are smooth, and the canonical morphism  $\mathcal{C}_2 \rightarrow \mathcal{D}_2$  is the stable model of  $X \rightarrow Y$ . Let  $\mathcal{Z}$  be the normalization of  $\mathcal{D}$  in  $K(\hat{X})$ . Let  $\Theta$  be an irreducible component of  $\mathcal{Z}_s$  lying over  $D_1$ . Then the separable closure of  $k(D_1)$  in  $k(\Theta)$  is Galois over  $k(D_1)$  (see [Ser68, I, § 7, Proposition 20]) and contains  $k(C_1)$ . Thus,  $k(\Theta)$  contains a subfield isomorphic to  $k(E)$ . In particular,  $p_a(\Theta) \geq 1$ . If  $\mathcal{C}_2 \rightarrow \mathcal{D}_2$  is equal to  $\hat{\mathcal{X}}/H \rightarrow \hat{\mathcal{X}}/G$ , then  $\mathcal{Z}$  dominates  $\hat{\mathcal{X}}$  because  $\mathcal{D}$  dominates  $\mathcal{D}_2$ . However,  $\Theta$  maps to a closed point of  $\hat{\mathcal{X}}_s$ , thus  $\hat{\mathcal{X}}$  cannot be semi-stable – contradiction.

*Remark 4.9.* If  $X$  has genus 1 and multiplicative reduction at some point of  $S$ , or if  $g(X) = 0$ , then over any ramified extension of  $S$ , there is no stable model of the identity map  $X \rightarrow X$ . The reason we take a ramified extension  $S'/S$  is that, when  $g(X) = 1$ ,  $X_{K'}$  has no minimal semi-stable model (see the proof of Lemma 1.5).

Let us give a characterization of the stable model.

DEFINITION 4.10. Let  $X_1 \rightarrow Y_1, X_2 \rightarrow Y_2$  be morphisms of schemes over some base scheme  $T$ . An *isomorphism* of the pairs  $(X_1 \rightarrow Y_1) \rightarrow (X_2 \rightarrow Y_2)$  is given by  $T$ -isomorphisms  $X_1 \rightarrow X_2, Y_1 \rightarrow Y_2$  such that the diagram

$$\begin{array}{ccc} X_1 & \longrightarrow & X_2 \\ \downarrow & & \downarrow \\ Y_1 & \longrightarrow & Y_2 \end{array}$$

is commutative. We denote by  $\text{Isom}_T((X_1 \rightarrow Y_1), (X_2 \rightarrow Y_2))$  the set of these isomorphisms. Now  $\text{Aut}_T(X \rightarrow Y)$  has an obvious meaning.

PROPOSITION 4.11. *Keep the hypothesis of Theorem 4.5 and suppose that  $g(X) \geq 2$ . Let  $\mathcal{X} \rightarrow \mathcal{Y}$  be a semi-stable model of  $X \rightarrow Y$ . Consider the following properties.*

- (i)  $\mathcal{X} \rightarrow \mathcal{Y}$  is the stable model of  $X \rightarrow Y$  over  $S$ .
- (ii) Let  $\Gamma$  be any irreducible component of  $\mathcal{Y}_s$  such that  $\text{deg } \omega_{\mathcal{Y}_s/k(s)}|_\Gamma \leq 0$ , then there exists an irreducible component  $\Theta$  of  $\mathcal{X}_s$  dominating  $\Gamma$  such that  $\text{deg } \omega_{\mathcal{X}_s/k(s)}|_\Theta > 0$ .
- (iii)  $\text{Aut}_{k(\bar{s})}(\mathcal{X}_{\bar{s}} \rightarrow \mathcal{Y}_{\bar{s}})$  is finite for all  $s \in S$ .

Then (i)  $\iff$  (ii)  $\implies$  (iii).

*Proof.* Looking at the proofs of Theorem 4.5 and Corollary 4.6, we see that  $X \rightarrow Y$  admits a stable model over  $S$ . Thus,  $\mathcal{X} \rightarrow \mathcal{Y}$  is stable if and only if it is a relatively minimal semi-stable model. Hence the equivalence (i)  $\iff$  (ii) is an immediate consequence of Lemma 2.13.

Suppose that condition (ii) is satisfied. Then the same condition holds over  $k(\bar{s})$  (with dualizing sheaves on  $\mathcal{X}_{\bar{s}}$  and  $\mathcal{Y}_{\bar{s}}$ ). Consider the natural inclusion

$$\text{Aut}_{k(\bar{s})}(\mathcal{X}_{\bar{s}} \rightarrow \mathcal{Y}_{\bar{s}}) \subseteq \text{Aut}_{k(\bar{s})}(\mathcal{X}_{\bar{s}}) \times \text{Aut}_{k(\bar{s})}(\mathcal{Y}_{\bar{s}}).$$

Note that the right-hand side is not a finite group in general. Let  $G$  be the subgroup (of finite index) of  $\text{Aut}_{k(\bar{s})}(\mathcal{X}_{\bar{s}})$  consisting in automorphisms which fix globally each irreducible component, and let  $H$  be the subgroup of  $\text{Aut}_{k(\bar{s})}(\mathcal{Y}_{\bar{s}})$  consisting of automorphisms  $\tau$  such that  $\tau|_\Gamma = \text{Id}$  for every irreducible component  $\Gamma$  with  $\text{deg } \omega_{\mathcal{Y}_{k(\bar{s})/k(\bar{s})}}|_\Gamma > 0$ . Then  $H$  is of finite index in  $\text{Aut}_{k(\bar{s})}(\mathcal{Y}_{\bar{s}})$  because  $\{\tau \in \text{Aut}_{k(\bar{s})}(\mathcal{Y}_{k(\bar{s})}) \mid \tau(\Gamma) = \Gamma\}$  is finite for any such  $\Gamma$ . Thus, it is enough to show that  $G' := \text{Aut}_{k(\bar{s})}(\mathcal{X}_{\bar{s}} \rightarrow \mathcal{Y}_{\bar{s}}) \cap (G \times H)$  is finite. Let  $I$  be the set of irreducible components  $\Theta$  of  $\mathcal{X}_{\bar{s}}$  such that  $\text{deg } \omega_{\mathcal{X}_{\bar{s}}/k(\bar{s})}|_\Theta > 0$ . Then  $G|_\Theta$  is finite for all  $\Theta \in I$ . Let  $(\sigma, \tau)$  be an element of  $\text{Ker}(G' \rightarrow G \times H \rightarrow \prod_{\Theta \in I} G|_\Theta)$ . Condition (ii) implies that  $\tau = \text{Id}$  on  $\mathcal{Y}_{\bar{s}}$ , hence  $\sigma \in \text{Aut}_{\mathcal{Y}_{\bar{s}}}(\mathcal{X}_{\bar{s}})$ . The latter is a finite group because  $\mathcal{X}_{\bar{s}} \rightarrow \mathcal{Y}_{\bar{s}}$  is a finite morphism. Since the projection map  $\text{Aut}_{k(\bar{s})}(\mathcal{X}_{\bar{s}} \rightarrow \mathcal{Y}_{\bar{s}}) \rightarrow \text{Aut}_{k(\bar{s})}(\mathcal{X}_{\bar{s}})$  is injective, the above kernel is finite. Hence,  $G'$  is finite.  $\square$

Remark 4.12. In general, property (iii) does not imply property (i). Let  $S$  be local, complete with algebraically closed residue field  $k$ . Let  $\pi : \mathbb{A}_k^1 \rightarrow \mathbb{A}_k^1$  be a finite separable cover with trivial automorphisms group  $\text{Aut}_k(\pi)$ . Then  $\pi$  extends to a finite cover  $\mathbb{P}_k^1 \rightarrow \mathbb{P}_k^1$  totally ramified at  $\infty$ . We can glue  $\pi$  with a finite separable cover  $C_2 \rightarrow D_2$  of smooth projective curves of genus  $\geq 2$  over  $k$  and obtain a finite cover  $C \rightarrow D$  with finite automorphisms group (see the construction in Remark 4.8) which lifts to a finite morphism of semi-stable curves  $\mathcal{C} \rightarrow \mathcal{D}$  over  $S$ . By the equivalence of properties (i) and (ii),  $\mathcal{C} \rightarrow \mathcal{D}$  is not stable.

Remark 4.13. We can see that Theorem 4.5 holds for finite morphisms of smooth projective marked curves  $(X, M) \rightarrow (Y, N)$ : let  $\mathcal{X}, \mathcal{Y}$  be respective models of  $X, Y$ , then after finite separable extension, there exists a stable marked hull  $\mathcal{X}' \rightarrow \mathcal{Y}'$  of  $\mathcal{X} \dashrightarrow \mathcal{Y}$ . Moreover, the formation of  $\mathcal{X}' \rightarrow \mathcal{Y}'$  commutes with flat base change. The proof is the same as for Theorem 4.5, except that we replace stable hull of a model by its stable marked hull, and we use Proposition 3.12 instead of Corollary 3.10.

The next lemma is a generalization of [LL99, Proposition 4.4(a)] (take  $N = \emptyset$ ).

LEMMA 4.14. *Let  $f : (X, M) \rightarrow (Y, N)$  be a finite morphism of connected smooth projective marked curves over  $K$ . Suppose that  $2g(Y) - 2 + |N| > 1$  and  $f^{-1}(N) = M$ . If  $(X, M)$  has a stable marked model  $\mathcal{X}$  over  $S$ , then  $(Y, N)$  has a stable marked model  $\mathcal{Y}$ , and  $\mathcal{X} \dashrightarrow \mathcal{Y}$  is a morphism.*

*Proof.* The stable marked model exists over  $S$  when  $X$  has semi-stable reduction over  $S$ . So the existence of  $\mathcal{X}$  implies that of  $\mathcal{Y}$  (see [LL99, Remark 4.8], when  $g(Y) \geq 1$ , the case  $g(Y) = 0$  is trivial). It remains to show that the rational map  $\mathcal{X} \dashrightarrow \mathcal{Y}$  is defined everywhere. Since the stable marked model commutes with flat base change (2.21), we can suppose that  $S$  is local with algebraically closed residue field and that  $\mathcal{W} := N(\mathcal{Y}, K(X))$  has a stable marked hull  $\mathcal{Z}$  over  $S$ . By definition,  $\mathcal{Z}$  dominates  $\mathcal{X}$ . We are going to show that  $\mathcal{Z} \rightarrow \mathcal{X}$  is an isomorphism, or, equivalently, that  $\mathcal{Z}$  is the stable marked hull of  $\mathcal{X}$ . Let  $\overline{M}$  denote the Zariski closure of  $M$  in  $\mathcal{Z}$ . It is enough to show that  $\deg \omega_{\mathcal{Z}/S}(\overline{M})|_{\Theta} > 0$  for all irreducible component  $\Theta$  of  $\mathcal{Z}_s$ . If  $\Theta$  is in the exceptional locus of  $\mathcal{Z} \rightarrow \mathcal{W}$ , then this is true because  $\mathcal{Z} \rightarrow \mathcal{W}$  is the stable marked hull. Suppose that  $\Theta$  is the strict transform of some irreducible component  $\Gamma$  of  $\mathcal{W}_s$ . Let  $\Delta$  be the image of  $\Gamma$  in  $\mathcal{Y}_s$ . Then every point  $y \in \Delta$  which is either a singular point of  $\mathcal{Y}_s$  or a specialization of  $N$  lifts to a point of  $\Theta$  which is either a singular point or a specialization of  $f^{-1}(N) = M$  (see [LL99, Proposition 4.4(b)] for singular points; use going-down property as in [LL99, Lemma 4.3] for specializations of  $N$ ). Since  $\mathcal{Y}$  is stable marked, this implies that  $\Theta$  contains at least three points of  $(\mathcal{Z}_s)_{\text{sing}} \cup \overline{M}_s$ , hence  $\deg \omega_{\mathcal{Z}/S}(\overline{M})|_{\Theta} > 0$ . □

Remark 4.15. It is known that, in general,  $f : X \rightarrow Y$  does not extend to a morphism of the respective minimal regular models of  $X$  and  $Y$  over  $S$  (see, e.g., [CES03, p. 333]). However, if  $g(Y) > 0$ , then  $f$  extends to a morphism of the respective minimal regular models with normal crossings, at least if  $k(s)$  is perfect for all closed points  $s \in S$ .

COROLLARY 4.16. *Let  $f : (X, M) \rightarrow (Y, N)$  be a finite morphism of geometrically connected smooth projective marked curves over  $K$ . Suppose that  $2g(Y) - 2 + |N| > 1$  and  $f^{-1}(N) = M$ . Then, after a finite separable extension of  $K$ ,  $f$  admits a stable marked model  $\mathcal{X} \rightarrow \mathcal{Y}$  over  $S$ . This construction commutes with flat base change.*

Remark 4.17. Let  $f : X \rightarrow Y$  be a finite separable morphism of smooth projective curves. A natural way to mark  $X$  and  $Y$  is to take  $M$  equal to the ramification locus of  $f$  and  $N$  equal to the branch locus. By definition  $f^{-1}(N) = M$ . Of course, in general,  $M$  is not contained in  $X(K)$ . However, if  $f$  is tamely ramified (e.g. if  $\text{char}(K) = 0$ ), then this becomes true over a finite separable extension of  $K$  (see [LL99, Lemma 3.3]). Moreover, if  $g(Y) \geq 2$ , or  $g(Y) = 1$  and  $f$  is not étale, or if  $g(Y) = 0$  and  $g(X) \geq 1$ , then  $2g(Y) - 2 + |N| > 1$ . So after again a finite separable extension of  $K$ , we have a canonical way to define a minimal semi-stable reduction of  $X \rightarrow Y$  in which the (horizontal) ramification and branch loci are finite unions of sections contained in the smooth locus. If  $(\deg f)!$  is invertible in  $\mathcal{O}_S$ , then  $\mathcal{X}$  and  $\mathcal{Y}$  are the respective stable marked models of  $X$  and  $Y$  (see [Moc95, § 3.11, second lemma]).

Remark 4.18. Let  $\hat{X}, G, H$  be as in Remark 4.8. Let  $\hat{X}$  be marked with its ramification locus over  $X$  (rational over  $K$  after a finite extension of  $K$ ). Let  $\hat{\mathcal{X}}$  be its stable marked model. If  $G$  has order prime to  $\text{char}(k)$ , then the stable marked model of  $X \rightarrow Y$  is equal to  $\hat{\mathcal{X}}/H \rightarrow \hat{\mathcal{X}}/G$  because both sides are, respectively, stable marked models of  $X$  and  $Y$  (see Remark 4.17). However, this is false if  $\text{char}(k)$  divides  $|G|$ . Let us go back to the example of Remark 4.8 and, moreover, let  $C_1 \rightarrow D_1$  be étale outside of  $y_1$ . Let  $\mathcal{C}' \rightarrow \mathcal{C}, \mathcal{D}' \rightarrow \mathcal{D}$  be the stable marked hulls of  $\mathcal{C}, \mathcal{D}$  (marked with horizontal



ramification/branch loci). Let  $\mathcal{C}' \rightarrow \mathcal{C}''$ ,  $\mathcal{D}' \rightarrow \mathcal{D}''$  be the contraction of (the strict transforms of)  $C_1$ ,  $D_1$ , respectively. Then it is easy to see that the stable marked model (4.16) of  $X \rightarrow Y$  is  $\mathcal{C}'' \rightarrow \mathcal{D}''$ . Similarly to Remark 4.8, we see that it is different from  $\hat{\mathcal{X}}/H \rightarrow \hat{\mathcal{X}}/G$ .

It remains to find a cover  $C_1 \rightarrow D_1$  as above. Consider the cover<sup>1</sup> defined by the extension  $k(D_1) = k(u) \rightarrow k(C_1) = k(u, v)$ , with  $v^{p^2+p} + v = u$ , where  $p = \text{char}(k) > 0$ . Then  $C_1 \rightarrow D_1$  is étale outside of the pole  $y_1$  of  $u$ . Let us show that the Galois closure  $k(E)$  has positive genus. Let  $t \in k(E)$  be such that  $t^{p^2+p} + t = u$  and  $t \neq v$ . Let  $w = t - v$ . Then  $w$  satisfies the equation  $(w^{p+1} - wt^p - w^p t)^p - w = 0$ . Hence,  $w = z^p$  where  $z = w^{p+1} - wt^p - w^p t$ . We have  $z = z^{p^2+p} - z^p t^p - z^{p^2} t$ , so

$$(t/z^p)^p + (t/z^p) = 1 - (1/z)^{p^2+p-1}.$$

This equation defines a  $p$ -cyclic cover  $E' \rightarrow \mathbb{P}_k^1$ , with conductor  $m = p^2 + p - 1$  at  $z = 0$ , and étale elsewhere. Hence,  $g(E) \geq g(E') = (p - 1)(m - 1)/2 \geq 2$ .

*Remark 4.19.* If we restrict ourselves to the category of regular models of  $X$ , then Theorem 2.3 still holds. More precisely, given any model  $\mathcal{X}$ , there exists (after finite separable extension of  $K$ ) a unique semi-stable and regular model dominating  $\mathcal{X}$  and minimal for this property. This model is just the minimal desingularization of the stable hull of  $\mathcal{X}$ . However, it does not commute with base change except when the normalization of  $\mathcal{X}$  is smooth. On the other hand, Theorem 4.5 is no longer true in the setting of regular models. In general, given a morphism of models  $\mathcal{X} \rightarrow \mathcal{Y}$ , there is no finite morphisms of regular models dominating  $\mathcal{X} \rightarrow \mathcal{Y}$ , even after any finite extension of  $K$  (see [LL99, Remark 6.5]).

ACKNOWLEDGEMENTS

I would like to thank D. Lorenzini for pointing out Coleman’s paper [Col03] and for interesting comments, Ch. Deninger and A. Werner for motivational discussions and Sylvain Maugeais for pointing out some mistakes in an earlier version. I also thank the referee for correcting a mistake in the proof of Proposition 4.11.

REFERENCES

BL85 S. Bosch and W. Lütkebohmert, *Stable reduction and uniformization of abelian varieties, I*, Math. Ann. **270** (1985), 349–379.  
 BLR90 S. Bosch, W. Lütkebohmert and M. Raynaud, *Néron models*, Ergeb. Math. Grenzgeb. (3), vol. 21 (Springer, Berlin, 1990).  
 BLR95 S. Bosch, W. Lütkebohmert and M. Raynaud, *Formal and rigid geometry, IV. The reduced fibre theorem*, Invent. Math. **119** (1995), 361–398.  
 Col03 R. Coleman, *Stable maps of curves*, Doc. Math., extra volume K. Kato (2003), 217–225.  
 CES03 B. Conrad, B. Edixhoven and W. Stein,  *$J_1(p)$  has connected fibers*, Doc. Math. **8** (2003), 331–408.  
 DW04 Ch. Deninger and A. Werner, *Vector bundles on  $p$ -adic curves and parallel transport*, Preprint (2004), arxiv.org/abs/math/0403516.  
 Epp73 H. P. Epp, *Eliminating wild ramification*, Invent. Math. **19** (1973), 235–249.  
 GR66 H. Grauert, R. Remmert, *Über die Methode der diskret bewerteten Ringe in der nicht-archimedischen Analysis*, Invent. Math. **2** (1966), 87–133.  
 Hen99 Y. Henrio, *Arbre de Hurwitz et automorphismes d’ordre  $p$  des disques et des couronnes  $p$ -adiques formels*, Thesis, Université Bordeaux 1 (1999).

<sup>1</sup>This example is given by Michel Matignon.

- Dej96 A. J. de Jong, *Smoothness, semi-stability and alterations*, Publ. Math. Inst. Hautes Études Sci. **83** (1996), 51–93.
- Dej97 A. J. de Jong, *Families of curves and alterations*, Ann. Inst. Fourier (Grenoble) **47** (1997), 599–621.
- Kuh03 F.-V. Kuhlmann, *A correction to: ‘Elimination of wild ramification’ by H. P. Epp*, Invent. Math. **153** (2003), 679–681.
- Liu02 Q. Liu, *Algebraic geometry and arithmetic curves*, Oxford Graduate Texts in Mathematics, vol. 6 (Oxford University Press, Oxford, 2002).
- Liu03 Q. Liu, *Reduction and lifting of finite covers of curves*, in *Proc. of the 2003 workshop on cryptography and related mathematics* (Chuo University, Tokyo, 2003), 161–180.
- LL99 Q. Liu, D. Lorenzini, *Models of curves and finite covers*, Compositio Math. **118** (1999), 61–102.
- Mau05 S. Maugeais, *On a compactification of a Hurwitz space in the wild case*, Preprint (2005).
- Moc95 S. Mochizuki, *The geometry of the compactification of the Hurwitz scheme*, Publ. Res. Inst. Math. Sci. **31** (1995), 355–441.
- Ray90 M. Raynaud,  *$p$ -groupes et réduction semi-stable des courbes*, in *The Grothendieck Festschrift*, vol. III, eds P. Cartier *et al.*, Progress in Mathematics, vol. 88 (Birkhäuser, Basel, 1990), 179–197.
- Ser68 J.-P. Serre, *Corps locaux* (Hermann, Paris, 1968).

Qing Liu [Qing.Liu@math.u-bordeaux1.fr](mailto:Qing.Liu@math.u-bordeaux1.fr)

CNRS, Laboratoire A2X, Institut de Mathématiques de Bordeaux, Université de Bordeaux I, 351, cours de la Libération, 33405 Talence, France