

Exploring Tonsil Pathology in PTEN Hamartoma Syndrome: A Cohort Study

Lauren M. Cairns*^{1,2}, Callum Findlay*^{3,4}, Nimeshi Jayakody⁴, Kwamena Amonoo-Kuofi⁴,
Nimesh N. Patel⁴, Katherine L. Lachlan^{1,5}

*Equal first authors

1. Wessex Clinical Genetics Service, University Hospital Southampton NHS Foundation Trust, UK.
2. Royal Berkshire Hospital, Royal Berkshire NHS Foundation Trust, UK.
3. Clinical and Experimental Sciences, Faculty of Medicine, University of Southampton, UK.
4. Department of Otolaryngology, Head and Neck Surgery, University Hospital Southampton NHS Foundation Trust, UK.
5. Human Genetics and Genomic Medicine, Faculty of Medicine, University of Southampton, UK.

Corresponding author: Katherine L. Lachlan

Katherine.Lachlan@uhs.nhs.uk

Wessex Clinical Genetics Service, University Hospital Southampton NHS Foundation Trust, Southampton, UK.

Introduction

PTEN hamartoma tumour syndrome (PHTS) comprises a heterogeneous autosomal dominant condition encompassing distinct clinical entities including Bannayan-Riley-Ruvalcaba syndrome, Cowden syndrome, Lhermitte-Duclos Disease, and macrocephaly-autism/developmental delay syndrome. Pathogenic variants in the PTEN (Phosphatase and tensin homolog deleted on chromosome 10) gene underlie the multifaceted PHTS¹⁻⁶. Marked by macrocephaly, developmental delay, mucocutaneous lesions, and an increased susceptibility to diverse cancers (breast, endometrial, thyroid, renal, and colon) PHTS presents a diagnostic challenge due to its variable and age-related manifestations^{4,7-11}.

Case reports have shown evidence that PHTS is associated with tonsillar pathology, in particular early-onset enlargement of tonsillar tissue. Various age of onset and severity of airway obstruction have been illustrated in the current literature. Mild tonsillar hypertrophy and a papilloma were found in a 2 year old, with the papilloma becoming larger and rougher after 6 months¹². A 3-year-old boy was noted to have hyperplasia of the pharyngeal tonsils¹³, while a 6 year old girl had more severe tonsillar hypertrophy resulting in sleep apnoea and subsequent tonsillectomy¹⁴ and an 11 year old boy developed open-mouth breathing from hypertrophic tonsils and adenoid vegetation¹⁵. Notably, an adult (55-year-old woman) was found to have airway obstruction during induction of general anaesthetic caused by multiple papillomas on the lingual tonsils, epiglottis and the surrounding structure. She was diagnosed with PHTS later post operatively¹⁶.

Despite these case report findings, studies examining tonsillar pathology within a specific PHTS patient cohort are absent. Our case series addresses this gap by reviewing the tonsil-related pathology in this distinct patient population, through rigorous examination of clinical records, diagnostic imaging, and histopathological data.

Methods

53 patients with *PTEN* mutations known to the Wessex Clinical Genetics Service were included in the study. The local electronic patient records of these patients were reviewed between April to June 2023. Information was extracted including the date and age of *PTEN* diagnosis, individual's genetic mutation, tonsil pharyngeal conditions, ENT operations, histology and post-operative course, focusing on tonsil issues and subsequent tonsillectomy details retrospectively.

Results

A cohort of 53 patients diagnosed with PHTS were included in this study. Among these patients, 28 (53%) were adults, while 25 (47%) were children. The gender distribution was 33 (62%) male and 20 (38%) female.

Tonsil related issues were prevalent in 31 (58%) of the PHTS patients, with 23 (74%) of them undergoing subsequent tonsillectomy (43% of the total cohort). The primary indication for tonsillectomy was obstructive sleep apnea (OSA), accounting for 10 (43%) cases, followed by recurrent tonsillitis in 4 (17%) cases, unknown indications in 4 (17%) cases, and both OSA and recurrent tonsillitis in 3 (13%) cases. Additionally, one patient exhibited asymmetrical tonsils, and another required emergency surgery due to acute airway obstruction.

Eight of the cohort (15%) exhibited tonsil issues but did not undergo surgical intervention. These individuals all presented with enlarged tonsils, with 5 (63%) having normal sleep study results, 2 displaying possible sleep apnea and having ongoing surveillance, and 1 showing no symptoms.

Regarding the timing of tonsil issues in relation to PHTS diagnosis, 14 (45%) patients experienced tonsil problems before the PHTS diagnosis, while 17 (55%) developed these issues subsequent to the diagnosis.

Tonsillectomy was the predominant surgical procedure, all patients who underwent surgery had this procedure, along with adenoidectomy in 20 (87%) cases. One adenoidectomy was a revision following a previous procedure that involved adenoidectomy and grommet insertion. The age range of PHTS patients undergoing tonsillectomy spanned from 1 to 27 years, with a mean age of 7.8 years. When the analysis was restricted to only include patients under 30 years at the time of data collection, there was an increase in the percentage of tonsil issues (71% 25/35) and tonsillectomies (49% 17/35) compared to the whole cohort (58% and 43% respectively). This reflects an age-related ascertainment bias likely due to older patients not having tonsil related diagnoses recorded.

No cases were severe enough for a post-op HDU bed. Histology information collection was limited due to surgeries taking place across different hospitals and years (1979 to 2020). Nonetheless, histology details were accessible for the 4 (17%) patients who underwent surgery in Southampton within the last decade, all indicating benign reactive lymphoid hyperplasia.

Among the PHTS patients who underwent tonsillectomies, 6 (26%) experienced post-surgery problems. These complications included ongoing loud snoring but with resolved apneas and normal sleep studies, tonsil remnants, hamartomas in the nasopharynx and tongue base and a single case of recurrent adenoidal hypertrophy necessitating a repeat adenoidectomy 6 years after the initial procedure.

Discussion

The present study offers significant insights into the relationship between PTEN Hamartoma Syndrome (PHTS) and tonsil pathology, shedding light on the prevalence, characteristics, and clinical implications of tonsil-related pathology within this patient population. Our findings underscore that tonsil related issues are a common problem in patients with PHTS that should be considered in their medical management.

Our study cohort consisted of 53 patients diagnosed with PHTS, exhibiting a gender distribution favouring males (62%) and encompassing a diverse age range spanning from children to adults. Tonsil issues emerged as a significant concern within our study cohort, affecting 58% of patients, much greater than the 11% incidence previously reported in school aged children¹⁷. These findings highlight the importance of recognising tonsil pathology as a potential aspect of PHTS that demands clinical attention. Moreover, among those with tonsil-related issues, 74% required tonsillectomy, a surgical intervention to remove the palatine tonsils for histological analysis or remove tissue causing upper airway obstruction. This accounted to 43% of the total cohort which is significantly more than the 13.6% of the general population undergoing tonsillectomy¹⁸. The primary indication for tonsillectomy was obstructive sleep apnea (OSA), a finding consistent with reports of increased OSA prevalence in individuals with PHTS. Importantly this OSA was never associated with peri-operative respiratory complications and HDU and ITU support was not required. Our study further delineated other indications for tonsillectomy, including recurrent tonsillitis. Notably, cases of asymmetrical tonsils and emergency surgical intervention for acute airway obstruction underscore the severe and sometimes urgent nature of tonsil-associated complications in PHTS.

Tonsillectomy was the principal surgical approach, often accompanied by adenoidectomy (87%). The inclusion of adenoidectomy is consistent with the need to address upper airway obstruction and sleep apnoea in individuals with PHTS. The rate of complications and revision surgery was not unexpected and is comparable to adenotonsillectomy in other groups.

Histological evaluation presented challenges due to the variety of hospitals and years during which surgeries were performed. Despite this limitation, histology information for a subset of patients undergoing surgery in Southampton within the last decade indicated benign

reactive lymphoid hyperplasia. This finding echoes previous reports of lymphoid hyperplasia associated with PHTS and underlines the need for detailed histopathological assessment to guide clinical decisions ¹³.

Our results reveal that tonsil related problems were identified and treated both before and after the diagnosis of PHTS was made. One patient in our cohort was diagnosed after the ENT surgeon performing the tonsillectomy recognised the macrocephaly and developmental delay and referred the patient to clinical genetics. This highlights the need for ENT surgeons to consider PHTS and be aware of its clinical manifestations when treating children for OSA secondary to tonsil hypertrophy.

Distinguishing patients with PTEN Hamartoma Syndrome (PHTS) from other patients requires a comprehensive approach that considers clinical, genetic, and diagnostic factors. As PHTS encompasses a diverse array of manifestations, including those affecting the head and neck region, distinguishing features can aid Ear, Nose, and Throat (ENT) surgeons in identifying individuals with PHTS. Here are some key considerations for ENT surgeons to differentiate patients with PHTS from others:

Clinical Phenotype: These may include macrocephaly, mucocutaneous lesions (trichilemmomas, oral papillomas), developmental delay, and specific cancer susceptibilities (breast, thyroid, endometrial). The presence of these features, especially when combined, should raise suspicion for PHTS.

Family History: Inquiring about a family history of PHTS-related conditions in this autosomal dominant condition may identify other family members with medical history consistent with PHTS prompting referral to consider genetic testing. If a pathogenic variant is identified in the family, predictive testing can be offered. This facilitates screening and discussion of risk reducing surgery, decreasing risk of malignancy.

Histopathological Findings: Skin biopsies of characteristic lesions, like trichilemmomas and oral papillomas, can provide histopathological evidence supporting the diagnosis of PHTS. Consultation with dermatopathologists can aid in recognising these distinctive features.

Genetic Testing: Genetic testing for mutations in the PTEN gene is a cornerstone in diagnosing PHTS. If clinical suspicion is high, patients should be referred to genetics specialists for appropriate genetic counseling and testing. Detection of pathogenic PTEN variants confirms the diagnosis of PHTS.

In summary, distinguishing patients with PTEN Hamartoma Syndrome from others requires a comprehensive approach that integrates clinical assessment, histopathology, and collaboration with specialists from various disciplines. A high index of suspicion, particularly when encountering a constellation of characteristic clinical features, can prompt timely diagnosis and appropriate management of individuals with PHTS.

Conclusion

In conclusion, our study highlights the crossover between PTEN Hamartoma Syndrome (PHTS) and Ear, Nose, and Throat (ENT) pathology. By reviewing a cohort of 53 PHTS

patients, we have found tonsil hyperplasia is a common condition in patients with PHTS and frequently causes obstructive sleep apnoea in the first decade of life.

The prevalence of tonsil issues in our study cohort emphasizes the need for targeted clinical evaluation for this. Clinical Geneticists and paediatricians caring for patients with PHTS should consider whether the patient has OSA and examine their tonsils. Conversely, ENT surgeons should consider whether children presenting with OSA and tonsillar hypertrophy also exhibit other signs of PHTS and consider referral for genetic testing.

Financial Support

This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

Competing Interests

The authors declare none.

Summary

- PTEN Hamartoma Tumour Syndrome (PHTS) encompasses a group of genetic disorders with varied clinical presentations (macrocephaly, developmental delay, mucocutaneous lesions and increased cancer susceptibility).
- Recent case reports have illustrated the occurrence of tonsil-related issues in PHTS.
- Tonsils issues were prevalent in 58% of the cohort.
- 43% of the PHTS cohort required tonsillectomy, compared to 13.6% of the general population.
- This study highlights the association of PHTS with tonsillar pathology throughout early life and its relevance for ENT surgeons.
- Early recognition and intervention are pivotal for managing sleep apnea and the associated health problems.

References

1. Arch EM, Goodman BK, Van Wesep RA, Liaw D, Clarke K, Parsons R, et al. Deletion of PTEN in a patient with Bannayan-Riley-Ruvalcaba syndrome suggests allelism with Cowden disease. *Am J Med Genet* 1997;**71**:489–93
2. Liaw D, Marsh DJ, Li J, Dahia PL, Wang SI, Zheng Z, et al. Germline mutations of the PTEN gene in Cowden disease, an inherited breast and thyroid cancer syndrome. *Nat Genet* 1997;**16**:64–7

3. Marsh DJ, Dahia PLM, Zheng Z, Liaw D, Parsons R, Gorlin RJ, et al. Germline mutations in PTEN are present in Bannayan-Zonana syndrome. *Nat Genet* 1997;**16**:333–4
4. Marsh DJ, Kum JB, Lunetta KL, Bennett MJ, Gorlin RJ, Ahmed SF, et al. PTEN Mutation Spectrum and Genotype-Phenotype Correlations in Bannayan-Riley-Ruvalcaba Syndrome Suggest a Single Entity With Cowden Syndrome. *Hum Mol Genet* 1999;**8**:1461–72
5. Nelen MR, Padberg GW, Peeters EAJ, Lin AY, Helm B van den, Frants RR, et al. Localization of the gene for Cowden disease to chromosome 10q22–23. *Nat Genet* 1996;**13**:114–6
6. Zhou X-P, Marsh DJ, Morrison CD, Chaudhury AR, Maxwell M, Reifemberger G, et al. Germline Inactivation of PTEN and Dysregulation of the Phosphoinositol-3-Kinase/Akt Pathway Cause Human Lhermitte-Duclos Disease in Adults. *The American Journal of Human Genetics* 2003;**73**:1191–8
7. Celebi JT, Ping XL, Zhang H, Remington T, Sulica VI, Tsou HC, et al. Germline PTEN mutations in three families with Cowden syndrome. *Exp Dermatol* 2000;**9**:152–6
8. DiLiberti JH. Inherited macrocephaly-hamartoma syndromes. *Am J Med Genet* 1998;**79**:284–90
9. Hendriks YMC, Verhallen JTCM, van der Smagt JJ, Kant SG, Hilhorst Y, Hoefsloot L, et al. Bannayan-Riley-Ruvalcaba syndrome: further delineation of the phenotype and management of PTEN mutation-positive cases. *Fam Cancer* 2003;**2**:79–85
10. Lachlan KL, Lucassen AM, Bunyan D, Temple IK. Cowden syndrome and Bannayan Riley Ruvalcaba syndrome represent one condition with variable expression and age-related penetrance: results of a clinical study of PTEN mutation carriers. *J Med Genet* 2007;**44**:579–85
11. Zori RT, Marsh DJ, Graham GE, Marliss EB, Eng C. Germline PTEN mutation in a family with Cowden syndrome and Bannayan-Riley-Ruvalcaba syndrome. *Am J Med Genet* 1998;**80**:399–402
12. Piccione M, Fragapane T, Antona V, Giachino D, Cupido F, Corsello G. PTEN hamartoma tumor syndromes in childhood: Description of two cases and a proposal for follow-up protocol. *Am J Med Genet A* 2013;**161**:2902–8
13. Heindl M, Hndel N, Ngeow J, Kionke J, Wittekind C, Kamprad M, et al. Autoimmunity, intestinal lymphoid hyperplasia, and defects in mucosal B-cell homeostasis in patients with PTEN hamartoma tumor syndrome. *Gastroenterology* W.B. Saunders: 2012;**142**
14. Abdul Latiff Zarina, Warouw Atmawidjaja Raymond, Raja Juanita Raja, Raja Lope Raja, Sharifah Azween Sharifah, Syed Omar Syed, et al. Bannayan Riley Ruvalcaba Syndrome. *Annals Academy of Medicine* 2010;**39**:578–9
15. Erkek E, Hizel S, Sanlı C, Erkek AB, Tombakoglu M, Bozdogan O, et al. Clinical and histopathological findings in Bannayan-Riley-Ruvalcaba syndrome. *J Am Acad Dermatol* 2005;**53**:639–43
16. Omote Keiichi, Kawamata Tomoyuki, Imaizumi Hitoshi, Namiki S Akiyoshi. Case of Cowden's Disease that Caused Airway Obstruction during Induction of Anesthesia [Internet]. *Anesthlesiol&y.* 1999
17. Kara CO, Ergin H, Koçak G, Kılıç İ, Yurdakul M. Prevalence of tonsillar hypertrophy and associated oropharyngeal symptoms in primary school children in Denizli, Turkey. *Int J Pediatr Otorhinolaryngol* 2002;**66**:175–9

18. Šumilo D, Nichols L, Ryan R, Marshall T. Incidence of indications for tonsillectomy and frequency of evidence-based surgery: a 12-year retrospective cohort study of primary care electronic records. *Br J Gen Pract* 2019;**69**:e33–41