

## Standard Paper

# PhyloKey: a novel method to rapidly and reliably identify species in complex, species-rich genera, and an opportunity for ‘non-molecular museomics’

Robert Lücking<sup>1</sup> , Bibiana Moncada<sup>1,2</sup>  and Manuela Dal Forno<sup>3,4</sup> 

<sup>1</sup>Botanischer Garten und Botanisches Museum Berlin, Freie Universität Berlin, 14195 Berlin, Germany; <sup>2</sup>Licenciatura en Biología, Universidad Distrital Francisco José de Caldas, Torre de Laboratorios, Herbario, Bogotá D.C., Colombia; Research Associate, Science & Education, The Field Museum, Chicago, IL 60605, USA; <sup>3</sup>Botanical Research Institute of Texas, Fort Worth Botanic Garden, Fort Worth, TX 76107, USA and <sup>4</sup>Research Associate, Department of Botany, Smithsonian Institution, National Museum of Natural History, Washington, DC 20560, USA

### Abstract

We present a novel identification tool called *PhyloKey*, based on the method of morphology-based, phylogenetic binning developed within the software package RAXML. This method takes a reference data set of species for which both molecular and morphological data are available, computes a molecular reference tree, maps the morphological characters on the tree, and computes weights based on their level of consistency versus homoplasy using maximum likelihood (ML) and maximum parsimony (MP). Additional units for which only morphological data are known are then binned onto the reference tree, calculating bootstrap support values for alternative placements. This approach is modified here to work as an identification tool which uses the same character coding approach as interactive keys. However, rather than identifying individual samples through a progressive filtering process when entering or selecting characters, query samples are binned in batch mode to all possible alternative species in the tree, with each placement receiving a bootstrap support adding to 100% for all alternative placements. In addition to the fact that, after scoring a character matrix, a large number of specimens can be identified at once in short time, all possible alternative identifications are immediately apparent and can be evaluated based on their bootstrap support values. We illustrate this approach using the basidiolichen genus *Cora*, which was recently shown to contain hundreds of species. We also demonstrate how the *PhyloKey* approach can aid the restudying of herbarium samples, adding further value to these collections and contributing with large quantitative data matrices to ‘non-molecular museomics’. Our analysis showed that *PhyloKey* identifies species correctly with as low as 50% of the characters sampled, depending on the nature of the reference tree and the character weighting scheme. Overall, a molecular reference tree worked best, but a randomized reference tree gave more consistent results, whereas a morphological reference tree performed less well. Surprisingly, even character weighting gave the best results, followed by parsimony weighting and then maximum likelihood weighting.

**Keywords:** biodiversity; integrative taxonomy; lichens; multi-access key; RAXML

(Accepted 29 May 2023)

### Introduction

Species are the basic units of biodiversity and important for all areas of basic and applied organismic research (Riddle & Hafner 1999; De Queiroz 2005; Wilkins 2017; Reydon 2019). As such, the most important instruments provided by taxonomists are identification tools (Sluys 2013; Lücking 2020). For most of the past 250 years, dichotomous, printed keys were the *status quo* for taxonomic identifications (Walter & Winterton 2007; Hagedorn *et al.* 2010). Such keys can be used without specific devices or software and they guide the user through the identification process, based on the fact that decision-making for the

human brain is facilitated by the existence of generally two alternatives laid out at each key couplet. The disadvantage of dichotomous keys lies in their fixed sequence and entry point (single-access), not allowing the selection of characteristic features that would immediately identify a particular taxon at hand.

With the advent of computing, machine-based, interactive keys became increasingly popular and are currently the standard in many areas (Edwards & Morse 1995; Dallwitz *et al.* 2006; Mayo *et al.* 2008; Nimis *et al.* 2012; Nimis & Martellos 2020; Murguía-Romero *et al.* 2021). These keys are based on a data matrix for known species with a set number of characters and character states coded in a specific manner, usually binary or in ordinal or categorical fashion. The same characters are scored for specimens to be identified and the identification is based on scores of agreement or similarity. The advantage of such keys is that they have flexible entry points (multi-access), which means that identification speed is increased by selecting individually

**Corresponding author:** Robert Lücking; Email: [r.luecking@bo.berlin](mailto:r.luecking@bo.berlin)

**Cite this article:** Lücking R, Moncada B and Dal Forno M (2023) *PhyloKey*: a novel method to rapidly and reliably identify species in complex, species-rich genera, and an opportunity for ‘non-molecular museomics’. *Lichenologist* 55, 181–192. <https://doi.org/10.1017/S0024282923000415>

© The Author(s), 2023. Published by Cambridge University Press on behalf of the British Lichen Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.



diagnostic characters for each specimen. In addition, many such applications offer a select set of best diagnostic characters based on initial character entry, thus guiding the character scoring process more effectively. The disadvantage lies in the necessary assembly of a full character matrix for all known species in a group, the required translation of a variety of characters into discrete scores (although some interactive keys also allow the use of morphometrics), and the need for access to a computer or compatible device (e.g. cell phone).

Organisms with underlying symmetry patterns also offer themselves for the use of image recognition for identification purposes, including plant identification applications (e.g. *LeafSnap*, *Pl@ntNet*, and *iNaturalist*; La Salle *et al.* 2009; Goëau *et al.* 2013; Joly *et al.* 2016). In addition, molecular identification using DNA barcoding loci is a trend focusing more on scientific applications (Seberg *et al.* 2003; Smith 2005; Schoch *et al.* 2012; DeSalle & Goldstein 2019; Lücking *et al.* 2020a). However, the latter approach is not broadly practicable, for example in community science, and it is often overlooked that a complete inventory of all species of a group, with correctly labelled barcoding sequences, is required before any molecular identification tool could work (Lücking *et al.* 2020b).

An advantage of molecular identification is the possibility of immediate feedback on the confidence of the identification, by means of phylogenetic assembly including statistical support through bootstrapping or posterior probabilities, such as achieved by sequence placement methods (Zhang *et al.* 2013; Carbone *et al.* 2017). Such statistical feedback is presently not available for non-molecular identification tools or for BLAST-based DNA barcoding, although it could theoretically be implemented for interactive keys. With dichotomous keys, users are often uncertain between two alternatives and a wrong turn will lead to a wrong identification or mismatch. Interactive keys seemingly avoid this problem by using discrete characters, but the problem is only shifted to the moment of correct character recognition. Some interactive tools, such as DELTA Intkey (Dallwitz *et al.* 2006), allow for a proportion of mismatches to bring up possible alternatives. However, the more complex and species-rich a genus, the more the user will become confused and uncertain about the accuracy of identification outcomes.

Here, we present a novel identification tool called *PhyloKey*, based on the method of morphology-based, phylogenetic binning (Berger *et al.* 2011; Lücking & Kalb 2018). This method takes a reference data set of species for which both molecular and morphological data are available, computes a molecular reference tree, maps the morphological characters on the tree and computes weights based on their level of consistency versus homoplasy using maximum likelihood (ML) and maximum parsimony (MP). Additional units for which only morphological data are known are then binned onto the reference tree, calculating bootstrap support values for alternative placements, an approach that can be modified into an identification tool.

We illustrate this approach using the basidiolichen genus *Cora*, which was recently shown to contain hundreds of species (Lücking *et al.* 2014, 2017; Dal Forno *et al.* 2022). Currently, around 265 species are distinguished using a combination of molecular and morphological data (Dal Forno *et al.* 2022). A total of 105 species has been formally described (Lücking *et al.* 2013, 2015a, 2017, 2020c; Vargas *et al.* 2014; Ariyawansa *et al.* 2015; Moncada *et al.* 2019) and 87 of these have molecular and comprehensive phenotype data available. *Cora* is an ideal model case to implement the *PhyloKey* approach, since it features only a small number of diagnostic characters compared to other


macrolichens and the characters are partly homoplastic, features that often lead to failure when using traditional dichotomous keys. We also demonstrate how the *PhyloKey* approach can help with restudying historical samples and thus contribute to ‘museomics’ by integrating them with molecular data.

It is with great pleasure that we dedicate this paper to our esteemed colleague and friend, Pier Luigi Nimis, on the occasion of his 70th birthday and his well-deserved retirement from a long and outstanding professional career. Besides his invaluable contributions to lichenology, particularly in Italy, Pier Luigi has greatly advanced the use of traditional and digital identification tools for lichens and other organisms.

## Material and Methods

Based on the studies by Lücking *et al.* (2014, 2017), we compiled three data sets of the genus *Cora* Fr. to illustrate and test *PhyloKey*: 1) a molecular alignment of the ITS fungal barcoding locus for 87 formally described and sequenced species plus two outgroup species of the genus *Corella* Vain. (‘alignment’ in fasta format; Supplementary Material File S1, available online); 2) a matrix of 20 characters for the 87 ingroup species (the same as analyzed in Dal Forno *et al.* (2022)), divided into one ecological (substratum), 11 phenotype (morphology, anatomy, chemistry; Fig. 1), and eight distributional characters (main distribution areas; ‘reference matrix’ in Phylip format; Table 1, Supplementary Material File S2, available online); and 3) a matrix of the same 20 characters for 398 samples to be identified (‘query matrix’ in Phylip format; Supplementary Material File S3, available online). For the latter, 200 test samples (‘samples’) were generated from the original data matrix by randomly selecting ten out of the 87 species (*C. applanata* B. Moncada *et al.*, *C. caliginosa* Holgado *et al.*, *C. campestris* Dal-Forno *et al.*, *C. crispoleslia* B. Moncada *et al.*, *C. davibogotana* Lücking *et al.*, *C. dewisanti* B. Moncada *et al.*, *C. dulcis* B. Moncada *et al.*, *C. fimbriata* L. Y. Vargas *et al.*, *C. pichinchensis* Paredes *et al.* and *C. soledavidia* Dal-Forno *et al.*), and 20 samples per species were generated by randomly deleting an increasing number of characters, from zero to 19, for each species. This data set was used to assess the effect of incomplete character sampling on the accuracy of species identification. In addition, we randomly selected 20 samples from herbarium collections held in B traditionally identified as *Cora pavonia* or *Dictyonema glabratum* (Table 2), to assess whether some of these would match described species and how potentially undescribed species would behave using this approach. We further added all 89 ingroup and outgroup species twice to the data set, once as reference and once for calibration purposes.

The molecular reference tree was computed from the molecular alignment through a maximum likelihood search with RAXML v. 8.2.8 (Stamatakis 2014), with non-parametric bootstrapping using 1000 replicates under a GTRGAMMA model. To test the effect of topology on the performance of the key, we also computed two additional trees. A morphological reference tree was built by subjecting the morphological matrix to a maximum likelihood search, also using RAXML v. 8.2.8 (Stamatakis 2014), with non-parametric bootstrapping using 1000 replicates under a MULTIGAMMA model. In addition, a random tree was generated in PAUP v. 4.0 b10 (Swofford 2003). The molecular and morphological reference trees included branch lengths (phylograms), whereas the random tree was used as a simple cladogram (‘reference trees’ in Newick format; Supplementary Material File S4, available online).



**Figure 1.** Selected characters and character states used to score *Cora* species and samples. For complete list of characters and states, see Table 1. A, sutures in *C. suturifera* Nugra *et al.* B, rugose surface in *C. auricleslia* B. Moncada *et al.* C, narrowly undulate surface in *C. celestino* B. Moncada *et al.* D, broadly undulate surface in *C. imi* Lücking *et al.* E, pitted surface in *C. elephas* Lücking *et al.* F, setose upper surface in *C. barbifera* B. Moncada *et al.* G, strigose upper surface in *C. hirsuta* (B. Moncada & Lücking) Moncada & Lücking. H, soredia in *C. hawksworthiana* Dal-Forno *et al.* I, viaduct-shaped upper cortex in *C. leslactu* Lücking *et al.* J, paraplectenchymatous upper cortex in *Corella melvinii* (Chaves *et al.*) Lücking *et al.* K, papillae in the lower medulla in *Cora haledana* Dal-Forno *et al.* L, adnate hymenophore in *C. soredata*. M, concentric hymenophore in *C. viliewoa* Lücking *et al.* N, cyphelloid hymenophore in *C. benitoana* B. Moncada *et al.* O, pigment (after rewetting) in *C. rubrosanguinea* Nugra *et al.* In colour online.

**Table 1.** Characters used to score *Cora* species and samples. For visual character definitions, see Fig. 1, and for additional details on character definitions, see Dal Forno *et al.* (2022).

Character	State 1	State 2	State 3	State 4	State 5
Substratum	saxicolous = 1	terricolous = 2	terrestrial = 3	epiphytic = 4	
Lobe size	small = 1	medium = 2	large = 3		
Sutures	absent = 0	short = 1	distinct = 2		
Colour	grey = 1	brown = 2	olive = 3	green = 4	blue = 5
Surface	even = 0	rugose = 1	pitted = 2	broadly undulate = 3	narrowly undulate = 4
Trichomes	absent = 0	felty = 1	setose = 2	strigose = 3	
Margin	glabrous = 0	pilose = 1	granular = 2	granular-pilose = 3	
Soredia	absent = 0	present = 1			
Cortex	viaduct-shaped = 1	collapsed = 2	compacted = 3	plettenchymatous = 4	paraplettenchymatous = 5
Papillae	absent = 0	present = 1			
Hymenophore	adnate = 1	concentric = 2	concentric-cyphelloid = 3	cyphelloid = 4	
Pigment	absent = 0	present = 1			
Central America	absent = 0	present = 1			
Caribbean	absent = 0	present = 1			
Galapagos	absent = 0	present = 1			
Northern Andes	absent = 0	present = 1			
Central Andes	absent = 0	present = 1			
Southern Andes	absent = 0	present = 1			
Brazil	absent = 0	present = 1			
Palaeotropics	absent = 0	present = 1			

Morphology-based phylogenetic binning is a two-step approach, first calculating the character weight vectors, using either maximum likelihood (ML) or maximum parsimony (MP), and then binning the samples to be identified onto the reference tree (Berger *et al.* 2011). For the weight vectors, we ran the vector analysis in RAXML v. 7.2.6 (Stamatakis *et al.* 2005) relating the matrix (in Phylip format) to each of the reference trees (in Newick format). The command line runs [raxmlHPC.exe -f u -m MULTIGAMMA -s matrix.phy -t reference.tre -n weight\_vector.txt] for the ML weight vector and is identical but with upper case U [-f U] for the MP weight vector. In addition, we employed a uniform weight vector and an arbitrary vector weighting character based on their ease of observation and distinctiveness (Table 3). For the latter, we assigned three weights: 100% for characters easy to observe, discrete or unique (e.g. soredia, hymenophore type, distribution in the Palaeotropics); 50% for characters difficult to observe, subtle, or continuous, or for ecological and most chorological characters (e.g. colour, lobe size, substratum); and 75% for characters that we considered intermediate in this respect (Table 3).

The matrix of samples to be identified requires inclusion of the species present in the reference tree, so the complete matrix contained 87 (known species) + 2 (outgroup) + 220 (samples) units. The samples were then binned onto each of the three reference trees, with three different weight vectors (even, MP, ML), in RAXML v. 7.2.6, with 1000 bootstrap replicates, using the

command line [raxmlHPC.exe -f v -m MULTIGAMMA -a weight\_vector.txt -s samples.phy -t reference.tre -n identification.txt -x 12345 -# 1000], which corresponds to the Evolutionary Placement Algorithm (EPA; Berger & Stamatakis 2010; Berger *et al.* 2011). This resulted in a total of nine combinations for the same 220 samples (three reference trees × three weight vectors). In addition, we tested the fourth (arbitrary) weight vector with the random tree.

For each of the randomly selected samples with decreasing number of characters binned and for each reference tree, we computed a combined score across the three weighting schemes as follows:

$$S_{\text{Ref}} = (N_{\text{MP}} \times B_{\text{MP}} + N_{\text{ML}} \times B_{\text{ML}} + N_{\text{even}} \times B_{\text{even}}) / 3000,$$

where  $S_{\text{Ref}}$  = combined score for each reference tree (molecular, morphological, random),  $N_{\text{MP}}$ ,  $N_{\text{ML}}$ ,  $N_{\text{even}}$  = number of correctly binned species, and  $B_{\text{MP}}$ ,  $B_{\text{ML}}$ ,  $B_{\text{even}}$  = mean bootstrap support for correctly binned species using MP, ML, and even weights.

The analytical output from RAXML includes a number of files in text format, two of which were used for visualization (all other output files can be discarded). One is the classification table, a tabular text file named 'RAXML\_classification', with the node placements of each query taxon and the corresponding bootstrap support values. This text file was adjusted for inspection in a spreadsheet editor, in this case Microsoft Excel, by globally

**Table 2.** Details of herbarium specimens held at B and traditionally identified as *Cora pavonia* or *Dictyonema glabratum*, used for the *PhyloKey* test.

Sample	Location	Barcode
Abrahamczyk s. n.	Mexico, Michoacán	B 60 0158981
Bach (et al.) 249	Bolivia, La Paz	B 60 0106364
Bach (et al.) 425	Bolivia, La Paz	B 60 0106366
Bach (et al.) 502	Bolivia, La Paz	B 60 0106368
Bach (et al.) 535	Bolivia, La Paz	B 60 0106369
Cleef (& Fernández-P.) 677	Colombia, Cauca	B 60 0146625
Cleef 2074	Colombia, Boyacá	B 60 0146627
Cleef 5232	Colombia, Cundinamarca	B 60 0146629
Florschuetz 3608a	Colombia, Cundinamarca	B 60 0146618
Follmann 35322	Chile, Región de Los Lagos	B 60 0160525
Hatschbach 52096	Brazil, Paraná	—
Krieger 13527	Brazil, Minas Gerais	B 60 0128817
LSE ( <i>Lichenes Selecti Exsiccati</i> ) 2445 Vivant s. n.	Guadeloupe	—
Rapp 581	USA, Florida	—
Sipman (& Aptroot) 19245	Guyana, Upper Mazaruni District	—
Sipman 31805	French Guiana	—
Sipman (et al.) 37684	El Salvador, Chalatenango	B 60 0106355
Sipman (et al.) 37777	El Salvador, Chalatenango	B 60 0106356
Steglich s. n.	Venezuela, Mérida	B 60 0164979
Welzen 1122	Costa Rica, San José	B 60 0164979

replacing spaces with tabulators (alternatively, the table can be opened in Excel using space as separator). The second file is the classification tree, originally in Newick format, named 'RAXML\_labelledTree', which can be opened in various tree viewing editors. Here, we used FigTree v. 1.4.4 (Rambaut 2018), for which the output file had to be adjusted as follows prior to opening: 1) globally replacing the string ':1.0]' with '[' and then ']' with ']:1.0' (which switched the order of branch length and node ID labels); 2) checking instances of identical terminal output names (which can occur as the bootstrap support values are added to the query names and alternative placements can have identical support). Such instances were then made unique by adding the suffix letters 'a', 'b', etc.

## Results

For the 'simulated' samples, the three different reference tree approaches resulted in overall similar outcomes in the individual results (Table 4; Supplementary Material File S5, available online) and the combined score (Fig. 2). In all three cases, samples with complete character sets were placed correctly, with a placement-support score (PSS) of 1.00 for the molecular and morphological reference trees and 0.99 for the random tree. Performance in terms of correct placement declined with increasing number of missing characters but resulted in very high PPS (0.95 or higher) down to three missing characters for the molecular and random

trees, and high PPS (0.70 or higher) down to ten missing characters for the molecular tree, four missing characters for the morphological tree, and eight missing characters for the random tree. The threshold of 0.50 PPS was reached for all trees at ten missing characters, followed by a strong drop in performance, and 15 or more missing characters resulted in a PPS of close to zero in all three approaches. The molecular reference tree performed best overall, followed by the random and the morphological reference trees. The random reference tree performed most consistently (Fig. 3), with the least amount of variation as a function of increasing number of missing characters, whereas both the molecular and the morphological trees had positive and negative spikes for particular proportions of missing characters (Fig. 2).

For the molecular reference tree, even and maximum likelihood (ML) weighting performed better than maximum parsimony (MP) weighting, both with nine out of ten correct placements with as many as ten missing characters; however, MP weighting was more consistent, with both even and ML weighting showing positive and negative spikes. The morphological reference tree showed a similar result between all three weighting approaches, whereas with the random tree, even weighting outperformed both MP and ML weighting (Table 4).

The test with herbarium samples, including all known species with a full character set each for calibration, placed the 89 known species correctly using even character weights, with 99.1% average bootstrap support; out of these, 80 species were placed correctly with 100% support, 85 species with  $\geq 95\%$ , and two species (*Cora squamiformis* Wilk *et al.* and *C. terricoleslia* Wilk *et al.*) with  $< 70\%$  (Supplementary Material File S5). Using maximum parsimony (MP) weights, 88 species were placed correctly, with an average support of 97.8%; of these, 54 had 100% support and 74 had  $> 95\%$  (none below 70%). Under maximum likelihood (ML) weighting, all 89 species were placed correctly, with an average of 96.6% support; 79 had 100% support, 81 had  $\geq 95\%$ , and three  $< 70\%$  (Supplementary Material File S5). Thus, overall performance was best with even weights (96% of species placed correctly with 95% support or higher), followed by ML weighting (91% of species) and MP weighting (83% of species).

Using the above results as expectation values for the outcome with the 20 herbarium samples, the latter were placed 50 times (out of a possible 60) with a known species: 17 times (out of 20) using even weight, with 90.8% average support, 15 times (out of 20) using MP weight, with 92.1% average support, and 18 times (out of 20) using ML weight (Supplementary Material File S6, available online), with 88.7% average support (Supplementary Material File S5). Based on expectation value from calibration with the known species, seven samples each were classified as 'no match' under even and MP weighting (more than 1% point difference with the expected value for the best scoring result), whereas under ML, ten samples were classified as 'no match'; for even weighting, 'no match' results varied between 94.3% and 56.1% support, for MP between 95.2% and 63.1%, and for ML between 94.7% and 50.7% (Supplementary Material File S5). Five out of the 20 samples were placed with a single species; of these, two included two internal node placements (unresolved) and one (*Steglich* s. n. from Venezuela) included one internal node placement (under MP weights), otherwise being placed with *C. dalehana* B. Moncada *et al.* from Colombia.

Samples Sipman 37684 and Sipman 37777 (both from El Salvador and with identical characters) were the only ones placed

**Table 3.** Character weights used in the different setups to bin the *Cora* samples onto the reference tree. For 'even', all characters were weighted equally. The MP and ML weights were derived from the corresponding weight vector algorithm implemented in RAxML, depending on the underlying reference tree. MP = maximum parsimony, ML = maximum likelihood, Mol = molecular reference tree, Mor = morphological reference tree, Ran = randomized reference tree. Note the differences in character weights between MP and ML approaches and between underlying reference trees.

Character	Even	MP Mol	MP Mor	MP Ran	ML Mol	ML Mor	ML Ran	Arbitrary
Substratum	100	24	11	5	100	100	54	50
Lobe size	100	22	67	12	100	100	0	50
Sutures	100	19	26	10	92	100	32	50
Colour	100	0	0	0	95	100	83	50
Surface	100	16	56	12	67	100	91	50
Trichomes	100	76	74	62	100	100	80	75
Margin	100	30	63	12	100	100	51	50
Soredia	100	76	85	65	100	100	67	100
Cortex	100	54	63	45	100	100	97	100
Papillae	100	41	63	42	59	100	60	100
Hymenophore	100	70	67	60	100	100	86	100
Pigment	100	100	100	90	100	100	78	100
Central America	100	57	59	55	94	100	45	50
Caribbean	100	100	100	100	100	100	10	50
Galapagos	100	100	100	100	85	100	15	75
Northern Andes	100	41	74	22	31	100	5	50
Central Andes	100	62	78	70	45	100	31	50
Southern Andes	100	97	96	97	100	100	43	50
Brazil	100	98	89	90	43	100	48	50
Paleotropics	100	100	100	100	100	100	68	100

three times (i.e. with each different weighting method) with the same known species (*C. barbulate* Lücking *et al.* from Costa Rica); they differ, however, in a few characters and do not represent that species but an undescribed taxon close to it. Nine further samples were placed with two different known species depending on the weighting approach; in five of these, additional placement at an unresolved, internal node was observed (three of these under MP weights). The remaining six samples were each placed with three different known species under each of the weighting approaches (Supplementary Material File S6).

Comparison of the character scores of the 20 herbarium samples revealed that two were conspecific with known species, namely a sample from southern Colombia (*Cleef* 677) with *Cora cuzcoensis* Holgado *et al.* from Peru, and a sample from Venezuela (*Steglich* s. n.) with *C. dalehana* from central Colombia. The remaining 18 samples represented presumably undescribed species; the number of ecological and morphological characters in which each of them differed from the most similar, known species, varied between one (three samples), two (five samples), three (eight samples), and four (two samples), out of 12 (not counting the eight distribution characters).

## Discussion


We introduced and tested *PhyloKey*, a novel method for batch-identification of specimens based on an underlying phenotype character matrix. *PhyloKey* is based on the approach of

phenotype-based phylogenetic binning (Berger & Stamatakis 2010; Berger *et al.* 2011), a tool used in integrative taxonomy of fungi (lichens), plants and animals to quantitatively integrate phenotype data with molecular phylogenies (Koch *et al.* 2012; Parnmen *et al.* 2012; Rivas Plata *et al.* 2012; Lücking *et al.* 2015b; Dohrmann *et al.* 2017; Buitrago *et al.* 2018; Lücking & Kalb 2018; Perlmutter *et al.* 2020; Badano *et al.* 2021; Černý & Natale 2022). In contrast to interactive keys, such as DELTA (Dallwitz *et al.* 2006), Xper2 (Ung *et al.* 2010) or Dryades KeyToNature (Nimis *et al.* 2012; Nimis & Martellos 2020), *PhyloKey* allows evaluation of identification results by means of bootstrap support values, thus providing a measure of reliability for individual placements, given all possible alternatives. An additional advantage of *PhyloKey* is the possibility to simultaneously bin a large number of previously scored specimens.

*PhyloKey* is comparable to other interactive identification tools in providing multi-access entry. However, in contrast to interactive keys, characters are not entered subsequently, not forcing the user to decide on the sequence of characters (in some tools, such as DELTA, guided by the identification process). In *PhyloKey*, all scored characters are evaluated simultaneously. Bootstrapping then reconstructs placements based on character subsets, thus assessing internal consistency of the sampled characters. Our test showed that using a molecular reference tree, specimens were mostly binned correctly and with support, with as little as ten out of 20 characters. For the random reference tree, the minimum number of characters required to bin most

**Table 4.** Results of the 'simulated' *Cora* test samples with increasing number of missing characters. MP = maximum parsimony; ML = maximum likelihood.

Sample	Molecular reference tree						Morphological reference tree						Random reference tree					
	Matches			Support			Matches			Support			Matches			Support		
	Even	MP	ML	Even	MP	ML	Even	MP	ML	Even	MP	ML	Even	MP	ML	Even	MP	ML
Complete	10	10	10	99	100	100	10	10	10	100	100	100	10	10	10	100	100	98
1 missing	9	9	9	99	95	100	8	9	8	100	100	100	10	10	10	99	100	98
2 missing	10	10	10	99	100	98	9	9	9	98	97	98	10	10	10	100	100	100
3 missing	10	10	10	99	100	98	9	9	9	100	97	100	10	10	9	100	100	100
4 missing	9	10	8	95	100	100	8	9	8	100	100	100	10	9	9	100	100	100
5 missing	8	8	7	93	89	88	7	7	7	97	100	97	9	9	8	100	99	100
6 missing	10	8	9	99	89	100	5	5	5	100	100	100	10	8	8	100	100	100
7 missing	6	7	5	71	76	72	7	6	7	94	100	94	6	6	6	99	100	100
8 missing	7	9	7	82	96	100	6	7	6	100	100	100	8	8	7	100	100	97
9 missing	6	7	6	66	76	77	4	3	4	100	100	100	7	5	6	100	100	100
10 missing	9	7	9	89	77	98	6	5	6	100	100	100	8	7	6	100	96	100
11 missing	4	4	4	69	67	81	3	3	3	100	100	100	3	4	4	100	93	100
12 missing	6	2	5	80	49	80	4	2	4	95	100	95	6	3	5	98	100	93
13 missing	2	1	2	50	35	49	3	2	3	100	100	100	2	1	3	99	100	90
14 missing	4	4	4	56	71	68	5	4	5	98	100	98	4	5	2	100	95	100
15 missing	2	3	3	52	47	69	1	1	1	99	83	99	2	2	2	100	94	100
16 missing	0	1	0	21	28	27	0	0	0	0	0	0	0	0	0	0	0	0
17 missing	0	0	0	4	4	3	1	1	1	82	87	82	2	0	1	80	0	70
18 missing	0	1	0	29	24	3	1	0	1	100	0	100	1	0	0	93	0	0
19 missing	0	0	0	4	7	17	0	0	0	0	0	0	0	0	0	0	0	0



**Figure 2.** Performance of the combined score (number of matches and mean bootstrap support) relative to the nature of the reference tree for *Cora* samples with decreasing number of sampled characters (e.g. minus\_01 means one less character out of the 20 scored and so forth, while minus\_19 means only one character was used). In colour online.


specimens correctly and with support was 12–14, and for the morphological reference tree was 16. Thus, scoring at least 70% of the characters resulted in correct binning with support in most cases under the various reference trees and weighting schemes. Surprisingly, even weighting performed consistently equal to or better than MP weighting and both methods performed better than ML weighting, consistent with the results of the binning approach by Berger *et al.* (2011) on the genera *Allographa* and *Graphis*, where MP was also found to outperform ML weighting. This is probably due to the nature of the underlying phenotype characters, which can hardly be forced into an evolutionary model and so the ML weighting approach is less intuitive than MP weighting.

Compared to a traditional dichotomous (single-access) key, *PhyloKey* requires a set of characters to be scored for each species to be identified, not knowing *a priori* which characters will eventually be critical for the identification. A traditional dichotomous

key will instead guide the user to observe specific characters, limited to those considered diagnostic at each step of the identification process. For instance, if a genus contains a single species with a unique pigment, in a dichotomous key that species will be easily keyed out first using just that one character. The time to observe all characters to establish a matrix is therefore longer when using *PhyloKey*, typically about twice as long (i.e. the mean time one would arrive at a species in the middle of a dichotomous key). For example, in the dichotomous key to *Cora* provided by Lücking *et al.* (2013), arriving at each species keyed out there would require the observation of between three and eight morpho-anatomical characters, on average about five. To use *PhyloKey*, about ten characters would have to be scored per specimen, not counting chorological characters (distribution), requiring about ten minutes per specimen (one minute per character). However, this increased amount of time is compensated in *PhyloKey* by the simultaneous identification of many specimens in batch mode. While the computation only takes seconds, running each specimen through a dichotomous key might typically require around five minutes. Therefore, for ten specimens, the total identification time using a dichotomous key would amount to 50 min (scoring of 5 characters each on average, at 1 min per character) + 50 min (average working time going with each specimen through the key) = 100 min, whereas in *PhyloKey* it would take 100 min (complete matrix scoring) + 0 min (key), so about the same time. Thus, on average, the advantages and disadvantages of both approaches regarding time balance each other out, with the difference that a matrix-based approach adds value by generating a lasting data set. Generally, a dichotomous key would work faster if the group in question is well known and has easily perceived diagnostic characters, whereas *PhyloKey* provides an advantage when the group in question is not well known (i.e. undescribed species are to be expected) and diagnostic characters are subtle. For instance, in the above example of a uniquely pigmented species within a genus, a dichotomous key would identify that species correctly only if it is indeed the only species with that character. If at least one other, unrecognized species with the same character existed, a dichotomous key would not alert the user to that possibility and since no other characters are used at that position in the key, the user may not be aware of potential deviations. In *PhyloKey*, as well as in other matrix-based identification tools such as DELTA Intkey, the resulting identity scores will tell the user whether there is a perfect match or whether there are deviations in one or more characters. A further, unique advantage of *PhyloKey* is found in obtaining a placement support value for each sample.

Character scoring in *PhyloKey* is no different to a scoring scheme required for an interactive multi-access key, such as DELTA Intkey, and so, apart from compatibility of import/export formats, no additional work is needed when implementing *PhyloKey* on a set of data originally prepared to be used in an interactive key. The required data matrices are not only interchangeable but can be used for many other downstream analyses, such as multivariate techniques or ancestral character state analysis on a phylogenetic reference tree (e.g. Parnmen *et al.* 2012). Comprehensive character scoring also forces the user to perform comparative observations, making the scoring process more objective and reliable, whereas dichotomous keys rely on *ad hoc* observations. One advantage of interactive keys over *PhyloKey* is that the evaluation of individual characters is transparent during the identification process and can be accompanied by guiding illustrations or imagery, such as in Dryades KeyToNature





**Figure 3.** Labelled classification tree resulting from phylogenetic binning of the simulated *Cora* samples onto the random reference tree under an even weighting scheme. Blue species names = correctly binned samples with decreasing number of randomly sampled characters; orange filled circles = incorrectly binned samples with low number of randomly sampled characters with  $\geq 70\%$  bootstrap support. For detailed tree, see Supplementary Material File S7 (available online). In colour online.

(Nimis *et al.* 2012; Nimis & Martellos 2020). This possibility may be more attractive to users who are not familiar with the diagnostic characters of a group in question. However, the scoring process in *PhyloKey* can also be accompanied by character illustrations, especially when set up for particular groups, as shown in the Material and Methods section above (Fig. 1).

One surprising result of our analysis was the good performance of the randomized reference tree. Normally, the binning method, and hence *PhyloKey*, would rely on a molecular reference tree to guarantee the best possible placement of a specimen or taxon based on phenotype data, due to the character weighting process. In lieu of a molecular reference tree (e.g. when only a small number of taxa within the target group have molecular data), a tree based on the phenotype characters seemed a viable alternative. However, in our analysis, such a morphological reference tree performed less well, probably because its internal topology is based on exactly the same characters that are being

used for the binning process. A randomized tree avoids this shortcoming and seems to have no negative effects on correct placement of individual taxa or specimens compared to a molecular reference tree, probably because the underlying relationships between taxa (backbone) do not affect the closest binning match. This finding offers a more universal use of the *PhyloKey* approach, by simply establishing a randomized reference tree when sufficient sequence data for a molecular reference tree are not available. We hypothesize that a randomized reference tree will work well if most of the species in a group, or at least the range of phenotypic variation of the group, are known and represented by the reference terminals; whereas a phenotype-based reference tree has the potential for better predictive placement of specimens representing unknown taxa, as placements also reflect the character composition of deeper nodes.

Looking at the performance of *PhyloKey* relative to previously unidentified herbarium samples, presumably undescribed species

obtained different placements under each weighting approach (including internal nodes), in each case with at least one strongly supported ‘mismatch’, that is, suggesting a known species as the result with 1% difference or less from the expected value. The only exceptions were the two samples *Sipman* 37684 and *Sipman* 37777, consistently placed with a single species (*C. barbulate*) but differing in three ecological and morphological characters, including the diagnostic medullary papillae. On the other hand, the two samples that presumably represented known species behaved differently. Sample *Steglich* s. n., representing *C. dalehana*, was placed twice with that species, under even and ML weights, whereas MP weighting placed the sample on an internal node. The sample deviated from *C. dalehana* in a single score, for the character ‘sutures’, scored as ‘1’ (short) in *C. dalehana* and ‘0’ (absent) in the sample, which we consider an ‘allowable’ deviation, as the sample was not sufficiently well developed. The deviation did not affect its inferred placement under even and ML weights but apparently did under MP weighting. In contrast, sample *Cleef* 677 (corresponding to *C. cuzcoensis*) was placed correctly under an even weighting scheme, on an internal node under ML weight, and incorrectly under MP weight. While this sample agreed with *C. cuzcoensis* in all ecological and morphological characters, it differed in the distribution grid (northern versus central Andes), which was included in the *PhyloKey* matrix and apparently caused the partially inconsistent placement. However, in both samples, even weighting was not sensitive to these scoring particularities and placed both correctly, whereas ML weighting resulted in one correct and one unresolved placement and MP weighting in one unresolved and one incorrect placement. In these cases, the *PhyloKey* approach makes the establishment of potentially new species more reliable, by simultaneously identifying their closest matches and hence avoiding overlooking potentially available names elsewhere, and also by allowing identification of the quantity and quality of character mismatches. In that sense, *PhyloKey* could not only be used as an identification tool but also as a quantitative tool to recognize potential new species, simultaneously highlighting the most similar known taxa and the number of differences to these. *PhyloKey* could therefore be a useful tool in ‘non-molecular museomics’, the quantitative assessment of phenotype characters and their integration with molecular data, by scoring a large number of herbarium samples and evaluating their placement on a reference tree. While DNA barcoding of older herbarium samples is partially feasible and has been shown to work in *Cora* (Dal Forno *et al.* 2022), it depends on the condition of the underlying sample and is often unsuccessful, so *PhyloKey* could be a non-molecular complement to this approach.

It should be noted that the phenotype-based phylogenetic binning approach provides an objective method to predict the phylogenetic placement of individuals in the absence of DNA sequence data; however, this prediction may not be accurate and depends on the number of taxa already sequenced and the phylogenetic signal of the scored characters. Usually, accuracy is obtained at genus or within-genus clade level, but not necessarily identifying the closest relative at species level (e.g. Berger *et al.* 2011; Perlmutter *et al.* 2020). One herbarium specimen tested here, *Rapp* 581 from the USA (Florida), recently described as *Cora timucua* Dal Forno *et al.* (Lücking *et al.* 2020c), showed slight differences in its placement when based on phenotype data versus DNA sequence data (Lücking *et al.* 2020c). Fortunately, the exact phylogenetic placement of an individual is of secondary importance in the *PhyloKey* application: as we could show,

individuals are generally placed correctly if a matching taxon is already in the reference matrix, whereas the recognition of potentially novel taxa does not depend on their precise phylogenetic placement.

### How to implement PhyloKey

The *PhyloKey* approach requires the following tools:

- a spreadsheet tool (Excel, Numbers, or similar)
- a text editor (Word, Editor, Wordpad, Notepad, Pages, BBEdit, or similar)
- RAxML (tested versions: 7.2.6, 8.2.0)
- tree viewing software (tested: FigTree v.1.4.4, see below).

The following data sets need to be established:

- a phylogenetic reference tree of a set of known taxa in Newick format (e.g. Supplementary Material File S4, available online); it can be based on actual molecular data (e.g. Supplementary File S1) or on phenotype data (e.g. Supplementary Material File S2) or can be assembled manually as a simple tree format assuming underlying ‘relationships’; the tree can contain branch lengths but these are not required
- a reference matrix of (diagnostically important) phenotype characters for the same taxa (terminals) used for the reference tree, with exactly the same names; the matrix can be established in a spreadsheet but must be converted into Phylip format prior to analysis (e.g. Supplementary Material File S2); note that unknown or missing data can be expressed using a ‘?’ sign
- a matrix of (diagnostically important) phenotype characters for a set of query specimens, in exactly the same format as the reference matrix; the matrix can be established in a spreadsheet but must be converted into Phylip format prior to analysis (e.g. Supplementary Material File S3); note that unknown or missing data can be expressed using a ‘?’ sign

The approach is implemented via the following steps:

*Step 1.* Computation of the weight vector(s) for the phenotype characters based on their distribution over the reference tree, either based on maximum parsimony (MP) or maximum likelihood (ML) or both; this is invoked in RAxML (e.g. v. 7.2.6) using the following command line (alternatively as Windows batch file ‘.bat’):

- raxmlHPC.exe -f U -m MULTIGAMMA -s matrix.phy -t reference.tre -n weight\_vector\_MP.txt (for the MP weight vector)
- raxmlHPC.exe -f u -m MULTIGAMMA -s matrix.phy -t reference.tre -n weight\_vector\_ML.txt (for the ML weight vector) where -f u (-f U) = algorithm, -m MULTIGAMMA = underlying evolutionary model, -s matrix.phy = matrix selection (matrix.phy = reference matrix in Phylip format), -t reference.tre = tree selection (reference.tre = reference tree in Newick format), and weight\_vector\_MP.txt/weight\_vector\_ML.txt = output of the weight vector (text format, a series of numbers between 0 and 100).

*Step 2.* Running the binning analysis; this is invoked in RAxML (e.g. v. 7.2.6) using the following command line (alternatively as Windows batch file ‘.bat’):

- raxmlHPC.exe -f v -m MULTIGAMMA -a weight\_vector\_ML.txt -s samples.phy -t reference.tre -n identification.txt -x 12345 -# 1000

where -f v = algorithm, -m MULTIGAMMA = underlying evolutionary model, -a weight\_vector\_ML.txt = selection of the weight vector, -s samples.phy = query matrix selection (samples.phy = query matrix of all samples to be analyzed, in Phylip format), -t reference.tre = tree selection (reference.tre = reference tree in Newick format), -n identification.txt = name designation for the various output files, -x 12345 = random number seed, and -# 1000 = number of bootstrap pseudoreplicates.


**Step 3.** Visualizing the classification table; open the output file named 'RAxML\_classification[...].txt' in a text editor and globally replace spaces with tabulators, then save and open with a spreadsheet editor, such as Microsoft Excel (alternatively, open original file in spreadsheet editor using space as separator). The table will display four columns: 1) sample name, 2) node ID of nearest placement, 3) bootstrap support value for placement (number of bootstrap replicates), and 4) branch length of the original reference tree for that node; edit classification table as desired (e.g. Supplementary Material File S5).

**Step 4.** Visualizing the classification tree (e.g. in FigTree); open the output file named 'RAxML\_labelledTree[...].txt' in a text editor; globally replacing the string ':1.0]' with '[' and then ']' with ':1.0]' (which switched the order of branch length and node ID labels); check instances of identical terminal output names and make them unique by adding the suffix letters 'a', 'b', etc.; open adjusted tree file in tree viewer (e.g. FigTree) and edit as desired, then export as PDF (e.g. Supplementary Material File S6).

**Step 5.** If desired, the taxonomic identities of the node labels in the output tree can be added to the classification table (e.g. Supplementary Material File S5 worksheet 'Herbarium').

**Step 6.** If desired, the original phenotype characters can be added to each query and ID label in the classification table, to assess the corresponding matching level (e.g. Supplementary Material File S5 worksheet 'Herbarium').

**Acknowledgements.** Harrie Sipman is thanked for his assistance in selecting the herbarium specimens tested here. MDF would like to acknowledge funding from NSF (PRFB 1609022) and the Smithsonian Institution (NMNH Peter Buck Postdoctoral Fellowship).

**Author ORCID.**  Robert Lücking, 0000-0002-3431-4636; Bibiana Moncada, 0000-0001-9984-2918; Manuela Dal Forno, 0000-0003-1838-1676.

**Supplementary Material.** The supplementary material for this article can be found at <https://doi.org/10.1017/S0024282923000415>.

## References

- Ariyawansa HA, Hyde KD, Jayasiri SC, Buyck B, Chethana KWT, Dai DQ, Dai YC, Daranagama DA, Jayawardena RS, Lücking R, *et al.* (2015) Fungal diversity notes 111–252 – taxonomic and phylogenetic contributions to fungal taxa. *Fungal Diversity* 75, 27–274.
- Badano D, Fratini M, Maugeri L, Palermo F, Pieroni N, Cedola A, Haug JT, Weiterschan T, Velten J, Mei M, *et al.* (2021) X-ray microtomography and

- phylogenomics provide insights into the morphology and evolution of an enigmatic Mesozoic insect larva. *Systematic Entomology* 46, 672–684.
- Berger SA and Stamatakis A (2010) Accuracy of morphology-based phylogenetic fossil placement under maximum likelihood. In *Proceedings of the ACS/IEEE International Conference on Computer Systems and Applications –AICCSA 2010, 1–19 May 2010*, New York, pp. 1–9.
- Berger SA, Stamatakis A and Lücking R (2011) Morphology-based phylogenetic binning of the lichen genera *Graphis* and *Allographa* (Ascomycota: Graphidaceae) using molecular site weight calibration. *Taxon* 60, 1450–1457.
- Buitrago A, MacDougal JM and Coca LF (2018) *Passiflora kumandayi* (Passifloraceae), a new species from the Colombian Andes in a new section within subgenus *Decaloba*. *Phytotaxa* 344, 13–23.
- Carbone I, White JB, Miadlikowska J, Arnold AE, Miller MA, Kauff F, U'Ren JM, May G and Lutzoni F (2017) T-BAS: Tree-Based Alignment Selector toolkit for phylogenetic-based placement, alignment downloads and metadata visualization: an example with the *Pezizomycotina* tree of life. *Bioinformatics* 33, 1160–1168.
- Černý D and Natale R (2022) Comprehensive taxon sampling and vetted fossils help clarify the time tree of shorebirds (*Aves*, *Charadriiformes*). *Molecular Phylogenetics and Evolution* 177, 107620.
- Dal Forno M, Lawrey JD, Moncada B, Bungartz F, Grube M, Schuettelpelz E and Lücking R (2022) DNA barcoding of fresh and historical collections of lichen-forming basidiomycetes in the genera *Cora* and *Corella* (Agaricales: Hygrophoraceae): a success story? *Diversity* 14, 284.
- Dallwitz MJ, Paine TA and Zurcher EJ (2006) *User's Guide to the DELTA System: a General System For Processing Taxonomic Descriptions*. Canberra: CSIRO Division of Entomology.
- De Queiroz K (2005) Ernst Mayr and the modern concept of species. *Proceedings of the National Academy of Sciences of the United States of America* 102 (Suppl. 1), 6600–6607.
- DeSalle R and Goldstein P (2019) Review and interpretation of trends in DNA barcoding. *Frontiers in Ecology and Evolution* 7, 302.
- Dohrmann M, Kelley C, Kelly M, Pisera A, Hooper JN and Reisch HM (2017) An integrative systematic framework helps to reconstruct skeletal evolution of glass sponges (*Porifera*, *Hexactinellida*). *Frontiers in Zoology* 14, 1–31.
- Edwards M and Morse DR (1995) The potential for computer-aided identification in biodiversity research. *Trends in Ecology and Evolution* 10, 153–158.
- Goëau H, Bonnet P, Joly A, Bakić V, Barbe J, Yahiaoui I, Selmi S, Carré J, Barthélémy D, Boujemaa N, *et al.* (2013) Pl@ntNet mobile app. In *Proceedings of the 21st ACM International Conference on Multimedia*, 21–25 October 2013, Barcelona, Spain, pp. 423–424.
- Hagedorn G, Press B, Hetzer S, Plank A, Weber G, von Mering S, Martellos S and Nimis PL (2010) A MediaWiki implementation of single-access keys. In Nimis PL and Vignes Lebbe R (eds), *Tools for Identifying Biodiversity: Progress and Problems*. Trieste: EUT-Edizioni Università di Trieste.
- Joly A, Bonnet P, Goëau H, Barbe J, Selmi S, Champ J, Dufour-Kowalski S, Affouard A, Carré J, Molino JF, *et al.* (2016) A look inside the Pl@ntNet experience. *Multimedia Systems* 22, 751–766.
- Koch MA, Kiefer M, German DA, Al-Shehbaz IA, Franzke A, Mummenhoff K and Schmickl R (2012) BrassiBase: tools and biological resources to study characters and traits in the *Brassicaceae* – version 1.1. *Taxon* 61, 1001–1009.
- La Salle J, Wheeler Q, Jackway P, Winterton S, Hobern D and Lovell D (2009) Accelerating taxonomic discovery through automated character extraction. *Zootaxa* 2217, 43–55.
- Lücking R (2020) Three challenges to contemporaneous taxonomy from a lichen-mycological perspective. *Megataxa* 1, 78–103.
- Lücking R and Kalb K (2018) Formal instatement of *Allographa* (Graphidaceae): how to deal with a hyperdiverse genus complex with cryptic differentiation and paucity of molecular data. *Herzogia* 31, 535–561.
- Lücking R, Dal-Forno M, Lawrey JD, Bungartz F, Rojas MEH, Marcelli MP, Moncada B, Morales EA, Nelsen MP, Salcedo L, *et al.* (2013) Ten new species of lichenized *Basidiomycota* in the genera *Dictyonema* and *Cora* (Agaricales: Hygrophoraceae), with a key to all accepted genera and species in the *Dictyonema* clade. *Phytotaxa* 139, 1–38.

- Lücking R, Dal-Forno M, Sikaroodi M, Gillevet PM, Bungartz F, Moncada B, Yáñez-Ayabaca A, Chaves JL, Coca LF and Lawrey JD (2014) A single macrolichen constitutes hundreds of unrecognized species. *Proceedings of the National Academy of Sciences of the United States of America* **111**, 11091–11096.
- Lücking R, Cáceres MES, Silva NG and Alves RJV (2015a) The genus *Cora* in the South Atlantic and the Mascarenes: two novel taxa and inferred biogeographic relationships. *Bryologist* **118**, 293–303.
- Lücking R, Mangold A, Rivas Plata E, Parnmen S, Kraichak E and Lumbsch HT (2015b) Morphology-based phylogenetic binning to assess a taxonomic challenge: a case study in *Graphidaceae* (*Ascomycota*) requires a new generic name for the widespread *Leptotrema wightii*. *Botanical Journal of the Linnean Society* **179**, 436–443.
- Lücking R, Dal Forno M, Moncada B, Coca LF, Vargas-Mendoza IY, Aptroot A, Arias LJ, Besal B, Bungartz F, Cabrera-Amaya DM, et al. (2017) Turbo-taxonomy to assemble a megadiverse lichen genus: seventy new species of *Cora* (*Basidiomycota: Agaricales: Hygrophoraceae*), honouring David Leslie Hawksworth's seventieth birthday. *Fungal Diversity* **84**, 139–207.
- Lücking R, Aime MC, Robertse B, Miller AN, Ariyawansa HA, Aoki T, Cardinali G, Crous PW, Druzhinina IS, Geiser DM, et al. (2020a) Unambiguous identification of fungi: where do we stand and how accurate and precise is fungal DNA barcoding? *IMA Fungus* **11**, 1–32.
- Lücking R, Truong BV, Huong DTT, Le NH, Nguyen QD, Nguyen VD, Von Raab-Straube E, Bollendorff S, Govers K and Di Vincenzo V (2020b) Caveats of fungal barcoding: a case study in *Trametes* s. lat. (*Basidiomycota: Polyporales*) in Vietnam reveals multiple issues with mislabelled reference sequences and calls for third-party annotations. *Willdenowia* **50**, 383–403.
- Lücking R, Kaminsky L, Perlmutter GB, Lawrey JD and Dal Forno M (2020c) *Cora timucua* (*Hygrophoraceae*), a new and potentially extinct, previously misidentified basidiolichen of Florida inland scrub documented from historical collections. *Bryologist* **123**, 657–673.
- Mayo SJ, Allkin R, Baker W, Blagoderov V, Brake I, Clark B, Govaerts R, Godfray C, Haigh A, Hand R, et al. (2008) Alpha e-taxonomy: responses from the systematics community to the biodiversity crisis. *Kew Bulletin* **63**, 1–16.
- Moncada B, Pérez-Pérez RE and Lücking R (2019) The lichenized genus *Cora* (*Basidiomycota: Hygrophoraceae*) in Mexico: high species richness, multiple colonization events, and high endemism. *Plant and Fungal Systematics* **64**, 393–411.
- Murguía-Romero M, Serrano-Estrada B, Ortiz E and Villaseñor JL (2021) Taxonomic identification keys on the web: tools for better knowledge of biodiversity. *Revista Mexicana de Biodiversidad* **92**, e923592.
- Nimis PL and Martellos S (2020) Towards a digital key to the lichens of Italy. *Symbiosis* **82**, 149–155.
- Nimis PL, Riccamboni R and Martellos S (2012) Identification keys on mobile devices: the *Dryades* experience. *Plant Biosystems* **146**, 783–788.
- Parnmen S, Lücking R and Lumbsch HT (2012) Phylogenetic classification at generic level in the absence of distinct phylogenetic patterns of phenotypical variation: a case study in *Graphidaceae* (*Ascomycota*). *PLoS ONE* **7**, e51392.
- Perlmutter GB, Rivas Plata E, LaGreca S, Aptroot A, Lücking R, Tehler A and Ertz D (2020) *Biatora akompsa* is revealed as a disjunct North American species of *Pentagenella* (*Opegraphaceae*) through molecular phylogenetic analysis and phenotype-based binning. *Bryologist* **123**, 502–516.
- Rambaut A (2018) *FigTree, version 1.4.4*. Institute of Evolutionary Biology, University of Edinburgh. [WWW resource] URL <http://tree.bio.ed.ac.uk/software/figtree> [Accessed 22 May 2023].
- Reydon TA (2019) Are species good units for biodiversity studies and conservation efforts? In Casetta E, Marques da Silva J and Vecchi D (eds), *From Assessing to Conserving Biodiversity*. Cham, Switzerland: Springer, pp. 167–193.
- Riddle BR and Hafner DJ (1999) Species as units of analysis in ecology and biogeography: time to take the blinders off. *Global Ecology and Biogeography* **8**, 433–441.
- Rivas Plata E, Lücking R and Lumbsch HT (2012) Molecular phylogeny and systematics of the *Ocellularia* clade (*Ascomycota: Ostropales: Graphidaceae*). *Taxon* **61**, 1161–1179.
- Schoch CL, Seifert KA, Huhndorf S, Robert V, Spouge JL, Levesque CA, Chen W, Bolchacova E, Voigt K, Crous PW, et al. (2012) Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. *Proceedings of the National Academy of Sciences of the United States of America* **109**, 6241–6246.
- Seberg O, Humphries CJ, Knapp S, Stevenson DW, Petersen G, Scharff N and Andersen NM (2003) Shortcuts in systematics? A commentary on DNA-based taxonomy. *Trends in Ecology and Evolution* **18**, 63–65.
- Sluys R (2013) The unappreciated, fundamentally analytical nature of taxonomy and the implications for the inventory of biodiversity. *Biodiversity and Conservation* **22**, 1095–1105.
- Smith VS (2005) DNA barcoding: perspectives from a 'Partnerships for Enhancing Expertise in Taxonomy' (PEET) debate. *Systematic Biology* **54**, 841–844.
- Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**, 1312–1313.
- Stamatakis A, Ludwig T and Meier H (2005) RAxML-III: a fast program for maximum likelihood-based inference of large phylogenetic trees. *Bioinformatics* **21**, 456–463.
- Swofford DL (2003) *PAUP\*: Phylogenetic Analysis Using Parsimony (and Other Methods), Version 4.0 b10*. Sunderland, Massachusetts: Sinauer Associates.
- Ung V, Dubus G, Zaragüeta-Bagils R and Vignes-Lebbe R (2010) Xper2: introducing e-taxonomy. *Bioinformatics* **26**, 703–704.
- Vargas LY, Moncada B and Lücking R (2014) Five new species of *Cora* and *Dictyonema* (*Basidiomycota: Hygrophoraceae*) from Colombia: chipping away at cataloging hundreds of unrecognized taxa. *Bryologist* **117**, 368–378.
- Walter DE and Winterton S (2007) Keys and the crisis in taxonomy: extinction or reinvention? *Annual Reviews of Entomology* **52**, 193–208.
- Wilkins J (2017) The reality of species: real phenomena not theoretical objects. In Joyce R (ed.), *The Routledge Handbook of Evolution and Philosophy*. London: Routledge, pp. 167–181.
- Zhang J, Kapli P, Pavlidis P and Stamatakis A (2013) A general species delimitation method with applications to phylogenetic placements. *Bioinformatics* **29**, 2869–2876.