

controlling for sources of prenatal stress. **RESULTS/ANTICIPATED RESULTS:** Global racism (total IRRS score) was significantly associated with birth weight when controlling for prenatal perceived stress and stressful life events within the last six months since study enrollment ($\beta = -16.7$, $p = .035$). Neonatal gestational age was associated with both global racism ($\beta = -0.03$, $p = .028$) and individual racism (IRRS'Individual' subscale score) ($\beta = -0.09$, $p = .032$) when controlling for prenatal perceived stress and stressful life events within the last six months since study enrollment. These results suggest that greater race-related stress contributes to lower weight and gestational age at birth in pregnant Black persons. **DISCUSSION/SIGNIFICANCE:** Future studies are necessary to determine the mechanisms by which race-related stress contributes to these adverse birth outcomes and to inform the development risk-assessment tools and interventions to mitigate the threat of race-related stress on adverse birth outcomes in high-risk populations.

196

COVID-19 Hospitalization and Death by Concentrated Racial/Ethnic and Economic Segregation: Los Angeles County, January 2020-June 2023*

Rebecca Fisher, Nina Harawa and Elizabeth Rose Mayeda
University of California Los Angeles

OBJECTIVES/GOALS: This study aims to assess associations over time between several area-based indices of concentration at the extremes and COVID-19 hospitalization and death in Los Angeles County, from January 2020 to June 2023. These measures reflect concentrations of racial/ethnic and economic segregation at the census tract level. **METHODS/STUDY POPULATION:** Indices of concentration at the extremes (ICEs) for economic segregation, racial/ethnic segregation, and a combination of the two were constructed for each Los Angeles County (LAC) census tract, using 2015-2019 ACS data. The index ranges from -1 to 1 and is the number of advantaged households (HH) minus the number of disadvantaged HH over the total number of HH measured. Economic segregation is HH income over \$100,000 vs. below \$25,000 per year and racial/ethnic segregation defined as White and Non-Hispanic vs. non-White or Hispanic HH. The distribution of index scores was divided into quintiles (Q1-Q5) for all LAC census tracts. Age-adjusted hospitalization and death rates were derived at the census tract level by quarter (QTR) based on Los Angeles County Department of Public Health surveillance data. **RESULTS/ANTICIPATED RESULTS:** Age-adjusted hospitalization and death rates were consistently higher across all quarters in Q1 (most deprived) vs. Q5 (most privileged) for all ICE measures. For ICE of economic segregation, the age-adjusted hospitalization and death rate ratios between Q1 and Q5 were 2.12 (range: 1.32 - 4.15; peak 2020 QTR2) and 2.02 (range: 1.46 - 3.21; peak 2021 QTR1), respectively. For ICE of racial segregation, the age-adjusted hospitalization and death rate ratio between Q1 and Q5 was 2.03 (range: 1.08 - 3.95; peak 2020 QTR3) and 1.77 (range: 1.03 - 2.80; peak 2021 QTR1). The ICE of economic/racial segregation combined was the highest, with averages of the age-adjusted hospitalization and death rate ratios between Q1 and Q5 being 2.26 (1.16 - 4.43; peak 2020 QTR2) and 1.99 (range: 1.22 - 3.32; peak 2021 QTR1). **DISCUSSION/SIGNIFICANCE:** This study assesses the impact of geographic segregation based on indices that quantify the concentration of both deprivation, privilege, and racial/ethnic group, demonstrating that segregation and economic deprivation are consistently associated with higher rates of age-adjusted hospitalization and death from COVID-19 in LAC.

197

Evaluating Disparities in Management of Solid Organ Injury in Children Treated at Pediatric vs. Adult Trauma Centers*

Carrinton Mauney, Charbel Chidiac and Mark Slidell
Johns Hopkins University School of Medicine

OBJECTIVES/GOALS: Non-operative management is preferred for pediatric solid organ injury (SOI). Despite this children treated at adult trauma centers (ATC) are more likely to be treated surgically and have worse outcomes than those treated at pediatric trauma centers (PTC). We hypothesize there are disparities by sex and race in management of pediatric SOI at ATC vs PTC. **METHODS/STUDY POPULATION:** Retrospective review of the National Trauma Data Bank (NTDB) from 2010-2018 was conducted. Inclusion criteria were age ≤ 18 and injury to spleen, liver or kidney. Outcomes at American College of Surgeons accredited ATC, PTC, and combined ATC/PTC trauma centers were evaluated. The primary outcome was operative management. Secondary outcomes include length of stay, and in-hospital complications. Multivariate logistical regression adjusting for race, sex, and insurance type will be performed. **RESULTS/ANTICIPATED RESULTS:** 40,111 children were treated for SOI from 2010-2018. 39.3% were treated at an ATC and 26.4% at a PTC. Of children treated at an ATC, 62% were White, 17% were Black, and 1% were Asian. Children treated at the PTC were 60% White, 20% Black, and 0.9% Asian. Primary insurance type was Medicaid for 33% of patients at an ATC and 39% at PTC. Median length of stay at ATC and PTC was 4 days (2-7) and 3 days (2-6) respectively. 3.85% of patients at ATC underwent splenectomy compared to 0.8% at PTC. It is anticipated that further analysis will demonstrate that ICU admission, transfusion, embolectomy, and other operative interventions will be more prevalent at ATC than PTC. Moreover, we anticipate that multivariate logistical regression will show the odds of receiving operative management at each center differ by race, sex and insurance type. **DISCUSSION/SIGNIFICANCE:** Initial analysis of the NTDB from 2010-2018 shows that children treated for SOI at ATC receive operative interventions more often than those treated at PTC. Elucidating disparities in SOI care is an important step towards minimizing the impact of these disparities and better allocating resources such that they may be eliminated.

198

Community & Recruitment Programs at OHSU: Leveraging a team science approach to ensure representative study populations through community engagement and recruitment

Laura Campbell, Meredith Zauflik and Melinda Davis
Oregon Health & Science University

OBJECTIVES/GOALS: Investigators are looking to integrate DE&I into trials, increasing study population diversity. In response, the Community and Recruitment programs of the Oregon Clinical and Translational Research Institute (OCTRI), built a collaborative, community-focused system for investigators interested in community engagement and recruitment. **METHODS/STUDY POPULATION:** Historically, the OCTRI Community and Recruitment programs operated independently. To build a community-focused support system for investigators, we began with the programs learning about each other's goals, values, and operations over a six-month period. Over the next two years, we then