

Review: nutritional and endocrine control of colostrogenesis in swine

H. Quesnel^{1†} and C. Farmer²

¹PEGASE, INRA, Agrocampus Ouest, 35590 Saint-Gilles, France; ²Agriculture and Agri-Food Canada, Sherbrooke R & D Centre, 2000 College St., Sherbrooke, QC, Canada J1M 0C8

(Received 9 July 2018; Accepted 30 November 2018)

Colostrum plays an essential role in ensuring the survival, growth and health of piglets by providing energy, nutrients, immunoglobulins, growth factors and many other bioactive components and cells. Both colostrum yield and composition are highly variable among sows, yet mechanisms and factors that regulate colostrogenesis are not fully known. Unlike sow milk yield, sow colostrum yield is not highly determined by litter size and suckling intensity but is largely driven by sow-related factors. Colostrum synthesis is under hormonal control, with prolactin and progesterone concentrations prepartum having, respectively, positive and negative influences on colostrum yield. Less is known about the endocrine control of the end of colostrogenesis in swine, which is characterized by the closure of tight junctions in the mammary epithelium and the cessation of transfer of immunoglobulin G (IgG) into lacteal secretions. Recent studies indicate that exogenous hormones may influence colostrogenesis. Inducing parturition by injecting prostaglandin F_{2α} on day 114 of gestation in combination with an oxytocin-like molecule reduced colostrum yield, and injection of prostaglandin F_{2α} alone either reduced colostrum yield or had no effect. Injecting a supraphysiological dose of oxytocin to sows in the early postpartum period delayed the tightening of mammary tight junctions, thereby prolonging the colostrum phase and increasing concentrations of IGF-I and IgG and IgA in early milk. The development of strategies to improve colostrum composition in swine through maternal feeding has been largely explored but very few attempts were made to increase colostrum yield. This is most likely because of the difficulty in measuring colostrum yield in swine. The fatty acid content of colostrum greatly depends on the amount of lipids provided in the sow diet during late gestation, whereas the fatty acid profile is largely influenced by the type of lipid being fed to the pregnant sow. Moreover, various ingredients that presumably have immuno-modulating effects (such as fish oil, prebiotics and probiotics) increased concentrations of IgG, IgA and/or IgM in sow colostrum when they were provided during the last weeks of gestation. Finally, there is some evidence that sow nutrition during late gestation may influence colostrum yield but this clearly warrants more research. This review emphasizes that although progress has been made in understanding the control of colostrogenesis in swine, and that strategies exist to manipulate fat and immunoglobulin contents of colostrum, ways to increase colostrum yield are still lacking.

Keywords: colostrum, hormone, immunoglobulin, nutrition, sow

Implications

An early and high intake of colostrum is a major determinant of piglet survival in the first days after birth and is essential for ensuring the health and growth of piglets at least until weaning. However, both colostrum yield and composition are highly variable among sows. Better knowledge of endocrine and nutritional factors that participate in this variability is needed to define new strategies to manage reproductive sows to increase yield and quality of colostrum.

Introduction

Colostrum intake is essential for piglet survival within the few days after birth and for piglet health and growth performance until weaning (Quesnel *et al.*, 2012). Long-term effects were also reported, with colostrum intake being positively associated with piglet weight at weaning, at onset and end of the fattening period and being negatively correlated with piglet mortality during the nursery period (Declerck *et al.*, 2016). Colostrum provides newborn piglets with the energy necessary for thermoregulation and body growth, with passive immunity needed for protection against pathogens, and with growth factors that stimulate growth and maturation of tissues and organs, especially the

[†] E-mail: helene.quesnel@inra.fr

gastrointestinal tract (Quesnel *et al.*, 2012). By transferring bioactive factors from the mother to the offspring, colostrum is crucial for the process of lactocrine signaling, which has an impact on the offspring much beyond the neonatal period. Bioactive components of colostrum include, for example, proteins, growth factors, peptides, oligosaccharides, fatty acid-derived molecules, steroids and microRNAs. The most striking example of a lactocrine-active factor is relaxin, which is a peptide hormone found in sow colostrum, that plays an important role in developmental programming of the female reproductive tract until puberty (Bartol *et al.*, 2017).

In swine, colostrum can be defined as the first secretion of the mammary glands excreted for approximately 24 h after the onset of parturition. In both swine and ruminants, the major difference between colostrum and milk resides in the high concentrations of colostral immunoglobulins (Ig), specifically immunoglobulins G (IgG) in swine (and IgG1 in ruminants). Immunoglobulins A, although more abundant in colostrum than in milk, are not specific to colostrum and are therefore also present in milk throughout lactation. The composition of sow colostrum, especially in terms of IgG content, is highly variable among sows (Quesnel, 2011). Sow colostrum yield is estimated by summing the colostrum intakes of each piglet in the litter; it thus depends on the prediction model used to estimate individual colostrum intake. Based on Devillers' model (Devillers *et al.*, 2004), colostrum yield averages 3.5 kg and ranges from <1 kg to >6.0 kg (Quesnel *et al.*, 2012). Based on Theil's model (Theil *et al.*, 2014a), it averages 6.0 kg with a range from 2.8 up to 8.5 kg (Vadmand *et al.*, 2015). Nevertheless, regardless of the method used to estimate colostrum yield, it is greatly variable. The amount of colostrum needed by an average piglet (weighing 1.4 kg at birth) was estimated at 250 g, or 180 g/kg birth weight (calculated with Devillers' model; Quesnel *et al.*, 2012). This amount would significantly reduce the risk of mortality, provide passive immunity and allow a slight weight gain. Based on this recommendation, Quesnel *et al.* (2012) estimated that at least one-third of sows do not produce enough colostrum to fulfill the needs of their litter.

The amount and composition of colostrum produced can be influenced by sow characteristics, such as endocrine status, nutritional status, immune status and level of stress, and by litter characteristics, especially vitality at birth. Environmental factors also could alter colostrum yield and composition through effects on the sows (e.g., heat stress is likely to reduce feed intake and cortisol concentration in sows). The complex interactions between these factors make it difficult to decipher their respective influence in the regulation of colostrogenesis. The present review will focus on the effects of sow endocrine status and nutrition because hormones play a major role in regulating colostrum synthesis, and nutrition could be a tool to manipulate colostrum yield and composition in herds.

When does colostrogenesis take place?

In the present review paper, colostrogenesis is defined as englobing both the synthesis of milk-specific constituents and the transfer of IgG into lacteal secretions. However, in order to efficiently investigate strategies to alter colostrogenesis in sows, it is imperative to know precisely when these two events occur. The synthesis of milk components takes place during late gestation, with the first lipid droplets being visible at 105 days of gestation (the gestational term of the sow is 115 days) (Kensinger *et al.*, 1986) and the synthesis of lactose starting not before 3 to 4 days before parturition (Hartmann *et al.*, 1984). The transfer of IgG from sow plasma to lacteal secretions slowly begins approximately 10 days before farrowing, increases 3 or 4 days before farrowing and is maximal on the day of farrowing (Huang *et al.*, 1992). This suggests that colostrum synthesis, when defined by IgG transfer, is not entirely finished before parturition in sows. This is somewhat different from what occurs in dairy cows, with IgG1 transfer beginning several weeks before parturition, peaking a week *prepartum* and ceasing abruptly before calving (Barrington *et al.*, 2001). Nevertheless, recent findings suggest that the transfer of IgG in dairy cows can persist at least until parturition (Gross *et al.*, 2014).

During colostrogenesis, tight junctions between mammary epithelial cells (MECs) are 'leaky,' thereby allowing the paracellular transfer of constituents from maternal plasma to the alveolar compartment, notably of hormones, growth factors and Ig. The uptake of IgG by mammary glands in pigs could also be mediated by a Fc-specific receptor, the neonatal Fc receptor (FcRn), as it is the case in ruminants (Barrington *et al.*, 2001). Expression of the FcRn protein was demonstrated in porcine mammary tissue in the *peripartum* period, from 3 days before up to 1 day after parturition (Schnulle and Hurley, 2003; Palombo *et al.*, 2018). However, the role of the FcRn in IgG transfer has not been demonstrated in swine. The uptake of IgA in a polymeric form (pIgA) could also be achieved via transcytosis mediated by the polymeric immunoglobulin receptor (pIgR), whose transcript was shown in sow's mammary tissue (Schnulle and Hurley, personal communication). The relative importance of the passive *v.* the active transport of IgG and IgA transfer into sow colostrum is not known.

The composition of colostrum gradually changes between the onset of parturition (T0) and 24 h later (T24; Foisnet *et al.*, 2010a). Lactose and fat contents increase while protein content decreases because of the drop in IgG and IgA concentrations (Theil *et al.*, 2014b). In parallel, concentrations of K increase while Na concentrations decrease, as well as the Na : K ratio (Quesnel, 2011). The decrease in the Na : K ratio is indicative of the development of tight junctions between cells in the mammary epithelium. Concentrations of IgG in colostrum at T24 are positively correlated with the Na : K ratio in colostrum at T24, whereas the drop in IgG concentrations in colostrum between T0 and T24 is negatively correlated to that same variable (Quesnel, 2011). These relations are consistent with the fact that the cessation of IgG

transfer into lacteal secretions and the closure of tight junctions are part of the process, marking the end of colostrumogenesis and the initiation of lactation.

Sow v. piglets, who drives?

In swine, it is well known that milk production is stimulated by the action of suckling from piglets, and thus depends on litter size and weight of piglets. Moreover, the estimation of colostrum yield actually corresponds to the consumption of colostrum by the litter. Therefore, it has long been argued that colostrum yield mainly depends on litter characteristics. However, there is now a large body of evidence that even though colostrum yield may be influenced by the global vitality of the litter at birth, it also largely depends on the capacity of the sow to produce colostrum (Farmer and Quesnel, 2009).

Endocrine control of colostrumogenesis

Role of endogenous hormones

In swine, the *prepartum* peak of prolactin is essential for colostrumogenesis (Farmer *et al.*, 1998) and is brought about by the drop in progesterone concentrations (Taverne *et al.*, 1982). In primiparous sows, delays in the decrease in progesterone and in the increase in prolactin relative to the onset of parturition were greatly associated with a reduced yield of colostrum (Foisnet *et al.*, 2010a). Furthermore, the relative concentrations of prolactin and progesterone during the *prepartum* period were recently shown to influence colostrum yield. Primiparous sows that had lower progesterone concentrations associated with a tendency for greater prolactin concentrations (high prolactin to progesterone ratio) during the 2 days before and the day of parturition produced more colostrum (+0.6 kg) than sows that had a low prolactin to progesterone ratio (Loisel *et al.*, 2015). There were no concomitant changes in the concentrations of macronutrients in colostrum, indicating an absence of dilution effect. The greater colostrum concentrations of progesterone at the onset of parturition were also negatively related with colostrum yield in sows from parities one to eight, with sows having concentrations of progesterone >4.5 ng/ml producing 1.6 kg less colostrum than those having progesterone concentrations of <4.5 ng/ml (Hasan *et al.*, 2018). It is however important to mention that even though lower *peripartum* concentrations of prolactin are observed in primiparous than in multiparous sows (Quesnel *et al.*, 2013), this does not necessarily translate into a lower production of colostrum. Indeed, significant discrepancies exist between studies on the effect of sow parity on colostrum yield (Table 1). Greater colostrum yield was reported in young sows compared with older sows (Decaluwé *et al.*, 2013; Quesnel, unpublished data), while Devillers *et al.* (2007) reported that sows of parities two and three tended to produce more colostrum than primiparous and older sows. Finally, Boonraungrod *et al.* (2018) recently reported that

multiparous sows produced more colostrum than primiparous sows.

Besides progesterone and prolactin, other hormones also undergo drastic changes *peripartum*. Among them, estrogens and glucocorticoids whose concentrations increase during the week and the day before parturition, respectively (Foisnet *et al.*, 2010a), are considered to be involved in the regulation of lactogenesis in rodents and ruminants. To the best of our knowledge, however, a role for estrogens and glucocorticoids in the control of colostrumogenesis has not yet been described in swine. Oxytocin, which is essential for fetus expulsion, peaks during parturition. The peak of oxytocin may be too late to participate in the regulation of colostrum nutrient synthesis. However, its amplitude and duration may influence the transfer of IgG into lacteal secretions by delaying the closure of tight junctions between mammary epithelial cells.

Finally, a link between colostrum yield and farrowing kinetics was reported in primiparous sows (Quesnel, 2011). The rhythm of piglet births during the onset of parturition was lower in sows with a low colostrum production (<3 kg) than in sows with a medium (3–4 kg) or high (>4 kg) production. It can be suggested that abnormalities in the endocrine status of sows in late pregnancy may affect the processes of both parturition and colostrumogenesis, and these two processes being partly regulated by common hormones, such as progesterone and maybe estrogens and glucocorticoids.

Colostrum yield was shown to be positively correlated with plasma concentrations of IGF-I in prepartal sows (Foisnet *et al.*, 2010a). Accordingly, it was suggested that the 20% increase in colostrum yield observed when sows were fed a fermented potato protein product for 7 days before expected farrowing could be due to greater IGF-I concentrations in sow plasma (Wavreille *et al.*, 2010). It can be hypothesized that IGF-I could stimulate lactogenesis because it can act as a mitogenic factor or an anabolic hormone that favors nutrient uptake by MEC, or both. However, these observations are insufficient to conclude for a stimulatory effect of IGF-I on colostrum production.

Table 1 Colostrum yield according to sow parity

Colostrum yield (kg) according to sow parity	n ¹	Reference
Parity 1 (3.4 kg) ^a < Parities 2 to 3 (4.3 kg) ^b	40	Devillers <i>et al.</i> , 2007
Parities 1 to 3 (3.7 kg) ^a > Parities 4 to 7 (2.8 kg) ^b	37	Decaluwé <i>et al.</i> , 2013
Parities 1 to 2 (3.7 kg) ^a > Parities 3 to 8 (3.2 kg) ^b	121	Quesnel, unpublished
Parity 1 (4.8 kg) ^a < Parities 2 to 5 (5.0 kg) ^b	183	Boonraungrod <i>et al.</i> , 2018

¹Number of sows.

^{ab}Average colostrum yield with different letters differed ($P < 0.05$).

Effects of exogenous hormones

Inducing parturition. A key factor for successful intensive farrowing management is farrowing supervision, hence natural and synthetic prostaglandin F₂α (PGF₂α) are commonly used in pig farms to induce parturition. Although one study reported that inducing parturition on day 114 of gestation (i.e., one day before the average term) decreased colostrum yield, other studies showed no significant effect (Table 2). Nevertheless, farrowing induction on day 113 of gestation induced transient alterations in plasma hormonal concentrations and colostrum composition (Foisnet *et al.*, 2011). Increases in prolactin and cortisol concentrations were observed 1 h after the injection of PGF₂α. Colostrum collected at the onset of parturition (T0) contained more lactose and less IgA when parturition was induced, whereas colostrum composition at T24 did not differ between sows with natural or induced farrowing (Foisnet *et al.*, 2011). Farrowing induction had no effect on colostral concentrations of IgG (Foisnet *et al.*, 2011; Otto *et al.*, 2017). It is apparent that if farrowing induction is performed not earlier than one day before the expected farrowing date (based on individual monitoring or on the average farrowing date in the herd), no negative impact on colostrum yield is expected. However, the effect of farrowing induction on IgA concentrations and on the subsequent occurrence of neonatal diarrhea warrants further investigation.

Preventing premature parturition. In a retrospective study, based on 61 000 farrowing records, a 10% occurrence of early parturition (<114 days) was observed (Vanderhaeghe *et al.*, 2011). Premature parturition may affect the birth weight, survival and growth of piglets and it interferes with the management procedures on the farm. It is thus important to prevent premature parturition in herds where its incidence is high. This can be achieved by administering an analog of progesterone (altrenogest) to maintain plasma concentrations of progesterone during the last few days of pregnancy. Administration of altrenogest from days 109 to 112 or 113 of gestation did not affect colostrum yield, likely because the treatment delayed farrowing and main hormonal changes without affecting the relative chronology of these changes (Foisnet *et al.*, 2010b). Such a treatment was tested on a

large scale in commercial herds, with no report of negative impact on piglet survival until weaning (Vanderhaeghe *et al.*, 2011), which is consistent with the lack of negative effect on colostrum yield.

Injecting oxytocin. As mentioned earlier, the composition of colostrum is affected by the status of tight junctions between mammary epithelial cells. Therefore, the ability to manipulate mammary tight junctions in the late colostral phase could allow Ig concentrations to be maintained at higher levels for a longer period of time, thereby increasing the quality of early milk. Injecting a supraphysiological dose of oxytocin to sows on day 2 of lactation (i.e., between 12 and 20 h after birth of the last piglet) increased the Na : K ratio (0.48 v. 0.28) and concentrations of IGF-I (78.1 v. 37.2 ng/ml), IgG (18.8 v. 6.6 mg/ml) and IgA (4.1 v. 2.7 mg/ml) in milk collected 8 h after the injection (Farmer *et al.*, 2017). The injection of oxytocin in the early *post-partum* period therefore delayed the occurrence of tightening of mammary tight junctions and prolonged the colostral phase, thereby having beneficial effects on the composition of early milk.

Effects of nutrition on colostrogenesis

The development of strategies to improve colostrum composition through maternal feeding has been largely explored, with a special focus on colostral fat and Ig because both are sensitive to nutritional changes. In contrast, very few attempts were made to increase colostrum yield, which is most likely due to the difficulty in measuring the amount of colostrum produced. Most recent attempts focused on maternal feeding during late gestation, when milk-specific constituents are synthesized. Alternatively, long-term strategies might aim at increasing mammary gland development. Although it is well known that milk yield depends on the number of mammary epithelial cells present at the onset of lactation (Head and Williams, 1991), it is not known whether this is also the case for colostrum production. Nutritional strategies to improve colostrum synthesis might also aim at managing sow body reserves and metabolic status during late pregnancy and this is covered in more

Table 2 Colostrum yield and gestation length in sows with farrowings induced on day 113 or 114 of gestation

Treatment	Colostrum yield, kg		Gestation length, days		n ¹	Reference
	Control	Induced ²	Control	Induced ²		
PGF ₂ α ³ analog on day 114	4.16	3.37**	113.3	114.6*	40 P1 to P6	Devillers <i>et al.</i> , 2007
PGF ₂ α ³ analog on day 113	4.47	3.95 [†]	113.7	114.0	47 P1	Foisnet <i>et al.</i> , 2011
PGF ₂ α ³ analog on day 114	4.81	4.66	–	–	96 P3 to P7	Otto <i>et al.</i> , 2017
PGF ₂ α ³ analog on day 114	4.84	4.90	114.8	114.7	130 P1 to P5	Boonraungrod <i>et al.</i> , 2018
PGF ₂ α ³ analog on day 114 + carbetocin ⁴	4.84	4.05*	114.8	115.1	133 P1 to P5	Boonraungrod <i>et al.</i> , 2018

¹Total number of sows and parity (P).

²Treatment effect: ***P* < 0.01, **P* < 0.05, [†]*P* < 0.10.

³Prostaglandin F₂α.

⁴Carbetocin is an oxytocin-like molecule with long-acting properties; it was injected 24 h after the PGF₂α analog.

Table 3 Nutritional manipulation of colostrum yield and effects on macrochemical composition of colostrum in sows

Treatment ¹	Duration	Effects on colostrum ²		n ³	Reference
		Yield	Composition		
Level of feeding 4.5 v. 1.5 kg/d	Day 108 to term	↗ [†]	↗ lactose* ↘ protein*	50 P2 to P7	Decaluwé <i>et al.</i> , 2014
Source or level of fat or dietary fiber Pectin residue ⁴	Mating to day 107	↗*	↗ lactose* ↘ protein* ↘ energy*	20	Theil <i>et al.</i> , 2014a
Various sources of fat ⁵	Day 108 to term	NS	↗ lactose* ⁶ ↘ DM ^{†6}	40	Theil <i>et al.</i> , 2014a
High dietary fiber ⁷ Various ingredients	Day 106 to term	NS	↗ fat*	29 P1	Loisel <i>et al.</i> , 2013
CLA ¹ , 1.3%	Day 108 to term	↘ [†]	↗ fat [†]	23 P1-P3	Krogh <i>et al.</i> , 2012
CLA ¹ , 0.75, 1.5 or 2.25%	Day 85 to term	NS	↘ fat	48 MP	Wu <i>et al.</i> , 2015
HMB ¹ , 2.0 g/d	Day 111 to term	–	↗ fat**	68 MP	Nissen <i>et al.</i> , 1994
HMB ¹ , 2.5 g/d	Day 108 to term	NS ⁸	No effect	16 P2	Flummer and Theil, 2012
Arginine, 25 g/d	Day 30 to weaning	NS	↘ lactose* ↗ protein [†] ↗ IGF-I [†]	21 MP	Krogh <i>et al.</i> , 2016
Yeast derivates, 2 g/kg diet	Gestation + lactation	↗*	↗ fat*	37 P1-P8	Hasan <i>et al.</i> , 2018

¹CLA: Conjugated linoleic acid; HMB: β-hydroxy β-methyl butyrate.

²Treatment effect: ***P* < 0.01, **P* < 0.05, [†]*P* < 0.10, NS: not significant.

³Total number of sows and parity (P, when known). MP: multiparous sows.

⁴A high-fiber diet formulated with pectin residue (32% DM, insoluble dietary fiber) was compared with a high-fiber diet rich in potato pulp (40% DM, soluble dietary fiber).

⁵Prefarrowing diets: standard (3% animal fat), 8% coconut oil, 8% sunflower oil, 8% fish oil or 4% fish oil + 4% octanoic acid.

⁶Effects of dietary supplements of 8% fish oil or 4% fish oil + 4% octanoic acid versus 8% coconut oil.

⁷A high-fiber diet with 23% insoluble and soluble dietary fiber was compared with a diet containing 13% insoluble and soluble dietary fiber.

⁸Colostrum yield per piglet tended to be increased (512 v. 434 g) but colostrum yield for the whole litter did not differ (5.3 v. 5.0 kg).

detail by Quesnel *et al.* (2015). However, the fact remains that the relation between sow metabolic state during gestation and colostrum yield is not clear, with discrepancies between findings.

Sow nutrition and colostrum yield

Manipulating circulating concentrations of prolactin. Given the effects of *prepartum* prolactin concentrations on colostrum yield, the development of strategies to influence prolactin concentrations through nutritional approaches is warranted. Silymarin, an extract from the milk thistle plant (*Silybum marianum*), is known for its hyperprolactinemic properties. Yet an attempt to induce hyperprolactinemia in sows before parturition by providing silymarin was not successful (Loisel *et al.*, 2014). To the best of our knowledge, there are no other known nutritional means to increase prolactin concentrations in swine.

Manipulating circulating concentrations of progesterone.

Circulating concentrations of progesterone can be influenced by the level of feeding. Indeed, in gilts and ewes, progesterone concentrations were reduced with a high feeding level, likely because of an increase in metabolic clearance rate (Banchero *et al.*, 2015). Supplementing ewes with energy during the last week of pregnancy accelerates the

decline in circulating progesterone that, in turns, increases colostrum yield (Banchero *et al.*, 2015). However, increasing sow feed intake for the last 2 weeks of pregnancy did not affect circulating concentrations of progesterone and piglet weight gain during the first 2 days *postpartum*, indicating that colostrum yield was unaltered (Miller *et al.*, 2004). Authors suggested that this absence of effect could be tied to the stimulatory effect of increased feeding level (via greater insulin and IGF-I concentrations) on the production of progesterone by corpora lutea, occurring concurrently with the increased clearance rate of progesterone. This problem may be overcome by imposing a high level of feeding for a shorter period of time at the beginning of luteolysis (i.e., 2 days before parturition) in order to accelerate the decrease in progesterone without stimulating its synthesis.

Manipulating colostrum yield. The few published experiments where nutritional attempts were made to stimulate colostrum yield used various strategies based on feeding level, source or level of fat and dietary fiber, and feeding specific nutrients (Table 3). A high feeding level (three times 1.5 kg daily compared with 1.5 kg/day) during the last week of gestation tended to increase colostrum yield from 3.5 to 4.0 kg (Decaluwé *et al.*, 2014). This effect may be related to the greater lactose content of colostrum from overfed sows,

lactose being the major osmotic factor that favors water transfer into lacteal secretions. Interestingly, Decaluwé *et al.* (2014) also showed that sow body condition had more impact on colostrum yield than feeding level. The highest colostrum yield was achieved when sows entered the farrowing unit with a moderate body condition (i.e., with a backfat thickness between 17 and 23 mm) and were fed the high plane of nutrition for a week before farrowing.

Theil *et al.* (2014a) combined two feeding strategies by providing various levels and sources of dietary fiber during most of gestation (from mating to day 107) and then various levels and sources of fat during the last week of gestation (Table 3). The gestation diet, but not the pre-farrowing diet, influenced colostrum yield, with sows fed pectin residue (rich in insoluble fiber) producing more colostrum than sows fed potato pulp (rich in soluble fiber; 6.2 v. 4.6 kg). It is not clear why sows fed pectin residue produced more colostrum but this was consistent with a greater lactose content in colostrum. However, when sows were fed sugar beet pulp (rich in soluble dietary fiber), their colostrum yield was intermediate compared with that of sows fed pectin residue or potato pulp, despite their higher lactose content in colostrum. Loisel *et al.* (2013) found no effect of high dietary fiber providing both soluble and insoluble fibers (23% of dietary fiber originating from a mixture of soybean hulls, wheat bran, undecorticated sunflower meal and sugar beet pulp) on colostrum yield when compared with a low fiber diet (13% of dietary fiber). Nevertheless, this high-fiber diet increased by nearly 60% the colostrum intake of low birth weight-piglets (<900 g) and was associated with lower piglet mortality until weaning. Clearly the effects of dietary fiber source on colostrum yield or composition and their underlying mechanisms need further attention.

The potential effects of various specific ingredients, namely conjugated linoleic acid (CLA), the leucine metabolite β -hydroxy β -methyl butyrate (HMB) and arginine, on colostrum yield were also investigated, with no significant effects being reported (Table 3). A trend for lower colostrum yield was reported in response to CLA supplementation during the last week of gestation (Krogh *et al.*, 2012).

Collectively, these data indicate that nutrition in late pregnancy is important for colostrum yield of sows, but that more investigations are needed to determine which nutrients or feeding strategies are most effective and which part of gestation (last week, last three weeks or last third) is most critical for colostrum production.

Sow nutrition and colostrum fat content

The effects of sow nutrition, encompassing feed intake, energy intake and source, and protein intake, on macrochemical composition of colostrum, have been extensively reviewed, as well as the effects of vitamins and minerals on microchemical composition (Farmer and Quesnel, 2009). The present review therefore only covers most recent findings, with a special focus on fat because it is by far the most labile macronutrient of colostrum. The fatty acid content of colostrum greatly depends on the amount of lipids provided in the

sow diet during late gestation, whereas the fatty acid profile is largely influenced by the type of lipid being fed to the pregnant sow (Farmer and Quesnel, 2009). Additional colostrum fat may be used, to a certain extent, as supplementary energy by the piglet and it may also increase lipid accretion, thereby enabling piglets to maintain a constant body temperature. Moreover, specific fatty acids are important for piglet maturation, such as polyunsaturated fatty acids (PUFA) which are known to play a role in brain development and function. Various studies were therefore carried out with the goal of increasing the amount of polyunsaturated fats in sow colostrum. This was successfully achieved via the inclusion of various fish oils or linseed oil in the diet of pregnant sows (Rooke *et al.*, 1998; 2001). Supplementation of soybean oil in the diet of sows during the last week of pregnancy also increased the fat content and the concentrations of polyunsaturated fatty acids in colostrum (Bai *et al.*, 2017). Dietary CLA also affects the fatty acid composition of colostrum fat in sows, increasing saturated fatty acids and decreasing monounsaturated fatty acids (Bee, 2000).

Besides the addition of fat to the maternal diet, providing additional dietary fiber in late gestation may be another way of increasing the fat content of colostrum. Loisel *et al.* (2013) demonstrated that feeding 23.4% compared with 13.3% dietary fiber to sows from day 106 of gestation until parturition increased colostrum fat from 8.3% in control sows to 10.7% in treated sows. Such an effect could not be attributed to dietary lipid intake or mobilization of body lipids since they did not differ significantly between sows fed the low and the high level of dietary fiber. This effect may be due to the fact that dietary fiber increases plasma concentrations of short chain fatty acids, including acetate which can be used by the mammary gland as precursors for *de novo* synthesis of lipids (Linzell *et al.*, 1969).

Colostrum fat content was also increased by supplementing the sow diet with the leucine metabolite HMB (2.0 g/day of Ca(HMB)₂) in the last 3 to 4 days before parturition (6.9 v. 4.9%; Nissen *et al.*, 1994). The mechanism of action of HMB is likely to be linked to its lipolytic effect, causing fat mobilization and elevated plasma concentrations of non-esterified fatty acids that can be used for *de novo* fat synthesis (Theil *et al.*, 2014b). However, Flummer and Theil (2012) could not reproduce this effect of HMB supplementation on colostrum composition (Table 3). Lastly, supplementing the sow diet with glutamine and glutamate can also affect colostrum fat concentration. Providing the two amino acids as Aminogut (1.5% w/w) during the last week of gestation increased colostrum fat by 60% (7.4 v.4.6%, Santos de Aquinos *et al.*, 2014). However, it is not known if glutamine or glutamate stimulate lipid synthesis or act as substrates for *de novo* lipogenesis.

Sow nutrition and immunoglobulin contents of colostrum

Numerous ingredients that presumably have immunomodulating effects were tested for their potential beneficial effects on the Ig concentrations of swine colostrum. These

Table 4 Nutritional manipulation of immunoglobulin concentrations in colostrum in sows, presumably through immunomodulating effects of active components¹

Treatment	Duration ²	Effect ³	References ⁴
Conjugated linoleic acid, 0.5%	8 days	↗ IgG	Bontempo <i>et al.</i> , 2004
Conjugated linoleic acid, 0.75, 1.5 or 2.25%	7 days	↗ IgG, IgA, IgM	Corino <i>et al.</i> , 2009
Conjugated linoleic acid, 0.5%	4 weeks	↗ IgG, IgA, IgM	Wu <i>et al.</i> , 2015
Shark liver oil	35 days	↗ IgG, = IgA	Mitre <i>et al.</i> , 2005
Phytogetic feed additive (source of essential oils)	1 week	↗ IgG, IgA	Wang <i>et al.</i> , 2008
Plant extract	1 week	↘ IgG, IgA	Illesley <i>et al.</i> , 2003
Saponins (plant extract)	3 weeks ⁵	= IgG, IgA	Illesley and Miller, 2005
Oregano essential oil	Whole gestation	= IgG, IgA	Ariza-Nieto <i>et al.</i> , 2011
Non-specific immunostimulants ⁶	4 to 6 weeks	↗ IgG	Krakowski <i>et al.</i> , 2002
Mannan oligosaccharides	2 weeks	↗ IgM; = IgG, IgA	Newman and Newman, 2001
Mannan oligosaccharides	3 weeks	↗ IgG, IgA, IgM	O'Quinn <i>et al.</i> , 2001
Fructo oligosaccharides	28 days	= IgG, ↗ IgA	Le Bourgot <i>et al.</i> , 2014
Astragalus polysaccharides	1 week	↗ IgG, IgM	Tan <i>et al.</i> , 2017
Yeast derivatives ⁷	Gestation	= IgG, IgA, IgM	Bass <i>et al.</i> , 2012
Yeast derivatives ⁸	Gestation	= IgG, IgA, IgM	Hasan <i>et al.</i> , 2018
Fermented liquid feed	2 weeks	↗ IgG, IgA, = IgM	Demečková <i>et al.</i> , 2003
<i>Saccharomyces cerevisiae boulardii</i>	21 days	↗ IgG, IgA	Guillou <i>et al.</i> , 2012
<i>Saccharomyces cerevisiae</i> ⁹	Gestation	↗ IgG, = IgA	Zanello <i>et al.</i> , 2013

¹Published in part by Farmer and Quesnel (2009).

²Duration of treatment before parturition.

³↗ increased immunoglobulin (Ig) concentrations; = no significant effect on Ig concentrations; ↘ reduced Ig concentrations.

⁴The list of references is given in Supplementary material S1.

⁵From days 72 to 93 of pregnancy.

⁶Factors that induce non-specific defense mechanisms of the organism toward the viral and bacterial pathogens.

⁷Whole extract derived from *P. guilliermondii*.

⁸Whole extract derived from *S. cerevisiae*.

⁹Two strains were tested: *S. cerevisiae* strain CNCM I-4407 and *S. cerevisiae* var. *boulardii* strain CNCM I-3799. A third strain (*S. cerevisiae* strain LC 10-05) had no effect on colostrum IgG and IgA.

ingredients belong to various families of components: specific fatty acids (such as CLA and fish oils), plant extracts (as essential oils), prebiotics (mannan oligosaccharides (MOS) or fructo oligosaccharides (FOS)) and probiotics (mainly *Saccharomyces cerevisiae boulardii*), or ingredients containing pre- or probiotics (fermented liquid feed). When fed to late pregnant sows, many of these ingredients increased IgG, IgA and (or) IgM concentrations in colostrum (Table 4).

As mentioned earlier, long-chain PUFA of the n-6 and n-3 series have received a lot of attention in sow nutrition because of their role in brain development. They may also have health-promoting properties through impacts on inflammatory and immune status. Conjugated linoleic acid which is the collective name for isomers of linoleic acid (an n-6 PUFA) was efficient in increasing Ig contents of colostrum, at various doses (Table 4). This effect was likely due to the increase in Ig concentrations in sow serum, itself potentially mediated through an increase in interleukin-2 synthesis in response to CLA supplementation (Wu *et al.*, 2015). The increase in colostrum IgG observed when the sow diet was supplemented with shark-liver oil was also associated with greater Ig concentrations in sow plasma. Shark-liver oil is a source of two bioactive lipids, the alkylglycerols and n-3 PUFA, and the increased plasmatic Ig production

by sows was mainly attributed to alkylglycerols. These supplementations improved the maternal transfer of passive immunity to the piglets, as indicated by greater IgG concentrations in piglet blood. Interestingly also, there are recent indications that feeding fish oil during the last month of gestation can affect the antioxidant capacity of sow colostrum (Shen *et al.*, 2015), yet much remains to be learned on this topic.

A wide range of beneficial properties are also attributed to plant-derived compounds, among them antimicrobial activity, antioxidant property and effects on the immune system. However, both positive and negative effects on colostrum Ig content were reported (Table 4).

Finally, probiotics and prebiotics receive considerable attention currently because of many potential beneficial effects on health. In short, probiotics are living organisms which colonize the gastrointestinal tract while prebiotics are fermented compounds that induce changes in gut microbiota. Effects on gut microbiota may be accompanied by beneficial effects on the gut immune system with consequences on the whole organism. Providing living bacteria (*Lactobacillus* strains in fermented liquid feed) or living cells (as *Saccharomyces cerevisiae*) or prebiotics MOS or FOS was shown to increase IgG or IgA concentrations in colostrum

(Table 4). In contrast, yeast extract provided throughout gestation had no effect on colostrum Igs. Interestingly, however, a recent study reported no effects of prebiotics supplementation on IgG, IgA or IgM content in sow colostrum but positive effects on both fat content and colostrum yield were observed (Hasan *et al.*, 2018). Multiparous sows were fed a control diet or a control diet supplemented with 2 g of yeast derivatives throughout pregnancy. This extract contained a mix of mannan oligosaccharides and polysaccharides, glucomannoproteins and betaglucans. Protein and lactose contents of colostrum were not affected, whereas fat content was increased by 21% and colostrum yield by 24%.

In conclusion, the most promising results on colostrum Ig content were obtained with CLA, prebiotics in the form of MOS and FOS, and living yeast cells, while the effects of plant extracts and yeast extracts are far less convincing.

Conclusions

Colostrum is crucial for the survival, development and health of piglets so that knowledge about mechanisms that regulate or influence colostrogenesis is essential. Although progress has been made in understanding the factors that regulate colostrum yield and composition, including the endocrine regulation and the impact of sow nutrition, several essential questions remain unanswered. Namely, when is colostrum synthesized precisely? What are the impacts of sow nutrition and metabolic status during late gestation? Does mammary development affect colostrum yield? Promising results on colostrum yield or quality were obtained by manipulating levels and sources of dietary fat and fiber during late gestation, or by supplementing sow diets with pre- and probiotics. In the future, nutritional strategies should be tested in commercial farms, using large numbers of animals, in order to investigate whether, beyond their effects on colostrum yield or composition, they affect piglet performance in terms of survival, growth, health and robustness at weaning.

Acknowledgement

None.

Declaration of interest

The authors declare that they have no competing interests.

Ethics statement

Not applicable.

Software and data repository resources

No data were deposited in an official repository.

Supplementary material

To view supplementary material for this article, please visit <https://doi.org/10.1017/S1751731118003555>

References

- Bai YS, Wang CQ, Zhao X, Shi BM and Shan AS 2017. Effects of fat sources in sow on the fatty acid profiles and fat globule size of milk and immunoglobulins of sows and piglets. *Animal Feed Science and Technology* 234, 217–227.
- Banchero GE, Milton JTB, Lindsay DR, Martin GB and Quintans G 2015. Colostrum production in ewes: a review of regulation mechanisms and of energy supply. *Animal* 9, 831–837.
- Barrington GM, McFadden TB, Huyler MT and Besser TE 2001. Regulation of colostrogenesis in cattle. *Livestock Production Science* 70, 95–104.
- Bartol FF, Wiley AA, George AE, Miller DJ and Bagnell CA 2017. Postnatal reproductive development and the lactocrine hypothesis. *Journal of Animal Science* 95, 2200–2210.
- Bee G 2000. Dietary conjugated linoleic acid alters adipose tissue and milk lipids of pregnant and lactating sows. *Journal of Nutrition* 130, 2292–2298.
- Boonraungrod N, Sutthiya N, Kumwan P, Tossakui P, Nuntapaitoon M, Muns R and Tummaruk P 2018. Control of parturition in swine using PGF₂α in combination with Carbetocin. *Livestock Science*, in press.
- Decaluwé R, Maes D, Cools A, Wuyts D, De Smet S, Marescau B, De Deyn PP and Janssens GPJ 2014. Effect of periparturient feeding strategy on colostrum yield and composition in sows. *Journal of Animal Science* 92, 3557–3567.
- Decaluwé R, Maes D, Declerck I, Cools A, Wuyts D, De Smet S and Janssens GPJ 2013. Changes in back fat thickness during late gestation predict colostrum yield in sows. *Animal* 7, 1999–2007.
- Declerck I, Dewulf J, Sarrazin S and Maes D 2016. Long-term effects of colostrum intake in piglet performance and mortality. *Journal of Animal Science* 94, 1633–1643.
- Devillers N, Farmer C, Le Dividich J and Prunier A 2007. Variability of colostrum yield and colostrum intake in swine. *Animal* 1, 1033–1041.
- Devillers N, van Milgen J, Prunier A and Le Dividich J 2004. Estimation of colostrum intake in the neonatal pig. *Animal Science* 78, 305–313.
- Farmer C, Lessard M, Knight CH and Quesnel H 2017. Oxytocin injections in the postparturient period affect mammary tight junctions in sows. *Journal of Animal Science* 95, 3532–3539.
- Farmer C, Robert S and Rushen J 1998. Bromocriptine given orally to periparturient or lactating sows inhibits milk production. *Journal of Animal Science* 76, 750–757.
- Farmer C and Quesnel H 2009. Nutritional, hormonal and environmental effects on colostrum in sows. *Journal of Animal Science* 87 (Suppl. 1), 56–65.
- Flummer C and Theil PK 2012. Effect of beta-hydroxy beta-methyl butyrate supplementation of sows in late gestation and lactation on sow production of colostrum and milk and piglet performance. *Journal of Animal Science* 90, 372–374.
- Foisnet A, Farmer C, David C and Quesnel H 2010a. Relationship between colostrum production by primiparous sows and sow physiology around parturition. *Journal of Animal Science* 88, 1672–1683.
- Foisnet A, Farmer C, David C and Quesnel H 2010b. Progestagen treatment during late pregnancy did not reduce colostrum yield in primiparous sows. *Journal of Animal Science* 88, 1684–1693.
- Foisnet A, Farmer C, David C and Quesnel H 2011. Farrowing induction induced transient alterations in prolactin concentrations and colostrum composition in primiparous sows. *Journal of Animal Science* 89, 3048–3059.
- Gross JJ, Kessler EC, Bjerre-Harpoth V, Dechow C, Baumrucker CR and Bruckmaier RM 2014. Periparturient progesterone and prolactin have little effect on the rapid transport of immunoglobulin G into colostrum of dairy cows. *Journal of Dairy Science* 97, 2923–2931.
- Hartmann PE, Whitely JL and Willcox DL 1984. Lactose in plasma during lactogenesis, established lactation and weaning in sows. *Journal of Physiology* 347, 453–463.
- Hasan S, Junnikkala S, Peltoniemi O, Paulin L, Lyyski A, Vuorenmaa J and Oliviero C 2018. Dietary supplementation with yeast hydrolysate in pregnancy influences colostrum yield and gut microbiota of sows and piglets after birth. *PLoS One* 13, e0197586.
- Head RH and Williams IH 1991. Mammarygenesis is influenced by pregnancy nutrition. In *Manipulating pig production III* (ed. ES Batterham), p 33. Australasian Pig Science Association, Werribee, VIC, Australia.

- Huang SC, Hu Z, Hasler-Rapacz J and Rapacz J 1992. Preferential mammary storage and secretion of immunoglobulin gamma (IgG) subclasses in swine. *Journal of Reproductive Immunology* 21, 15–28.
- Kensinger RS, Collier RJ and Bazer FW 1986. Ultrastructural changes in porcine mammary tissue during lactogenesis. *Journal of Anatomy* 145, 49–59.
- Krogh U, Flummer C, Jensen SK and Theil PK 2012. Colostrum and milk production of sows is affected by dietary conjugated linoleic acid. *Journal of Animal Science* 90, 366–368.
- Krogh U, Pksbjerg N, Purup S, Ramackers P and Theil PK 2016. Colostrum and milk production in multiparous sows fed supplementary arginine during gestation and lactation. *Journal of Animal Science* 94, 22–25.
- Linzell JL, Mephram TB, Annison EF and West CE 1969. Mammary metabolism in lactating sows: arteriovenous differences of milk precursors and the mammary metabolism of [¹⁴C]glucose and [4 C]acetate. *British Journal of Nutrition* 23, 319–332.
- Loisel F, Farmer C, Ramaekers P and Quesnel H 2013. Effect of high dietary fiber during late pregnancy on sow physiology, colostrum production, and piglet performance. *Journal of Animal Science* 91, 5269–5279.
- Loisel F, Farmer C, Ramaekers P and Quesnel H 2014. Colostrum yield and piglet growth during lactation are related to gilt metabolic and hepatic status pre-partum. *Journal of Animal Science* 92, 2931–2941.
- Loisel F, Farmer C, van Hees H and Quesnel H 2015. Relative prolactin to progesterone concentrations around farrowing influence sow colostrum yield. *Domestic Animal Endocrinology* 53, 35–41.
- Miller HM, Foxcroft GR and Aherne FX 2004. Increasing feed intake in late gestation does not affect plasma progesterone concentration in the sow. *Theriogenology* 62, 1618–1626.
- Nissen S, Faidley TD, Zimmerman DR, Izard R and Fisher CT 1994. Colostral milk fat percentage and pig performance are enhanced by feeding the leucine metabolite beta-hydroxy-beta-methyl butyrate to sows. *Journal of Animal Science* 72, 2331–2337.
- Otto MA, Machado AP, Moreira LP, Bernardi ML, Coutinho ML, Vaz IS Jr, Wentz I and Bortolozzo FP 2017. Colostrum yield and litter performance in multiparous sows subjected to farrowing induction. *Reproduction in Domestic Animals* 52, 749–755.
- Palombo V, Looor JJ, D'Andrea M, Vailati-Riboni M, Shazad K, Krogh U and Theil PK 2018. Transcriptional profiling of swine mammary gland during the transition from colostrumogenesis to lactogenesis using RNA sequencing. *BMC Genomics* 19, 322.
- Quesnel H 2011. Colostrum production by sows: variability of colostrum yield and immunoglobulin G concentrations. *Animal* 5, 1546–1553.
- Quesnel H, Farmer C and Devillers N 2012. Colostrum intake: influence on piglet performance and factors of variation. *Livestock Science* 145, 105–114.
- Quesnel H, Farmer C and Theil PK 2015. Colostrum and milk production. In *The gestating and lactating sow* (ed. C Farmer), pp. 173–192. Wageningen Academic Publishers, Wageningen, the Netherlands.
- Quesnel H, Ramaekers P, van Hees H and Farmer C 2013. Short communication: relations between peripartum concentrations of prolactin and progesterone in sows and piglet growth in early lactation. *Canadian Journal of Animal Science* 93, 109–112.
- Rooke JA, Bland IM and Edwards SA 1998. Effect of feeding tuna oil or soyabean oil as supplements to sows in late pregnancy on piglet tissue composition and viability. *British Journal of Nutrition* 80, 273–280.
- Rooke JA, Sinclair AG, Edwards SA, Cordoba R, Pkiyach R, Penny PC, Penny P, Finch AM and Horgan GW 2001. The effect of feeding salmon oil to sows throughout pregnancy on pre-weaning mortality of piglets. *Animal Science* 73, 489–500.
- Santos de Aquino R, Dutra Junior WM, Manso HECC, MansoFilho HC, Kutschenko M, Nogueira ET and Watford M 2014. Glutamine and glutamate (AminoGut) supplementation influences sow colostrum and mature milk composition. *Livestock Science* 169, 112–117.
- Schnulle PM and Hurley WL 2003. Sequence and expression of the FcRn in the porcine mammary gland. *Veterinary Immunology and Immunopathology* 91, 227–231.
- Shen Y, Wan H, Zhu J, Fang Z, Che L, Xu S, Lin Y, Li J and Wu D 2015. Fish oil and olive oil supplementation in late pregnancy and lactation differentially affect oxidative stress and inflammation in sows and piglets. *Lipids* 50, 647–658.
- Taverne M, Bevers M, Bradshaw JMC, Dieleman SJ, Willemsse AH and Porter DG 1982. Plasma concentrations of prolactin, progesterone, relaxin, and oestradiol-17 β in sows with progesterone, bromocriptine or indomethacin during late pregnancy. *Journal of Reproduction and Fertility* 65, 85–96.
- Theil PK, Flummer C, Hurley WL, Kristensen NB, Labouriau R and Sørensen MT 2014a. Mechanistic model to predict colostrum intake based on deuterium oxide dilution technique and impact of of gestation and prepartum diets on piglet intake and sow yield of colostrum. *Journal of Animal Science* 92, 5507–5519.
- Theil P, Lauridsen C and Quesnel H 2014b. Neonatal piglet survival: Impact of sow nutrition around parturition on fetal glycogen deposition and production and composition of colostrum and transient milk. *Animal* 8, 1021–1030.
- Vadmand CN, Krogh U, Hansen CF and Theil PK 2015. Impact of sow and litter characteristics on colostrum yield, time for onset of lactation, and milk yield of sows. *Journal of Animal Science* 93, 2488–2500.
- Vanderhaeghe C, Dewulf J, Jourquin J, De Kruijff A and Maes D 2011. Incidence and Prevention of Early Parturition in Sows. *Reproduction in Domestic Animals* 46, 428–433.
- Wavreille J, Planchon V, Renaville R, Forier R, Agneessens R and Bartiaux-Thill N 2010. Effet d'un produit fermenté de pomme de terre dans l'aliment de la truie en maternité sur les performances des truies et des porcelets allaités. *Journées de la Recherche Porcine* 42, 133–134.
- Wu F, Li PL, Bai LL, Liu H, Lai CH, Thacker PA and Wang FL 2015. Responses in colostrum production and immunoglobulin concentrations to conjugated linoleic acid fed to multiparous sows during late gestation. *Animal Feed Science and Technology* 210, 200–208.