

## Phyto-oestrogen content of berries, and plasma concentrations and urinary excretion of enterolactone after a single strawberry-meal in human subjects

W. M. Mazur<sup>1</sup>, M. Uehara<sup>1</sup>, K. Wähälä<sup>2</sup> and H. Adlercreutz<sup>1\*</sup>

<sup>1</sup>*Folkhälsan Institute for Preventive Medicine, Nutrition and Cancer, and Department of Clinical Chemistry, University of Helsinki, PO Box 60, FIN-00014 Helsinki, Finland*

<sup>2</sup>*Department of Chemistry, PO Box 55, FIN-00014 University of Helsinki, Finland*

(Received 15 February 1999 – Revised 1 September 1999 – Accepted 14 September 1999)

Quantitative data on phyto-oestrogen, particularly lignan, content in edible plants are insufficient. We, therefore, measured isoflavonoids and lignans in nine edible berries using an isotope dilution gas chromatography–mass spectrometry method for foods and found substantial concentrations of the lignan secoisolariciresinol (1.39–37.18 mg/kg DM), low amounts of matairesinol (0–0.78 mg/kg DM) and no isoflavones. To determine pharmacokinetics and urinary excretion pattern of the mammalian lignan enterolactone derived from plant lignans, a study with human subjects was conducted. Five healthy women and two men consumed, after a 72 h period of a phyto-oestrogen-free regimen, a single strawberry-meal containing known amounts of plant lignans. Basal and post-meal blood and urine samples were collected at short intervals. The samples were analysed using time-resolved fluoroimmunoassay of enterolactone. The meal increased plasma concentration of enterolactone after 8–24 h and in urine in the 13–24 h and 25–36 h urine collections. High individual variability of the metabolic response was observed. Enterolactone excreted in the urine collected throughout the 48 h post-meal yielded on average 114 % of the plant lignans consumed. It is concluded that berries containing relatively high concentrations of plant lignans contribute to plasma and urinary levels of mammalian enterolactone in human subjects.

### Berries: Phyto-oestrogens: Kinetics

Epidemiological investigations in human subjects have provided evidence linking diet to incidence of cancer (e.g. breast, prostate cancer) and other degenerative disorders (e.g. CHD, postmenopausal osteoporosis). The detection and identification in human subjects of lignans and isoflavonoids, plant-derived oestrogenic compounds abundant in plasma of subjects living in areas with low cancer incidence, have challenged studies on their metabolic pathways and role in human health and disease.

Despite laboratory research efforts revealing interesting properties of phyto-oestrogens as inhibitors of carcinogenic and atherosclerotic processes, there is a dearth of studies on their presence in terms of quantity and quality in the human diet. Very little is known about the fate of lignans in common foods after ingestion in human subjects. There is a single study in rats, however, in which food lignan content was measured and then urinary levels of mammalian lignans were measured following ingestion (Landström *et al.* 1998).

As early as in the middle of the 1980s, it was suggested

(Adlercreutz *et al.* 1986a, 1987, 1988) that berries and fruits containing plant lignans may be the precursors of mammalian enterolactone (ENL) and enterodiol (END). This was based on correlations found between dietary intake of fibre from berries and fruits, and urinary excretion of mammalian lignans END and ENL. In order to verify this suggestion we developed the isotope dilution gas chromatography–mass spectrometry method (ID-GC–MS) for food samples and analysed both lignans and isoflavones in some edible berries. The present paper describes for the first time concentrations of plant oestrogens in nine different types of berries.

The urinary excretion of lignans (Fig. 1) has been the most frequently studied aspect of their metabolism (Adlercreutz *et al.* 1982, 1986b, 1988, 1991a). Urinary lignan response to consumption of linseed, which contains over 200-fold more of the ENL precursor secoisolariciresinol (SECO) (Mazur *et al.* 1996) than berries, has been investigated in some long-term studies in urine (Shultz *et al.*

**Abbreviations:** ID-GC–MS–SIM, isotope dilution gas chromatography–mass spectrometry in the selected-ion-monitoring mode; END, enterodiol; ENL, enterolactone; MAT, matairesinol; SECO, secoisolariciresinol.

\* **Corresponding author:** Professor Herman Adlercreutz, fax +358 9 61585633, email herman.adlercreutz@helsinki.fi




Fig. 1. Structures of plant and mammalian lignans.

1991; Lampe *et al.* 1994; Cunnane *et al.* 1995) and faeces (Kurzer *et al.* 1995). Our previous investigations on lignans (and isoflavonoids) in plasma (Adlercreutz *et al.* 1993, 1994) allowed us to assess normal concentrations of these phyto-oestrogens in Finnish omnivorous and vegetarian women who consumed their habitual diets. Morton *et al.* (1994) reported plasma levels of lignans (and isoflavonoids) for postmenopausal women consuming a traditional diet supplemented with linseed, soyabean flour or clover sprouts. Brzezinski *et al.* (1997), studying the effects of a phyto-oestrogen-rich diet on menopausal symptoms in women, examined basal and end-of-study levels of phyto-oestrogens in serum without measuring the dietary content of plant oestrogens. In none of these investigations have the basal and post-meal levels of lignans in blood and urine of human subjects been measured after a challenge with food with known lignan-precursor concentration. In the present investigation we studied the kinetics of ENL for 48 h after a strawberry meal in human subjects.

## Materials and methods

### Subjects

Healthy subjects ( $n = 7$ ) were recruited to the experiment based on an oral interview with the project leader. All but one man (H.A.) were employed by the Helsinki University Department of Clinical Chemistry and were technicians, or medical or chemistry students. None were taking any medication and none had had any antibiotics during 3 months preceding the sampling period. The subjects' ages ranged from 23 to 55 (mean 38.1) years and BMI from 18.8 to 26.0 (mean 23.3) kg/m<sup>2</sup>.

### Berry samples

The berries studied were: strawberry (*Fragaria × ananassa*), blackberry (*Rubus fruticosus*), raspberry (*Rubus idaeus*) and cloudberry (*Rubus chamaemorus*) of the family

*Rosaceae*; blackcurrant (*Ribes nigrum*), redcurrant (*Ribes rubrum*) of the family *Grossulariaceae*; and cranberry (*Vaccinium oxycoccos*), lingonberry (*Vaccinium vitis-idaea*), bilberry (*Vaccinium myrtillus*) and blueberry (*Vaccinium corymbosum*) of the family *Ericaceae*. The berries were purchased from commercial sources. Strawberries for consumption by the subjects of the study (5 kg) were brought from a farm in central Finland. These strawberries were divided into ten portions of 500 g (wet weight) and frozen at  $-20^{\circ}$  until consumed. Before phyto-oestrogen analysis, the berry samples were dried in the oven ( $45-50^{\circ}$ , 4–5 d) and pulverized manually with a mortar. DM was determined with a Moisture Analyzer 40 (Sartorius AG, Göttingen, Germany).

### Chemicals, standards and instruments

The berries were analysed using our ID-GC–MS in the selected-ion-monitoring mode (SIM) method for the determination of lignans and isoflavonoids in food samples. All chemicals and details of the synthesis of standards and the validity of the method have been presented (Mazur *et al.* 1996). Time-resolved fluoro-immunoassay of plasma and urine ENL was performed as described in detail by Adlercreutz *et al.* (1998a) using the following instruments, all purchased from Wallac, Turku, Finland: a Victor 1420 multilabel counter and Wallac 1420 Victor software version 1.0 for data analysis. Microtitration strips were washed and shaken with 1296-026 DELFIA platewasher and 1296-003 DELFIA plateshaker respectively. Radioactivity counting was performed with a LKB Model 1217 Rackbeta Liquid Scintillation Counter. White microtitration strips coated with goat anti-rabbit IgG were also products of Wallac.

### Study design

The study design was approved by the ethics committee of the Helsinki University Central Hospital. Detailed written dietary instructions and a list of forbidden food items were given to the participants at least 1 month before starting the experiment. The subjects were requested to exclude from their diet any food items containing phyto-oestrogens for 3 d (72 h) before the consumption of the berry meal and during the collection of plasma and urine samples (the following 48 h). The participants avoided most vegetables and fruits, legumes, seeds and nuts, berries, grains, whole-grain products, and tea, beer and wine. Consumption of certain wheat products (white bread, pasta), milk products (milk, yoghurt, cheese), meat and fish, a few vegetables (e.g. potato) and fruits (e.g. an apple, a pear) was allowed so that, in principle, the intake of vitamins and macronutrients was unchanged. The subjects began this restricted (zero) diet 3 d before the experiment (D1) and followed it throughout the experiment (D5). On day 3 (D3), starting after the first morning void, participants collected one 24 h urine sample ('zero' urine), including the first morning void on day 4 (D4). Daily urine was collected into 2 litre plastic bottles containing 2 g ascorbic acid and stored at  $+4^{\circ}$  until the collection was completed. Total urine volume was measured and portions were stored at  $-20^{\circ}$  until analysis. On D4 at

08.00 hours, a vascular catheter was introduced to a median cubital (or basilic) vein and the first blood sample ('zero' plasma, four 10 ml tubes) was taken. Thereafter subjects consumed 500 g strawberries. Following the strawberry meal the 24 h urine sample was collected in four portions: 4 h+4 h+4 h+12 h. Blood samples (4×10 ml) were taken through the catheter at the following intervals: 0.5 h, 1 h, 2 h, 4 h, 8 h after the berry meal. The last blood sample was taken on the morning of the next day (D5) (24 h after the meal). After centrifugation of the tubes, plasma (15–20 ml) was separated and stored at –20° until analysis. On D5 urine sample collection was continued in two portions (12 h+12 h) per d. Total urine collection obtained in four (D4) and two (D5) portions was measured and stored at –20° until analysis.

The subject K.L. collected D5 urine mistakenly in one portion, 25–48 h, instead of in two portions (25–36 h and 37–48 h).

#### Analytical methods

*Isotope dilution gas chromatography–mass spectrometry in the selected-ion-monitoring mode method for food samples.* The method for the determination of lignans, isoflavones and coumestrol in berry samples is described briefly here. Dried and powdered berry samples were hydrolysed with an enzyme mixture (*Helix pomatia*; Bio Septra Inc., Marlborough, MA, USA) and acid (HCl), extracted with diethyl ether and the extract subsequently was purified on DEAE- and QAE-Sephadex columns (Pharmacia Fine Chemicals, Uppsala, Sweden). Formononetin, daidzein, biochanin A, genistein (isoflavones) and coumestrol, as well as SECO and matairesinol (MAT) (lignans) were derivatized to form trimethylsilyl ethers and analysed by the ID-GC–MS–SIM method (Mazur *et al.* 1996) using synthesized deuterated internal standards for the correction of losses during the procedure. All assays were carried out in duplicate or quadruplicate.

*Time-resolved fluoroimmunoassay methods for enterodiol in plasma and urine.* The analytical procedure is briefly as follows (Adlercreutz *et al.* 1997, 1998a,b). After addition of the radioactive internal standard <sup>3</sup>H-oestradiol

glucuronide (only for plasma), hydrolysis reagent (0.2 U/ml β-glucuronidase and 2 U/ml sulfatase in 0.1 mol/l acetate buffer, pH 5.0) was added to the plasma or urine sample which were mixed and then incubated overnight at 37°. For plasma samples diethyl ether was used to extract free (unconjugated) ENL after the hydrolysis; for urine no extraction was used. Then the hydrolysed and extracted plasma or urine samples in assay buffer (Tris-HCl buffer containing 5 g BSA/l, pH 7.8) were pipetted into prewashed goat anti-rabbit IgG microstrips. Simultaneously anti-serum (dilution 1 : 250 000) and Eu-labelled ENL (dilution 1 : 400 000) in the assay buffer were added into the microstrips. After incubating and shaking the strips slowly on a DELFIA plate shaker (Wallac) at room temperature for 90 min, the strips were washed with a DELFIA plate washer (Wallac). Subsequently DELFIA enhancement solution (Wallac) was added to each well and the strips were shaken slowly for an additional 5 min. Fluorescence was read in the DELFIA Victor multilabel counter (Wallac). Results were calculated according to the formula:

$$\text{Plasma ENL} = \text{concentration (read)}$$

$$\times 1/\text{recovery} \times \text{dilution factor,}$$

$$\text{Urine ENL} = \text{concentration (read)} \times \text{dilution factor.}$$

#### Statistics

Values in the text were expressed as means with pooled standard errors. After one-way ANOVA, the data were assessed using Duncan's multiple range test. Differences were considered significant at  $P < 0.05$ . All the statistical analyses were performed with the SPSS statistical software package (Version 6.1J, SPSS, Chicago, IL, USA).

#### Results

The lignan concentrations in eight types of berries grown in Finland are shown in Table 1. None of the isoflavonoids in these samples was measurable with our ID-GC–MS–SIM technique, mainly due to their low concentrations and some interfering compounds at these very low (trace) levels. However, relatively high levels of the lignan SECO were

**Table 1.** Phyto-oestrogen content of berries (μg/kg DM)\*

Berry	Daidzein	Genistein	SECO†	Matairesinol
<i>Rosaceae</i>				
Blackberry ( <i>Rubus fruticosus</i> )	Tr	Tr	37180	225
Strawberry ( <i>Fragaria × ananassa</i> )	Tr	Tr	15046	781
Cloudberry ( <i>Rubus chamaemorus</i> )	0	0	2030	0
Raspberry ( <i>Rubus idaeus</i> )	0	Tr	1390	0
<i>Ericaceae</i>				
Lingonberry ( <i>Vaccinium vitis-idaea</i> )	0	0	15100	0
Cranberry ( <i>Vaccinium oxycoccos</i> )	0	0	10540	0
Blueberry ( <i>Vaccinium corymbosum</i> )	0	0	8350	0
<i>Grossulariaceae</i>				
Blackcurrant ( <i>Ribes nigrum</i> )	0	0	3880	95
Redcurrant ( <i>Ribes rubrum</i> )	0	0	1653	0

SECO, secoisolariciresinol; Tr, trace (present in trace amounts; over 50–60% of detection limit).

\* Analysis by isotope dilution gas chromatography–mass spectrometry in the selected-ion-monitoring mode.

† Total amount of anhydrosecoisolariciresinol and SECO.

**Table 2.** Enterolactone in plasma (nmol/l) and urine (nmol/l) before and after consumption of the strawberry meal  
(Values for individual subjects and mean values with pooled standard errors)

Time of plasma collection (h after strawberry meal)	Urine collection Time (h)	Subject	Subject							Mean	SEM
			A.S.	I.W.	W.M.	T.N.	H.A.	R.T.	K.L.		
0		Plasma	10.6	8.3	1.7	11.0	4.7	13.5	22.4	10.3*	2.5
	Day 3; 24	Urine	686.8	849.3	303.6	608.9	2413.1	2470.3	3128.4	1494.3	429.2
0.5		Plasma	7.9	11.7	1.1	12.1	4.2	11.3	21.0	9.9*	2.4
	Day 4; 0–4	Urine	697.8	782.7	99.8	523.0	440.3	1447.2	2324.6	902.2	283.6
1		Plasma	6.9	11.3	1.5	11.5	4.0	11.7	24.1	10.1*	2.8
	Day 4; 5–8	Urine	719.3	659.0	586.5	941.8	251.6	786.7	2293.6	891.2	247.2
2		Plasma	8.8	9.1	1.5	10.4	4.2	10.3	19.8	9.2*	2.2
	Day 4; 9–12	Urine	776.5	1000.4	587.3	458.5	1049.3	1351.7	7071.6	1756.5	893.1
4		Plasma	13.5	11.3	0.9	10.4	3.2	8.6	17.4	9.3*	2.2
	Day 4; 13–24	Urine	627.2	2742.2	816.0	1438.9	1889.7	4363.6	1237.3	1873.6	493.3
8		Plasma	14.1	13.4	4.0	15.5	3.3	9.2	25.9	12.2	2.9
	Day 5; 1–12	Urine	673.0	3973.0	698.6	1339.5	771.0	2296.8	1246.6	1571.2	454.2
24		Plasma	16.5	50.0	2.4	25.9	8.1	10.8	30.2	20.6	6.2
	Day 5; 13–24	Urine	1230.2	1797.3	562.6	1238.9	1716.7	1984.6	Nd	1421.7	212.4

Nd, not determined (the sample was not collected).

Mean values were significantly different from that of the sample taken 24 h after the strawberry meal: \*  $P < 0.05$ .


found in all the berries, ranging from 1.39 mg/kg (raspberry) to 37.2 mg/kg (blackberry) (3.9 and 104.3  $\mu\text{mol/kg}$  DM respectively). Blackberry contained the highest concentrations of SECO and total plant lignans. Strawberry (the most common Finnish variety) contained 15.1 mg/kg DM, (42.4  $\mu\text{mol/kg}$ ). MAT, the immediate precursor of ENL, was found only in three berries (strawberry, blackberry and blackcurrant) at low concentrations. The strawberries used in the study (500 g, 77.5 g DM) contained 3.3  $\mu\text{mol}$  (11.7 mg on a DM basis) of SECO and 0.17  $\mu\text{mol}$  (0.61 mg on a DM basis) MAT.

The results for ENL concentrations in plasma and urine of each subject are shown in Table 2. Mean plasma ENL values before and during the experiment are displayed in Fig. 2.

In the seven subjects studied, ENL plasma concentrations recorded in the basal plasma samples taken before consumption of strawberries ranged from 1.7 to 22.4 nmol/l

(mean 10.3 (SEM 2.5) nmol/l). At the four next measurements, 0.5, 1, 2 and 4 h after the berry meal, plasma ENL levels slightly decreased, ranging from 1.1 to 21.0 nmol/l (mean 9.9 (SEM 2.4) nmol/l), 1.5 to 24.1 nmol/l (mean 10.1 (SEM 2.8) nmol/l), 1.5 to 19.8 nmol/l (mean 9.2 (SEM 2.2) nmol/l) and 0.9 to 17.4 nmol/l (mean 9.3 (SEM 2.2) nmol/l) respectively. The analysis of plasma ENL in two subsequent plasma samples taken on D4 and D5 indicated an increase (statistically significant for values at 24 h; 8 h: 4.0–25.9 nmol/l, mean 12.2 (SEM 2.9) nmol/l; 24 h: 2.4–30.2 nmol/l, mean 20.6 (SEM 6.2) nmol/l). ENL concentrations recorded in the subject K.L.'s plasma (all collections but at 24 h) were higher than those of the other subjects.

The urinary excretion of ENL was calculated for 4 h collections (nmol/4 h). A 24 h urine collection on D3, corrected for 4 h period, yielded results for the subjects ranging from 70.8 nmol/4 h (subject W.M.) to 1053.2 nmol/4 h (subject K.L.) (mean 517.5 (SEM 132.3) nmol/4 h). ENL



**Fig. 2.** Enterolactone concentrations in plasma of the subjects ( $n=7$ ) after consumption of a strawberry meal. Values are means with standard errors indicated by vertical bars. Mean values were significantly different from that of the sample taken 24 h after the strawberry meal: \*  $P < 0.05$ .

**Table 3.** Individual urinary excretion (nmol/24 h) and recovery (%) of enterolactone (ENL)

Subject	Time of urine collection			Total ENL excreted post-challenge	Recovery (%)
	Day 3; 24 h (pre-challenge)	Day 4; 1–24 h	Day 5; 25–48 h		
A.S.	1339	1844	1817	2322	56.4
I.W.	2820	6600	14265	18045	438.0
W.M.	425	920	785	1280	31.1
T.N.	1796	2558	2745	3507	85.1
H.A.	4344	1664	1732	-948	-23.0
R.T.	4694	4839	4792	4947	119.8
K.L.	6319	6558	3491	3730	90.5
Mean					114.0

excretion measured in the urine collected in portions within the four next 4 h periods (D4: 0–4 h, 5–8 h, 9–12 h and 13–24 h) after the strawberry meal varied individually between the subjects (range 83.8–906.6 nmol/4 h, 105.6–1032.9 nmol/4 h, 114.5–1909.4 nmol/4 h and 205.4–903.2 nmol/4 h respectively); however, there was a tendency for an increase (mean 551.9 (SEM 132.2) nmol/4 h, 434.1 (SEM 103.1) nmol/4 h, 643.3 (SEM 233.8) nmol/4 h, and 646.6 (SEM 189.9) nmol/4 h respectively). In the next collection, 25–36 h, mean ENL excretion increased and was the highest recorded during the experimental period (range 139.7–3125.5 nmol/4 h, mean 926.3 (SEM 390.5) nmol/4 h). ENL excretion decreased in the urine collected at 37–48 h in all the subjects except for K.L. (the sample not collected) and A.S. (range 121.9–1629.6, mean 543.0 (SEM 227.7) nmol/4 h).

Urinary ENL excretion per d (nmol/24 h) differed among the subjects (Table 3). The recovery of plant lignans SECO and MAT determined as urinary ENL varied from -23 to 438% (mean 114%) for the subjects. Two subjects, however, showed very different yields: H.A. excreted 23% less ENL in urine than calculated by the intake, whilst I.W.'s ENL recovery in urine was almost 5-fold higher (438.0%) than judged by the amount of plant lignans ingested.

When the ENL values in plasma with respect to sampling periods were analysed statistically, the 24 h mean ENL concentration was significantly higher than the mean concentrations at 0, 0.5, 1, 2, 4 h ( $P < 0.05$ ). With respect to the ENL values in urine, no significances were found between the different periods due to high interindividual variations.

## Discussion

The quantitative results for isoflavones in berries, showing very low content of these phenols in berries, are consistent with our earlier findings in various fruits and vegetables (Mazur, 1998; Mazur & Adlercreutz, 1998) as well as with earlier reports by others (Price & Fenwick, 1985; Jones *et al.* 1989; Dewick, 1993; Reinli & Block, 1996). With regard to lignans, our present study is the first describing plant lignan SECO and MAT levels in berries. The results for the three berry families *Rosaceae*, *Ericaceae* and *Grossulariaceae* varied. Blackberry and strawberry of the *Rosacea* family contained the highest amounts of lignans whereas in another

member of this family, the raspberry, the lowest amount of SECO was detected. When compared with other edible plant foods analysed until now (Mazur, 1998; Mazur & Adlercreutz, 1998), berries are an important source of the lignans in our diet.

The present pilot study was primarily designed to evaluate whether and how rapidly a single meal, relatively rich in lignans, would affect plasma levels and urinary excretion in subjects consuming a phyto-oestrogen-free diet. In order to follow lignan metabolic changes in the biological fluids immediately after a berry meal we collected short-time urine portions and took blood at short intervals within the first 48 h post-challenge. We chose to determine plasma and urinary ENL because ENL is the chief mammalian lignan in human subjects and we have developed specific, sensitive and economical time-resolved fluoroimmunoassays for it. Strawberry was selected for the experimental meal due to its popularity in the Finnish diet, and its relatively high lignan content.

The results of our present study led to certain observations on lignan metabolism in human subjects. The first observation is that a single strawberry meal, demonstrated to contain relatively high amounts of plant lignan SECO, contributed to elevated plasma ENL levels and increased the urinary excretion of this mammalian lignan (statistically significant in 24 h post-meal plasma). The metabolic response, however, was different in different subjects. Maximal mean ENL value measured in the 24 h plasma post-challenge was accompanied by the highest mean urinary ENL in D5 25–36 h urine of the subjects. From the data it is obvious that it takes about 8 h for the lignan precursors to reach the colon where conversion to ENL takes place, reaching its maximum about 25–36 h after the meal.

The second comment concerns the high individual variation of ENL values in plasma and urine, before and during the experiment. Although all the subjects ingested a fixed dose of the lignans, there were substantial variations between the subjects in the post-challenge plasma levels and urinary excretion of the mammalian ENL. The between-individual variability of ENL levels in plasma and urine, before and during the experiment, is probably due to differences in individual lignan metabolism and absorption depending on the gastrointestinal microflora (Adlercreutz, 1998). The gut microflora produces enzymes catalysing

metabolic conversion of SECO, the major lignan in strawberry, to END through dehydroxylation and then demethylation (Borriello *et al.* 1985). END is then oxidized by facultative bacteria to yield ENL. ENL also can be synthesized directly from MAT, however, strawberry (and other berries) contains minimal amounts of this lignan. The conversion of plant lignans to mammalian lignans depends on the presence of facultative bacteria in the intestine, as evidenced by studies with rats and human subjects (Axelson & Setchell, 1981; Setchell *et al.* 1982; Pettersson *et al.* 1996).

There is a large body of evidence that our diet influences our gastrointestinal microflora (Rowland *et al.* 1985). The Finnish (except for W.M.) participants of the study have been consuming a lignan-rich diet for many years. Their everyday diet consists predominantly of cereals and grain (e.g. rye bread), milk and yogurt products, vegetables, seeds, and berries (all these are sources of lignans). This dietary pattern must therefore affect their microflora and influence mammalian lignan production from plant precursors (Adlercreutz & Mazur, 1997; Adlercreutz, 1998; Mazur & Adlercreutz, 1998). Daily intake of these foods is a natural stimulation of the production of mammalian lignans; the fibre increases the bacterial mass. Interestingly the subject K.L., a vegetarian, does not consume any meat and meat products at all; she had the highest ENL concentrations in pre- and particularly post-meal plasma and urine which could reflect her high metabolic response to the berry meal. High individual variation in phyto-oestrogen excretion in response to isoflavonoid- or lignan-rich plant food consumption has been reported by other investigators (Adlercreutz *et al.* 1991b; Lampe *et al.* 1994; Morton *et al.* 1994; Kelly *et al.* 1995; Karr *et al.* 1997; Watanabe *et al.* 1998). The subjects in these studies, however, had been treated with a much greater dose of phyto-oestrogens for a longer period of time.

The urinary recovery of ENL differed only slightly from amounts estimated by its precursors in the strawberry meal. Considering lignan metabolism and ENL recovery one should bear in mind that substantial amounts of lignans are excreted by the faecal route and the amounts are similar or only slightly lower in faeces than in urine (Bannwart *et al.* 1987) (e.g. faecal excretion of lignans in nine omnivorous women: ENL 1510 nmol/24 h, END 147.7 nmol/24 h and MAT 22.3 nmol/24 h (Adlercreutz *et al.* 1995)). This would mean that the total recovery of lignans in urine and faeces is probably higher than the intake of lignans from the strawberries. Furthermore, the time-resolved fluoroimmunoassay for urinary ENL gives about 30% higher values than if assayed by GC-MS (H Adlercreutz and M Uehara, unpublished results), probably due to the presence of other lignan metabolites cross-reacting with the antiserum. As microflora is an indispensable factor involved in the constant process of mammalian lignan production, the bacteria seem to be loaded with lignans and could themselves supply 5–10% ENL recovered in urine (basal plasma and urinary ENL levels, even after a long period of restricted diet, did not reach 0 value). Xu *et al.* (1995) in their study on bioavailability of soyabean isoflavones concluded that the breakdown of isoflavones by the microflora in the gut determines the recovery of the compounds.

In summary, berries contain different, relatively high amounts of the lignan SECO, but not isoflavones, and therefore contribute to elevated plasma levels and increased urinary excretion of mammalian lignan ENL. A supplement of 500 g lignan-rich strawberries increased plasma ENL concentrations 24 h post-challenge (observed as higher urinary excretion in D5 25–36 h collection) (statistically significant for mean values in plasma). Our pilot study showed clearly high individual variability of metabolic response to a single dose of lignan-rich berries.

### Acknowledgements

The authors would like to thank Adile Samaletdin and Ritva Takkinen for skillful technical assistance. One author (W. M.) thanks the Graduate School in Steroid Research for a scholarship. The analytical work was supported by NIH grant no. 2 R01 CA 56289-04.

### References

- Adlercreutz H (1998) Evolution, nutrition, intestinal microflora, and prevention of cancer: a hypothesis. *Proceedings of the Society for Experimental Biology and Medicine* **217**, 241–246.
- Adlercreutz H, Fotsis T, Bannwart C, Hämäläinen E, Bloigu S & Ollus A (1986a) Urinary estrogen profile determination in young Finnish vegetarian and omnivorous women. *Journal of Steroid Biochemistry* **24**, 289–296.
- Adlercreutz H, Fotsis T, Bannwart C, Wähälä K, Brunow G & Hase T (1991a) Isotope dilution gas chromatographic–mass spectrometric method for the determination of lignans and isoflavonoids in human urine, including identification of genistein. *Clinica Chimica Acta* **199**, 263–278.
- Adlercreutz H, Fotsis T, Bannwart C, Wähälä K, Mäkelä T, Brunow G & Hase T (1986b) Determination of urinary lignans and phytoestrogen metabolites, potential antiestrogens and anticarcinogens, in urine of women on various habitual diets. *Journal of Steroid Biochemistry* **25**, 791–797.
- Adlercreutz H, Fotsis T, Heikkinen R, Dwyer JT, Woods M, Goldin BR & Gorbach SL (1982) Excretion of the lignans enterolactone and enterodiol and of equol in omnivorous and vegetarian women and in women with breast cancer. *Lancet* **2**, 1295–1299.
- Adlercreutz H, Fotsis T, Kurzer MS, Wähälä K, Mäkelä T & Hase T (1995) Isotope dilution gas chromatographic mass spectrometric method for the determination of unconjugated lignans and isoflavonoids in human feces, with preliminary results in omnivorous and vegetarian women. *Analytical Biochemistry* **225**, 101–108.
- Adlercreutz H, Fotsis T, Lampe J, Wähälä K, Mäkelä T, Brunow G & Hase T (1993) Quantitative determination of lignans and isoflavonoids in plasma of omnivorous and vegetarian women by isotope dilution gas chromatography–mass spectrometry. *Scandinavian Journal of Clinical and Laboratory Investigation* **53**, 5–18.
- Adlercreutz H, Fotsis T, Watanabe S, Lampe J, Wähälä K, Mäkelä T & Hase T (1994) Determination of lignans and isoflavonoids in plasma by isotope dilution gas chromatography–mass spectrometry. *Cancer Detection and Prevention* **18**, 259–271.
- Adlercreutz H, Höckerstedt K, Bannwart C, Bloigu S, Hämäläinen E, Fotsis T & Ollus A (1987) Effect of dietary components, including lignans and phytoestrogens, on enterohepatic circulation and liver metabolism of estrogens, and on sex hormone binding globulin (SHBG). *Journal of Steroid Biochemistry* **27**, 1135–1144.

- Adlercreutz H, Höckerstedt K, Bannwart C, Hämäläinen E, Fotsis T & Bloigu S (1988) Association between dietary fiber, urinary excretion of lignans and isoflavonic phytoestrogens, and plasma non-protein bound sex hormones in relation to breast cancer. In *Progress in Cancer Research and Therapy*, vol. 35: *Hormones and Cancer 3*, pp. 409–412 [F Bresciani, RJB King, ME Lippman and J-P Raynaud, editors] New York, NY: Raven Press.
- Adlercreutz H, Honjo H, Higashi A, Fotsis T, Hämäläinen E, Hasegawa T & Okada H (1991b) Urinary excretion of lignans and isoflavonoid phytoestrogens in Japanese men and women consuming a traditional Japanese diet. *American Journal of Clinical Nutrition* **54**, 1093–1100.
- Adlercreutz H, Lapcik O, Hampl R, Wähälä K, Al-Maharik N, Wang G-J & Mikola H (1997) Immunoassay of phytoestrogens in human plasma. *Symposium on Phytoestrogen Research Methods, Tucson, Arizona, USA*. Abstr.
- Adlercreutz H & Mazur W (1997) Phyto-oestrogens and western diseases. *Annals of Medicine* **29**, 95–120.
- Adlercreutz H, Wang G-J, Lapcik O, Hampl R, Wähälä K, Mäkelä T, Lusa K, Talme M & Mikola H (1998a) Time-resolved fluoroimmunoassay for plasma enterolactone. *Analytical Biochemistry* **265**, 208–215.
- Adlercreutz H, Wang G-J, Uehara M, Lapcik O, Al-Maharik N, Mäkelä T, Mikola H, Hampl R & Wähälä K (1998b) Immunoassays of phytoestrogens in human plasma. *The Cost 916 Workshop on Phytoestrogens: Exposure, Bioavailability, Health Benefits and Safety Concerns. April 17–18, 1998, Doorwerth, The Netherlands*, pp. 23–28 [S Bausch-Goldbolm, S Kardinaal and F Serra, editors]. European Communities.
- Axelsson M & Setchell KDR (1981) The excretion of lignans in rats – evidence for an intestinal bacterial source for this new group of compounds. *FEBS Letters* **123**, 337–342.
- Bannwart C, Adlercreutz H, Wähälä K, Brunow G & Hase T (1987) Identification of the isoflavonic phytoestrogens formononetin and dihydrodaidzein in human urine. In *International Symposium on Applied Mass Spectrometry in the Health Sciences. Abstracts*, p. 169. Barcelona: Fira de Barcelona, Palau de Congressos.
- Borriello SP, Setchell KDR, Axelsson M & Lawson AM (1985) Production and metabolism of lignans by the human faecal flora. *Journal of Applied Bacteriology* **58**, 37–43.
- Brzezinski A, Adlercreutz H, Shaoul R, Rosler A, Shmueli A, Tanos V & Schenker JG (1997) Short-term effects of phytoestrogen-rich diet on postmenopausal women. *Menopause – The Journal of the North American Menopause Society* **4**, 89–94.
- Cunnane SC, Hamadeh MJ, Liede AC, Thompson LU, Wolever TM & Jenkins DJ (1995) Nutritional attributes of traditional flaxseed in healthy young adults. *American Journal of Clinical Nutrition* **61**, 62–68.
- Dewick PM (1993) Isoflavonoids. In *The Flavonoids: Advances in Research Since 1986*, pp. 117–217 [JB Harborne, editor]. London: Chapman & Hall.
- Jones AE, Price KR & Fenwick GR (1989) Development of high-performance liquid chromatographic method for the analysis of phytoestrogens. *Journal of the Science of Food and Agriculture* **46**, 357–364.
- Karr SC, Lampe JW, Hutchins AM & Slavin JL (1997) Urinary isoflavonoid excretion in humans is dose dependent at low to moderate levels of soy-protein consumption. *American Journal of Clinical Nutrition* **66**, 46–51.
- Kelly GE, Joannou GE, Reeder AY, Nelson C & Waring MA (1995) The variable metabolic response to dietary isoflavones in humans. *Proceedings of the Society for Experimental Biology and Medicine* **208**, 40–43.
- Kurzer MS, Lampe JW, Martini MC & Adlercreutz H (1995) Fecal lignan and isoflavonoid excretion in premenopausal women consuming flaxseed powder. *Cancer Epidemiology, Biomarkers and Prevention* **4**, 353–358.
- Lampe JW, Martini MC, Kurzer MS, Adlercreutz H & Slavin JL (1994) Urinary lignan and isoflavonoid excretion in premenopausal women consuming flaxseed powder. *American Journal of Clinical Nutrition* **60**, 122–128.
- Landström M, Zhang J-X, Hallmans G, Åman P, Bergh A, Damber JE, Mazur W, Wähälä K & Adlercreutz H (1998) Inhibitory effects of soy and rye diets on the development of Dunning R3327 prostate adenocarcinoma in rats. *Prostate* **36**, 151–161.
- Mazur WM (1998) Phytoestrogens in food. In *Baillière's Clinical Endocrinology and Metabolism: Phyto-oestrogens*, vol. 12, no. 4, pp. 729–742 [H Adlercreutz, editor]. London: Baillière Tindall.
- Mazur WM & Adlercreutz H (1998) Naturally occurring oestrogens in food. *Journal of Pure and Applied Chemistry* **70**, 1759–1776.
- Mazur WM, Fotsis T, Wähälä K, Ojala S, Salakka A & Adlercreutz H (1996) Isotope dilution gas chromatographic–mass spectrometric method for the determination of isoflavonoids, coumestrol, and lignans in food samples. *Analytical Biochemistry* **233**, 169–180.
- Morton MS, Wilcox G, Wahlqvist ML & Griffiths K (1994) Determination of lignans and isoflavonoids in human female plasma following dietary supplementation. *Journal of Endocrinology* **142**, 251–259.
- Petersson D, Åman P, Knudsen KEB, Lundin E, Zhang JX, Hallmans G, Harkonen H & Adlercreutz H (1996) Intake of rye bread by ileostomists increases ileal excretion of fiber polysaccharide components and organic acids but does not increase plasma or urine lignans and isoflavonoids. *Journal of Nutrition* **126**, 1594–1600.
- Price KR & Fenwick GR (1985) Naturally occurring oestrogens in foods – A review. *Food Additives and Contaminants* **2**, 73–106.
- Reinli K & Block G (1996) Phytoestrogen content of foods – a compendium of literature values. *Nutrition and Cancer* **26**, 123–148.
- Rowland IR, Mallett AK & Wise A (1985) The effect of diet on the mammalian gut flora and its metabolic activities. *Critical Reviews in Toxicology* **16**, 31–103.
- Setchell KDR, Lawson AM, Borriello SP, Adlercreutz H & Axelsson M (1982) Formation of lignans by intestinal microflora. In *Colonic Carcinogenesis: Falk Symposium 31*, pp. 93–97 [RA Malt and RCN Williamson, editors]. Lancaster: MTP Press.
- Shultz TD, Bonorden WR & Seaman WR (1991) Effect of short-term flaxseed consumption on lignan and sex hormone metabolism in men. *Nutrition Research* **11**, 1089–1100.
- Watanabe S, Yamaguchi M, Sobue T, Takahashi T, Miura T, Arai Y, Mazur W, Wähälä K & Adlercreutz H (1998) Pharmacokinetics of soybean isoflavones in plasma, urine and feces of men after ingestion of 60g baked soybean powder (kinako). *Journal of Nutrition* **128**, 1710–1715.
- Xu X, Harris KS, Wang HJ, Murphy PA & Hendrich S (1995) Bioavailability of soybean isoflavones depends upon gut microflora in women. *Journal of Nutrition* **125**, 2307–2315.