

The effect of duration of manual restraint during blood sampling on plasma cortisol levels in calves

G Stilwell G^{*†}, C de Carvalho R[‡], MS Lima[†] and DM Broom[§]

[†] Centro de Investigação Interdisciplinar em Sanidade Animal, Departamento de Clínica, Faculdade de Medicina Veterinária, UTL. Av. Universidade Técnica, Alto da Ajuda, 1300-477 Lisbon, Portugal

[‡] Clínica VetMondego, 3140 Montemor-o-velho, Portugal

[§] Centre for Animal Welfare and Anthrozoology, Department of Veterinary Medicine, University of Cambridge, Madingley Road, Cambridge CB3 0ES, UK

* Contact for correspondence and requests for reprints: stilwell@fmv.utl.pt

Abstract

Many studies on stress and pain rely, solely or mainly, on plasma cortisol assessment. Confounding factors, such as handling, may cause a release of cortisol making the interpretation of the results difficult. We looked at the influence of duration of restraint on the plasma cortisol levels of one-to-two month old calves. Forty-three calves were divided into four groups according to the interval between restraint and blood sampling: i) Group 0, immediate blood-sampling; ii) Group 0.5M, half a minute restraint; iii) Group 1M, one minute restraint and iv) Group 2M, two minutes restraint. The only increase in plasma cortisol, compared with all the other groups, was seen with blood sampling after two minutes of restraint. This study provides evidence to suggest that cortisol released as a result of handling stress is not evident if blood sampling is carried out within one minute of restraining calves.

Keywords: animal welfare, calves, cortisol, handling, restrain, stress

Introduction

Various hormones (eg ACTH, glucocorticoids, catecholamines, prolactin, etc) play a role in the stress response of animals (for a review see Matteri *et al* 2000). It is well established that glucocorticoid production, following the activation of the hypothalamic-pituitary-adrenal (HPA) axis, is part of an emergency response (Broom & Zanella 2004) intended to defend the organism against stressful conditions (Möstl & Palme 2002). It is for these reasons that cortisol is medically termed 'the stress hormone'.

The concentration of cortisol in blood is widely used as an indicator of stress, although caution in the interpretation of results is advised because an increase does not occur with every type of stressor (Broom & Johnson 2000) and because a wide variety of stressors can activate the HPA system (Molony & Kent 1997; Broom & Johnson 2000). In farm animals, examples of these are: weaning (Hickey *et al* 2003); social isolation (review by Cockram 2004); transport (Crookshank *et al* 1979; Grigor *et al* 2004); social mixing (Arthington *et al* 2003); novelty (van Reenen *et al* 2005); restraint and handling (Ewbank *et al* 1992) and multiple venipuncture (Hopster *et al* 1999). See also review by Lane (2006).

Sample collection, which often involves confinement and handling of animals, may be, in itself, stressful and may

confound the results of studies (Cook *et al* 2000). Some authors have used in-dwelling catheters applied some time before the study begins but this implies regular flushing of catheters and may cause discomfort or infection. Also, this method does not preclude all handling. Non-invasive measurements of cortisol (milk, saliva, urine and faeces) may reduce or prevent this disadvantage but only provide information on overall levels and, thus, is more useful for studies on chronic stress. For the main disadvantages of non-invasive measurements of cortisol, see Lane (2006). So, to validate blood cortisol as an indicator of stress or pain caused by a particular procedure, all redundant effects should be eliminated (Cook *et al* 2000).

Calf blood cortisol is useful for the evaluation of stress and pain after routine farm procedures, such as disbudding, tail-docking and castration (Molony *et al* 1995; Morisse *et al* 1995; McMeekan *et al* 1998; Faulkner *et al* 2000; Schreiner & Ruegg 2002; Sutherland *et al* 2002; Stafford *et al* 2003; Stilwell & Lima 2004, 2008). Although all studies use control groups that are not subjected to the procedures, the effect of duration between restraint and blood sampling on blood cortisol levels of young calves should be taken on account. This study was designed, therefore, to measure the effect of time between first restraint and blood sampling, on plasma cortisol levels of young calves.

Table 1 Mean (\pm SD) blood cortisol levels of calves restrained for differing periods of time.

Time to sampling (Group)	n	Age (days)	Cortisol (nmol l ⁻¹)
Immediate	10	47 \pm 11	11.71 \pm 7.97 ^a
0.5M	9	46 \pm 13	18.39 \pm 13.85 ^a
1M	10	54 \pm 10	16.34 \pm 14.33 ^a
2M	13	49 \pm 11	40.12 \pm 31.10 ^b

Different superscripts indicate significant differences ($P < 0.05$).

Materials and methods

The study site

This study was carried out at a large cattle rearing unit which receives between 100 and 200 young calves each month from dairy farms. The great majority are Holstein-Friesian, but some crossbreeds are seen (Holstein \times Limousin and Holstein \times Belgian Blue). Transport distances from farms of origin range from 2 to 200 km and on arrival all calves are put in individual boxes and receive an electrolyte solution.

Animals are fed twice a day with a commercial milk-replacer in individual buckets. Water and concentrate are available all day. New straw is added every three-to-four days, but bed material is only completely removed when the calf is weaned and moved to group paddocks. Calves have close contact with herdspeople at feeding, adding of bedding and during twice daily individual visual monitoring. Weaning occurs at approximately two months of age.

Study animals

Forty-three male Holstein-Friesian calves were included in the study. They were housed in the same building along four rows and, although age varied between 31 and 67 days (Table 1), all calves were milk fed.

The animals underwent systematic allocation to different groups. Starting at one end of the first row every four calves were distributed to the following four groups, according to the time between entering the individual pen and blood sampling: i) Group 0, immediate blood sampling; ii) Group 0.5M, 0.5 min restraint; iii) Group 1M, one minute restraint and iv) Group 2M, two minute restraint. Restraint was carried out by squeezing the calf gently against the pen wall with a knee while holding the head with one hand. This was done by an experienced veterinary surgeon and no excessive force was needed with any of the animals. A second person, five metres from the pen, measured the time and advised when venipuncture and blood sampling should be done. One calf that should have been included in Group 0.5M was excluded because of signs of illness. The last three calves of the last row were included in Group 2M. There were no age differences between groups.

Blood samples (7 ml) were taken into a heparinised tube by left jugular venipuncture. Blood was immediately

centrifuged and frozen (-20°C). Cortisol was assayed in duplicate and measured by a validated solid radioimmunoassay, without extraction, using commercial kits (Coat-A-Count®, Diagnostic Product Corporation, Los Angeles, CA, USA). The inter-assay coefficient of variation for cortisol was 5.5% for the level of $1\mu\text{g dl}^{-1}$ and 1.9% for the level of $5\mu\text{g dl}^{-1}$.

Statistical analysis

The distributions and variance of the data were shown not to be normally distributed by Levene and Shapiro-Wilks tests. Significant differences between the four groups were then determined by the Mann Whitney *U*-test following a Kruskal-Wallis, one-way analysis of variance. Computer software SPSS version 14.0 was used for the analysis.

Results

The results (Table 1) showed a significant difference between Group 2M and Group 0 ($P = 0.002$), Group 0.5M ($P = 0.03$) and Group 1M ($P = 0.021$). Individual variation in blood cortisol levels was very large within each group but especially in the 2M group. Within each group we also compared cortisol levels of animals that were younger than the mean age with those that were older than the mean age and found no differences (data not shown in table).

Discussion and conclusions

The question, ‘how long after an animal has been stressed by handling and restraint will the cortisol response be evident?’, has not been answered for young calves used for studies on pain associated with disbudding, dehorning, tail docking and castration.

Hopster *et al* (1999) found that initial collection within one minute of restraint did not alter baseline cortisol in dairy cows, but repeated venipuncture at 15 min intervals caused an increase in cortisol in primiparous cows less accustomed to handling. Our study used young dairy calves that had been accustomed, since birth, to human proximity and contact. Although restraint was easy and the animals did not show any evidence of distress, we did show that handling alone does cause a significant cortisol response, even in very young calves that were used to human contact. However, we also showed that cortisol levels are not affected if blood sampling is done immediately after restraint (up to one minute of restraint, at least). This suggests that when studies on distress and pain in calves are carried out, non-treated control groups may give reliable information on baseline plasma cortisol levels, providing that blood sampling is carried out by an experienced operator and takes place within one minute of first handling and restraint.

Acknowledgements

We would like to thank Manuel Barata for providing animals for the study and Professor Dr Luísa Mateus at the Faculdade de Medicina Veterinária for cortisol measuring and valuable advice. Finally, we would like to thank Marta Vacas de Carvalho for advice regarding the statistical analysis.

References

- Arthington JD, Eicher SD, Kunkle WE and Martin FG** 2003 Effect of transportation and co-mingling on the acute-phase protein response, growth, and feed intake of newly weaned beef calves. *Journal of Animal Science* 81: 1120-1125
- Broom DM and Johnson KG** 2000 *Assessing Welfare: Short term Responses* pp 87-110. Kluwer Academic Publishers: Dordrecht, The Netherlands
- Broom DM and Zanella AJ** 2004 Brain measures which tell us about animal welfare. *Animal Welfare* 135: 41-45
- Cockram MS** 2004 A review of behavioural and physiological responses of sheep to stressors to identify potential behavioural signs of distress. *Animal Welfare* 13: 283-291
- Cook CJ, Mellor DJ, Harris PJ, Ingram JR and Matthews LR** 2000 Hands-on and hands-off measurement of stress. In: Moberg GP and Mench JA (eds) *The Biology of Animal Stress* pp 123-146. CABI Publishing: Wallingford, UK
- Crookshank HR, Elissalde MH, White RG, Clanton DC and Smalley HE** 1979 Effect of transportation and handling of calves upon blood serum composition. *Journal Animal Science* 48: 430-435
- Ewbank R, Parker MJ and Mason CW** 1992 Reactions of cattle to head-restraint at stunning: a practical dilemma. *Animal Welfare* 1: 55-64
- Faulkner PM and Weary DM** 2000 Reducing pain after dehorning in dairy calves. *Journal of Dairy Science* 83: 2037-2041
- Grigor PN, Cockram MS, Steele WB, McIntyre J, Williams CL, Leushuisb IE and van Reenen CG** 2004 A comparison of the welfare and meat quality of veal calves slaughtered on the farm with those subjected to transportation and lairage. *Livestock Production Science* 91: 219-228
- Hopster H, Joop TN, van der Werf JT, Erkens JH and Blokhuis HJ** 1999 Effects of repeated jugular puncture on plasma cortisol concentrations in loose-housed dairy cows. *Journal of Animal Science* 77: 708-714
- Hickey MC, Drennan M and Earley B** 2003 The effect of abrupt weaning of suckler calves on the plasma concentrations of cortisol, catecholamines, leukocytes, acute-phase proteins and *in vitro* interferon-gamma production. *Journal of Animal Science* 81: 2847-2855
- Lane J** 2006 Can non-invasive glucocorticoid measures be used as reliable indicators of stress in animals? *Animal Welfare* 15: 331-342
- Matteri RL, Carroll JA and Dyer CJ** 2000 Neuroendocrine responses to stress. In: Moberg GP and Mench JA (eds) *The Biology of Animal Stress* pp 43-76. CABI Publishing: Wallingford, UK
- McMeekan CM, Stafford KJ, Mellor DJ, Bruce RA, Ward RN and Gregory NG** 1998 Effects of regional analgesia and/or a non-steroidal anti-inflammatory analgesic on the acute cortisol response to dehorning in calves. *Research in Veterinary Science* 64: 147-150
- Molony V, Kent JE and Robertson IS** 1995 Assessment of acute and chronic pain after different methods of castration of calves. *Applied Animal Behaviour Science* 46: 33-48
- Molony V and Kent JE** 1997 Assessment of acute pain in farm animals using behavioural and physiological measurements. *Journal of Dairy Science* 75: 266-272
- Morisse JP, Cotte JP and Huonnic D** 1995 Effect of dehorning on behaviour and plasma cortisol responses in young calves. *Applied Animal Behaviour Science* 43: 239-247
- Möstl E and Palme R** 2002 Hormones as indicators of stress. *Domestic Animal Endocrinology* 23: 67-74
- Schreiner DA and Ruegg PL** 2002 Responses to tail docking in calves and heifers. *Journal of Dairy Science* 85: 3287-3296
- Stafford KJ, Mellor DJ, Todd SE, Ward RN and McMeekan CM** 2003 The effect of different combinations of lignocaine, ketoprofen, xylazine and tolazoline on the acute cortisol response to dehorning in calves. *New Zealand Veterinary Journal* 51(5): 219-226
- Stilwell G, Lima MS, Nunes T and Capitão E** 2004 Effect of three different methods of dehorning on plasma cortisol levels and behaviour of calves. *Proceedings of the XXIII World Buiatrics Congress*. 11-16 July 2004. Quebec, Canada
- Stilwell G, Lima MS and Broom DM** 2008 The effects of two non-steroidal anti-inflammatory drugs and of epidural injection on plasma cortisol and behaviour of calves castrated using a Burdizzo clamp. *American Journal of Veterinary Research* 69(6): 744-750
- Sutherland MA, Mellor DJ, Stafford KJ and Gregory NG** 2002 Cortisol responses to dehorning of calves given a 5-h local anaesthetic regimen plus phenylbutazone, ketoprofen, or adrenocorticotropic hormone prior to dehorning. *Research in Veterinary Science* 73: 115-123
- van Reenen CG, O'Connell NE, van der Werf JTN, Korte SM, Hopster H, Jones RB and Blokhuis HJ** 2005 Responses of calves to acute stress: individual consistency and relations between behavioural and physiological measures. *Physiology & Behaviour* 85: 557-570