

Erratum

Effect of midazolam, dexamethasone and their combination on the prevention of nausea and vomiting following strabismus repair in children

W. Riad*, R. Altaf†, A. Abdulla†, H. Oudan‡

*King Khaled Eye Specialist Hospital, Department of Anaesthesia; King Saud University, Faculty of Medicine, Departments of †Anesthesia, ‡Ophthalmology, Riyadh, Saudi Arabia

doi: 10.1017/S0265021507000166

The Publishers and Editors would like to apologise to the authors and readers for the inadvertent transposition of data columns two and three in Table 1 in the article by Riad *et al.* Effect of midazolam, dexamethasone and their combination on

the prevention of nausea and vomiting following strabismus repair in children. *European Journal of Anaesthesiology* 2007; 24: 697–701. The correct version of Table 1 is shown below.

Table 1. Patient characteristics and clinical data.

	Placebo	Dexamethasone	Midazolam	Midazolam + Dexamethasone
Sex (M/F)	14/11	15/10	12/13	11/14
Age (yr)	6.7 (2.9)	7.2 (2.4)	8.3 (3.6)	8.3 (3.9)
Weight (kg)	23 (13)	24 (15)	27 (12)	28 (11)
No. of muscles				
1	4	4	10	11
2	19	20	15	9
3 or 4	2	1	0	5
Anaesthesia time (min)	60 (18)	58 (23)	46 (13)	57 (26)
Recovery time (min)	15 (2.1)	17 (1.7)	24 (1.9)*	23 (2.5)*
Number of patients with oculocardiac reflex requiring atropine (%)	3	2	2	0

* $P < 0.05$.

Data expressed as a mean (SD) or number of patients.