

306

Intraoral Spectroscopy for the Identification and Study of Molar Hypomineralization

Sarah Boyer¹, Ray Jurado², Ashlee Cosantino², Azza Ahmed³ and Derk Joester⁴

¹Northwestern University; ²Lurie Children's Hospital; ³University of Illinois Chicago School of Dentistry and ⁴Northwestern University, Material Science and Engineering

OBJECTIVES/GOALS: Molar hypomineralization (MH) is a highly prevalent dental disease that leads to rapid enamel decay even with preventative measures. While the harm is apparent, the etiology of the condition is not. Further, MH is difficult to study due to limited intact extracted teeth. Therefore, we need a method to study MH non-destructively and intraorally. **METHODS/STUDY POPULATION:** Recent work has shown excess proteins not commonly found in enamel are present in teeth with MH. This is theorized to be due to disruption and infiltration in the cells that form enamel leading to leftover protein and under mineralized enamel. We hypothesize these proteins have specific spectroscopic signatures that can be detected using light. We further hypothesize that modern fiber optic probes can provide a method for non-destructive, pain-free, and rapid intraoral examination of MH. Initially, Micro Raman spectroscopy was used to detect specific vibrational bands associated with organics in teeth with MH followed by the collection of spectra of teeth with MH and healthy enamel using a spectrophotometer. These spectra were examined for any obvious differences. **RESULTS/ANTICIPATED RESULTS:** Currently 12 teeth were collected, and micro-computed tomography reconstructions confirmed location in 3D of MH lesions. Micro Raman of a MH-affected tooth revealed clear organic associated Amide I and III vibrational bands when compared to a synthetic hydroxyapatite (mineral in enamel) powder standard. We determined the wavelength of light that can be used to detect spectral differences between healthy enamel and teeth with MH. The next steps include optimization of the protocol of the intraoral spectrometer with the determined wavelength for implementation in clinic to allow for collection of spectra without the need for tooth extraction. **DISCUSSION/SIGNIFICANCE:** We hope that this work will lead to advancements in our understanding of the mechanism of MH as well as act as a proof of concept for a MH diagnostic tool. In the long term, the goal is to ideally lead to improvements in dental health care, decrease dental costs, and improve overall quality of life for the children with this condition.

307

Revealing Candidate Inherited Retinal Disease Candidates via Genome-wide Screening of Knockout Mice

Benjamin Yang, Andy Shao, Mustafa Jundi, Michael Shea, Cameron Ylagan and Ala Moshiri
University of California, Davis

OBJECTIVES/GOALS: The goal of this study is to reveal strong candidate genes for inherited retinal diseases (IRDs) in humans to better understand the mechanisms behind IRD development and reveal potential therapeutic targets. We hope these findings will help improve our understanding of IRDs and, subsequently, diagnostic

accuracy and prognosis for IRD patients. **METHODS/STUDY POPULATION:** The goal of the International Mouse Phenotyping Consortium is to identify the function of all protein-coding genes in the mouse genome via generation and phenotypic characterization of single gene knockout (KO) mice. Using this database, we identified all KO strains associated with abnormal retinal phenotypes, removed all RNA coding genes and pseudogenes, converted to human orthologues, and conducted a literature search for existing research regarding candidate IRD genes and retinal function/abnormalities. A similar process was used for RetNet genes. Subsequently, we performed bioinformatics analysis, including functional annotation (e.g. panther-db), pathway analysis (e.g. KEGG), and string-db to visualize known and predicted protein-protein interactions between the two data sets. **RESULTS/ANTICIPATED RESULTS:** Analysis of the IMPC database revealed, out of 8481 phenotyped genes, 572 unique protein coding genes were associated with 14 categories of retinal abnormalities such as abnormal retinal vasculature and abnormal retinal thickness, 377 of which have never been associated with retinal pathology in humans or mice. Pathway analysis of the IMPC database highlighted a general metabolism pathway as well as PI3K-Akt and MAPK pathways, not found in RetNet pathway results. Unique clusters from functional annotation clustering of the IMPC include DNA methylation and protein ubiquitination. Visualization of protein-protein interactions in string-db between the IMPC (mouse) and RetNet (human) revealed 4 clusters of interest with gold standard RetNet IRD proteins interacting with candidate IMPC IRD proteins. **DISCUSSION/SIGNIFICANCE:** IMPC analysis revealed 572 candidate IRD genes, 377 of which are novel with no existing independent research related to the retina outside of the IMPC. Bioinformatic analysis reveals 4 strong clusters of interest through string-db where a gold standard RetNet gene interacts with candidate IRD genes as well as many functional pathways of interest.

308

Targeting Cdk8 to improve Ischemic Fracture Healing

Christina Capobianco, Michelle Song, Jeanna Schmanski, Alexis Donneys, Karen Kessell, Tristan Maerz and Kurt Hankenson
University of Michigan

OBJECTIVES/GOALS: There are 1.78 million bone fractures globally each year, and 46% of fractures that have accompanying vascular damage (ischemia) will not heal without surgical intervention. Using single cell RNAseq we identified Cdk8 as a gene upregulated under ischemic fracture. Our work seeks to inhibit Cdk8 to assess its potential as a therapeutic target. **METHODS/STUDY POPULATION:** Most bone injuries heal through a cartilage intermediate that requires mesenchymal progenitor cells (hMSCs) to become cartilage forming chondrocytes. hMSCs underwent pelleted 3D chondrogenic differentiation in the presence of Cdk8 inhibitor, Senexin B. Chondrogenic gene expression was assessed via gene analysis of Aggrecan, Collagen II, and Collagen X. Content of sulfated glycosaminoglycans (sGAGs) was quantified through DMMB analysis. With IACUC approval, C57Bl/6 WT mice underwent femoral artery isolation and resection to create an ischemic environment prior to a transverse tibia fracture. Mice underwent