

reaction, PCR) of resistance genes using primers for *bla*TEM, *bla*CTX-M, and *bla*SHV. PCR-based replicon typing (PBRT) was conducted to type resistance plasmids carrying ESBL genes. **Results:** During the study period, 30 *E. coli* isolates with reduced susceptibility to third-generation cephalosporins (ie, ceftazidime, cefotaxime, and ceftriaxone) were detected in hospitalized patients. These isolates stemmed from blood culture (66.7%), wound swabs (13.3%), urine (13.3%), and drainage content (6.7%). Alongside complete resistance to  $\beta$ -lactam antimicrobial agents, they were also characterized by high resistance to gentamicin (93.3%) and ciprofloxacin (96.7%), whereas 23.3% of isolates were also resistant to ertapenem. Most isolates harbored both *bla*CTX-M and *bla*TEM genes concurrently (46.7%), while solitary *bla*TEM and *bla*CTX-M genes were found in 33.3% and 20% of these isolates, respectively. The presence of SHV  $\beta$ -lactamases was not found in any of the isolates. PBRT revealed a wide array of diverse plasmid groups, with most of the isolates harboring different combinations; however, 80% of isolates were characterized by plasmid incompatibility group B/O (IncB/O). **Conclusions:** We detected increased frequency of both TEM and CTX-M type  $\beta$ -lactamases in *E. coli* isolates from a single-hospital setting, with significant consequences for further treatment approaches. The high prevalence of broad- and extended-spectrum  $\beta$ -lactamase producers tends to prompt an increased carbapenem use (potentially resulting in increased resistance to carbapenems); thus, this type of analytical work should become a standard approach (where possible) in hospital centers in our country and worldwide.

**Funding:** No

**Disclosures:** None

*Antimicrobial Stewardship & Healthcare Epidemiology* 2021;1(Suppl. S1):s71–s72

doi:10.1017/ash.2021.139

#### **Presentation Type:**

Poster Presentation

**Subject Category:** MRSA/VRE

#### **Hospital Onset Vancomycin-Resistant Enterococci Infections in a Northern California Tertiary-Care Center From 2019 to 2020**

Stephanie Rasmussen and Sarah Waldman

**Background:** Understanding the epidemiology and risk factors for nosocomial infections with vancomycin-resistant enterococci (VRE) is necessary for the prevention and control of VRE infections in the hospital setting. We sought to determine the incidence of nosocomial infections of VRE and to ascertain predictors associated with nosocomial infection. **Methods:** In this retrospective cohort study, data were collected from patients with VRE infection from January 2019 to December 2020 at a tertiary-care center in northern California. VRE infections were designated as hospital onset (HO) if the specimen was collected >3 days after hospital admission or community onset (CO) if the specimen was collected  $\leq$ 3 days after admission. Associations between HO infections with time, unit, and specimen were identified. **Results:** Over the 2-year period, 214 unique VRE infections were identified in hospitalized patients; 115 infections were CO and 99 were HO. HO-VRE were associated most frequently with stay in medical/telemetry units (68%), followed by oncology–transplant units (15%) and ICUs (12%). There were  $\sim$ 4.7 and  $\sim$ 3.6 HO-VRE infections per month in 2019 and 2020, respectively. No differences were identified between HO-VRE infections in 2 medical units regarding glycopeptide and cephalosporin use in those units. The sources of VRE infections were urinary 45%, bloodstream 15%, stool 10%, and other 30%. Of the 45 infections in urine, 51% were identified from catheters (Foley and straight) and 27% were identified from clean-catch urine. Interestingly, 71% of patients with VRE identified from urine did not report urinary tract infection (UTI) symptoms at the time of collection. Urine was most often collected for urinalysis and culture from patients with nonspecific symptoms such as fever, leukocytosis, hypotension, tachycardia, or altered mental status. All urine collected from patients who reported UTI symptoms grew >100,000 CFU/mL in culture, while only 75% of cultures from patients without symptoms grew >100,000 CFU/mL. The most common antibiogram was resistance to

ampicillin, cefazolin, levofloxacin, minocycline, penicillin, tetracycline, and/or vancomycin (42% of cases) and susceptibility to both daptomycin and linezolid (60% of cases). **Conclusions:** HO-VRE infections were frequently identified with urinary sources and were often associated with catheter use. However, the frequent lack of concurrent UTI symptoms suggests VRE colonization rather than infection in many cases. Understanding the epidemiology and risk factors for HO-VRE infections is essential for developing infection prevention protocols to reduce the incidence of those infections.

**Funding:** No

**Disclosures:** None

*Antimicrobial Stewardship & Healthcare Epidemiology* 2021;1(Suppl. S1):s72

doi:10.1017/ash.2021.140

#### **Presentation Type:**

Poster Presentation

**Subject Category:** Occupational Health

#### **Experience with Bloodborne Pathogen Exposures Among Operation Room Nurses in Korea: A Qualitative Study**

JaHyun Kang and Hyunsook Lee

**Background:** Operation room (OR) nurses are at a high risk for bloodborne pathogen (BBP) exposures because they are constantly in contact with blood and body fluid (BBF) through surgical sites, and they frequently use sharp surgical instruments and sutures during surgeries. We explored occupational experiences with BBP exposures among OR nurses in South Korea. **Methods:** With an institutional review board's approval, this qualitative research was performed with 12 OR nurses who had worked for >3 months and had experienced BBP exposures. Based on the main research question "How is the experience of BBP exposure among OR nurses?" the semistructured questions for in-depth interview were prepared. Narrative data were collected through 1-hour individual interviews from June to September 2020 and were analyzed using a thematic analysis method. **Results:** The average age of the participants was 34.4 years with an average 9.75 years of clinical experience. The main theme extracted was "The nurses are alerted to their safety after experiencing the aftereffects and emotional trauma from BBP exposures," with 4 subthemes and 14 concepts. The first subtheme, "OR nurses risking exposure to BBF," included (1) hurried doctors and hasty nurses; (2) sharp surgical instruments everywhere; (3) deprioritized self-protection due to ongoing surgery; (4) inattentive to BBF risk; and (5) uncomfortable, foggy goggles and retracted safety devices. The second subtheme, "BBP exposures occurred in a flash," included (1) sharp injury occurred in a split second; (2) temporarily treat sharp wounds while hiding frightened feelings; and (3) OR nurses concentrated on surgery by suppressing anxiety and sharp wounds. The third subtheme, "Burdened time that could be overwhelmed alone," included (1) BBP exposures moments that I wish to reverse; (2) anger over dangerous environments and the turmoil of anxiety; (3) exhausted body and facial discoloration due to taking post-exposure prophylaxis; and (4) exposure to BBP that I want to hide from family and friends. The fourth subtheme, "Voice for everyone's safety," included (1) establishing a safety culture, which requires everyone's efforts, and (2) necessity of practical resources for decreasing BBP exposures. **Conclusions:** The participating OR nurses felt that they were working with a high risk to BBF exposures and called for institutional interventions to reduce the risks, including surgeons' attentive collaboration. Korean hospitals should make greater efforts to establish safety culture in ORs and to provide repeated, tailored education to prevent BBF exposures. They should also supply high-quality protective equipment and safety-engineered devices for OR nurses.

**Funding:** No

**Disclosures:** None

*Antimicrobial Stewardship & Healthcare Epidemiology* 2021;1(Suppl. S1):s72

doi:10.1017/ash.2021.141