

Original Article

Cite this article: Wright SL *et al* (2024). EMDR v. other psychological therapies for PTSD: a systematic review and individual participant data meta-analysis. *Psychological Medicine* 54, 1580–1588. <https://doi.org/10.1017/S0033291723003446>

Received: 17 April 2023

Revised: 25 October 2023

Accepted: 1 November 2023

First published online: 4 January 2024

Keywords:

EMDR; individual participant data meta-analysis; moderators; PTSD; systematic review; treatment

Corresponding author:

Simonne Lesley Wright;
Email: s.l.wright@vu.nl

EMDR v. other psychological therapies for PTSD: a systematic review and individual participant data meta-analysis

Simonne Lesley Wright^{1,2} , Eirini Karyotaki² , Pim Cuijpers² ,
Jonathan Bisson³ , Davide Papola⁴ , Anke Witteveen² ,
Sharain Suliman¹ , Georgina Spies¹ , Khodabakhsh Ahmadi⁵ ,
Liuva Capezzani^{6,7}, Sara Carletto⁸ , Thanos Karatzias⁹ , Claire Kullack¹⁰,
Jonathan Laugharne¹¹ , Christopher William Lee¹² , Mirjam J. Nijdam^{13,14} ,
Miranda Olff^{13,14} , Luca Ostacoli⁸ , Soraya Seedat^{1,15} 
and Marit Sijbrandij² 

¹South Africa PTSD Research Programme of Excellence, Department of Psychiatry, Faculty of Medicine & Health Sciences, Stellenbosch University, Cape Town, South Africa; ²Department of Clinical, Neuro- and Developmental Psychology, World Health Organization Collaborating Center for Research and Dissemination of Psychological Interventions, Amsterdam Public Health Research Institute, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands; ³Division of Psychological Medicine and Clinical Neurosciences, Cardiff University, Cardiff, UK; ⁴Department of Neuroscience, Biomedicine, and Movement Sciences, Section of Psychiatry, WHO Collaborating Centre for Research and Training in Mental Health and Service Evaluation, University of Verona, Verona, Italy; ⁵Behavioral Sciences Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran; ⁶The International Institute for Psychoanalytic Research and Training of Health Professionals (IIPRTHP), Rome, Italy; ⁷International School for Psychotherapy (SIPSI), Rome, Italy; ⁸Department of Clinical and Biological Sciences, University of Turin, Turin, Italy; ⁹School of Health & Social Care, Edinburgh Napier University, Edinburgh, Scotland; ¹⁰Pax Centre, West Leederville, Australia; ¹¹Faculty of Health and Medical Sciences, University of Western Australia, Perth, Australia; ¹²Faculty of Health and Medical Sciences, The University of Western Australia, Crawley, Australia; ¹³Department of Psychiatry & Amsterdam Public Health, Amsterdam University Medical Center location University of Amsterdam, Amsterdam, The Netherlands; ¹⁴ARQ National Psychotrauma Center, Diemen, The Netherlands and ¹⁵South African Medical Research Council Unit on the Genomics of Brain Disorders, Department of Psychiatry, Stellenbosch University, Cape Town, South Africa

Abstract

Background. This systematic review and individual participant data meta-analysis (IPDMA) examined the overall effectiveness of eye movement desensitization and reprocessing (EMDR) in reducing posttraumatic stress disorder (PTSD) symptoms, achieving response and remission, and reducing treatment dropout among adults with PTSD compared to other psychological treatments. Additionally, we examined available participant-level moderators of the efficacy of EMDR.

Methods. This study included randomized controlled trials. Eligible studies were identified by a systematic search in PubMed, Embase, PsycINFO, PTSDpubs, and CENTRAL. The target population was adults with above-threshold baseline PTSD symptoms. Trials were eligible if at least 70% of study participants had been diagnosed with PTSD using a structured clinical interview. Primary outcomes included PTSD symptom severity, treatment response, and PTSD remission. Treatment dropout was a secondary outcome. The systematic search retrieved 15 eligible randomized controlled trials (RCTs); 8 of these 15 were able to be included in this IPDMA (346 patients). Comparator treatments included relaxation therapy, emotional freedom technique, trauma-focused cognitive behavioral psychotherapies, and REM-desensitization.

Results. One-stage IPDMA found no significant difference between EMDR and other psychological treatments in reducing PTSD symptom severity ($\beta = -0.24$), achieving response ($\beta = 0.86$), attaining remission ($\beta = 1.05$), or reducing treatment dropout rates ($\beta = -0.25$). Moderator analyses found unemployed participants receiving EMDR had higher PTSD symptom severity at the post-test, and males were more likely to drop out of EMDR treatment than females.

Conclusion. The current study found no significant difference between EMDR and other psychological treatments. We found some indication of the moderating effects of gender and employment status.

distressing mental images (exposure to the traumatic event), the patient's emotional arousal is interrupted by employing another stimulus (bilateral stimulation and interruption of attention) which is assumed to lead to a reduction in arousal and distress (Jeffries & Davis, 2013; Rothbaum, Astin, & Marsteller, 2005).

Since EMDR's introduction, several mechanistic hypotheses have been proposed to explain the effects of bilateral stimulation in EMDR (Landin-Romero, Moreno-Alcazar, Pagani, & Amann, 2018). One of them is the adaptive information processing (AIP) model (Shapiro & Lialiotis, 2011). This suggests that EMDR involves a re-setting of the system that processes and stores events during stressful situations, which reduces distress and negative emotions triggered by traumatic experiences (Shapiro, 2001, 2006). Another theory to explain the effects of bilateral stimulation is the working memory theory, which proposes that by taxing the working memory, eye movements permanently reduce the vividness and emotionality of aversive memories (van den Hout & Engelhard, 2012). This theory has been supported by laboratory studies in healthy individuals (Mertens, Lund, & Engelhard, 2021). Finally, the orientating response model suggests that bilateral stimulation triggers an investigatory reflex, reducing negative emotions and enhancing awareness, facilitating exploratory behavior, and potentially improving cognitive processes (Barrowcliff, Gray, Freeman, & MacCulloch, 2004). However, a recent meta-analysis that included dismantling studies comparing EMDR with and without eye movements, found no benefit of eye movements, casting doubt about the superiority of EMDR to trauma-focused treatments without eye movements, such as exposure therapy or cognitive behavioral therapy with a trauma-focus (CBT-TF; Cuijpers, Veen, Sijbrandij, Yoder, & Cristea, 2020).

Most international guidelines for the treatment of PTSD recommend the use of either CBT-TF or EMDR as first-line treatments for adults with PTSD (Hamblen et al., 2019; Phelps et al., 2022; VA/DoD Clinical Practice Guideline, 2023). These guidelines are based on evidence-based research and provide recommendations to optimize patient treatment. Past meta-analyses (MA) have found EMDR to significantly improve PTSD symptom severity at post-test assessment (Bisson & Olff, 2021; Cuijpers et al., 2020; Cusack et al., 2016; Lewis, Roberts, Andrew, Starling, & Bisson, 2020a). Consistent with other disorders (Papola et al., 2022), large effect sizes have been reported when comparing EMDR for PTSD to wait-list-control (WLC) groups, and smaller effects when compared to treatment-as-usual groups and other active treatment groups. While the effectiveness of EMDR against WLC has been established, it is unclear what the additive benefit of the eye movements are. Some researchers argue that the eye movements are unnecessary, while others argue that they have an added advantage (van den Hout & Engelhard, 2012).

Currently, very little is known about moderators and predictors of EMDR treatment outcomes. Age, gender, baseline severity of PTSD, depression, and anxiety were not significantly associated with PTSD symptoms after EMDR treatment (Capezzani et al., 2013). Similarly, gender did not significantly influence treatment effects in a later study (Ter Heid, Mooren, van de Schoot, de Jongh, & Keber, 2016). In the same study, participants who did not have refugee status had a greater reduction in PTSD symptoms compared to those with refugee status (Ter Heid et al., 2016). There is literature suggesting that veterans with PTSD respond less to trauma-focused treatments in general, and to

EMDR specifically (Haagen, Ter Heide, Mooren, Knipscheer, & Kleber, 2017).

However, the methodological quality of the studies may play a role in these comparisons. There is also a great deal of inconsistency in the literature concerning the influence of specific moderators and predictors of psychotherapy outcomes in PTSD in general. No significant associations with treatment outcomes have been found for factors such as age (Ivarsson et al., 2014; Karatzias et al., 2007; Lewis et al., 2017), gender (Blanchard et al., 2003; Galovski, Blain, Mott, Elwood, & Houle, 2012; Haagen et al., 2017; Karatzias et al., 2007; Lewis et al., 2017), marital status (Karatzias et al., 2007), employment status (Ivarsson et al., 2014; Karatzias et al., 2007), therapy type (Karatzias et al., 2007), time since trauma (Ehlers et al., 2003; Karatzias et al., 2007), type of trauma (Karatzias et al., 2007), and psychiatric comorbidity (Cloitre, Koenen, Cohen, & Han, 2002; Rizvi, Vogt, & Resick, 2009).

Some individual psychotherapy studies have found that higher education (Lewis et al., 2017), higher levels of guilt (Rizvi et al., 2009), and therapeutic alliance (Cloitre et al., 2002) were associated with a better PTSD treatment response. Additionally, there is some evidence suggesting that comorbid psychiatric disorders reduce the beneficial effects of treatment on PTSD outcomes (Hagenaars, van Minnen, & Hoogduin, 2010; van Minnen, Wessel, Dijkstra, & Roelofs, 2002).

A long-standing issue in the field is that randomized controlled trials (RCT) and study-level (also known as aggregate or traditional) meta-analysis (MA) often lack sufficient statistical power to identify significant moderators of treatment effect (Gurung, Ellard, Mistry, Patel, & Underwood, 2015). This may be the reason for the gap in the current literature when it comes to moderators and predictors of EMDR. An individual participant data meta-analysis (IPDMA) synthesizes raw participant-level data from multiple related studies to answer a specific set of research questions. This can be done using a one-stage or two-stage approach (Riley, Tierney, & Stewart, 2021). The simpler and more utilized two-stage approach uses the participant-level data to calculate aggregate data in each trial separately, and then combines the aggregate data in a univariate MA model. This is similar to a study-level MA. On the other hand, the one-stage approach analyses participant-level data from all the trials in a single step using a generalized linear mixed model that accounts for the clustering of participants within trials. These two approaches tend to give very similar results when the same assumptions and estimation methods are used. However, when the number of trials included in the IPDMA is small then a one-stage approach is more exact (Riley et al., 2021). Using an IPDMA approach, we can maximize statistical power to detect more precise effects and explore participant-level characteristics as moderators of treatment outcomes.

By gaining insight into potential moderators and predictors of the effectiveness of EMDR, we may have better precision to identify patients who would benefit the most from EMDR. This is important since EMDR is highly protocolized, relatively straightforward to administer, and requires shorter episodes of imaginal trauma exposure (Nijdam, Gersons, Reitsma, de Jongh, & Olff, 2012; Schubert & Lee, 2009) in comparison to CBT-TF psychotherapies. The aims of this study were to (1) investigate the effectiveness of EMDR in reducing PTSD symptom severity, achieving treatment response, attaining PTSD remission, and reducing treatment dropout rates in comparison to another psychological treatment among adults with PTSD, and (2) explore

potential sociodemographic, clinical, and intervention-related moderators of EMDR treatment effects among adults with PTSD.

Methods

This IPDMA was reported in compliance with the Preferred Reporting Items for Systematic Review and Meta-Analyses (PRISMA) IPD Statement (Stewart *et al.*, 2015). We registered the study on PROSPERO (CRD42020138638). Additional information can also be found in the protocol paper (Wright *et al.*, 2022).

Eligibility criteria

Study inclusion was limited to RCTs comparing EMDR to active treatments. Active treatments included other psychological treatments (e.g. relaxation therapy, emotional freedom technique, and trauma-focused cognitive behavioral psychotherapies). We excluded studies comparing EMDR to WLC and treatment-as-usual groups. Studies were included if participants were adults (18 years or older) with above-threshold PTSD symptoms based on any established self-report scale or a clinical diagnosis of PTSD. Within each study, at least 70% of the participants were required to have a clinical diagnosis of PTSD according to any version of the DSM or ICD criteria. Only studies published in English were included.

Study identification and selection process

We used an existing database of psychological treatments for PTSD that was created by the Cardiff University Traumatic Stress Research Group (CUTSRG) to perform a systematic review and MA for the treatment guidelines of the International Society for Traumatic Stress Studies (ISTSS). A search was conducted by the Cochrane Collaboration Cochrane with the same inclusion criteria including all studies published until May 2018 (Bisson, Andrew, Roberts, Cooper, & Lewis, 2013; Bisson *et al.*, 2019; Lewis *et al.*, 2020a). We updated the search using the same search strategy (see online Appendix 1). The search strategy included the screening of major bibliographic databases such as PubMed, Embase, PsycINFO, PTSDpubs, and CENTRAL. We also screened past systematic reviews for additional articles. Our updated searches included studies published between the May 1, 2018 and the 11 January 2021 (see online Appendix 2).

Two researchers (SLW and DP) independently screened the titles and abstracts for the initial update (1 May 2018 till the 13 May 2019). Titles and abstracts for the second update (1 January 2019 till the 11 January 2021) were also screened by two researchers (SLW and ABW). The same researchers screened the full text of studies that possibly met the inclusion criteria. Senior members of the review team (MS and GS) resolved any uncertainties regarding study inclusion.

Data collection

Authors of eligible trials were contacted to request the use of their participant-level datasets. At least six additional reminder emails were sent at intervals of two to three weeks if no response was received. If a response had not been received at this point, an additional two authors were contacted (when possible). If after this no response was received, we attempted to reach out to the corresponding authors on ResearchGate and LinkedIn. If a response

was still not received, the dataset was considered unavailable along with those where the corresponding author declined to share data. In cases where the author was able to provide their participant-level dataset, data collection and storage were conducted in alignment with the European General Data Protection Regulation (Regulation (EU) 2016/679). Participant-level datasets were anonymized before sharing and were stored in an existing encrypted, password-protected folder at Vrije Universiteit Amsterdam. For data protection purposes, we sent the link for the encrypted folder to a different email address than the one we used for the password. Our data collection commenced at the end of 2019 as planned, however it was extended to two and a half years because an insufficient number of datasets had been collected by the end of 2020.

Primary and secondary outcomes

The primary outcome data included PTSD symptom severity, treatment response, and PTSD remission. The secondary outcome was treatment dropout. PTSD symptom severity scores were measured using PTSD measures and clinical interviews. Studies included in this IPDMA used self-report measures like the Impact of Event Scale (total score ranging from 0 to 75; Horowitz, Wilner, & Alvarez, 1979), Impact of Event Scale-Revised (IES-R; total score ranging from 0 to 88; Weiss & Marmar, 1997), PTSD Checklist (PCL; total score ranging from 0 to 80; Blanchard, Jones-Alexander, Buckley, & Forneris, 1996), and the Mississippi Scale for PTSD (MPTSD; total score ranging from 35 to 175; Keane, Malloy, & Fairbank, 1984). We used self-report measures because we wanted to use the same type of assessment across studies. All available studies provided self-report PTSD severity outcomes but not all studies provided PTSD severity outcome data based on clinical interviews. Treatment response was defined as a 50% reduction in baseline PTSD symptom scores (Karyotaki *et al.*, 2017). Participants were considered in PTSD remission if they no longer had a formal diagnosis of PTSD at the post-test. In line with the definition used by CUTSRG for ISTSS' treatment guidelines, we considered a participant who left the study before the post-test assessment as a treatment dropout (Lewis, Roberts, Gibson, & Bisson, 2020b). Assessment time points included baseline and post-test. In addition, we requested available baseline sociodemographic, clinical, and intervention-related variables in order to investigate their effect on treatment outcomes (when there was sufficient data to do so). Moderators included gender; age; relationship status; partner or no partner (married/cohabitating or divorced/widowed/separated/single); marital status (yes or no); tertiary education (yes or no); employed (yes or no); using psychotropic medication (yes/no); above threshold depression (yes/no); anxiety threshold (yes or no); chronic (duration of PTSD symptoms > 3 months; yes or no), and comorbid psychiatric disorder (yes or no).

Risk of bias assessment

Studies were assessed for risk of bias using Cochrane Risk of Bias 2 Tool (RoB 2; Higgins, Savović, Page, Elbers, & Sterne, 2022). Each study was assessed independently by two researchers (SW and SS) to determine whether there was a risk for bias related to bias arising from the following domains: randomization process (D1), deviations from intended intervention (D2), and measurement of the outcome (D4). We did not include domains 3 and 5 (missing outcome data and selection of the reported result respectively) because missing values were addressed in

our analyses, and bias related to the selection of reported outcomes because we had access to the participant-level datasets (Karyotaki et al., 2023). Each domain comprises signaling questions which lead to a domain-level judgement on the risk of bias. All uncertainties were cleared up through deliberation by SW and SS with the assistance of an additional member of the review team.

Data analysis

Data analysis was conducted in Stata 17 (StataCorp, 2021). We combined all individual data sets into a merged data set, using a generic standardized protocol for integrating participant-level datasets (Stewart et al., 2015). Study-level variables were included for both the available and unavailable participant-level datasets, which were extracted from the studies' available documentation, such as publications and trial registries.

Study-level MA

We conducted a study-level MA to examine the differences across the studies that provided participant-level data and the studies for which participant-level data were unavailable. The difference between the studies that did and did not provide data was compared in a subgroup analysis. Heterogeneity was examined by calculating I^2 indicating heterogeneity as a percentage with 25% as low, 50% as moderate, and 75% as high (Higgins & Thompson, 2002). The 95% confidence intervals (CI) around I^2 were calculated using the non-central chi-squared-based approach in the heterogi module of Stata (Ioannidis, Patsopoulos, & Evangelou, 2007; Orsini, Bottai, Higgins, & Buchan, 2005). We examined publication bias by visually inspecting the funnel plot, using the trim and fill procedure and Egger's test of funnel plot asymmetry (Duval & Tweedie, 2000; Egger, Davey Smith, Schneider, & Minder, 1997).

IPDMA

PTSD total scores were standardized by transforming each study's PTSD severity original (raw) scores to z scores before combining the individual datasets into the master dataset. Missing outcome data for PTSD symptom severity at post-test were estimated using multiple imputation under the missing-at-random assumption (miimputemvnl in STATA software, version 17; StataCorp, 2021). This method generated 20 imputed data sets using data on baseline PTSD symptom severity scores. These newly imputed datasets included the observed and the imputed standardized PTSD symptom scores for missing values. They were analyzed separately using the selected model, and the results were averaged according to Rubin's rules (Riley, Lambert, & Abo-Zaid, 2010).

One-stage IPDMA

In a one-stage IPDMA, we merged all participant-level data from all studies with participants clustered within studies. One-stage IPDMA yields more precise and less biased estimates of effect, maximizes the statistical power, and accounts for parameter correlation (Debray, Moons, Abo-Zaid, Koffijberg, & Riley, 2013; Stewart & Parmar, 1993). We calculated the standardized β coefficient for the examined comparisons. This estimate indicates how many S.D.s the dependent variable changes per S.D. increase in the predictor variable. Thus, the higher the β the greater the effect of the predictor variable on the dependent variable, although there is

no association among the variables if the β is 0. Using a one-stage approach, we analyzed the effect of the interventions on PTSD symptom severity at the end of treatment with a multilevel mixed-effects linear regression using a random intercepts model with a random effect for each trial and fixed effects for the intervention and symptom severity, using STATA's mixed command. Post-test PTSD scores were used as the dependent variable and trial arm condition (EMDR *v.* control) as the independent variable while controlling for baseline PTSD symptom severity. We analyzed the effect of the interventions on treatment response at post-test using a multilevel mixed-effects logistic regression (using a random intercepts model with a random effect for each trial and fixed effects for the intervention and PTSD symptom severity, using STATA's melogit command). Response (yes or no) was the dependent variable, and treatment condition was the independent variable. This was repeated for PTSD remission and dropout.

Two-stage IPDMA

The two-stage approach uses participant-level data to derive aggregate data (such as treatment effect estimates) in each trial separately, and then combines the aggregate data in a study-level MA model. We ran a two-stage IPDMA using STATA's ipdmetan command.

Moderator analyses

We tested whether available demographic and clinical characteristics (gender, age, relationship status, marital status, completed some form of tertiary education, employment status, chronic PTSD status, use of psychotropics, above threshold depression, and presence of comorbidity) moderated the effect of EMDR on PTSD outcomes. Not all included studies reported data on the selected moderators. We included moderator analyses when the variables were reported by three or more studies. To examine moderators, we added the interaction between each potential moderator and treatment outcome on PTSD severity into a multilevel mixed-effects linear regression model. We similarly added the interaction between each potential moderator and treatment response into a multilevel mixed-effects logistic regression model. Each potential moderator was included in a separate model as the main effect.

Sensitivity analysis

Sensitivity analyses were conducted to examine the effect of EMDR compared to CBT-TF. We also compared the effect of EMDR to other psychotherapies, while excluding CBT-TF.

Results

Study selection and participant-level data obtained

The systematic literature search resulted in 15 eligible articles. We were able to obtain participant-level data from eight studies, comprising 346 participants (Ahmadi, Hazrati, Ahmadzadeh, & Noohi, 2015; Capezzani et al., 2013; Carletto et al., 2016; Devilly & Spence, 1999; Karatzias et al., 2011; Laugharne et al., 2016; Lee, Gavriel, Drummond, Richards, & Greenwald, 2002; Nijdam et al., 2012) and these were included in the analyses reported herein. Seven eligible datasets were unavailable and could not be included in this IPDMA (Boterhoven-De Haan et al., 2020; Carlson, Chemtob, Rusnak, Hedlund, & Muraoka,

1998; Power *et al.*, 2002; Rothbaum *et al.*, 2005; Taylor *et al.*, 2003; Ter Heid *et al.*, 2016; Vaughan *et al.*, 1994). Of the unavailable studies, corresponding authors reported that two were lost, three did not respond to the study invitations, one indicated that consent issues precluded data sharing, and one study was still in progress. While the main results had been published (Boterhoven-De Haan *et al.*, 2020), the authors decided not to share their data because the additional papers were still being written up. See online Appendix 3 for a summary of study characteristics for the unavailable studies.

Study and participant characteristics

The eight studies evaluated EMDR against the following treatments: one relaxation therapy (Carletto *et al.*, 2016), one REM-Desensitization (Ahmadi *et al.*, 2015), one emotional freedom technique (Karatzias *et al.*, 2011), one Prolonged Exposure (CBT-TF [PE]) (Laugharne *et al.*, 2016), one brief eclectic psychotherapy (CBT-TF [BEP]) (Nijdam *et al.*, 2012), and three CBT-TF (Unspecified) (Capezzani *et al.*, 2013; Devilly & Spence, 1999; Lee *et al.*, 2002). Both the EMDR and comparator treatments had significant improvements in overall PTSD symptom severity at the post-test assessment in comparison to their group baseline scores.

All eight EMDR studies used the standard EMDR protocol (Shapiro, 1989). The included studies were conducted in the following countries: Iran (1), Netherlands (1), Australia (3), Italy (2), and Scotland (1). All the included EMDR interventions were conducted in person and in a one-on-one format (see online Appendix 4). Additionally, all participants had a diagnosis of PTSD at the baseline assessment. The mean (s.d.) age of participants was 38.61 (11.90) years. 204 (59.13%) of 345 were female, 125 (51.02%) of 245 were married or cohabitating, and 101 (41.22%) were single. 109 (53.17%) of 205 had completed some form of tertiary education, 133 (53.63%) of 248 had no comorbid diagnosis at the baseline assessment, 95 (41.67%) of 228 were stable on psychotropic medication, and 261 (95.96%) of 272 had chronic PTSD at the baseline assessment (see online Appendix 5). The mean (s.d.) baseline PTSD symptom score was 53.78 (10.63; 95% CI 50.76–56.80) on the Impact of Event Scale (Horowitz *et al.*, 1979), 68.88 (20.28; 95% CI 66.07–71.69) on the Impact of Event Scale-Revised (IES-R; Weiss & Marmar, 1997); 57.61 (10.76; 95% CI 54.96–60.25) on the PCL (Blanchard *et al.*, 1996), and 148.43 (11.23; 95% CI 143.31–153.54) on Keane's Post-Traumatic Stress Disorder Scale from the Minnesota Multiphasic Personality Inventory (MMPI-K; Keane *et al.*, 1984) in the respective studies. Finally, 94 (33.22%) out of the 283 participants dropped out of treatment before the post-test assessment.

Risk of bias assessment

Online Appendix 6 presents the RoB2 ratings for the studies included in this IPDMA. One of the eight studies scored *some concerns* on D1, risk of bias arising from the randomization process because the information about the randomization methods was limited to a statement that the study was randomized. Additionally, masking participants is difficult to achieve in psychotherapy research, which resulted in three studies being rated as having *some concerns*. None of the included studies were rated as being at a *high risk* of bias on any of the domains.

Results of study-level MA

Fifteen studies compared EMDR with another psychological treatment. The results of the study-level MA of all 15 included studies revealed no significant difference in PTSD symptom severity between EMDR and the comparator interventions at the post-test assessment $g = -0.091$, 95% CI -0.33 to 0.15 , $p = 0.462$. Heterogeneity was moderate, $I^2 = 55.79\%$. There was no significant difference between the outcome findings of studies included in the present IPDMA and studies with unavailable data, $p = 0.87$. See online Appendix 7. Based on a visual inspection of the funnel plot of standard error Hedges' g , it is unlikely that publication bias is present in this MA (see online Appendix 8).

Primary outcomes

IPDMA: PTSD symptom severity

Online Appendix 9 presents the main and moderator results of a one-stage IPDMA on PTSD symptom severity at the post-test. A one-stage IPDMA found no significant difference between EMDR and comparator interventions on PTSD symptom severity, $\beta = -0.24$, 95% CI -0.62 to 0.14 , $p = 0.210$, n (studies) = 270 (8) in the completer analysis. The full sample one-stage IPDMA analysis based on imputed PTSD severity outcome data, $\beta = -0.20$, 95% CI -0.52 to 0.12 , $p = 0.217$, n (studies) = 339 (8), and the two-stage yielded a similar result to the one-stage IPDMA completer analysis, $g = -0.20$, 95% CI -0.55 to 0.14 , $p = 0.251$.

Baseline PTSD symptom severity was found to be a significant predictor of post-test PTSD severity in the one-stage completer analysis ($\beta = 0.43$; $p = 0.000$), and the imputed full sample analysis, $\beta = 0.43$, $p = 0.000$. More specifically, higher baseline PTSD symptom severity was associated with higher post-test PTSD severity.

In the completer analysis only, employment status significantly moderated the relationship between therapy type and post-test PTSD symptom severity. More specifically, unemployed participants who received EMDR reported significantly higher PTSD symptom severity at the post-test than employed participants who received EMDR, $\beta = 0.80$, $p = 0.019$. None of the other participant-level variables (sociodemographic, clinical, and intervention-related characteristics) significantly moderated PTSD symptom severity after EMDR treatment in the completer or imputed analyses (see online Appendix 9).

IPDMA: treatment response

In one-stage analysis, no significant difference in the effect of EMDR compared with other psychological treatments was found for PTSD treatment response, $\beta = 0.86$, 95% CI -0.03 to 1.74 , $p = 0.057$, n (studies) = 270(8). See online Appendix 10. The OR was 2.36. The two-stage analysis MA also found no significant difference in effect between EMDR and other psychological treatments for PTSD treatment response, $\beta = 0.52$, 95% CI -0.42 to 1.46 , $p = 0.278$. The OR was 1.68.

Employment status significantly moderated the relationship between therapy type and PTSD treatment response. More specifically, unemployed participants who received EMDR were significantly less likely to have responded to treatment at post-test than the employed participants who received EMDR, $\beta = -0.63$, $p = 0.005$.

None of the other sociodemographic, clinical, and intervention-related characteristics of participants was significantly associated with treatment response (see online Appendix 11).

IPDMA: PTSD remission

In a one-stage IPDMA, no significant difference in effect between EMDR and other psychological treatments for PTSD remission at post-test were found, $\beta = 1.05$, 95% CI -0.11 to 2.22 , $p = 0.075$, n (studies) = 199(5). See online Appendix 11. The OR was 2.87. The two-stage analysis found a significant effect of EMDR compared with other psychological treatments for PTSD remission at post-test, $g = 1.00$, 95% CI 0.14 – 1.87 , $p = 0.023$. The OR was 2.73. There was insufficient data to run a moderator analysis.

Secondary outcome

IPDMA: treatment dropout

One-stage IPDMA on treatment dropout found no significant difference in effect between EMDR and other psychological treatments at post-test, $\beta = -0.25$, 95% CI -0.79 to 0.29 , $p = 0.369$, n (studies) = 283 (6). See online Appendix 12. The OR was 0.78. The two-stage analysis found no significant difference in the effect of EMDR over controls for PTSD treatment dropout at post-test, $\beta = -0.19$, 95% CI -0.83 to 0.45 , $p = 0.553$. The OR was 0.82.

Gender significantly moderated the relationship between therapy type and PTSD dropout. More specifically, male participants in EMDR groups were significantly more likely to drop out of EMDR treatment than female participants in EMDR groups, $\beta = 0.23$, $p = 0.028$. None of the other sociodemographic, clinical, and intervention-related characteristics of participants was significantly associated with treatment dropout (see online Appendix 12).

Sensitivity analysis

We re-ran the one-stage IPDMA, including only the CBT-TF comparison groups. We found no significant difference in effect between EMDR and CBT-TF comparison groups, $\beta = -0.18$, 95% CI -0.75 to 0.38 , $p = 0.525$, n (studies) = 180 (5).

We re-ran the one-stage IPDMA, excluding the CBT-TF comparison groups. We found no significant difference in effect between EMDR and the other comparison groups, $\beta = -0.29$, 95% CI -0.82 to 0.25 , $p = 0.294$, n (studies) = 90 (3).

Discussion

To the best of our knowledge, this is the first IPDMA to explore moderators of EMDR for adults with PTSD using individual participant-level data. One of the strengths of the present study was the statistical power to detect statistically significant moderators compared with study-level MA and published RCTs aimed at investigating the efficacy of EMDR for adults with PTSD. This IPDMA made it possible to investigate available participant-level moderators, such as employment status and gender.

In line with past research, the current study found no significant difference between EMDR and other psychological treatments on PTSD severity, treatment response, or treatment dropout in either the one- or two-stage IPDMA (Lewis et al., 2020a). Despite the proposed different mechanisms of action, no significant difference in efficacy was found between EMDR and CBT-TF comparison groups. When we removed the CBT-TF comparison groups from the analysis, there was still no significant difference. The current study did not provide support for the notion that psychotherapies with eye movements

are more effective in treating PTSD than those without eye movements.

It is important to note that all the psychological comparator treatment groups were found to be effective in treating PTSD when interpreting these findings. We found no significant difference between EMDR and the psychological treatment control groups on PTSD remission at post-test in the one-stage analysis. However, a significant main effect in favor of EMDR was found in the two-stage analysis on PTSD remission. Considering the small sample sizes of the included trials, the one-stage IPDMA result is most likely the more accurate reflection of the true effect.

In line with previous research, baseline PTSD was a significant predictor of post-test PTSD symptom severity (Taylor et al., 2003). Specifically, higher baseline PTSD symptom severity was associated with higher post-test PTSD symptom severity. An earlier study found that higher PTSD baseline scores on PTSD self-report measures were associated with better treatment outcomes on self-report PTSD questionnaires (Karatzias et al., 2007). Overall, this was not the case in this aggregated set of trials. While there are distinct differences between EMDR and the other TF therapies, our finding suggests they are equally efficacious at treating PTSD symptoms. These findings are in line with past study-level MA (Lewis et al., 2020a).

Our moderator analysis was exploratory in nature and based on available sociodemographic, clinical, and intervention-related variables available in the obtained databases. Results from our completer moderator analysis found unemployed participants who received EMDR reported significantly higher PTSD symptom severity at post-test than employed participants who received EMDR. Similarly, we found unemployed participants who received EMDR were significantly less likely to respond to treatment by post-test than the participants who received EMDR and were employed.

Past research supports our current findings. Unemployed participants were found to be more likely to suffer from higher levels of mental health problems including PTSD (Bosmans & van der Velden, 2018; McKee-Ryan, Song, Wanberg, & Kinicki, 2005). In a longitudinal study among employed and unemployed trauma-exposed participants, unemployed participants continued to experience higher levels of mental health problems even years after exposure (Bosmans & van der Velden, 2018). Research has attributed the benefits of employment as income, status, relationships, and esteem (Chen, Westman, & Hobfoll, 2015; Paul & Batinic, 2010). Unemployed participants might be more socially isolated in comparison to their employed counterparts or have more severe or further advanced symptoms (Nijdam, Vermetten, & McFarlane, 2023) resulting in less beneficial PTSD outcomes at post-test. Furthermore, it is possible that financial concerns and associated psychological distress could distract or hinder the recovery process in unemployed participants. Another explanation for poorer outcomes for unemployed participants may be that unemployed participants may more often be engaged in compensation seeking procedures that may cause additional distress. Indeed, a direct relationship was found between compensation-related distress and higher PTSD symptom severity (O'Donnell et al., 2015).

We also found that male participants who received EMDR were significantly more likely to drop out of treatment than female participants who received EMDR. Both brain and behavior differences in men and women may explain why men were found to be more likely to drop out of EMDR treatment in comparison to female participants (Olf, 2017). In a recent survey among

Australian males who attended mental health services, various reasons for drop-out were self-reported, among which was a lack of connection with the therapist, the sense that therapy lacked progress and the cost/inconvenience related with attending therapy sessions (Seidler et al., 2021). Thus, it is crucial that studies examine strategies to make interventions more attractive and acceptable for males, to prevent drop-out.

To our knowledge, this study is the first MA to use individual participant-level data to examine moderators of EMDR for adults with PTSD. Among the strengths of the present study was its higher power to detect statistically significant moderators compared with study-level MA or any of the current RCTs aimed at investigating the efficacy of EMDR for adults with PTSD. The use of an IPDMA made it possible to investigate participant-level moderators (such as employment status and gender). Based on a visual inspection of the funnel plot of standard error Hedges' g , it is unlikely publication bias is present in this MA.

Several limitations of our IPDMA should be mentioned. First, the small sample sizes of the included studies, and consequently the total number of participants included in this IPDMA, limited our ability to detect certain moderators. Second, our findings are at risk of availability bias because we could not access data from seven eligible studies. However, the results of the study-level MA indicated no significant difference between the studies included in the present IPDMA and studies with unavailable data. Six of the 15 studies were published more than 20 years ago (≤ 2003). Only two of these six older studies were available for this IPDMA (Deville & Spence, 1999; Lee et al., 2002). Additionally, we could not examine several variables that could potentially influence EMDR treatment response, such as symptom duration or the number of treatment sessions attended because we did not have sufficient studies reporting these variables to conduct these analyses. Two studies were excluded from the dropout analysis because only the completer data were provided (Ahmadi et al., 2015; Carletto et al., 2016). Furthermore, most of the participants had chronic PTSD. Therefore, the current findings can only be generalized to patients with chronic PTSD.

Considering that EMDR is highly protocolized, and is relatively straightforward to administer, it may be a more cost and resource-effective treatment option to implement in areas with limited human resources. In the only systematic review to compare the relative cost-effectiveness of different PTSD treatments, EMDR was found to be the most cost-effective (Mavranzouli et al., 2020). However, further studies are needed in this area, in particular, large international RCTs.

With the increased use of secondary analyses and IPDMA, researchers are strongly urged to anonymize and store their data (in a usable format) for long-term use. In terms of the FAIR data principles, this not only improves scientific integrity but also prevents us from overlooking important discoveries. Data sharing, compiling and storage have become much faster and easier. By increasing the sample sizes in the EMDR effectiveness trials, reducing risk of bias, and increasing the number of RCTs statistical power can be increased which could improve our precision in detecting clinically relevant moderators of treatment outcomes. An update of this IPDMA in the future may have greater statistical power to provide further insight into moderating effects of participant-level factors on PTSD treatment response. Further research on the mechanisms underlying the efficacy of EMDR is needed, as well as whether bilateral stimulation offers additional therapeutic benefit. This requires adequately powered studies with dismantling designs. It is also

necessary to carry out well-designed randomized controlled trials in large samples.

In sum, this is the first IPDMA to have examined the effect of EMDR in comparison to other psychological treatments and explore what individual-level characteristics moderate PTSD treatment outcomes. Overall, no significant difference was found between EMDR and other effective psychological treatments in terms of PTSD outcomes. The use of IPDMA allowed us to examine the moderators of EMDR in a more powerful manner. It is vital that studies investigate approaches to make interventions more attractive and acceptable for males, in order to reduce drop-out. Findings from this IPDMA suggest that particular consideration should be taken when administering EMDR to unemployed individuals who may have extra challenges that could influence their treatment course.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0033291723003446>.

Funding statement. This research is funded by the NRF-NUFFIC scholarship, grant number 115977.

Competing interests. None.

References

- Ahmadi, K., Hazrati, M., Ahmadizadeh, M., & Noohi, S. (2015). REM desensitization as a new therapeutic method for post-traumatic stress disorder: A randomized controlled trial. *Acta Medica Indonesiana*, 47(2), 111–119. Retrieved from <http://www.inaactamedica.org/archives/2015/26260553.pdf>
- Barrowcliff, A. L., Gray, N. S., Freeman, T. C. A., & MacCulloch, M. J. (2004). Eye-movements reduce the vividness, emotional valence and electrodermal arousal associated with negative autobiographical memories. *The Journal of Forensic Psychiatry & Psychology*, 15(2), 325–345. doi:10.1080/14789940410001673042
- Bisson, J. I., Andrew, M., Roberts, N., Cooper, R., & Lewis, C. (2013). Psychological therapies for chronic post-traumatic stress disorder (PTSD) in adults (review). *Cochrane Database of Systematic Reviews*, 2013(12), CD003388. doi:10.1002/14651858.CD003388.pub4
- Bisson, J. I., Berliner, L., Cloitre, M., Forbes, D., Jensen, T. K., Lewis, C., ... Shapiro, F. (2019). The international society for traumatic stress studies new guidelines for the prevention and treatment of posttraumatic stress disorder: Methodology and development process. *Journal of Traumatic Stress*, 32(4), 475–483. doi:10.1002/jts.22421
- Bisson, J. I., & Olf, M. (2021). Prevention and treatment of PTSD: The current evidence base. *European Journal of Psychotraumatology*, 12(1), 1824381. doi:10.1080/20008198.2020.1824381
- Blanchard, E. B., Hickling, E. J., Devineni, T., Veazey, C. H., Galovski, T. E., Mundy, E., ... Buckley, T. C. (2003). A controlled evaluation of cognitive behavioural therapy for posttraumatic stress in motor vehicle accident survivors. *Behaviour Research and Therapy*, 41(1), 79–96. doi:10.1016/s0005-7967(01)00131-0
- Blanchard, E. B., Jones-Alexander, J., Buckley, T. C., & Forneris, C. A. (1996). Psychometric properties of the PTSD checklist (PCL). *Behaviour Research and Therapy*, 34(8), 669–673. doi:10.1016/0005-7967(96)00033-2
- Bosmans, M. W. G., & van der Velden, P. G. (2018). The effect of employment status in postdisaster recovery: A longitudinal comparative study among employed and unemployed affected residents. *Journal of Traumatic Stress*, 31(3), 460–466. doi:10.1002/jts.22282
- Boterhoven-De Haan, K. L., Lee, C. W., Fassbinder, E., van Es, S. M., Menninga, S., Meewisse, M. L., ... Arntz, A. (2020). Imagery rescripting and eye movement desensitisation and reprocessing as treatment for adults with post-traumatic stress disorder from childhood trauma: Randomised clinical trial. *The British Journal of Psychiatry: The Journal of Mental Science*, 217(5), 609–615. doi:10.1192/bjp.2020.158
- Capezzani, L., Ostacoli, L., Cavallo, M., Carletto, S., Fernandez, I., Solomon, R., ... Cantelmi, T. (2013). EMDR And CBT for cancer patients: Comparative study

- of effects on PTSD, anxiety, and depression. *Journal of EMDR Practice and Research*, 7, 134–143. doi:10.1891/1933-3196.7.3.134
- Carletto, S., Borghi, M., Bertino, G., Oliva, F., Cavallo, M., Hofmann, A., ... Ostacoli, L. (2016). Treating post-traumatic stress disorder in patients with multiple sclerosis: A randomized controlled trial comparing the efficacy of eye movement desensitization and reprocessing and relaxation therapy. *Frontiers in Psychology*, 7, 526. doi:10.3389/fpsyg.2016.00526
- Carlson, J., Chemtob, C., Rusnak, K., Hedlund, N., & Muraoka, M. (1998). Eye movement desensitization and reprocessing (EDMR) treatment for combat-related posttraumatic stress disorder. *Journal of Traumatic Stress*, 11(1), 32–33. doi:10.1023/A:1024448814268
- Chen, S., Westman, M., & Hobfoll, S. E. (2015). The commerce and crossover of resources: Resource conservation in the service of resilience. *Stress and Health*, 31, 95–105. doi:10.1002/smi.2574
- Cloitre, M., Koenen, K., Cohen, L., & Han, H. (2002). Skills training in affective and interpersonal regulation followed by exposure: A phase-based treatment for PTSD related to childhood abuse. *Journal of Consulting and Clinical Psychology*, 70, 1067–1074. doi:10.1037//0022-006x.70.5.1067
- Cuijpers, P., Veen, S. C. V., Sijbrandij, M., Yoder, W., & Cristea, I. A. (2020). Eye movement desensitization and reprocessing for mental health problems: A systematic review and meta-analysis. *Cognitive Behaviour Therapy*, 49(3), 165–180. doi:10.1080/16506073.2019.1703801
- Cusack, K., Jonas, D. E., Forneris, C. A., Wines, C., Sonis, J., Middleton, J. C., & Gaynes, B. N. (2016). Psychological treatments for adults with post-traumatic stress disorder: A systematic review and meta-analysis. *Clinical Psychology Review*, 43, 128–141. doi:10.1016/j.cpr.2015.10.003
- Debray, T. P., Moons, K. G., Abo-Zaid, G. M., Koffijberg, H., & Riley, R. D. (2013). Individual participant data meta-analysis for a binary outcome: One-stage or two-stage? *PLoS One*, 8(4), e60650. doi:10.1371/journal.pone.0060650
- Devilly, G., & Spence, S. (1999). The relative efficacy and treatment distress of EMDR and a cognitive-behavior trauma treatment protocol in the amelioration of posttraumatic stress disorder. *Journal of Anxiety Disorders*, 13(1–2), 131–157. doi:10.1016/s0887-6185(98)00044-9
- Duval, S., & Tweedie, R. (2000). Trim and fill: A simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics*, 56(2), 455–463. doi:10.1111/j.0006-341x.2000.00455.x
- Egger, M., Davey Smith, G., Schneider, M., & Minder, C. (1997). Bias in meta-analysis detected by a simple, graphical test. *British Medical Journal*, 315(7109), 629–634. doi:10.1136/bmj.315.7109.629
- Ehlers, A., Clark, D. M., Hackmann, A., McManus, F., Fennell, M., Herbert, C., & Mayou, R. (2003). A randomized controlled trial of cognitive therapy, a self-help booklet, and repeated assessments as early interventions for post-traumatic stress disorder. *Archives of General Psychiatry*, 60(10), 1024–1032. doi:10.1001/archpsyc.60.10.1024
- Galovski, T. E., Blain, L. M., Mott, J. M., Elwood, L., & Houle, T. (2012). Manualized therapy for PTSD: Flexing the structure of cognitive processing therapy. *Journal of Consulting and Clinical Psychology*, 80(6), 968–981. doi:10.1037/a0030600
- Gurung, T., Ellard, D. R., Mistry, D., Patel, S., & Underwood, M. (2015). Identifying potential moderators for response to treatment in low back pain: A systematic review. *Physiotherapy*, 101(3), 243–251. doi:10.1016/j.physio.2015.01.006
- Haagen, J. F., Ter Heide, F. J., Mooren, T. M., Knipscheer, J. W., & Kleber, R. J. (2017). Predicting post-traumatic stress disorder treatment response in refugees: Multilevel analysis. *The British Journal of Clinical Psychology*, 56(1), 69–83. doi:10.1111/bjc.12121
- Hagenaars, M. A., van Minnen, A., & Hoogduin, K. A. (2010). The impact of dissociation and depression on the efficacy of prolonged exposure treatment for PTSD. *Behaviour Research and Therapy*, 48(1), 19–27. doi:10.1016/j.brat.2009.09.001
- Hamblen, J. L., Norman, S. B., Sonis, J. H., Phelps, A. J., Bisson, J. I., Nunes, V. D., ... Schnurr, P. P. (2019). A guide to guidelines for the treatment of post-traumatic stress disorder in adults: An update. *Psychotherapy (Chic)*, 56(3), 359–373. doi:10.1037/pst0000231
- Higgins, J. P., & Thompson, S. G. (2002). Quantifying heterogeneity in a meta-analysis. *Statistics in Medicine*, 21(11), 1539–1558. doi:10.1002/sim.1186
- Higgins, J. P. T., Savović, J., Page, M. J., Elbers, R. G., & Sterne, J. A. C. (2022). Chapter 8: Assessing risk of bias in a randomized trial. In J. P. T. Higgins, J. Thomas, J. Chandler, M. Cumpston, T. Li, M. J. Page, & V. A. Welch (Eds.), *Cochrane handbook for systematic reviews of interventions* (2nd ed.). Chichester, UK: John Wiley & Sons.
- Horowitz, M., Wilner, N., & Alvarez, W. (1979). Impact of event scale: A measure of subjective stress. *Psychosomatic Medicine*, 41(3), 209–218. doi:10.1097/00006842-197905000-00004
- Ioannidis, J. P., Patsopoulos, N. A., & Evangelou, E. (2007). Uncertainty in heterogeneity estimates in meta-analyses. *British Medical Journal*, 335(7626), 914–916. doi:10.1136/bmj.39343.408449.80
- Ivarsson, D., Blom, M., Hesser, H., Carlbring, P., Enderby, P., Nordberg, R., & Andersson, G. (2014). Guided internet-delivered cognitive behavior therapy for post-traumatic stress disorder: A randomized controlled trial. *Internet Interventions*, 1, 33–40. doi:10.1016/j.invent.2014.03.002
- Jeffries, F. W., & Davis, P. (2013). What is the role of eye movements in eye movement desensitization and reprocessing (EMDR) for post-traumatic stress disorder (PTSD)? A review. *Behavioural and Cognitive Psychotherapy*, 41(3), 290–300. doi:10.1017/S1352465812000793
- Karatzias, A., Power, K., McGoldrick, T., Brown, K., Buchanan, R., Sharp, D., & Swanson, V. (2007). Predicting treatment outcome on three measures for post-traumatic stress disorder. *European Archives of Psychiatry and Clinical Neuroscience*, 257(1), 40–46. doi:10.1007/s00406-006-0682-2
- Karatzias, T., Power, K., Brown, K., McGoldrick, T., Begum, M., Young, J., ... Adams, S. (2011). A controlled comparison of the effectiveness and efficiency of two psychological therapies for posttraumatic stress disorder: Eye movement desensitization and reprocessing vs. Emotional freedom techniques. *The Journal of Nervous and Mental Disease*, 199(6), 372–378. doi:10.1097/NMD.0b013e31821cd262
- Karyotaki, E., Riper, H., Twisk, J., Hoogendoorn, A., Kleiboer, A., Mira, A., ... Cuijpers, P. (2017). Efficacy of self-guided internet-based cognitive behavioral therapy in the treatment of depressive symptoms: A meta-analysis of individual participant data. *The Journal of the American Medical Association*, 74(4), 351–359. doi:10.1001/jamapsychiatry.2017.0044
- Karyotaki, E., Sijbrandij, M., Purgato, M., Acarturk, C., Lakin, D., Bailey, D., ... Barbui, C. (2023). Self-help plus for refugees and asylum seekers: An individual participant data meta-analysis. *BMJ Mental Health*, 26(1), e300672. doi:10.1136/bmjment-2023-300672
- Keane, T. M., Malloy, P. F., & Fairbank, J. A. (1984). Empirical development of an MMPI subscale for the assessment of combat-related posttraumatic stress disorder. *Journal of Consulting and Clinical Psychology*, 52(5), 888–891. doi:10.1037//0022-006x.52.5.888
- Landin-Romero, R., Moreno-Alcazar, A., Pagani, M., & Amann, B. L. (2018). How does eye movement desensitization and reprocessing therapy work? A systematic review on suggested mechanisms of action. *Frontiers in Psychology*, 9, 1395. doi:10.3389/fpsyg.2018.01395
- Laugharne, J., Kullack, C., Lee, C. W., McGuires, T., Brockman, S., Drummond, P. D., & Starkstein, S. (2016). Amygdala volumetric change following psychotherapy for posttraumatic stress disorder. *The Journal of Neuropsychiatry and Clinical Neurosciences*, 28(4), 312–318. doi:10.1176/appi.neuropsych.16010006
- Lee, C., Gavriel, H., Drummond, P., Richards, J., & Greenwald, R. (2002). Treatment of PTSD: Stress inoculation training with prolonged exposure compared to EMDR. *Journal of Clinical Psychology*, 58, 1071–1089. doi:10.1002/jclp.10039
- Lewis, C., Roberts, N. P., Andrew, M., Starling, E., & Bisson, J. I. (2020a). Psychological therapies for post-traumatic stress disorder in adults: Systematic review and meta-analysis. *European Journal of Psychotraumatology*, 11(1), 1729633. doi:10.1080/20008198.2020.1729633
- Lewis, C., Roberts, N. P., Gibson, S., & Bisson, J. I. (2020b). Dropout from psychological therapies for post-traumatic stress disorder (PTSD) in adults: Systematic review and meta-analysis. *European Journal of Psychotraumatology*, 11(1), 1709709. doi:10.1080/20008198.2019.1709709
- Lewis, C. E., Farewell, D., Groves, V., Kitchiner, N. J., Roberts, N. P., Vick, T., & Bisson, J. I. (2017). Internet-based guided self-help for posttraumatic stress disorder (PTSD): Randomized controlled trial. *Depression and Anxiety*, 34(6), 555–565. doi:10.1002/da.22645
- Mavranzeouli, I., Megnin-Viggars, O., Grey, N., Bhutani, G., Leach, J., Daly, C., ... Pilling, S. (2020). Cost-effectiveness of psychological treatments for post-traumatic stress disorder in adults. *PLoS One*, 15(4), e0232245. doi:10.1371/journal.pone.0232245

- McKee-Ryan, F. M., Song, Z., Wanberg, C. R., & Kinicki, A. J. (2005). Psychological and physical well-being during unemployment: A meta-analytic study. *Journal of Applied Psychology, 90*, 53–76. doi:10.1037/0021-9010.90.1.53
- Mertens, G., Lund, M., & Engelhard, I. M. (2021). The effectiveness of dual-task interventions for modulating emotional memories in the laboratory: A meta-analysis. *Acta Psychologica, 220*, 103424. doi:10.1016/j.actpsy.2021.103424
- Nijdam, M. J., Gersons, B. P., Reitsma, J. B., de Jongh, A., & Olff, M. (2012). Brief eclectic psychotherapy v. eye movement desensitization and reprocessing therapy for post-traumatic stress disorder: Randomised controlled trial. *The British Journal of Psychiatry, 200*(3), 224–231. doi:10.1192/bjp.bp.111.099234
- Nijdam, M. J., Vermetten, E., & McFarlane, A. C. (2023). Toward staging differentiation for posttraumatic stress disorder treatment. *Acta Psychiatrica Scandinavica, 147*(1), 65–80. doi:10.1111/acps.13520
- O'Donnell, M. L., Grant, G., Alkemade, N., Spittal, M., Creamer, M., Silove, D., ... Studdert, D. M. (2015). Compensation seeking and disability after injury: The role of compensation-related stress and mental health. *Journal of Clinical Psychiatry, 76*(8), e1000–e1005. doi:10.4088/JCP.14m09211
- Olff, M. (2017). Sex and gender differences in post-traumatic stress disorder: An update. *European Journal of Psychotraumatology, 8*(sup4), 1. doi:10.1080/20008198.2017.1351204
- Orsini, N., Bottai, M., Higgins, J., & Buchan, I. (2005). *Heterogi: Stata module to quantify heterogeneity in a meta-analysis* (software program; Statistical Software Components S449201). Statistical software components. Boston: Boston College Department of Economics.
- Papola, D., Ostuzzi, G., Tedeschi, F., Gastaldon, C., Purgato, M., Del Giovane, C., ... Barbui, C. (2022). Comparative efficacy and acceptability of psychotherapies for panic disorder with or without agoraphobia: Systematic review and network meta-analysis of randomised controlled trials. *The British Journal of Psychiatry, 221*(3), 507–519. doi:10.1192/bjp.2021.148
- Paul, K. I., & Batinic, B. (2010). The need for work: Jahoda's latent functions of employment in a representative sample of the German population. *Journal of Organizational Behavior, 31*, 45–64. doi:10.1002/job.622
- Phelps, A. J., Lethbridge, R., Brennan, S., Bryant, R. A., Burns, P., Cooper, J. A., ... Silove, D. (2022). Australian guidelines for the prevention and treatment of posttraumatic stress disorder: Updates in the third edition. *Australian & New Zealand Journal of Psychiatry, 56*(3), 230–247. doi:10.1177/00048674211041917
- Power, K., McGoldrick, T., Brown, K., Buchanan, R., Sharp, D., Swanson, V., & Karatzias, T. (2002). A controlled comparison of eye movement desensitization and reprocessing versus exposure plus cognitive restructuring versus waiting list in the treatment of post-traumatic stress disorder. *Clinical Psychology and Psychotherapy, 9*, 229–318. doi:10.1002/cpp.341
- Riley, R. D., Lambert, P. C., & Abo-Zaid, G. (2010). Meta-analysis of individual participant data: Rationale, conduct, and reporting. *British Medical Journal, 340*, c221. doi:10.1136/bmj.c221
- Riley, R. D., Tierney, J. F., & Stewart, L. A. (2021). *Individual participant data meta-analysis: A handbook for healthcare research*. Chichester: Wiley.
- Rizvi, S. L., Vogt, D. S., & Resick, P. A. (2009). Cognitive and affective predictors of treatment outcome in cognitive processing therapy and prolonged exposure for posttraumatic stress disorder. *Behaviour Research and Therapy, 47*(9), 737–743. doi:10.1016/j.brat.2009.06.003
- Rothbaum, B. O., Astin, M. C., & Marsteller, F. (2005). Prolonged exposure versus eye movement desensitization and reprocessing (EMDR) for PTSD rape victims. *Journal of Traumatic Stress, 18*(6), 607–616. doi:10.1002/jts.20069
- Schubert, S. J., & Lee, C. (2009). Adult PTSD and its treatment with EMDR: A review of controversies, evidence, and theoretical knowledge. *Journal of EMDR Practice and Research, 3*, 117–132. doi:10.1891/1933-3196.3.3.117
- Seidler, Z. E., Wilson, M. J., Kealy, D., Oliffe, J. L., Ogradniczuk, J. S., & Rice, S. M. (2021). Men's dropout from mental health services: Results from a survey of Australian men across the life span. *American Journal of Men's Health, 15*(3), 15579883211014776. doi:10.1177/15579883211014776
- Shapiro, F. (1989). Eye movement desensitization: A new treatment for post-traumatic stress disorder. *Journal of Behavior Therapy and Experimental Psychiatry, 20*, 211–217. doi:10.1016/0005-7916(89)90025-6
- Shapiro, F. (2001). *Eye movement desensitization and reprocessing (EMDR): Basic principles, protocols, and procedures*. New York, NY: Guilford Press.
- Shapiro, F. (2006). *New notes on adaptive information processing*. Hamden, CT: EMDR Humanitarian Assistance Programs.
- Shapiro, F., & Liliotis, D. (2011). EMDR and the adaptive information processing model: Integrative treatment and case conceptualization. *Clinical Social Work Journal, 39*(2), 191–200. doi:10.1007/s10615-010-0300-7
- StataCorp. (2021). *Stata statistical software: Release 17*. College Station, TX: StataCorp LLC.
- Stewart, L. A., Clarke, M., Rovers, M., Riley, R. D., Simmonds, M., Stewart, G., ... PRISMA-IPD Development Group. (2015). Preferred reporting items for systematic review and meta-analyses of individual participant data: The PRISMA-IPD statement. *The Journal of the American Medical Association, 313*(16), 1657–1665. doi:10.1001/jama.2015.3656
- Stewart, L. A., & Parmar, M. K. (1993). Meta-analysis of the literature or of individual patient data: Is there a difference? *Lancet, 341*(8842), 418–422. doi:10.1016/0140-6736(93)93004-k
- Taylor, S., Thordarson, D. S., Maxfield, L., Fedoroff, I. C., Lovell, K., & Ogradniczuk, J. (2003). Comparative efficacy, speed, and adverse effects of three PTSD treatments: Exposure therapy, EMDR, and relaxation training. *Journal of Consulting and Clinical Psychology, 71*, 330–338. doi:10.1080/00332747.2017.1286892
- Ter Heid, F. J., Mooren, T. M., van de Schoot, R., de Jongh, A., & Keber, R. J. (2016). Eye movement desensitization and reprocessing therapy v. Stabilisation as usual for refugees: Randomised controlled trial. *British Journal of Psychiatry, 209*(4), 311–318. doi:10.1192/bjp.bp.115.167775
- VA/DoD Clinical Practice Guideline. (2023). *Management of posttraumatic stress disorder and acute stress disorder work group*. Washington, DC: U.S. Government Printing Office.
- van den Hout, M. A., & Engelhard, I. M. (2012). How does EMDR work? *Journal of Experimental Psychopathology, 3*(5), 724–738. doi:10.5127/jep.028212
- van Minnen, A., Wessel, I., Dijkstra, T., & Roelofs, K. (2002). Changes in PTSD patients' narratives during prolonged exposure therapy: A replication and extension. *Journal of Traumatic Stress, 15*(3), 255–258. doi:10.1023/A:1015263513654
- Vaughan, K., Armstrong, M. S., Gold, R., O'Connor, N., Jenneke, W., & Tarrier, N. (1994). A trial of eye movement desensitization compared to image habituation training and applied muscle relaxation in post-traumatic stress disorder. *Journal of Behavior Therapy and Experimental Psychiatry, 25*, 283–291. doi:10.1016/0005-7916(94)90036-1
- Weiss, D. S., & Marmar, C. R. (1997). The impact of event scale-revised. In J. P. Wilson, & T. M. Keane (Eds.), *Assessing psychological trauma and PTSD: A practitioner's handbook* (pp. 399–411). New York: Guilford Press.
- Wright, S. L., Karyotaki, E., Bisson, J. I., Cuijpers, P., Papola, D., Witteveen, A. B., ... Sijbrandij, M. (2022). Protocol for individual participant data meta-analysis of interventions for post-traumatic stress. *British Medical Journal Open, 12*(2), e054830. doi:10.1136/bmjopen-2021-054830