








Biological Sciences

Impacts of ocean acidification on the palatability of two Antarctic macroalgae and the consumption of a grazer

Hannah E. Oswalt¹, Margaret O. Amsler¹, Charles D. Amsler¹, James B. McClintock¹ and Julie B. Schram²

¹Department of Biology, University of Alabama at Birmingham, Birmingham, AL, USA and ²Department of Natural Sciences, University of Alaska Southeast, Juneau, AK, USA

Abstract

Increases in atmospheric CO₂ have led to more CO₂ entering the world's oceans, decreasing the pH in a process called 'ocean acidification'. Low pH has been linked to impacts on macroalgal growth and stress, which can alter palatability to herbivores. Two common and ecologically important macroalgal species from the western Antarctic Peninsula, the unpalatable *Desmarestia menziesii* and the palatable *Palmaria decipiens*, were maintained under three pH treatments: ambient (pH 8.1), near future (7.7) and distant future (7.3) for 52 days and 18 days, respectively. Discs of *P. decipiens* or artificial foods containing extracts of *D. menziesii* from each treatment were presented to the amphipod *Gondogeneia antarctica* in feeding choice experiments. Additionally, *G. antarctica* exposed to the different treatments for 55 days were used in a feeding assay with untreated *P. decipiens*. For *D. menziesii*, extracts from the ambient treatment were eaten significantly more by weight than the other treatments. Similarly, *P. decipiens* discs from the ambient and pH 7.7 treatments were eaten more than those from the pH 7.3 treatment. There was no significant difference in the consumption by treated *G. antarctica*. These results suggest that ocean acidification may decrease the palatability of these macroalgae to consumers but not alter consumption by *G. antarctica*.

Keywords: Chemical defence; climate change; herbivory; ocean acidification; seaweed

(Received 3 June 2024; revised 23 October 2024; accepted 9 December 2024)

Introduction

Human-derived emissions from the combustion of fossil fuels, the production of cement and deforestation have led to a drastic increase in atmospheric CO₂ (Malhi *et al.* 2002, Heede 2014, Paraschiv & Paraschiv 2020). In 2019 alone, ~98.2 Mt CO₂ were added to the atmosphere daily (Liu *et al.* 2020). The culmination of decades of emissions since the Industrial Revolution has increased atmospheric CO₂ concentrations by ~50% from 280 ppm in 1750 to a peak of 420 ppm in 2024 (Joos & Spahni 2008, Lan & Keeling 2024).

One consequence of increased atmospheric CO₂ concentrations is ocean acidification (OA). Approximately a third of atmospheric CO₂ enters the ocean (Sabine *et al.* 2004, Gruber *et al.* 2019), where it reacts with seawater in a series of steps that ultimately release free hydrogen ions, thereby lowering pH. As a result of this gaseous uptake, the average ocean's surface pH has decreased by 0.1 pH units and is predicted to decrease a further 0.4 pH units over the next 80 years (IPCC 2022).

Changes in abiotic factors from climate change, particularly decreased ocean pH, can have a direct impact on the physiology and behaviour of marine organisms (Kroeker *et al.* 2013, Kindinger *et al.* 2022). These changes not only impact species individually but

also impact the interactions between species, such as macroalgae-herbivore interactions. For example, OA can affect the growth rate (Johnson *et al.* 2014), tissue strength (Kinnby *et al.* 2021), nutritional quality (Duarte *et al.* 2016, Fieber & Bourdeau 2021) and putative chemical defences of macroalgae (Swanson & Fox 2007, Fieber & Bourdeau 2021, Kinnby *et al.* 2021). Similarly, OA can impact herbivores by altering growth (Sheppard Brennan *et al.* 2010, Bhuiyan *et al.* 2022), survival (Lopes *et al.* 2019, Park *et al.* 2020), development (Kurihara & Shirayama 2004, Watson *et al.* 2009, Place & Smith 2012) and reproduction (Kurihara & Shirayama 2004, Morita *et al.* 2010, Borges *et al.* 2018) and by shifting energy demands (Melzner *et al.* 2011, Saba *et al.* 2012, Ramajo *et al.* 2016, Lim & Harley 2018).

The consequences of OA on one species can impact the relationship between macroalgae and herbivores. For example, grazing pressure from invertebrates can become greater if energy demands on invertebrates increase from compensatory strategies in response to changing pH (Saba *et al.* 2012, Leung *et al.* 2018). Alternatively, grazing pressure could be altered from changes in the macroalgae, but the specific type of change is heavily dependent on the species and how the macroalga is being affected. For example, changes in nutritional quality or qualitative or quantitative changes in the production of chemical defences have been documented (Swanson & Fox 2007, Duarte *et al.* 2016, Fieber & Bourdeau 2021).

A significant portion of previous OA work has focused on calcifying species and communities (e.g. Andersson *et al.* 2011, Leung *et al.* 2022, Page *et al.* 2022), mainly due to the negative impacts of OA on the availability of carbonate and the alteration

Corresponding author: Hannah E. Oswalt; Email: heoswalt@uab.edu

Cite this article: Oswalt, H. E., Amsler, M. O., Amsler, C. D., McClintock, J. B., & Schram, J. B. 2025. Impacts of ocean acidification on the palatability of two Antarctic macroalgae and the consumption of a grazer. *Antarctic Science*, 1–10. <https://doi.org/10.1017/S095410202400052X>

of seawater chemistry (Doney *et al.* 2009, Hurd *et al.* 2009) that reduce these organisms' ability to calcify. However, focusing on one functional group of macroalgae cannot provide a full picture of OA consequences on algae. Macroalgae are taxonomically and metabolically diverse (Li-Beisson *et al.* 2019, Tully 2019, Schubert *et al.* 2023, Guiry 2024), making it difficult to generalize regarding how they will respond as a whole to climate change based on a few studied species.

Fleshy macroalgal species are expected to be more resilient to increased CO₂ concentrations compared to calcified species because the former do not experience dissolution and have access to more CO₂ for photosynthesis (Koch *et al.* 2013), which has been supported by modelling (Schlenger *et al.* 2021). However, experiments have shown that the consequences of OA appear to be species-specific. Previous studies have found increases in net growth for some, but not all, of the fleshy macroalgae held under different conditions of OA (Johnson *et al.* 2014, Kram *et al.* 2016, Paine *et al.* 2023). In some cases, there have been contrasting results in net growth with OA conditions. For example, Swanson & Fox (2007) found that CO₂ levels predicted for 2100 resulted in a positive effect on the growth of the kelp *Nereocystis luetkeana* but a negative effect on the growth of the kelp *Saccharina latissima*.

In addition to changes in net growth, shifts in chemical composition and photosynthesis as a response to lowered pH have also been observed in macroalgae (Garcia-Sanchez *et al.* 1994, Mercado *et al.* 1999, Gordillo *et al.* 2001, Duarte *et al.* 2016). Changes in chemical composition can alter the overall nutritional quality of macroalgae and can impact herbivore preference (Duarte *et al.* 2016). Furthermore, these changes in nutritional content can impact the consumers of algae (Urabe *et al.* 2003, Schoo *et al.* 2013, Sudo & Yoshida 2021, Sepúlveda *et al.* 2024). For example, the amphipod *Ampithoe tarasovi*, the sea urchin *Tetrapygus niger* and the sea snail *Tegula atra* experience reduced growth when fed a diet of lower-quality macroalgae (Sudo & Yoshida 2021, Sepúlveda *et al.* 2024).

High-latitude environments, such as the those of the western Antarctic Peninsula region (WAP; including islands to the west and north of the continental peninsula), are particularly vulnerable to decreased seawater pH due to the low buffering capacity associated with cold waters (Sabine *et al.* 2004, Nissen *et al.* 2024). The WAP is characterized by its large macroalgal forests that dominate shallow (< 70 m) benthic communities (Wiencke *et al.* 2014). These macroalgal communities contribute standing biomass levels comparable to temperate kelp forests, often covering more than 80% of the ocean benthos (Wiencke *et al.* 2014), and they play a vital role in nutrient cycling, nutrient retention and benthic food webs (Dunton 2001, Quartino & Boraso de Zaiuso 2008, Wiencke & Amsler 2012, Iken *et al.* 2023). The dominant macroalgae in these communities are large, perennial browns from the family Desmarestiaceae, while the understory is dominated by smaller red macroalgae (Wiencke *et al.* 2014). In addition to producing large amounts of biomass, many of the macroalgae in these near-shore communities also produce chemical defences that protect them from herbivory (Amsler *et al.* 2020). Most of this macroalgal community is chemically defended, including all of the large browns and many of the common reds, with the notable exception of *Palmaria decipiens* (Amsler *et al.* 2005, Aumack *et al.* 2010). Despite their chemical defences, these macroalgae can serve as a food resource after they die and their former chemical defences break down (Wiencke & Amsler 2012).

One unique aspect of the WAP is the incredibly large number of mesograzers found in the macroalgal communities. Macroalgae

shelter large mesograzer assemblages that mostly consist of amphipods with large numbers of gastropods and smaller numbers of isopods, copepods and ostracods (Iken 1999, Huang *et al.* 2007, Schram *et al.* 2016, Amsler *et al.* 2022). The density of amphipods on macroalgae in this region can be extremely abundant, with densities reaching as high as 300 000 and 26 000 individuals m² on the benthos in stands of *Desmarestia menziesii*, a dominant brown macroalga, and the red macroalga *Plocamium* sp., respectively (Huang *et al.* 2007, Amsler *et al.* 2008). These densities are approximately one to three times higher than reported amphipod densities in tropical and temperate regions (Amsler *et al.* 2014).

In the present study, we assessed how the palatability of two common macroalgae, *Desmarestia menziesii* J. Agardh and *Palmaria decipiens* (Reinsch) R. W. Ricker, is impacted by reduced seawater pH. These two species represent a chemically defended macroalga (*D. menziesii*) and a palatable macroalga (*P. decipiens*) that occur commonly in the community. Thalli of both species were exposed to three different pH treatments simulating ambient conditions (approximately pH 8.1), near-future conditions for 2100 (pH 7.7; IPCC 2022) and distant-future conditions (pH 7.3; IPCC 2022) and then used in a feeding assay with the common amphipod *Gondogeneia antarctica* (Chevreux). Additionally, we also conducted a feeding assay with *G. antarctica* exposed to the same pH treatments to determine whether there is an impact of pH on the amphipod's total consumption. *Gondogeneia antarctica* was selected because it is one of the most common amphipod species in this region (Huang *et al.* 2007) and is a very commonly studied WAP amphipod (e.g. Amsler *et al.* 2020, Park *et al.* 2020, Ahn *et al.* 2021). We hypothesized that the palatability of *P. decipiens* would increase while the palatability of *D. menziesii* would either increase or decrease due to reported increases in photosynthetic capacity and growth in several fleshy macroalgae species. Furthermore, we hypothesized that *G. antarctica* would demonstrate higher consumption at lowered pH due to a possible increase in energetic demands from compensatory strategies.

Materials and methods

Collection of macroalgae and amphipods

The shallow-water brown macroalga *D. menziesii* was collected from five sites near the US Antarctic Research Program's Palmer Station on Anvers Island on the WAP. Scuba divers collected *D. menziesii* from Palmer Station pier (S64°46.477', W64°03.274'), Hero Inlet (S64°46.569', W64°02.874'), Bonaparte Point (S64°46.745', W64°02.559'), Gamage Point (S64°46.432', W64°03.368') and Hermit Island (S64°47.880', W64°00.438'). *Desmarestia menziesii* samples were collected between 4 and 11 m on 19–27 December 2019 (Year 1) by cutting a ~85 g or larger section from an individual of *D. menziesii* with a knife and then placing the section into a fine mesh bag. *Palmaria decipiens* was collected between 2 and 8 m from Amsler Island (S64°46.487', W64°03.262') on 14 March 2023 (Year 2) by collecting small rocks with macroalgae attached and placing them in a mesh bag.

The collection bags were immediately placed in a 19 l bucket filled with seawater and transported to Palmer Station. *Desmarestia menziesii* was rinsed repeatedly with seawater in a series of fine mesh bags submerged in a tank of seawater to dislodge associated grazers. After the final rinse, the macroalgae were inspected to remove any remaining grazers by hand. The sections of *P. decipiens* were placed in a sorting tank. Each macroalga was visually inspected for grazers. Any grazers found were removed.

Amphipods for the experiments were collected between 9 January 2023 and 18 February 2023 (Year 2) from multiple locations in the near vicinity of Palmer Station using the methods described by Huang *et al.* (2007). Briefly, macroalgal-associated crustaceans were collected by gently prying *D. menziesii* off the substrata with a knife then carefully floating the macroalga into a fine mesh bag to avoid disturbing the associated grazers. After being transported back to station, the grazers were removed from *D. menziesii* as described above. Then, the mesograzers were placed in a sorting table aquarium to separate *G. antarctica* from the rest of the mesograzers. A haphazard subset of these amphipods was used in the OA experiment. Another subset of the amphipods was held in an ambient flow-through tank in groups of ~200 individuals in bottles fitted with mesh windows to allow for water flow until they were used in the macroalgae feeding experiments.

Experimental set-up

A nearly identical experimental set-up was created for the Year 1 and Year 2 experiments. The experimental set-up consisted of 24 white plastic buckets of 19 l in volume placed on a submerged platform in two adjacent aquarium tanks (2.5 m diameter and 1 m depth; 3800 l) in the aquarium facility at Palmer Station (see Fig. S1). The facility was maintained on a 24 h light cycle in Year 1 and an ambient light cycle in Year 2, consistent with light availability at the time of collection. Both tanks were plumbed with ambient flow-through seawater to create a flow-through seawater bath to maintain an ambient temperature in all replicates. The tops of the buckets were elevated above the water level of the flow-through seawater.

Filtered seawater collected and flowed out of a common, constant head tank into a water distributor located in the centre of each water bath. Water flowed from the distributor to 12 mixing reservoirs arranged in a circle around the distributor. Along with an inflow water pipe, mixing reservoirs also contained a pH probe, a tube with mixed air and compressed CO₂ and a tube with compressed air alone to monitor and control the pH. After the seawater was adjusted to the desired pH in the mixing reservoirs (as described below), the seawater flowed into the experimental buckets. Finally, water flowed out of the bucket into the flow-through seawater bath.

There were eight buckets per pH treatment (ambient (8.1), 7.7 and 7.3). Seawater pH treatment levels were based on the average pH recorded for Palmer Station during collection (~8.1) and levels for the predicted near and distant future (IPCC 2022). The pH of the buckets was monitored throughout the experiment. Buckets in both decreased pH treatments were individually regulated with an automated pH monitoring system (AquaMedic, USA; Schoenrock *et al.* 2016, Schram *et al.* 2016). The target pH for each bucket was maintained by bubbling an air-CO₂ mixture or air alone into the mixing reservoirs as needed. The air-CO₂ mixture was created by combining ratios of air and CO₂ with a Multi-tube Gas Proportioning rotameter (Omega Engineering, Inc., Stamford, CT, USA) connected to a CO₂ cylinder and multiple air pumps.

The eight buckets for each pH treatment were randomly assigned within each water bath. Each water bath contained half of the replicates for each treatment to control for differences in light availability due to water bath location. The buckets had clear Plexiglass covers to reduce gas exchange with the atmosphere (Schram *et al.* 2016). In Year 2, some of the bucket lids were outfitted with screen mesh to equalize light levels inside of the

buckets. The control seawater pH replicates received ambient air only.

Seawater samples from a quarter of the experimental buckets were collected for pH measurements (determined spectrophotometrically) and total alkalinity (TA; determined by potentiometric titration; Dickson *et al.* 2007) daily. This sampling method resulted in all buckets being tested over a 4 day cycle. In addition, the pH in all buckets was monitored at least once daily with a calibrated pH electrode.

In Year 1, seven thalli of *D. menziesii* were cut into three parts (one additional individual died and could not be used as planned). Each part of the macroalga was distributed into each of the pH treatments so that one individual was placed in the ambient, near-future and far-future treatments for comparison. After the macroalgae were placed in the buckets, the pH was slowly decreased in the lower pH treatments over a 28 h period. The experiment ran for a total of 52 days. At the end of the experiment, *D. menziesii* were removed from the experimental buckets and rinsed with seawater to remove epiphytes and grazers. Each macroalga was visually inspected to ensure epiphytes and grazers were fully removed. Each sample was then frozen in a -20°C freezer until processed.

In Year 2, individuals of *P. decipiens* were split into three parts. One third of each macroalga was distributed to each of the pH treatments. The macroalgae were placed inside clear plastic beakers with window screening mesh secured to the top of the beaker. This allowed for water flow while also protecting the macroalgae from grazing from amphipods in a concurrent experiment. The macroalgae were exposed to their respective pH treatments for 18 days. This macroalga had a shorter exposure period compared to *D. menziesii* because it is a pseudoperennial species with a relatively short-lived upright blade portion (Wiencke 1990, Weykam *et al.* 1997) and needed to be exposed to the treatments and used in the feeding assay before it started to senesce.

One hundred *G. antarctica* were placed into each of the experimental buckets in Year 2 to be used in a feeding rate assay. The amphipods were allowed to acclimatize to the experimental buckets for a few days before the pH in the non-ambient treatments was slowly decreased over the course of 52 h. The amphipods were fed small scrap pieces of *P. decipiens* and were also provided with plastic artificial plants seeded with diatoms from an outdoor tank and replaced with new, diatom-seeded artificial plants regularly. The amphipods were exposed to their respective pH treatments for 55 days.

Carbonate chemistry determination

Seawater pH was determined on the total hydrogen scale (pH_T) using a Perkin Elmer UV/VIS Spectrometer Lambda 40P in Year 1 and a Perkin Elmer UV/VIS Lambda 365 in Year 2 after the addition of the pH-sensitive indicator dye m-cresol purple (SOP 6b; Dickson *et al.* 2007). Seawater TA for the samples was determined with open-cell potentiometric titration (SOP 3b; Dickson *et al.* 2007) using a Mettler-Toldeo T50 open-cell titrator equipped with a pH probe (Model DGi 115-SC). Samples were siphoned into a jacketed beaker plumbed to a water bath (Neslab RTE-7 Circulating Bath, Thermo Scientific, USA) to maintain a constant temperature during titration (20°C). Mettler-Toledo LabX® software was used to record titrant volumes in real time.

Carbonate chemistry parameters were calculated based on pH, TA, temperature and salinity data. In Year 1, CO₂calc software (Robbins *et al.* 2010) with CO₂ constants provided by Roy *et al.*

(1993) and a KHSO_4 acidity constant from Dickson (1990) was used for carbonate calculations as directed by Roy *et al.* (1993). In Year 2, the *R* package (R Core Team 2020) *seacarb* by Gattuso *et al.* (2022) with constants from Lueker *et al.* (2000) and Perez & Fraga (1987) was used for carbonate calculations as directed in *seacarb*. The temperature was recorded during sample collection and the salinity was measured with a Seabird 45 MicroTSG from the Palmer Station Waterwall (Palmer Station Instrument Technician 2023).

Preparation of *Desmarestia menziesii* discs

The chemical defences from each sample of *D. menziesii* from Year 1 were extracted using three changes of 1:1 CH_2Cl_2 :methanol. The resulting extracts from each solvent change were combined and then filtered. The solvents were removed by evaporation using a rotary evaporator. The resulting dry lipophilic extracts were weighed to calculate the extract yield.

Artificial food was created using 2.2% alginate with 5% *Cladophora repens* powder, a palatable, sympatric green alga (Amsler *et al.* 2005). Macroalgal extracts were dissolved in a minimum volume of methanol and dried on the algal powder with a rotary evaporator (Hay *et al.* 1994).

The extracts from *D. menziesii* are strongly unpalatable at their natural concentrations. A preliminary feeding trial was run to determine what concentration of extract would deter feeding but also allow feeding rates to be measured. Extracts from several different *D. menziesii* were combined and then diluted to 50% of natural concentration. The extract-coated *C. repens* powder was placed in a 30 mm plastic Petri dish and mixed thoroughly with a cold alginate solution and food colouring (to help visually distinguish each treatment). Then, the mixture was gelled using cold 1 M CaCl_2 . Finally, the artificial food rounds were cut into smaller discs with a cork borer to ~10 mm in diameter and 2 mm in thickness. Control artificial foods were created using the same methods without the addition of the *D. menziesii* extract. Artificial foods for the *D. menziesii* experiment were prepared using these methods. All experimental extracts were diluted to 50% of their natural concentrations.

To examine the palatability of the macroalgal extract concentration, *G. antarctica*, one of the most common species of amphipods in the macroalgal-associated community, was given a choice of artificial food discs in a feeding assay following the methods of Amsler *et al.* (2005). Twenty amphipods were haphazardly placed into nine 250 ml bottles filled with ambient seawater. The bottles contained an extract disc and a control disc. Paired bottles without amphipods served as autogenic controls for mass changes in the discs. Feeding preference was determined by calculating wet mass change between the paired discs from the experimental and control bottles at the end of assay.

Preparation of *Palmaria decipiens* discs

Two small discs were cut adjacent to each other from each section of *P. decipiens* from the experimental buckets. Each disc was labelled with a color-coded thread tied through the centre of the disc. The discs were allowed to soak in 250 ml plastic bottles for 4 h, then the water in the bottles was changed before the feeding assay to remove compounds released from the macroalgae when they were injured from the cutting of the discs (cf. McDowell *et al.* 2014).

Feeding assays

To examine the palatability of the macroalgal discs from each pH treatment, 20 non-treated amphipods were haphazardly placed into seven 250 ml bottles for *D. menziesii* and eight such bottles for *P. decipiens* filled with ambient seawater. The bottles contained a disc from each of the pH treatments (ambient, pH 7.7 and pH 7.3) from the same initial individual that was split at the beginning of the experiment. Paired bottles without amphipods served as controls for autogenic changes in the discs. Feeding preference was determined by calculating wet mass change between the paired discs from the experimental and autogenic control bottles at the end of assay.

To test the feeding rate of amphipods exposed to lowered pH, 10 *G. antarctica* from each experimental bucket were placed in a 250 ml bottle filled with water from that bucket to maintain a constant pH. Two discs from one *P. decipiens* were cored from adjacent spots on the algal blade. One disc was placed in the bottle with the amphipods and the other was placed in a paired bottle without amphipods to serve as an autogenic control. The feeding assay ran for 12 h. Amphipods were returned to their experimental buckets and the discs were weighed. Consumption was determined by calculating wet mass change between the paired discs from the experimental and autogenic control bottles at the end of assay.

Statistical analyses

Statistical analyses were performed with *R* v4.0.2 (R Core Team 2020) using the *ez* (Lawrence 2016), *car* (Fox & Weisberg 2019) and *mvnrmtest* packages (Jarek 2012). Normality and homogeneity of variance were tested using the Shapiro-Wilks test and Levene's test, respectively. The consumption amount from each assay was corrected using the autogenic controls. The consumption from the extract concentration experiment was compared using a *t*-test. Extract yields from the pH treatments were compared using an analysis of variance (ANOVA). A one-sample *t*-test was used to test whether amphipods were eating macroalgal discs with 50% extract concentration. The feeding assays comparing the pH treatments were compared using an ANOVA for the *G. antarctica* assay and a repeated-measures ANOVA for the three-choice feeding assays because the choices in such an assay are not independent of each other (Roa 1992). A Greenhouse-Geisser correction was utilized for *P. decipiens* because it did not meet the assumption of sphericity (Greenhouse & Geisser 1959, Armstrong 2017). Pairwise comparisons using a Bonferroni correction were utilized as a *post hoc* test for the repeated-measures ANOVA. The cut-off for statistical significance was set to $P \leq 0.05$.

Results

Seawater parameters during the experiments are summarized in Table I. Mean (\pm SD) pH values of the ambient and lowered pH treatments were very similar between Year 1 and Year 2. Data for each parameter (pH, TA, salinity, dissolved inorganic carbon (DIC), etc.) on each day of the experiment are available at the US Antarctic Program Data Center (see 'Details of data deposit' statement below). TA, temperature and salinity remained similar between the pH treatments across both years.

The amphipod *G. antarctica* showed a significant preference for the control artificial foods over the foods that contained 50% natural concentrations of *D. menziesii* extracts ($t_{16} = 2.4068$, $P = 0.0285$). On average, macroalgal discs containing *D. menziesii*

Table 1. Carbonate chemistry of the ambient and lowered pH treatments (mean \pm SD). Seawater parameters ($n = 8$) are calculated from total alkalinity (TA; $\mu\text{mol kg}^{-1}$ seawater, spectrophotometric pH_T (mean \pm SD), temperature ($^{\circ}\text{C}$) and salinity (ppt). Calculated parameters included pCO_2 (μatm) and saturation states of aragonite (Ω_{arg}) and calcite (Ω_{cal}).

	Year 1			Year 2		
	Ambient	pH 7.7	pH 7.3	Ambient	pH 7.7	pH 7.3
pH_T	8.07 \pm 0.07	7.69 \pm 0.11	7.32 \pm 0.08	8.02 \pm 0.06	7.73 \pm 0.08	7.35 \pm 0.08
TA	2284 \pm 78	2261 \pm 42	2245 \pm 64	2243 \pm 17	2248 \pm 29	2247 \pm 15
Temperature	2.45 \pm 0.34	2.46 \pm 0.34	2.46 \pm 0.33	2.61 \pm 0.40	2.65 \pm 0.43	2.65 \pm 0.42
Salinity	32.97 \pm 0.29	32.96 \pm 0.29	32.96 \pm 0.29	33.46 \pm 0.20	33.48 \pm 0.21	33.48 \pm 0.21
pCO_2	367 \pm 48	947 \pm 221	2245 \pm 409	408 \pm 48	856 \pm 145	2123 \pm 363
DIC	2145 \pm 56	2243 \pm 41	2343 \pm 72	2121 \pm 30	2218 \pm 37	2334 \pm 32
Ω_{arg}	1.62 \pm 0.37	0.73 \pm 0.21	0.32 \pm 0.07	1.45 \pm 0.22	0.79 \pm 0.16	0.34 \pm 0.08
Ω_{cal}	2.59 \pm 0.58	1.17 \pm 0.33	0.50 \pm 0.11	2.31 \pm 0.35	1.25 \pm 0.26	0.54 \pm 0.12

DIC = dissolved inorganic carbon.




Figure 1. Feeding preference of *Gondogeneia antarctica* grazing on macroalgal discs containing ground tissue of the macroalga *Cladophora repens* and extract of the macroalga *Desmarestia menziesii* ($n = 9$). Bars correspond to mean consumption \pm SE. The asterisk indicates a significant difference ($P < 0.05$) in a t -test.

extract were eaten 36% less than control discs (Fig. 1). The consumption of macroalgal discs with *D. menziesii* extract was found to be significantly higher than 0 using a one-sample t -test, meaning amphipods were eating discs with the extract ($t_8 = 4.746$, $P = 0.0015$).

Extract yields for *D. menziesii* ranged from 22.63 to 48.32 mg extract/g alga and did not significantly differ between the pH treatments. A preference for macroalgal discs containing extract from *D. menziesii* exposed to ambient seawater over extracts from macroalgae exposed to pH 7.7 and 7.3 was observed in *G. antarctica* (Fig. 2). A repeated-measures ANOVA showed a significant impact of pH on *D. menziesii* palatability ($F_{2,12} = 3.927$, $P = 0.0488$). Artificial food pellets made with extracts from the ambient pH treatment were consumed $\sim 38\%$ more than those made with extracts from macroalgae maintained at reduced pH (Bonferroni pairwise comparison, $P < 0.05$).

Gondogeneia antarctica had a strong preference for *P. decipiens* kept under ambient pH or pH 7.7 compared to macroalgae exposed to pH 7.3 ($F_{1,132,7.924} = 6.839$, $P = 0.0286$; Fig. 2). On average, pairwise comparisons revealed that consumption was two-fold lower in the pH 7.3 treatment compared to the other two treatments ($P < 0.05$).

Gondogeneia antarctica that were exposed to the different pH treatments for 8 weeks exhibited no significant difference in total




Figure 2. Feeding preference of the amphipod *Gondogeneia antarctica* grazing on macroalgae exposed to different pH treatments. **a.** Consumption of discs containing ground tissue of the macroalga *Cladophora repens* and extracts of the macroalga *Desmarestia menziesii* ($n = 7$). **b.** Consumption of discs of the macroalga *Palmaria decipiens* ($n = 8$). Bars correspond to mean consumption \pm SE. Letters indicate significant groups ($P < 0.05$) in a repeated-measures analysis of variance.

consumption of untreated *P. decipiens* (Fig. 3). Amphipods in each of the treatments consumed ~ 25 mg of *P. decipiens* over the 12 h of the experiment.

Discussion

The macroalgae *D. menziesii* and *P. decipiens* and the amphipod *G. antarctica* were exposed to three pH treatments simulating




Figure 3. Total consumption of discs of the macroalga *Palmaria decipiens* by the amphipod *Gondogeneia antarctica* kept under different pH treatments for 8 weeks ($n = 8$). Bars correspond to mean consumption \pm SE.

current ambient conditions (pH 8.1), end-of-the-century conditions (pH 7.7; IPCC 2022) and distant-future conditions (pH 7.3; IPCC 2022). These macroalgae were selected to represent a common unpalatable (*D. menziesii*) and palatable (*P. decipiens*) species (Amsler *et al.* 2005, Aumack *et al.* 2010). We found a reduction in palatability to (non-treated) *G. antarctica* in both *D. menziesii* extracts and *P. decipiens* discs when exposed to decreased pH. However, when pH-treated *G. antarctica* were offered untreated, identical foods we found no significant difference in total consumption between *G. antarctica* from any of the treatments.

Previous studies have shown that OA can have a direct impact on the physiology and behaviour of marine organisms (Ericson *et al.* 2010, Kroeker *et al.* 2013, Ramajo *et al.* 2016), especially in consumers (e.g. Kurihara & Shirayama 2004, Lim & Harley 2018, Park *et al.* 2020, Anand *et al.* 2021). In addition to these potential direct impacts on a species, the interactions between species, such as macroalgae-herbivore interactions, can also be altered. OA-induced changes in the growth and chemical composition of macroalgae can alter consumer preference for specific macroalgal species (Fieber & Bourdeau 2021). Additionally, differences in tolerance to low pH can shift entire ecosystem macroalgal composition, which can change the abundance of specific food items for consumers (Hall-Spencer *et al.* 2008, Harvey *et al.* 2021).

Fleshy macroalgae are generally more resistant to OA compared to calcifying species and can sometimes benefit from a decrease in pH (Hall-Spencer *et al.* 2008, Johnson *et al.* 2014, Kram *et al.* 2016). However, resistance to OA does not mean that these species are unaffected by this process. Changes in macroalgal palatability have been observed in several macroalgae when exposed to OA conditions (Swanson & Fox 2007, Poore *et al.* 2013, Duarte *et al.* 2016, Fieber & Bourdeau 2021). For example, the palatability of *Fucus vesiculosus* has been found to increase when exposed to low-pH treatments in summer through winter (Raddatz *et al.* 2017). Additionally, Poore *et al.* (2013) found that the palatability of *Sargassum linearifolium* was not impacted by decreases in pH alone. However, decreases in pH were able to mitigate reductions in palatability from increases in temperature.

Stress from changes in abiotic factors such as salinity, temperature and pH can alter secondary metabolite production in algae (Kolackova *et al.* 2023). Alterations in secondary metabolites used as chemical defences can impact the overall palatability of macroalgae. The response can range from decreased, increased or

unaffected chemical defence production between different species (Swanson & Fox 2007, Rich *et al.* 2018, Fieber & Bourdeau 2021). The kelp *Laminaria setchellii* and several species of the green algal genus *Halimeda* experience a decrease in phenolic and terpene content, respectively, when exposed to decreased pH (Campbell *et al.* 2014, Fieber & Bourdeau 2021). However, the amount of phlorotannins, a putative chemical defence, have been shown to increase in some kelps when exposed to high-CO₂ treatments (Swanson & Fox 2007).

In our study, *D. menziesii* extracts from an ambient pH were preferred over extracts from the pH 7.7 and pH 7.3 treatments. Since the extracts were lipophilic and therefore did not contain other desirable components to consumers, such as proteins, this preference for ambient-treatment extracts demonstrates that the potency of the extracts increased with decreased pH. These results suggest that OA could delay when the carbon from *D. menziesii* would become available to consumers in macroalgal communities along the WAP. The brown macroalga *D. menziesii* commonly dominates or co-dominates shallow-water communities down to 20 m in this region of the WAP (Amsler *et al.* 1995, 2024). Although this macroalga is typically not consumed while it is living, this macroalga and other sympatric species are important food resources for grazers when their chemical defences break down after death (Wiencke & Amsler 2012). However, an increase in the amount or the potency of the chemical defences in *D. menziesii* could increase the amount of time for which the macroalga would need to be dead before the chemical defences would break down sufficiently to become palatable to consumers, lengthening the time for their carbon to become available to higher trophic levels. This concept is supported by the findings that high CO₂ exposure and larger putative chemical defence levels have both been correlated with longer decay times in other brown macroalgae (Swanson & Fox 2007).

In addition to changes in chemical defences, shifts in chemical composition that lower nutritional quality have also been observed in OA experiments (e.g. Gordillo *et al.* 2001, Duarte *et al.* 2016). Decreases in protein and fatty acid concentrations have been observed in several macroalgal species when exposed to CO₂ enrichment (Garcia-Sanchez *et al.* 1994, Mercado *et al.* 1999, Gordillo *et al.* 2001, Rossoll *et al.* 2012, Duarte *et al.* 2016). However, these changes in composition vary with corresponding photosynthesis or growth changes for particular species. For example, decreases in protein content and increases in C:N ratios are correlated with lower growth and photosynthesis rates in *Porphyra leucostricta* and *Gracilaria tenuistipitata* (Garcia-Sanchez *et al.* 1994, Mercado *et al.* 1999). Gordillo *et al.* (2001), however, found similar composition changes but increases in growth and photosynthesis in *Ulva rigida* with CO₂ enrichment. These reductions in nutritional quality can result in lower palatability to consumers. Duarte *et al.* (2016) found that exposure to high CO₂ reduced protein content and organic matter in the brown macroalga *Durvillaea antarctica*. As a result, the amphipod *Orchestoidea tuberculata* preferred *D. antarctica* from the ambient CO₂ treatment three-fold more than macroalgae from the higher CO₂ treatment (Duarte *et al.* 2016).

Our study found that the palatability of *P. decipiens* decreases by half when exposed to a pH 7.3 treatment for 18 days. This decrease in palatability could be attributed to a decrease in nutritional quality of the macroalga or to the production of an unpalatable metabolite as a response to stress. The nutritional quality of primary producers can influence ecosystem trophic structure by determining herbivory pressure, consumer to producer biomass

ratios and the rate at which energy is cycled through the food web (Cebrian *et al.* 2009). *Palmaria decipiens* is one of the few palatable macroalgae in these communities (Amsler *et al.* 2020). A reduction in nutritional quality could lead to heavier herbivory pressure on this macroalga if grazers must consume more macroalgae to compensate for low-nutrition food items (Duarte *et al.* 2016). Furthermore, reductions in nutritional quality can impact consumers through shifting biochemical compositions, decreasing egg production and reducing growth rates in some invertebrates (Urabe *et al.* 2003, Rossoll *et al.* 2012, Schoo *et al.* 2013).

Previous studies have shown that invertebrates can be susceptible to OA through alterations in reproduction (Kurihara & Shirayama 2004, Ericson *et al.* 2010, Morita *et al.* 2010), development (Kurihara & Shirayama 2004, Watson *et al.* 2009, Place & Smith 2012) and growth (Sheppard Brennard *et al.* 2010). In addition to these changes, the energy demands of invertebrates can also shift as organisms reallocate available energy to different processes as a response to low pH exposure (Pan *et al.* 2015). One common response of invertebrates to higher energy demands is increasing food intake (Saba *et al.* 2012, Ramajo *et al.* 2016). Saba *et al.* (2012) found that Antarctic krill increase their feeding rates greater than three-fold and increase their metabolic rates when exposed to high CO₂ concentrations. In some situations, having access to greater food amounts can mitigate some of the negative effects of OA exposure. For example, Melzner *et al.* (2011) found that inner-shell corrosion in the mussel *Mytilus edulis* was not observed in moderate CO₂ treatments when given a high volume of food. Similarly, the hard coral *Acropora cervicornis* can utilize increased feeding to mitigate decreases in calcification and growth from high CO₂ exposure (Towle *et al.* 2015). In contrast to other studies, Ramajo *et al.* (2016) found that juvenile scallops exposed to high CO₂ had significantly higher metabolic, growth, calcification and feeding rates compared to when exposed to ambient CO₂. However, scallops with a low food supply exposed to the same pCO₂ altered the composition of their shell-periostracum layer.

Amphipods utilize calcium carbonate to create their exoskeletons (Luquet 2012), making them susceptible to increased energetic requirements under OA to maintain their calcification. However, in the present study, we found no significant difference in the total consumption of *G. antarctica* of food discs with decreasing pH. This result aligns with the reported robustness of arthropod feeding, as a whole, to OA compared to more susceptible groups, including echinoderms and molluscs (Clements & Darrow 2018). Other OA studies found that the larvae of the Dungeness crab, *Metacarcinus magister*, and the Pacific green shore crab, *Hemigrapsus oregonensis*, showed no difference in feeding rates on prey items when exposed to lowered pH (Christmas 2013). Similar trends in feeding rates on algae have also been observed in some krill, shrimp, amphipods and copepods (Arnberg *et al.* 2013, Poore *et al.* 2013, Li *et al.* 2015, Cooper *et al.* 2016, Glandon & Miller 2017, McLaskey *et al.* 2019). However, while measuring feeding responses can be a tool for estimating energy acquisition (Clements & Darrow 2018), feeding cannot measure all processes that might be impacted by OA. For example, there can be changes in respiration, fatty acid accumulation efficiency and development rates that are not reflected in consumption rates (Arnberg *et al.* 2013, Li *et al.* 2015, McLaskey *et al.* 2019). Consequently, any such impacts on *G. antarctica* could have gone undetected in our study. In addition, *P. decipiens* is one of the most protein-rich macroalgae in this region (Peters *et al.* 2005), making it possible that the untreated *P. decipiens* used as food in the experiment could have provided

enough nutrition to mask possible differences in energy demands of *G. antarctica*.

The results of the present study show that OA negatively affected the palatability of the unpalatable *D. menziesii* and the palatable *P. decipiens* but did not impact the feeding of *G. antarctica*. These changes in palatability are probably occurring because of changes in either nutritional quality or secondary metabolite production. Alterations in either of these components could lead to major impacts for the WAP food web, such as delaying when an unpalatable macroalga enters the detrital food web or decreasing the nutritional quality of a palatable macroalga for consumers. Overall, our results suggest that the indirect effects of OA on food items can be more impactful than the direct effects on amphipods along the WAP.

Supplementary material. To view supplementary material for this article, please visit <http://doi.org/10.1017/S095410202400052X>.

Details of data deposit. Data are available at the US Antarctic Program Data Center: <https://www.usap-dc.org/view/project/p0010193>.

Acknowledgements. The authors gratefully acknowledge the science and logistical support staff of Antarctic Support Contract for their help and support of the US Antarctic Program. Addie Knight and Jami de Jesus provided valuable field assistance. Stacy Krueger-Hadfield and two reviewers provided constructive comments on earlier drafts of the manuscript.

Financial support. This research was supported by National Science Foundation award OPP- 1848887 from the Antarctic Organisms and Ecosystems Program.

Competing interests. The authors declare none.

Author contributions. HEO, CDA, JBM and JBS designed the study. HEO, MOA, CDA and JBS collected the macroalgae. HEO, MOA and CDA collected the amphipods. All authors conducted the experiment. HEO analysed the data, prepared the figures and wrote the first draft of the manuscript. All authors contributed to editing the final manuscript.

References

- AHN, I.-Y., ELIAS-PIERA, F., HA, S.-Y., ROSSI, S. & KIM, D.-U. 2021. Seasonal dietary shifts of the gammarid amphipod *Gondogeneia antarctica* in a rapidly warming fjord of the West Antarctic Peninsula. *Journal of Marine Science and Engineering*, **9**, 10.3390/jmse9121447.
- AMSLER, C.D., MCCLINTOCK, J.B. & BAKER, B.J. 2008. Macroalgal chemical defenses in polar marine communities. In AMSLER, C.D., ed., *Algal chemical ecology*. Berlin: Springer Berlin Heidelberg, 91–103.
- AMSLER, C.D., MCCLINTOCK, J.B. & BAKER, B.J. 2014. Chemical mediation of mutualistic interactions between macroalgae and mesograzers structure unique coastal communities along the western Antarctic Peninsula *Journal of Phycology*, **50**, 10.1111/jpy.12137.
- AMSLER, C.D., MCCLINTOCK, J.B. & BAKER, B.J. 2020. Chemical mediation of Antarctic macroalga-grazer interactions. In GÓMEZ, I. & HUOVINEN, P., eds, *Antarctic seaweeds: diversity, adaptation and ecosystem services*. Cham: Springer International Publishing, 339–363.
- AMSLER, C.D., ROWLEY, R.J., LAUR, D.R., QUETIN, L.B. & ROSS, R.M. 1995. Vertical distribution of Antarctic peninsular macroalgae: cover, biomass and species composition. *Phycologia*, **34**, 10.2216/i0031-8884-34-5-424.1.
- AMSLER, C.D., AMSLER, M.O., HEISER, S., MCCLINTOCK, J.B., IKEN, K., GALLOWAY, A.W.E. & KLEIN, A.G. 2024. Vertical distribution of brown and red macroalgae along the central western Antarctic Peninsula. *Botanica Marina*, **67**, 10.1515/bot-2023-0085.
- AMSLER, C.D., MILLER, L.R., EDWARDS, R.A., AMSLER, M.O., ENGL, W., MCCLINTOCK, J.B. & BAKER, B.J. 2022. Gastropod assemblages associated with *Himantothallus grandifolius*, *Sarcopeltis antarctica* and other subtidal macroalgae. *Antarctic Science*, **34**, 10.1017/S0954102022000153.

- AMSLER, C.D., IKEN, K., MCCLINTOCK, J.B., AMSLER, M., PETERS, K., HUBBARD, J., et al. 2005. Comprehensive evaluation of the palatability and chemical defenses of subtidal macroalgae from the Antarctic Peninsula. *Marine Ecology Progress Series*, **294**, 10.3354/meps294141.
- ANAND, M., RANGESH, K., MARUTHUPANDY, M., JAYANTHI, G., RAJESWARI, B. & PRIYA, R.J. 2021. Effect of CO₂ driven ocean acidification on calcification, physiology and ovarian cells of tropical sea urchin *Salmacis virgulata* - a microcosm approach. *Heliyon*, **7**, 10.1016/j.heliyon.2021.e05970.
- ANDERSSON, A.J., MACKENZIE, F.T. & GATTUSO, J.-P. 2011. Effects of ocean acidification on benthic processes, organisms, and ecosystems. In GATTUSO, J.-P. & HANSSON, L. eds, *Ocean acidification*. Oxford: Oxford University Press, 121–153.
- ARMSTRONG, R.A. 2017. Recommendations for analysis of repeated-measures designs: testing and correcting for sphericity and use of MANOVA and mixed model analysis. *Ophthalmic and Physiological Optics*, **37**, 10.1111/opo.12399.
- ARNBERG, M., CALOSI, P., SPICER, J.I., TANDBERG, A.H.S., NILSEN, M., WESTERLUND, S. & BECHMANN, R.K. 2013. Elevated temperature elicits greater effects than decreased pH on the development, feeding and metabolism of northern shrimp (*Pandalus borealis*) larvae. *Marine Biology*, **160**, 10.1007/s00227-012-2072-9.
- AUMACK, C.F., AMSLER, C.D., MCCLINTOCK, J.B. & BAKER, B.J. 2010. Chemically mediated resistance to mesoherbivory in finely branched macroalgae along the western Antarctic Peninsula. *European Journal of Phycology*, **45**, 10.1080/09670260903171668.
- BHUIYAN, M.K.A., RODRÍGUEZ, B.M., BILLAH, M.M., PIRES, A., FREITAS, R. & CONRADI, M. 2022. Effects of ocean acidification on the biochemistry, physiology and parental transfer of *Ampelisca brevicornis* (Costa, 1853). *Environmental Pollution*, **293**, 10.1016/j.envpol.2021.118549.
- BORGES, F.O., FIGUEIREDO, C., SAMPAIO, E., ROSA, R. & GRILO, T.F. 2018. Trans-generational deleterious effects of ocean acidification on the reproductive success of a keystone crustacean (*Gammarus locusta*). *Marine Environmental Research*, **138**, 10.1016/j.marenvres.2018.04.006.
- CAMPBELL, J., CRAFT, J., MUEHLEHNER, N., LANGDON, C. & PAUL, V. 2014. Responses of calcifying algae (*Halimeda* spp.) to ocean acidification: implications for herbivores. *Marine Ecology Progress Series*, **514**, 10.3354/meps10981.
- CEBRIAN, J., SHURIN, J.B., BORER, E.T., CARDINALE, B.J., NGAL, J.T., SMITH, M.D. & FAGAN, W.F. 2009. Producer nutritional quality controls ecosystem trophic structure. *PLoS ONE*, **4**, 10.1371/journal.pone.0004929.
- CHRISTMAS, A.-M.F. 2013. *Effects of ocean acidification on dispersal behavior in the larval stage of the Dungeness crab and the Pacific green shore crab*. PhD thesis. Western Washington University, 72 pp.
- CLEMENTS, J.C. & DARROW, E.S. 2018. Eating in an acidifying ocean: a quantitative review of elevated CO₂ effects on the feeding rates of calcifying marine invertebrates. *Hydrobiologia*, **820**, 10.1007/s10750-018-3665-1.
- COOPER, H.L., POTTS, D.C. & PAYTAN, A. 2016. Metabolic responses of the North Pacific krill, *Euphausia pacifica*, to short- and long-term pCO₂ exposure. *Marine Biology*, **163**, 10.1007/s00227-016-2982-z.
- DICKSON, A.G. 1990. Standard potential of the reaction and the standard acidity constant of the ion HSO⁴⁻ in synthetic sea water from 273.15 to 318.15 K. *Journal of Chemical Thermodynamics*, **22**, 10.1016/0021-9614(90)90074-Z.
- DICKSON, A.G., SABINE, C.L. & CHRISTIAN, J.R., eds. 2007. *Guide to best practices for ocean CO₂ measurements*. Sidney, BC: North Pacific Marine Science Organization, 191 pp.
- DONEY, S.C., FABRY, V.J., FEELY, R.A. & KLEYPAS, J.A. 2009. Ocean acidification: the other CO₂ problem. *Annual Review of Marine Science*, **1**, 10.1146/annurev.marine.010908.163834.
- DUARTE, C., LÓPEZ, J., BENÍTEZ, S., MANRÍQUEZ, P.H., NAVARRO, J.M., BONTA, C.C., et al. 2016. Ocean acidification induces changes in algal palatability and herbivore feeding behavior and performance. *Oecologia*, **180**, 10.1007/s00442-015-3459-3.
- DUNTON, K.H. 2001. δ¹⁵N and δ¹³C measurements of Antarctic Peninsula fauna: trophic relationships and assimilation of benthic seaweeds. *American Zoologist*, **41**, 10.1093/icb/41.1.99.
- ERICSON, J.A., LAMARE, M.D., MORLEY, S.A. & BARKER, M.F. 2010. The response of two ecologically important Antarctic invertebrates (*Sterechnus neumayeri* and *Parborlasia corrugatus*) to reduced seawater pH: effects on fertilisation and embryonic development. *Marine Biology*, **157**, 10.1007/s00227-010-1529-y.
- FIEBER, A.M. & BOURDEAU, P.E. 2021. Elevated pCO₂ reinforces preference among intertidal algae in both a specialist and generalist herbivore. *Marine Pollution Bulletin*, **168**, 10.1016/j.marpolbul.2021.112377.
- FOX, J. & WEISBERG, S. 2019. An R Companion to Applied Regression. Retrieved from <https://socialsciences.mcmaster.ca/jfox/Books/Companion/>
- GARCIA-SANCHEZ, M.J., FERNANDEZ, J.A. & NIELL, X. 1994. Effect of inorganic carbon supply on the photosynthetic physiology of *Gracilaria tenuistipitata*. *Planta*, **194**, 10.1007/BF00201034.
- GATTUSO, J.-P., EPITALON, J.-M., LAVIGNE, H. & ORR, J. 2022. *Seacarb*: seawater carbonate chemistry. Retrieved from <https://CRAN.R-project.org/package=seacarb>
- GLANDON, H.L. & MILLER, T.J. 2017. No effect of high pCO₂ on juvenile blue crab, *Callinectes sapidus*, growth and consumption despite positive responses to concurrent warming. *ICES Journal of Marine Science*, **74**, 10.1093/icesjms/fsw171.
- GORDILLO, F.J.L., NIELL, F.X. & FIGUEROA, F.L. 2001. Non-photosynthetic enhancement of growth by high CO₂ level in the nitrophilic seaweed *Ulva rigida* C. Agardh (Chlorophyta). *Planta*, **213**, 10.1007/s004250000468.
- GREENHOUSE, S.W. & GEISSER, S. 1959. On methods in the analysis of profile data. *Psychometrika*, **24**, 10.1007/BF02289823.
- GRUBER, N., CLEMENT, D., CARTER, B.R., FEELY, R.A., VAN HEUVEN, S., HOPPEMA, M., et al. 2019. The oceanic sink for anthropogenic CO₂ from 1994 to 2007. *Science*, **363**, 10.1126/science.aau5153.
- GUIRY, M.D. 2024. How many species of algae are there? A reprise. Four kingdoms, 14 phyla, 63 classes and still growing. *Journal of Phycology*, **60**, 10.1111/jpy.13431.
- HALL-SPENCER, J.M., RODOLFO-METALPA, R., MARTIN, S., RANSOME, E., FINE, M., TURNER, S.M., et al. 2008. Volcanic carbon dioxide vents show ecosystem effects of ocean acidification. *Nature*, **454**, 10.1038/nature07051.
- HARVEY, B.P., KON, K., AGOSTINI, S., WADA, S. & HALL-SPENCER, J.M. 2021. Ocean acidification locks algal communities in a species-poor early successional stage. *Global Change Biology*, **27**, 10.1111/gcb.15455.
- HAY, M.E., KAPPEL, Q.E. & FENICAL, W. 1994. Synergisms in plant defenses against herbivores: interactions of chemistry, calcification, and plant quality. *Ecology*, **75**, 10.2307/1939631.
- HEEDE, R. 2014. Tracing anthropogenic carbon dioxide and methane emissions to fossil fuel and cement producers, 1854–2010. *Climatic Change*, **122**, 10.1007/s10584-013-0986-y.
- HUANG, Y.M., AMSLER, M.O., MCCLINTOCK, J.B., AMSLER, C.D. & BAKER, B.J. 2007. Patterns of gammaridean amphipod abundance and species composition associated with dominant subtidal macroalgae from the western Antarctic Peninsula. *Polar Biology*, **30**, 10.1007/s00300-007-0303-1.
- HURD, C.L., HEPBURN, C.D., CURRIE, K.I., RAVEN, J.A. & HUNTER, K.A. 2009. Testing the effects of ocean acidification on algal metabolism: considerations for experimental designs. *Journal of Phycology*, **45**, 10.1111/j.1529-8817.2009.00768.x.
- IKEN, K. 1999. Feeding ecology of the Antarctic herbivorous gastropod *Laevilacunaria antarctica* Martens. *Journal of Experimental Marine Biology and Ecology*, **236**, 10.1016/S0022-0981(98)00199-3.
- IKEN, K., AMSLER, C.D., GORMAN, K.B., KLEIN, A.G., GALLOWAY, A.W.E., AMSLER, M.O., et al. 2023. Macroalgal input into the coastal food web along a gradient of seasonal sea ice cover along the western Antarctic Peninsula. *Marine Ecology Progress Series*, **718**, 10.3354/meps14388.
- IPCC. 2022. *Climate change 2022: impacts, adaptation and vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge and New York: Cambridge University Press, 10.1017/9781009325844.
- JAREK, S. 2012. *mvnormtest*: normality test for multivariate variables. Retrieved from <https://CRAN.R-project.org/package=mvnormtest>
- JOHNSON, M.D., PRICE, N.N. & SMITH, J.E. 2014. Contrasting effects of ocean acidification on tropical fleshy and calcareous algae. *PeerJ*, **2**, 10.7717/peerj.411.
- JOOS, F. & SPEAHN, R. 2008. Rates of change in natural and anthropogenic radiative forcing over the past 20,000 years. *Proceedings of the National Academy of Sciences of the United States of America*, **105**, 10.1073/pnas.0707386105.

- KINDINGER, T.L., TOY, J.A. & KROEKER, K.J. 2022. Emergent effects of global change on consumption depend on consumers and their resources in marine systems. *Proceedings of the National Academy of Sciences of the United States of America*, **119**, 10.1073/pnas.2108878119.
- KINNBY, A., WHITE, J.C.B., TOTH, G.B. & PAVIA, H. 2021. Ocean acidification decreases grazing pressure but alters morphological structure in a dominant coastal seaweed. *PLoS ONE*, **16**, 10.1371/journal.pone.0245017.
- KOCH, M., BOWES, G., ROSS, C. & ZHANG, X.-H. 2013. Climate change and ocean acidification effects on seagrasses and marine macroalgae. *Global Change Biology*, **19**, 10.1111/j.1365-2486.2012.02791.x.
- KOLACKOVA, M., JANOVA, A., DOBESOVA, M., ZVALOVA, M., CHALOUPEK, P., KRYSOFOVA, O., *et al.* 2023. Role of secondary metabolites in distressed microalgae. *Environmental Research*, **224**, 10.1016/j.envres.2023.115392.
- KRAM, S.L., PRICE, N.N., DONHAM, E.M., JOHNSON, M.D., KELLY, E.L.A., HAMILTON, S.L. & SMITH, J.E. 2016. Variable responses of temperate calcified and fleshy macroalgae to elevated pCO₂ and warming. *ICES Journal of Marine Science*, **73**, 10.1093/icesjms/fsv168.
- KROEKER, K.J., KORDAS, R.L., CRIM, R., HENDRIKS, I.E., RAMAJO, L., SINGH, G.S., *et al.* 2013. Impacts of ocean acidification on marine organisms: quantifying sensitivities and interaction with warming. *Global Change Biology*, **19**, 10.1111/gcb.12179.
- KURIHARA, H. & SHIRAYAMA, Y. 2004. Effects of increased atmospheric CO₂ on sea urchin early development. *Marine Ecology Progress Series*, **274**, 10.3354/meps274161.
- LAN, X. & KEELING, R. 2024. Trends in atmospheric carbon dioxide (CO₂). *Global Monitoring Laboratory*. Retrieved from <https://gml.noaa.gov/ccgg/trends/data.html>
- LAWRENCE, M.A. 2016. *ez*: easy analysis and visualization of factorial experiments. R package version 4.4-0. Retrieved from <https://CRAN.R-project.org/package=ez>
- LEUNG, J.Y.S., ZHANG, S. & CONNELL, S.D. 2022. Is ocean acidification really a threat to marine calcifiers? A systematic review and meta-analysis of 980+ studies spanning two decades. *Small*, **18**, 10.1002/sml.202107407.
- LEUNG, J.Y.S., NAGELKERKEN, I., RUSSELL, B.D., FERREIRA, C.M. & CONNELL, S.D. 2018. Boosted nutritional quality of food by CO₂ enrichment fails to offset energy demand of herbivores under ocean warming, causing energy depletion and mortality. *Science of the Total Environment*, **639**, 10.1016/j.scitotenv.2018.05.161.
- LI, W., HAN, G., DONG, Y., ISHIMATSU, A., RUSSELL, B.D. & GAO, K. 2015. Combined effects of short-term ocean acidification and heat shock in a benthic copepod *Tigriopus japonicus* Mori. *Marine Biology*, **162**, 10.1007/s00227-015-2722-9.
- LI-BEISSON, Y., THELEN, J.J., FEDOSEJEVS, E. & HARWOOD, J.L. 2019. The lipid biochemistry of eukaryotic algae. *Progress in Lipid Research*, **74**, 10.1016/j.plipres.2019.01.003.
- LIM, E.G. & HARLEY, C.D.G. 2018. Caprellid amphipods (*Caprella* spp.) are vulnerable to both physiological and habitat-mediated effects of ocean acidification. *PeerJ*, **6**, 10.7717/peerj.5327.
- LIU, Z., CIAIS, P., DENG, Z., LEI, R., DAVIS, S.J., FENG, S., *et al.* 2020. Near-real-time monitoring of global CO₂ emissions reveals the effects of the COVID-19 pandemic. *Nature Communications*, **11**, 10.1038/s41467-020-18922-7.
- LOPES, A.R., BORGES, F.O., FIGUEIREDO, C., SAMPAIO, E., DINIZ, M., ROSA, R. & GRILO, T.F. 2019. Transgenerational exposure to ocean acidification induces biochemical distress in a keystone amphipod species (*Gammarus locusta*). *Environmental Research*, **170**, 10.1016/j.envres.2018.12.040.
- LUEKER, T.J., DICKSON, A.G. & KEELING, C.D. 2000. Ocean pCO₂ calculated from dissolved inorganic carbon, alkalinity, and equations for K₁ and K₂: validation based on laboratory measurements of CO₂ in gas and seawater at equilibrium. *Marine Chemistry*, **70**, 10.1016/S0304-4203(00)00022-0.
- LUQUET, G. 2012. Biomineralizations: insights and prospects from crustaceans. *ZooKeys*, **176**, 10.3897/zookeys.176.2318.
- MALHI, Y., MEIR, P. & BROWN, S. 2002. Forests, carbon and global climate. *Philosophical Transactions of the Royal Society of London. Series A: Mathematical, Physical and Engineering Sciences*, **360**, 10.1098/rsta.2002.1020.
- MCDOWELL, R.E., AMSLER, C.D., DICKINSON, D.A., MCCLINTOCK, J.B. & BAKER, B.J. 2014. Reactive oxygen species and the Antarctic macroalgal wound response. *Journal of Phycology*, **50**, 10.1111/jpy.12127.
- MCLASKEY, A.K., KEISTER, J.E., SCHOO, K.L., OLSON, M.B. & LOVE, B.A. 2019. Direct and indirect effects of elevated CO₂ are revealed through shifts in phytoplankton, copepod development, and fatty acid accumulation. *PLoS ONE*, **14**, 10.1371/journal.pone.0213931.
- MELZNER, F., STANGE, P., TRÜBENBACH, K., THOMSEN, J., CASTIES, I., PANKNIN, U., *et al.* 2011. Food supply and seawater pCO₂ impact calcification and internal shell dissolution in the blue mussel *Mytilus edulis*. *PLoS ONE*, **6**, 10.1371/journal.pone.0024223.
- MERCADO, J.M., JAVIER, F., GORDILLO, L., XAVIER NIELL, F. & FIGUEROA, F.L. 1999. Effects of different levels of CO₂ on photosynthesis and cell components of the red alga *Porphyra leucosticta*. *Journal of Applied Phycology*, **11**, 10.1023/A:1008194223558.
- MORITA, M., SUWA, R., IGUCHI, A., NAKAMURA, M., SHIMADA, K., SAKAI, K. & SUZUKI, A. 2010. Ocean acidification reduces sperm flagellar motility in broadcast spawning reef invertebrates. *Zygote*, **18**, 10.1017/S0967199409990177.
- NISSEN, C., LOVENDUSKI, N.S., BROOKS, C.M., HOPPEMA, M., TIMMERMANN, R. & HAUCK, J. 2024. Severe 21st-century ocean acidification in Antarctic Marine Protected Areas. *Nature Communications*, **15**, 10.1038/s41467-023-44438-x.
- PAGE, H.N., BAHR, K.D., CYRONAK, T., JEWETT, E.B., JOHNSON, M.D. & MCCOY, S.J. 2022. Responses of benthic calcifying algae to ocean acidification differ between laboratory and field settings. *ICES Journal of Marine Science*, **79**, 10.1093/icesjms/fsab232.
- PAINE, E.R., BRITTON, D., SCHMID, M., BREWER, E.A., DIAZ-PULIDO, G., BOYD, P.W. & HURD, C.L. 2023. No effect of ocean acidification on growth, photosynthesis, or dissolved organic carbon release by three temperate seaweeds with different dissolved inorganic carbon uptake strategies. *ICES Journal of Marine Science*, **80**, 10.1093/icesjms/fsac221.
- PALMER STATION INSTRUMENT TECHNICIAN. 2023. Palmer Station Water-wall data, 2016-present (ongoing). Retrieved from <https://amrddata.ssec.wisc.edu/dataset/palmer-station-waterwall-data>
- PAN, T.-C.F., APPLEBAUM, S.L. & MANAHAN, D.T. 2015. Experimental ocean acidification alters the allocation of metabolic energy. *Proceedings of the National Academy of Sciences of the United States of America*, **112**, 10.1073/pnas.1416967112.
- PARASCHIV, S. & PARASCHIV, L.S. 2020. Trends of carbon dioxide (CO₂) emissions from fossil fuels combustion (coal, gas and oil) in the EU Member States from 1960 to 2018. *Energy Reports*, **6**, 10.1016/j.egy.2020.11.116.
- PARK, S., AHN, I.-Y., SIN, E., SHIM, J. & KIM, T. 2020. Ocean freshening and acidification differentially influence mortality and behavior of the Antarctic amphipod *Gondogeneia antarctica*. *Marine Environmental Research*, **154**, 10.1016/j.marenvres.2019.104847.
- PEREZ, F.F. & FRAGA, F. 1987. Association constant of fluoride and hydrogen ions in seawater. *Marine Chemistry*, **21**, 16 10.1016/0304-4203(87)90036-3.
- PETERS, K.J., AMSLER, C.D., AMSLER, M.O., MCCLINTOCK, J.B., DUNBAR, R.B. & BAKER, B.J. 2005. A comparative analysis of the nutritional and elemental composition of macroalgae from the western Antarctic Peninsula. *Phycologia*, **44**, 10.2216/0031-8884(2005)44[453:ACAOTN]2.0.CO;2.
- PLACE, S.P. & SMITH, B.W. 2012. Effects of seawater acidification on cell cycle control mechanisms in *Strongylocentrotus purpuratus* embryos. *PLoS ONE*, **7**, 10.1371/journal.pone.0034068.
- POORE, A.G.B., GRABA-LANDRY, A., FAVRET, M., SHEPPARD BRENNAND, H., BYRNE, M. & DWORJANYN, S.A. 2013. Direct and indirect effects of ocean acidification and warming on a marine plant-herbivore interaction. *Oecologia*, **173**, 10.1007/s00442-013-2683-y.
- QUARTINO, M.L. & BORASO DE ZAIXSO, A.L. 2008. Summer macroalgal biomass in Potter Cove, South Shetland Islands, Antarctica: its production and flux to the ecosystem. *Polar Biology*, **31**, 10.1007/s00300-007-0356-1.
- R CORE TEAM. 2020. R: a language and environment for statistical computing. R Foundation for Statistical Computing. Retrieved from <https://www.R-project.org/>
- RADDATZ, S., GUY-HAIM, T., RILOV, G. & WAHL, M. 2017. Future warming and acidification effects on anti-fouling and anti-herbivory traits of the brown alga *Fucus vesiculosus* (Phaeophyceae). *Journal of Phycology*, **53**, 10.1111/jpy.12473.
- RAMAJO, L., MARBÀ, N., PRADO, L., PERON, S., LARDIES, M.A., RODRIGUEZ-NAVARRO, A.B., *et al.* 2016. Biomineralization changes with food supply

- confer juvenile scallops (*Argopecten purpuratus*) resistance to ocean acidification. *Global Change Biology*, **22**, 10.1111/gcb.13179.
- RICH, W.A., SCHUBERT, N., SCHLÄPFER, N., CARVALHO, V.F., HORTA, A.C.L. & HORTA, P.A. 2018. Physiological and biochemical responses of a coralline alga and a sea urchin to climate change: implications for herbivory. *Marine Environmental Research*, **142**, 10.1016/j.marenvres.2018.09.026.
- ROA, R. 1992. Design and analysis of multiple-choice feeding-preference experiments. *Oecologia*, **89**, 509–515.
- ROBBINS, L.L., HANSEN, M.E., KLEYPAS, J.A. & MEYLAN, S.C. 2010. *CO2calc*: a user-friendly seawater carbon calculator for Windows, Max OS X, and iOS (iPhone). Retrieved from <https://pubs.usgs.gov/publication/ofr20101280>
- ROSSOLL, D., BERMÚDEZ, R., HAUSS, H., SCHULZ, K.G., RIEBESELL, U., SOMMER, U. & WINDER, M. 2012. Ocean acidification-induced food quality deterioration constrains trophic transfer. *PLoS ONE*, **7**, 10.1371/journal.pone.0034737.
- ROY, R.N., ROY, L.N., VOGEL, K.M., PORTER-MOORE, C., PEARSON, T., GOOD, C.E., *et al.* 1993. The dissociation constants of carbonic acid in seawater at salinities 5 to 45 and temperatures 0 to 45°C. *Marine Chemistry*, **44**, 10.1016/0304-4203(93)90207-5.
- SABA, G.K., SCHOFIELD, O., TORRES, J.J., OMBRES, E.H. & STEINBERG, D.K. 2012. Increased feeding and nutrient excretion of adult Antarctic krill, *Euphausia superba*, exposed to enhanced carbon dioxide (CO₂). *PLoS ONE*, **7**, 10.1371/journal.pone.0052224.
- SABINE, C.L., FEELY, R.A., GRUBER, N., KEY, R.M., LEE, K., BULLISTER, J.L., *et al.* 2004. The oceanic sink for anthropogenic CO₂. *Science*, **305**, 10.1126/science.1097403.
- SCHLENGER, A.J., BEAS-LUNA, R. & AMBROSE, R.F. 2021. Forecasting ocean acidification impacts on kelp forest ecosystems. *PLOS ONE*, **16**, 10.1371/journal.pone.0236218.
- SCHOENROCK, K.M., SCHRAM, J.B., AMSLER, C.D., MCCLINTOCK, J.B., ANGUS, R.A. & VOHRA, Y.K. 2016. Climate change confers a potential advantage to fleshy Antarctic crustose macroalgae over calcified species. *Journal of Experimental Marine Biology and Ecology*, **474**, 10.1016/j.jembe.2015.09.009.
- SCHOO, K.L., MALZAHN, A.M., KRAUSE, E. & BOERSMA, M. 2013. Increased carbon dioxide availability alters phytoplankton stoichiometry and affects carbon cycling and growth of a marine planktonic herbivore. *Marine Biology*, **160**, 10.1007/s00227-012-2121-4.
- SCHRAM, J.B., AMSLER, M.O., AMSLER, C.D., SCHOENROCK, K.M., MCCLINTOCK, J.B. & ANGUS, R.A. 2016. Antarctic crustacean grazer assemblages exhibit resistance following exposure to decreased pH. *Marine Biology*, **163**, 10.1007/s00227-016-2894-y.
- SCHUBERT, N., ALVAREZ-FILIP, L. & HOFMANN, L.C. 2023. Systematic review and meta-analysis of ocean acidification effects in *Halimeda*: implications for algal carbonate production. *Climate Change Ecology*, **4**, 10.1016/j.ecochg.2022.100059.
- SEPÚLVEDA, F., QUIJÓN, P.A., QUINTANILLA-AHUMADA, D., VARGAS, J., ALDANA, M., FERNÁNDEZ, M., *et al.* 2024. Cross-examining the influence of upwelling and seaweed quality on herbivores' feeding behavior and growth. *Marine Environmental Research*, **193**, 10.1016/j.marenvres.2023.106288.
- SHEPPARD BRENNAND, H., SOARS, N., DWORJANYN, S.A., DAVIS, A.R. & BYRNE, M. 2010. Impact of ocean warming and ocean acidification on larval development and calcification in the sea urchin *Tripneustes gratilla*. *PLoS ONE*, **5**, 10.1371/journal.pone.0011372.
- SUDO, H. & YOSHIDA, G. 2021. Effects of a reduction in algal nitrogen content on survival, growth, and reproduction of an herbivorous amphipod. *Journal of Experimental Marine Biology and Ecology*, **539**, 10.1016/j.jembe.2021.151543.
- SWANSON, A.K. & FOX, C.H. 2007. Altered kelp (Laminariales) phlorotannins and growth under elevated carbon dioxide and ultraviolet-B treatments can influence associated intertidal food webs. *Global Change Biology*, **13**, 10.1111/j.1365-2486.2007.01384.x.
- TOWLE, E.K., ENOCHS, I.C. & LANGDON, C. 2015. Threatened Caribbean coral is able to mitigate the adverse effects of ocean acidification on calcification by increasing feeding rate. *PLoS ONE*, **10**, 10.1371/journal.pone.0139398.
- TULLY, B.J. 2019. Metabolic diversity within the globally abundant Marine Group II Euryarchaea offers insight into ecological patterns. *Nature Communications*, **10**, 10.1038/s41467-018-07840-4.
- URABE, J., TOGARI, J. & ELSER, J.J. 2003. Stoichiometric impacts of increased carbon dioxide on a planktonic herbivore. *Global Change Biology*, **9**, 10.1046/j.1365-2486.2003.00634.x.
- WATSON, S.-A., SOUTHGATE, P.C., TYLER, P.A. & PECK, L.S. 2009. Early larval development of the Sydney rock oyster *Saccostrea glomerata* under near-future predictions of CO₂. *Journal of Shellfish Research*, **28**, 10.2983/035.028.0302.
- WEYKAM, G., THOMAS, D.N. & WIENCKE, C. 1997. Growth and photosynthesis of the Antarctic red algae *Palmaria decipiens* (Palmariales) and *Iridaea cordata* (Gigartinales) during and following extended periods of darkness. *Phycologia*, **36**, 10.2216/i0031-8884-36-5-395-1.
- WIENCKE, C. 1990. Seasonality of red and green macroalgae from Antarctica - a long-term culture study under fluctuating Antarctic daylengths. *Polar Biology*, **10**, 10.1007/BF00239371.
- WIENCKE, C. & AMSLER, C.D. 2012. Seaweeds and their communities in polar regions. In WIENCKE, C. & BISCHOF, K., eds. *Seaweed biology. Ecological Studies*. Berlin: Springer, 265–291.
- WIENCKE, C., AMSLER, C.D. & CLAYTON, M.N. 2014. Macroalgae. In DE BROYER, C., KOUUBI, P., GRIFFITHS, H.J., RAYMOND, B., D'UDEKEM D'ACOS, C., VAN DE PUTTE, A., eds. *Biogeographic atlas of the Southern Ocean*. Cambridge: Scientific Committee on Antarctic Research, 66–73.