



COMPOSITIO MATHEMATICA

Integral canonical models for Spin Shimura varieties

Keerthi Madapusi Pera

Compositio Math. **152** (2016), 769–824.

[doi:10.1112/S0010437X1500740X](https://doi.org/10.1112/S0010437X1500740X)



FOUNDATION
COMPOSITIO
MATHEMATICA



LONDON
MATHEMATICAL
SOCIETY
EST. 1865



Integral canonical models for Spin Shimura varieties

Keerthi Madapusi Pera

ABSTRACT

We construct regular integral canonical models for Shimura varieties attached to Spin and orthogonal groups at (possibly ramified) primes $p > 2$ where the level is not divisible by p . We exhibit these models as schemes of ‘relative PEL type’ over integral canonical models of larger Spin Shimura varieties with good reduction at p . Work of Vasiu–Zink then shows that the classical Kuga–Satake construction extends over the integral models and that the integral models we construct are canonical in a very precise sense. Our results have applications to the Tate conjecture for K3 surfaces, as well as to Kudla’s program of relating intersection numbers of special cycles on orthogonal Shimura varieties to Fourier coefficients of modular forms.

Introduction

The objects of study in this paper are certain Shimura varieties attached to GSpin and special orthogonal groups. More precisely, we will consider the \mathbb{Q} -algebraic stack $\mathrm{Sh}_{K_{\max}} := \mathrm{Sh}_{K_{\max}}(G, X)$, where $G = \mathrm{GSpin}(V, Q)$ is attached to a quadratic space (V, Q) over \mathbb{Q} of signature $(n, 2)$ and X is the space of oriented negative definite planes in $V_{\mathbb{R}}$. The level sub-group $K_{\max} \subset G(\mathbb{A}_f)$ is attached to a lattice $L \subset V$: it is the intersection of $G(\mathbb{A}_f)$ with $C(L \otimes \widehat{\mathbb{Z}})^{\times}$, the unit group of the Clifford algebra of $L \otimes \widehat{\mathbb{Z}}$. The image of K_{\max} in $\mathrm{SO}(V)(\mathbb{A}_f)$ is the *discriminant kernel*: the largest sub-group of $\mathrm{SO}(L)(\widehat{\mathbb{Z}})$ that acts trivially on the discriminant L^{\vee}/L . Here, $L^{\vee} \subset V$ is the dual lattice for L .

Our results can be summarized by the following theorem.

THEOREM. *Assume that L has square-free discriminant. Then, over $\mathbb{Z}[\frac{1}{2}]$, $\mathrm{Sh}_{K_{\max}}$ admits a regular canonical model with a regular compactification.*

The precise meaning of the word ‘canonical’ is explained in §4. In the body of the paper, we will isolate a prime $p > 2$ and work with finite quasi-projective covers $\mathrm{Sh}_K \rightarrow \mathrm{Sh}_{K_{\max}}$, where $K \subset K_{\max}$ is of the form $K_p K^p$ with $K_p = G(\mathbb{Q}_p) \cap C(L \otimes \mathbb{Z}_p)^{\times}$ and $K^p \subset G(\mathbb{A}_f^p)$ is a small enough compact open sub-group. We will then build canonical models over $\mathbb{Z}_{(p)}$ for the tower of such covers.

The basic idea of the proof is quite simple and involves exhibiting the model as the solution to a ‘relative PEL’ problem¹ over the smooth integral model of a larger Shimura variety attached to a self-dual lattice. To explain this, we work over $\mathbb{Z}_{(p)}$ for some odd prime p . When the lattice

Received 4 December 2012, accepted in final form 19 December 2014, published online 7 December 2015.

2010 Mathematics Subject Classification 11G18 (primary).

Keywords: Shimura varieties, integral canonical models, Spin groups.

This work was partially supported by NSF Postdoctoral Research Fellowship DMS-1204165.

This journal is © **Foundation Compositio Mathematica** 2015.

¹ We thank an anonymous referee for pointing us to [Vas14], where Vasiu has introduced the same terminology in a different context.

L is self-dual at a prime p , K_p is hyperspecial and the results of [Kis10] already give us a smooth integral canonical model over $\mathbb{Z}_{(p)}$ (compactifications are dealt with in [MaP15]). In general, we can exhibit L as a sub-lattice of a bigger lattice \tilde{L} that is self-dual at p and is such that the associated quadratic space (\tilde{L}, \tilde{Q}) over \mathbb{Q} again has signature $(m, 2)$ for some $m \in \mathbb{Z}_{\geq 0}$. If $\text{Sh}_{\tilde{K}}$ is the Shimura variety attached to a \tilde{L} and a compact open $\tilde{K} \subset \tilde{G}(\mathbb{A}_f^p)$ containing K , this allows us to exhibit Sh_K as an intersection of divisors in $\text{Sh}_{\tilde{K}}$. Let $\mathcal{S}_{\tilde{K}}$ be the integral canonical model for $\text{Sh}_{\tilde{K}}$ over $\mathbb{Z}_{(p)}$. We show that the divisors have a moduli interpretation and we use this interpretation to define models for them as schemes over $\mathcal{S}_{\tilde{K}}$. We then construct our regular models as intersections (over $\mathcal{S}_{\tilde{K}}$) of the integral models of these divisors.

The moduli interpretation of the divisors can be described as follows: the classical Kuga-Satake construction, combined with Kisin’s construction, gives us a natural polarized abelian scheme $(\tilde{A}^{\text{KS}}, \tilde{\lambda}^{\text{KS}})$ over $\mathcal{S}_{\tilde{K}}$. Let \tilde{L}_B be the $\mathbb{Z}_{(p)}$ -local system over $\mathcal{S}_{\tilde{K}, \mathbb{C}}$ attached to $\tilde{L}_{\mathbb{Z}_{(p)}}$. Let \tilde{H}_B be the first Betti cohomology of $\tilde{A}_{\mathbb{C}}^{\text{KS}}$ over $\mathcal{S}_{\tilde{K}, \mathbb{C}}$ with coefficients in $\mathbb{Z}_{(p)}$. Then the construction allows us to view \tilde{L}_B as a sub-local system of $\tilde{H}_B \otimes \tilde{H}_B^{\vee}$. Given an $\text{Sh}_{\tilde{K}}$ -scheme T , we say that an endomorphism f of \tilde{A}_T^{KS} is *special* if over $T_{\mathbb{C}}$ it induces a section of \tilde{L}_B . Then the divisors mentioned above can be viewed as the loci where \tilde{A}^{KS} inherits certain special endomorphisms of fixed degree. So, we obtain a moduli interpretation for $\text{Sh}_K \rightarrow \text{Sh}_{\tilde{K}}$: it is the locus over which \tilde{A}^{KS} inherits a certain family of special endomorphisms. For an analytic viewpoint of all this in the case of orthogonal Shimura varieties, cf. [Kud97].

Let \tilde{H}_{cris} be the first crystalline cohomology of $\tilde{A}_{\mathbb{F}_p}^{\text{KS}}$ over $\mathcal{S}_{\tilde{K}, \mathbb{F}_p}$. Then Kisin’s work provides us with a canonical sub-crystal $\tilde{L}_{\text{cris}} \subset \tilde{H}_{\text{cris}} \otimes \tilde{H}_{\text{cris}}^{\vee}$ attached to the quadratic space \tilde{L} . This allows us to give a definition of specialness in characteristic p as well: for any $\mathcal{S}_{\tilde{K}, \mathbb{F}_p}$ -scheme T , an endomorphism f of \tilde{A}_T^{KS} is *special* if its crystalline realization is a section of \tilde{L}_{cris} at every point of T . We can patch together the two notions of specialness to get the notion of a special endomorphism of \tilde{A}^{KS} in general.

The moduli interpretation for $\text{Sh}_K(G, X)$ as a scheme over $\text{Sh}_{\tilde{K}}(\tilde{G}, \tilde{X})$ can now be extended over $\mathbb{Z}_{(p)}$ to obtain a natural integral model $\mathcal{S}_K(G, X)$. To study its local properties, we study the problem of deforming special endomorphisms of \tilde{A}^{KS} , using ideas of Deligne [Del81] and Ogus [Ogu79]. These methods help us show that the quadric M_G^{loc} of isotropic lines in L is an étale local model for $\mathcal{S}_K(G, X)$. In the case where the discriminant of L is square-free, M_G^{loc} is regular, and this completes the construction.

The above construction actually works for general lattices L , but then it only gives us access to a certain open locus of the desired integral model, which we denote by $\mathcal{S}_K^{\text{pr}}(G, X)$: this is still modeled by M_G^{loc} . In general, the special fiber of this open locus will miss some very important parts of the expected ‘true’ integral model. For instance, in the situation where L is maximal, and p is a prime such that p^2 divides the discriminant of L , $\mathcal{S}_K^{\text{pr}}(G, X)$ will entirely *exclude* the part of the supersingular locus with maximal Artin invariant;² cf. § 6.27 for a discussion of this phenomenon. In particular, for $n = 3$, where our Shimura variety is closely related to Siegel threefolds with parahoric level structure, our theory does *not* recover the integral models of Chai and Norman [CN90] and Görtz [Gör03]: the simple local models that we describe are insufficient for this purpose. We expect to fix this gap in the theory in future work.

Along these lines, we should note that, for other low values of n , there are direct, moduli-theoretic ways to construct integral models. For the case of Shimura curves, cf. [KRY06], and for

² We thank the referee for pointing out many subtleties arising in this situation.

the case of certain Hilbert–Blumenthal surfaces, cf. [DP94, KR99]. Moreover, there is work in progress by Kisin and Pappas, which will generalize the methods of [Kis10] to construct integral models of Shimura varieties of abelian type with general parahoric level at p . However, the simple and direct nature of our construction seems to be quite useful in applications, which include the Tate conjecture for K3 surfaces in odd characteristic [MaP14], and also forthcoming work, in collaboration with F. Andreatta, E. Goren and B. Howard, on the arithmetic intersection theory over these Shimura varieties.

In the body of the paper, we also concurrently consider the case of orthogonal Shimura varieties. These are Shimura varieties of abelian type, but of a particularly simple nature: they are finite étale quotients of GSpin Shimura varieties, and we can easily deduce results for them from corresponding ones for their GSpin counterparts.

Notational conventions

For any prime ℓ , ν_ℓ will be the ℓ -adic valuation satisfying $\nu_\ell(\ell) = 1$. We will use \mathbb{A}_f to denote the ring of finite adèles over \mathbb{Q} , and $\widehat{\mathbb{Z}} \subset \mathbb{A}_f$ will be the pro-finite completion of \mathbb{Z} . Given a rational prime p , \mathbb{A}_f^p will denote the ring of prime-to- p finite adèles; that is, the restricted product $\prod'_{\ell \neq p} \mathbb{Q}_\ell$. Moreover, $\widehat{\mathbb{Z}}^p \subset \mathbb{A}_f^p$ will be the closure of \mathbb{Z} . Unless otherwise specified, the letter k will always represent a perfect field of characteristic p . Given such a k , $W(k)$ will denote its ring of Witt vectors, and $\sigma : W(k) \rightarrow W(k)$ will be the canonical lift of the Frobenius automorphism of k . For any group G , \underline{G} will denote the locally constant étale sheaf (over a base that will be clear from context) with values in G .

1. Clifford algebras and Spin groups

We will quickly summarize some standard facts about Clifford algebras and the GSpin group. A good reference is [Bou07, § 9]; cf. also [Bas74].

1.1 Let R be a commutative ring in which 2 is invertible, and let (L, Q) be a quadratic space over R : by this we mean a projective R -module L of finite rank equipped with a quadratic form $Q : L \rightarrow R$. We will denote by $[\cdot, \cdot]_Q : L \otimes L \rightarrow R$ the associated symmetric bi-linear form, its relation with Q being given by the formula $[v, w]_Q = Q(v + w) - Q(v) - Q(w)$.

Let $C := C(L)$ be the associated Clifford algebra over R . It is equipped with an embedding $L \hookrightarrow C$, which is universal for maps $f : L \rightarrow B$ into associative R -algebras B satisfying $f(v)^2 = Q(v)$. The algebra C is equipped with a natural $\mathbb{Z}/2\mathbb{Z}$ -grading $C = C^+ \oplus C^-$, so that C^+ is a sub-algebra of C .

1.2 Suppose now that Q is *non-degenerate*: that is, it induces an isomorphism $L \xrightarrow{\cong} L^\vee$. In this situation, we will also call the module L itself *self-dual*, especially when working over discrete valuation rings.

Then we can use the Clifford algebra to define a reductive group scheme $\mathrm{GSpin}(L, Q)$ over R . For any R -algebra S , we have

$$\mathrm{GSpin}(L, Q)(S) = \{x \in (C_S^+)^{\times} : x(L_S)x^{-1} = L_S\}.$$

It is a central extension of the special orthogonal group $\mathrm{SO}(L, Q)$ by \mathbb{G}_m , and there is a canonical character, the *spinor norm*, $\nu : \mathrm{GSpin}(L, Q) \rightarrow \mathbb{G}_m$. The spinor norm is defined as follows: there is a canonical anti-involution $*$ on C : it is the unique anti-automorphism that restricts to

the identity on L ; cf. [Bou07, § 9], where it is termed ‘*l’antiautomorphisme principal*’. For any $x \in \mathrm{GSpin}(L, Q)(S)$, we set $\nu(x) = x^*x$. It follows from [Bas74, 3.2.1] that this is indeed an element of $S^\times \subset (C_S^+)^{\times}$.

The group scheme $\mathrm{GSpin}(L, Q)$ acts naturally on C via left multiplication, and we will denote the resulting representation by H . The right multiplication action of C on itself endows H with a $\mathrm{GSpin}(L, Q)$ -equivariant right C -module structure, and the grading on C endows it with a $\mathrm{GSpin}(L, Q)$ -stable $\mathbb{Z}/2\mathbb{Z}$ -grading $H = H^+ \oplus H^-$. The action of L on C via left multiplication provides us with a $\mathrm{GSpin}(L, Q)$ -equivariant embedding $L \hookrightarrow \mathrm{End}_C(H)$.

1.3 Let $\mathrm{GL}_R^+(H) \subset \mathrm{GL}_R(H)$ be the sub-group³ of automorphisms that preserve the grading on H : this is the group of units in the algebra $\mathrm{End}_R^+(H)$ of graded endomorphisms of H . Let $\mathrm{U}(H) \subset \mathrm{GL}_R^+(H)$ be the centralizer of the right C -action. Then, for any R -algebra S , we have $\mathrm{U}(H)(S) = (\mathrm{End}_C^+(H_S))^{\times} = (C_S^+)^{\times}$. By definition, $\mathrm{GSpin}(L, Q)$ is a sub-group of $\mathrm{U}(H)$.

The pairing $[\varphi_1, \varphi_2] = (1/2^n) \mathrm{Tr}(\varphi_1 \circ \varphi_2)$ on $\mathrm{End}_R(H)$ is symmetric, non-degenerate and restricts to the pairing $[\cdot, \cdot]_Q$ on $L \subset C$. Choose an R -basis e_1, \dots, e_m of L and let $A \in \mathrm{GL}_m(R)$ be the matrix whose inverse is $([e_i, e_j]_Q)$. We will use the basis $\{e_i\}$ to identify L with R^m and hence to view A as an automorphism of L . Consider the endomorphism $\pi : \mathrm{End}_R(H) \rightarrow \mathrm{End}_R(H)$ given, for $\varphi \in \mathrm{End}_R(H)$, by

$$\pi(\varphi) = \sum_i [\varphi, e_i] \cdot Ae_i.$$

LEMMA 1.4.

- (i) π is an idempotent endomorphism of $\mathrm{End}_R(H)$ with image $L \subset \mathrm{End}_R(H)$.
- (ii) π is the unique projector onto L satisfying

$$\ker \pi = \{\varphi \in \mathrm{End}_R(H) : [\varphi, v] = 0 \text{ for all } v \in L\}.$$

In particular, it is independent of the choice of basis.

- (iii) $\mathrm{GSpin}(L, Q) \subset \mathrm{U}(H)$ is the stabilizer of $\pi : \mathrm{End}_R(H) \rightarrow \mathrm{End}_R(H)$.

Proof. For (i), we note that

$$\pi(\pi(\varphi)) = \sum_{i,k} [\varphi, e_i] \cdot [Ae_i, e_k]_Q \cdot Ae_k = \sum_{i,k} [\varphi, e_i] \cdot \delta_{i,k} \cdot Ae_k = \pi(\varphi).$$

An easy computation also shows that, for $1 \leq k \leq m$, $\pi(e_k) = e_k$, so that the image of π is precisely L .

(ii) is clear.

The stabilizer in $\mathrm{U}(H)$ of π must preserve L under the conjugation action on $\mathrm{End}_R(H)$; it must therefore be contained in $\mathrm{GSpin}(L, Q)$. On the other hand, for any $g \in \mathrm{GSpin}(L, Q)$, since g must preserve L , $g\pi g^{-1}$ is again a projector onto L whose kernel agrees with that of π . Hence, we must have $g\pi g^{-1} = \pi$. This shows (iii). \square

Remark 1.5. In the above situation, suppose that $R = \mathbb{Z}_{(p)}$ and only that $(L_{\mathbb{Q}}, Q)$ is non-degenerate. If we choose a basis $\{e_1, \dots, e_m\}$ for L , the corresponding matrix A belongs to $\mathrm{GL}(L_{\mathbb{Q}})$ and $A \cdot L \subset L_{\mathbb{Q}}$ is precisely the dual lattice L^{\vee} . Therefore, the image of $\mathrm{End}_{\mathbb{Z}_{(p)}}(H)$ under π is exactly $L^{\vee} \subset L_{\mathbb{Q}}$.

³ For the rest of the section, the word ‘group’ will mean ‘ R -group scheme’.

1.6 As shown in [Bou07, §9, Théorèmes 2 and 3], C is an Azumaya algebra over its center $Z(C)$. Here, $Z(C) = R$, if m is even, and is finite étale of rank 2 over R , if m is odd. Therefore, there exists an R -linear *reduced trace map* $\text{Trd} : C \rightarrow R$ such that the pairing $(x, y) \mapsto \text{Trd}(xy)$ is a non-degenerate symmetric bilinear form on C . For any $\delta \in C^\times$ such that $\delta^* = -\delta$, the form $\psi_\delta(x, y) = \text{Trd}(x\delta y^*)$ defines an R -valued symplectic form on H .

LEMMA 1.7. For δ as above, the line $[\psi_\delta]$ in $\text{Hom}(H \otimes H, R)$, spanned by the symplectic form ψ_δ , is preserved by $\text{GSpin}(L, Q)$. The similitude character of $\text{GSpin}(L, Q)$ obtained from its action on this line agrees with the spinor norm. \square

DEFINITION 1.8. We will need one further piece of notation: for any pair of positive integers (r, s) , we set

$$H^{\otimes(r,s)} = \underbrace{H^\vee \otimes \cdots \otimes H^\vee}_r \otimes \underbrace{H \otimes \cdots \otimes H}_s.$$

We will also use this notation for objects in other tensor categories without comment.

Note that we can now think of π as an element of $H^{\otimes(2,2)}$.

1.9 Let $G_0 = \text{SO}(L, Q)$. Since there exists a central isogeny $G \rightarrow G_0$ of reductive groups, there is a bijective correspondence between parabolic sub-groups of G and those of G_0 . We want to make this correspondence explicit on the level of linear algebra for certain parabolic sub-groups. To each isotropic sub-space $L^1 \subset L$, we can attach the parabolic sub-group $P_0 \subset G_0$ that stabilizes L^1 . We get a decreasing filtration

$$0 = F^2L \subset F^1L = L^1 \subset F^0L = (L^1)^\perp \subset F^{-1}L = L.$$

Since L^1 is isotropic, we have a canonical embedding of R -algebras

$$\wedge^\bullet L^1 \hookrightarrow C \hookrightarrow \text{End}_R(H).$$

If $N \subset \text{End}_R(H)$, write $\text{im } N$ for the union of the images in H of the endomorphisms in N . Similarly, write $\text{ker } N$ for the intersection of the kernels of the elements of N . Then we have, for every integer $i = 0, \dots, r + 1$,

$$\text{im}(\wedge^i L^1) = \text{ker}(\wedge^{r-i+1} L^1).$$

Moreover, $\text{im}(\wedge^i L^1) \subset \text{im}(\wedge^{i-1} L^1)$. So, we can define a descending filtration $F^\bullet H$ on H by

$$F^i H = \text{im}(\wedge^i L^1).$$

Suppose that $\mu_0 : \mathbb{G}_m \rightarrow G_0$ is a co-character splitting $F^\bullet L$. It gives rise to a splitting

$$L = L^1 \oplus L^0 \oplus L^{-1}$$

with $F^i L = \bigoplus_{j \geq i} L^j$ and where $\mu_0(z)$ acts on L^i via z^i . In particular, L^{-1} is another isotropic direct summand of L that pairs non-degenerately with L^1 .

Take the increasing filtration $E_i H = \text{ker}(\wedge^{i+1} L^{-1}) = \text{im}(\wedge^{r-i} L^{-1})$ and set $H^i = E_i H \cap F^i H$. One easily checks that this is a splitting of $F^\bullet H$.

Let $\mu : \mathbb{G}_m \rightarrow \text{GL}(H)$ be the co-character that acts via z^i on H^i . By construction, H^i is C -stable and one can check easily that $\mu(\mathbb{G}_m)$ preserves the grading on H . So, μ must factor through $\text{U}(H)$. Furthermore, we find that, if $v \in L$ and $i = 0, 1, \dots, r$, then

$$v \cdot H^i \subset \begin{cases} H^{i+1} & \text{if } v \in L^1, \\ H^i & \text{if } v \in L^0, \\ H^{i-1} & \text{if } v \in L^{-1}. \end{cases}$$

This shows that

$$\mu(z)v\mu(z)^{-1} = \begin{cases} z \cdot v & \text{if } v \in L^1, \\ v & \text{if } v \in L^0, \\ z^{-1} \cdot v & \text{if } v \in L^{-1}. \end{cases}$$

In other words, μ factors through G and is a lift of μ_0 . This shows the following result.

PROPOSITION 1.10. *The parabolic sub-group $P \subset G$ lifting $P_0 \subset G_0$ is the stabilizer in G of $F^\bullet H$. \square*

2. Local models

We fix a prime $p > 2$ for the rest of this paper.

2.1 Suppose that (M, q) is a non-degenerate quadratic space over a field κ and let $N \subset M$ be a sub-space. Set

$$\mathfrak{a}_N = \{f \in \text{Hom}(N, M) : [f(v), w]_q + [v, f(w)]_q = 0 \text{ for all } v, w \in N\}.$$

LEMMA 2.2. *The natural map $\text{Lie}(\text{SO}(M)) \rightarrow \mathfrak{a}_N$ induced by restriction to N is surjective.*

Proof. We will first consider three special cases.

- *q restricted to N is also non-degenerate:* in this case, we have an orthogonal decomposition $M = N \perp N^\perp$. So, given $f \in \mathfrak{a}_N$, we can extend it to the element of $\text{Lie}(\text{SO}(M))$ which restricts to 0 on N^\perp .
- *N is isotropic:* in this case, we can find a splitting $M = N \oplus (N \oplus N')^\perp \oplus N'$, where $N' \subset M$ is also isotropic and pairs non-degenerately with N , and the restriction of q to $(N \oplus N')^\perp$ is non-degenerate. Given $f \in \mathfrak{a}_N$, we can write it in the form $f_1 \oplus f_2 \oplus f_3$, where $f_1 : N \rightarrow N$, $f_2 : N \rightarrow (N \oplus N')^\perp$ and $f_3 : N \rightarrow N'$. The duality between N' and N allow us to view f_1^\vee as a map $N' \rightarrow N'$. Similarly, f_2^\vee can be viewed as a map $(N \oplus N')^\perp \rightarrow N'$, and f_3^\vee as a map $N \rightarrow N'$. The condition that $f \in \mathfrak{a}_N$ simply means that $f_3^\vee = -f_3$. We can now extend f to an element $X \in \text{Lie}(\text{SO}(N))$, which restricts to $-f_2^\vee$ on $(N \oplus N')^\perp$ and $-f_1^\vee$ on N' .
- *N contains $\perp N$ and the image of f lies in N^\perp :* again, we can find a splitting $M = N \oplus N'$ with $N = N^\perp \oplus (N^\perp \oplus N')^\perp$. Here, N' is isotropic and pairs non-degenerately with N^\perp . Now, f is of the form $f_1 \oplus 0 \oplus 0$, where $f_1 : (N^\perp \oplus N')^\perp \rightarrow N^\perp$. Extend this to the element $X \in \text{Lie}(\text{SO}(N))$, which restricts to $-f_1^\vee : N' \rightarrow (N^\perp \oplus N')^\perp$ on N' and to 0 on N .

In general, let $N_0 \subset N$ be the radical. Since N_0 is isotropic, by the second case above, we can find $X_1 \in \text{Lie}(\text{SO}(M))$ such that $X_1|_{N_0} \equiv f|_{N_0}$. Therefore, replacing f with $f - X_1|_N$, we can assume that $f|_{N_0} \equiv 0$. In this case, the image of f must lie within N_0^\perp . Let $\bar{f} : N/N_0 \rightarrow N_0^\perp/N_0$ be the induced map. Now, N/N_0 is a non-degenerate sub-space of the non-degenerate quadratic space N_0^\perp/N_0 . Therefore, from the first case treated above, we can find $\bar{X}_2 \in \text{Lie}(\text{SO}(N_0^\perp/N_0))$ such that $\bar{X}_2|_{N/N_0} \equiv \bar{f}$. Lift \bar{X}_2 to an element $X_2 \in \text{Lie}(\text{SO}(M))$ that stabilizes N_0 . Then, replacing f by $f - X_2|_N$, we can assume that the image of f lands in N_0 . Extend f to any map $\tilde{f} : N_0^\perp \rightarrow N_0$. Then, by the third case above, we can find $X \in \text{Lie}(\text{SO}(M))$ such that $X|_{N_0^\perp} \equiv \tilde{f}$. \square

2.3 Suppose that we are given a discrete valuation ring \mathcal{O} of mixed characteristic $(0, p)$ with residue field k , and a quadratic space (M, q) over \mathcal{O} such that $M[p^{-1}]$ is non-degenerate over the fraction field $\mathcal{O}[p^{-1}]$. The quadratic form q endows the discriminant $\text{disc}(M) = M^\vee/M$ with a natural non-degenerate $\mathcal{O}[p^{-1}]/\mathcal{O}$ -valued quadratic form \bar{q} .

We say that M is *maximal* if it is maximal among \mathcal{O} -lattices in $M[p^{-1}]$ on which q takes values in \mathcal{O} .

LEMMA 2.4.

- (i) *The following are equivalent.*
 - (a) *M is maximal;*
 - (b) *$\text{disc}(M)$ is a k -vector space and the form \bar{q} is anisotropic.*
- (ii) *Suppose that (\tilde{M}, \tilde{q}) is a self-dual quadratic space over \mathcal{O} containing (M, q) isometrically as a direct summand. Then M is maximal if and only if $M^\perp \subset \tilde{M}$ is maximal.*

Proof. First, we claim that M is maximal only if $\text{disc}(M)$ is a k -vector space. Otherwise, if $\pi \in \mathcal{O}$ is a uniformizer, we can find $m \in M^\vee$ such that $\pi \cdot m \notin M$ but $\pi^2 \cdot m \in M$. But then $M + \langle \pi \cdot m \rangle$ is a lattice bigger than M on which q is \mathcal{O} -valued.

To prove (i), we can now assume that $\text{disc}(M)$ is a k -vector space. Then the assignment $M' \mapsto M'/M$ sets up a bijection between the following two sets.

- Lattices $M' \subset M[p^{-1}]$ containing M and on which q takes values in \mathcal{O} .
- Sub-spaces of $\text{disc}(M)$ that are isotropic for \bar{q} .

From this, the claimed equivalence is clear.

For the final assertion, note that the identification $\tilde{M} \xrightarrow{\sim} \tilde{M}^\vee$ induced by \tilde{q} gives us canonical isomorphisms

$$\text{disc}(M) \xleftarrow{\sim} \frac{\tilde{M}}{M + M^\perp} \xrightarrow{\sim} \text{disc}(M^\perp).$$

This gives us an isomorphism $\text{disc}(M) \xrightarrow{\sim} \text{disc}(M^\perp)$ that preserves quadratic forms up to sign. We now conclude the proof from (i). □

2.5 Let (L, Q) be a quadratic space over $\mathbb{Z}_{(p)}$ such that $(L_\mathbb{Q}, Q)$ is non-degenerate. Suppose that we are given a self-dual quadratic space (\tilde{L}, \tilde{Q}) over $\mathbb{Z}_{(p)}$ admitting (L, Q) as a direct summand. Set $\Lambda = L^\perp \subset \tilde{L}$. Set $\tilde{G} = \text{GSpin}(\tilde{L}, \tilde{Q})$ and let $G \subset \tilde{G}$ be the closed sub-group such that, for any $\mathbb{Z}_{(p)}$ -algebra R , we have

$$G(R) = \{g \in \tilde{G}(R) : g|_{\Lambda_R} \equiv 1\}.$$

Note that the central embedding $\mathbb{G}_{m, \mathbb{Q}} \hookrightarrow G_\mathbb{Q} = \text{GSpin}(L_\mathbb{Q}, Q)$ is induced from an embedding $\mathbb{G}_m \hookrightarrow G$. Let G_0 (respectively \tilde{G}_0) be the $\mathbb{Z}_{(p)}$ -group scheme G/\mathbb{G}_m (respectively \tilde{G}/\mathbb{G}_m), so that $G_{0, \mathbb{Q}} = \text{SO}(L_\mathbb{Q}, Q)$.

LEMMA 2.6. *For any flat $\mathbb{Z}_{(p)}$ -algebra R , we have*

$$\begin{aligned} G(R) &= G(R_\mathbb{Q}) \cap C(L)_R^\times \subset C(L)_{R_\mathbb{Q}}^\times, \\ G_0(R) &= \{g \in G_0(R_\mathbb{Q}) : gL_R = L_R \text{ and } g \text{ acts trivially on } L_R^\vee/L_R\}. \end{aligned}$$

Proof. The obvious inclusion $L \hookrightarrow \tilde{L} \hookrightarrow C(\tilde{L})$ gives rise to a canonical map $C(L) \rightarrow C(\tilde{L})$. We claim that this sets up an identification:

$$C(L)^+ = \{z \in C(\tilde{L})^+ : vz = zv \in C(\tilde{L}) \text{ for all } v \in \Lambda\}. \tag{2.6.1}$$

Since both sides of this purported identity are saturated $\mathbb{Z}_{(p)}$ -sub-modules of $C(\tilde{L})^+$, it is enough to show that

$$C(L_{\mathbb{Q}})^+ = \{z \in C(\tilde{L}_{\mathbb{Q}})^+ : vz = zv \in C(\tilde{L}_{\mathbb{Q}}) \text{ for all } v \in \Lambda\}. \tag{2.6.2}$$

This is easily checked using the orthogonal decomposition $\tilde{L}_{\mathbb{Q}} = L_{\mathbb{Q}} \perp \Lambda_{\mathbb{Q}}$.

The description of $G(R)$ for a flat $\mathbb{Z}_{(p)}$ -algebra R is now clear, since both $G(R_{\mathbb{Q}}) \cap C(L)_{\mathbb{R}}^{\times}$ and $G(R)$ can be identified with the set of elements of $\tilde{G}(R)$ that commute with the left multiplication action of Λ on $C(\tilde{L})_R$.

Proving the corresponding description of $G_0(R)$ amounts to checking the following thing: suppose that we have $g \in G_0(R_{\mathbb{Q}})$ such that $gL_R = L_R$. Then

$$g\tilde{L}_R = \tilde{L}_R \Leftrightarrow g \text{ acts trivially on } L_R^{\vee}/L_R.$$

Indeed, suppose that $g\tilde{L}_R = \tilde{L}_R$. Given any $v \in L_R^{\vee}$, there exists $w \in \Lambda_R^{\vee}$ such that the element $v + w$ of $\tilde{L}_{R_{\mathbb{Q}}} = L_{R_{\mathbb{Q}}} \oplus \Lambda_{R_{\mathbb{Q}}}$ lies in \tilde{L}_R . Therefore, since g acts trivially on Λ_R , we have

$$gv - v = g(v + w) - (v + gw) = g(v + w) - (v + w) \in \tilde{L}_R \cap L_{R_{\mathbb{Q}}} = L_R.$$

This shows that g acts trivially on L_R^{\vee}/L_R .

On the other hand, suppose that g acts trivially on L_R^{\vee}/L_R . Every element $\tilde{v} \in \tilde{L}_R$ can be written uniquely in the form $v + w$ for some $v \in L_R^{\vee} \subset L_{R_{\mathbb{Q}}}$ and $w \in \Lambda_R^{\vee} \subset \Lambda_{R_{\mathbb{Q}}}$. We now have

$$g\tilde{v} = g(v + w) = gv + w = (gv - v) + v + w = (gv - v) + \tilde{v} \in \tilde{L}_{R_{\mathbb{Q}}}.$$

But, by our hypothesis on g , $gv - v \in L_R \subset \tilde{L}_R$. Therefore, $g\tilde{v} \in \tilde{L}_R$ for all $\tilde{v} \in \tilde{L}_R$. □

LEMMA 2.7. *Set*

$$\mathfrak{a}_{\Lambda} = \{f \in \text{Hom}(\Lambda, \tilde{L}) : [f(v), w]_{\tilde{Q}} + [v, f(w)]_{\tilde{Q}} = 0 \text{ for all } v, w \in \Lambda\}.$$

Then we have a short exact sequence of finite free $\mathbb{Z}_{(p)}$ -modules

$$0 \rightarrow \text{Lie } G_0 \rightarrow \text{Lie } \tilde{G}_0 \xrightarrow{X \mapsto X|_{\Lambda}} \mathfrak{a}_{\Lambda} \rightarrow 0.$$

Proof. The only non-obvious part is the surjectivity on the right-hand side. By Nakayama's lemma, it suffices to prove this after tensoring with \mathbb{F}_p . But then it follows from (2.2). □

LEMMA 2.8. *Write $\iota_0 : \Lambda \hookrightarrow \tilde{L}$ for the natural embedding. Let R be a $\mathbb{Z}_{(p)}$ -algebra. Suppose that we have another isometric embedding $\iota : \Lambda_R \hookrightarrow \tilde{L}_R$ onto a local direct summand of \tilde{L}_R . Then the functor \mathcal{P}_{ι} on R -algebras given by*

$$\mathcal{P}_{\iota}(B) = \{g \in \tilde{G}_0(B) : g \circ \iota_0 = \iota\}$$

is represented by a scheme that is affine, smooth and faithfully flat over R .

Proof. It is clear that \mathcal{P}_ι is represented by a closed sub-scheme of $\tilde{G}_{0,R}$. Moreover, Witt’s extension theorem [Bou07, §4, Theorem 1] shows that $\mathcal{P}_\iota(\kappa)$ is non-empty for any residue field κ of R . To finish the proof of the proposition, it suffices to show that \mathcal{P}_ι is formally smooth over R .

Let B be an R -algebra, let $I \subset B$ be a square-zero ideal and let $B_0 = B/I$. Suppose that we have $g_0 \in \mathcal{P}_\iota(B_0)$: we want to find $g \in \mathcal{P}_\iota(B)$ mapping to g_0 . Choose any $\tilde{g} \in \tilde{G}_0(B)$ lifting g_0 . Consider the assignment on Λ_B given by $v \mapsto v - \tilde{g}^{-1}\iota(v)$. This factors through a map $U : \Lambda_{B_0} \rightarrow I \otimes_{B_0} \tilde{L}_{B_0}$:

$$\Lambda_B \rightarrow \Lambda_{B_0} \xrightarrow{U} I \otimes_{B_0} \tilde{L}_{B_0} \xrightarrow{\cong} I \otimes_B \tilde{L}_B \hookrightarrow \tilde{L}_B.$$

For $v, w \in \Lambda$, we have

$$\begin{aligned} [U(v), w]_{\tilde{Q}} + [v, U(w)]_{\tilde{Q}} &= [v - \tilde{g}^{-1}\iota(v), w]_{\tilde{Q}} + [v, w - \tilde{g}^{-1}\iota(w)]_{\tilde{Q}} \\ &= [v - \tilde{g}^{-1}\iota(v), w - \tilde{g}^{-1}\iota(w)]_{\tilde{Q}} + [v, w]_{\tilde{Q}} - [\tilde{g}^{-1}\iota(v), \tilde{g}^{-1}\iota(w)]_{\tilde{Q}} \\ &= [v - \tilde{g}^{-1}\iota(v), w - \tilde{g}^{-1}\iota(w)]_{\tilde{Q}} = 0. \end{aligned}$$

Here, the last equality holds because $I^2 = 0$.

Therefore, $U \in I \otimes_{\mathbb{Z}_{(p)}} \mathfrak{a}_\Lambda$ and, by (2.7), we can find $X \in I \otimes_{\mathbb{Z}_{(p)}} \text{Lie}(\tilde{G}_0)$ such that $X|_{\Lambda_{R_0}} \equiv U$. Then $g = \tilde{g} \circ (1 - X)$ is an element of $\mathcal{P}_\iota(B)$ that lifts g_0 . \square

PROPOSITION 2.9. *The group schemes G and G_0 are smooth and faithfully flat over $\mathbb{Z}_{(p)}$. In particular, up to (unique) isomorphism, they do not depend on the choice of self-dual quadratic space \tilde{L} containing L .*

Proof. It is enough to show that G_0 is smooth over $\mathbb{Z}_{(p)}$. This follows from (2.8) with $R = \mathbb{Z}_{(p)}$ and $\iota = \iota_0$.

Take $R = \mathbb{Z}_{(p)}^{nr}$ to be a strict henselization of $\mathbb{Z}_{(p)}$. That G and G_0 are independent of the choice of \tilde{L} is immediate from (2.6) and the following assertion [BT84, 1.7.6]: given a smooth \mathbb{Q} -scheme X , a smooth $\mathbb{Z}_{(p)}$ -model \mathfrak{X} for X is determined up to unique isomorphism by its set of R -valued points $\mathfrak{X}(R) \subset X(R_{\mathbb{Q}})$. \square

2.10 Let M_G^{loc} be the $\mathbb{Z}_{(p)}$ -scheme such that, for every $\mathbb{Z}_{(p)}$ -algebra R , we have

$$M_G^{\text{loc}}(R) = \{\text{Isotropic lines } F^1 L_R \subset L_R\}^4$$

LEMMA 2.11. *Let $N \subset L_{\mathbb{F}_p}$ be the radical. Set $r = \dim L$, $t = \dim N$ and $s = r - t - 1$. We will assume that $N \neq L_{\mathbb{F}_p}$ or, equivalently, $s \geq 0$.*

- (i) M_G^{loc} is flat and projective of relative dimension $r - 2$ over $\mathbb{Z}_{(p)}$.
- (ii) The singular locus of $M_{G, \mathbb{F}_p}^{\text{loc}}$ consists of lines contained in N and so can be identified with $\mathbb{P}(N)$. It has co-dimension s in $M_{G, \mathbb{F}_p}^{\text{loc}}$.⁵
- (iii) $M_{G, \mathbb{F}_p}^{\text{loc}}$ is a local complete intersection (lci) variety. It is reduced if and only if $s \geq 1$. It is normal if and only if $s \geq 2$, and smooth if and only if $t = 0$.

⁴ A line is an R -sub-module that is locally a direct summand of rank 1.

⁵ Here, $\mathbb{P}(N)$ denotes the space of lines in N .

Proof. Part (i) is a direct consequence of the hypothesis that $N \neq L_{\mathbb{F}_p}$. Part (ii) is an easy fact about quadrics. Finally, $M_{G, \mathbb{F}_p}^{\text{loc}}$, being a quadric, is an lci variety. The remainder of (iii) follows from (ii) and standard criteria for reducedness and normality. \square

We now recall some definitions and results from [VZ10].

DEFINITION 2.12. A regular local $\mathbb{Z}_{(p)}$ -algebra R with maximal ideal \mathfrak{m} is *quasi-healthy* if it is faithfully flat over $\mathbb{Z}_{(p)}$ and if every abelian scheme over $\text{Spec } R \setminus \{\mathfrak{m}\}$ extends uniquely to an abelian scheme over $\text{Spec } R$.

A regular $\mathbb{Z}_{(p)}$ -scheme X is *healthy* if it is faithfully flat over $\mathbb{Z}_{(p)}$ and if, for every open sub-scheme $U \subset X$ containing $X_{\mathbb{Q}}$ and all generic points of $X_{\mathbb{F}_p}$, every abelian scheme over U extends uniquely to an abelian scheme over X . It is *locally healthy* if, for every point $x \in X_{\mathbb{F}_p}$ of co-dimension at least 2, the complete local ring $\widehat{\mathcal{O}}_{X,x}$ is quasi-healthy.

Remark 2.13.

- Any regular, flat $\mathbb{Z}_{(p)}$ -scheme of dimension at most 1 is trivially healthy.
- By faithfully flat descent, a regular local ring R is quasi-healthy whenever its completion \widehat{R} is quasi-healthy.
- If X is locally healthy, then it is healthy. Indeed, suppose that $U \subset X$ is as in the definition of ‘healthy’ above; the complement $X \setminus U$ lies entirely in the special fiber and has co-dimension at least 2 in X . The claim follows by using ascending noetherian induction on the co-dimension of $X \setminus U$, and repeatedly using quasi-healthiness of the local rings of X .

We do not know if the converse holds.

THEOREM 2.14 (Vasiu–Zink). *Let R be a regular local, faithfully flat $\mathbb{Z}_{(p)}$ -algebra of dimension at least 2.*

- (i) *Suppose that there exists a faithfully flat complete local R -algebra \widehat{R} that admits a surjection $\widehat{R} \rightarrow W[[T_1, T_2]]/(p-h)$, where $h \in (T_1, T_2)W[[T_1, T_2]]$ is a power series that does not belong to the ideal $(p, T_1^p, T_2^p, T_1^{p-1}T_2^{p-1})$. Then R is quasi-healthy.*
- (ii) *Let $\mathfrak{m}_R \subset R$ be the maximal ideal and suppose that $p \notin \mathfrak{m}_R^p$. Then R is quasi-healthy.*
- (iii) *If R is a formally smooth complete local $\mathbb{Z}_{(p)}$ -algebra, then R is quasi-healthy.*

Proof. See [VZ10, Theorem 3 and Corollary 4]. \square

2.15 Suppose that (L, Q) is maximal. The isomorphism $pL^\vee/pL \xrightarrow{\cong} \text{disc}(L)$ allows us to identify $\text{disc}(L)$ with the radical $N \subset L_{\mathbb{F}_p}$. By (2.4), $\text{disc}(L)$ with its induced form \overline{Q} has to be an anisotropic quadratic space over \mathbb{F}_p . This implies that $t = \dim N \leq 2$. If $t = 2$, $\text{disc}(L)_{\mathbb{F}_{p^2}} = N_{\mathbb{F}_{p^2}}$ admits two lines that are \overline{Q} -isotropic. These lines, since every sub-space of $N_{\mathbb{F}_{p^2}}$ is Q -isotropic, can be viewed as points in $M_G^{\text{loc}}(\mathbb{F}_{p^2})$. We will call these points *irregular*.

PROPOSITION 2.16. *If $t \leq 1$, then M_G^{loc} is regular and locally healthy. If $t = 2$, then the same assertion holds for the complement of the irregular points defined above.*

Proof. If $t = 0$, then M_G^{loc} is smooth over $\mathbb{Z}_{(p)}$ and the result is immediate from Theorem 2.14.

So, we assume that $t \geq 1$. The assertions in the proposition can be checked at the complete local rings of M_G^{loc} at points valued in algebraically closed fields of characteristic p . As such, it suffices to prove them after base change to $\mathbb{Z}_{p^2} = W(\mathbb{F}_{p^2})$. Now, we can find a basis for $\mathbb{Z}_{p^2} \otimes L$

so that the quadratic form has the shape

$$\sum_{i=1}^{r-1} X_i^2 + pY^2 \quad \text{if } t = 1, \quad \sum_{i=1}^{r-2} X_i^2 + pYZ \quad \text{if } t = 2.$$

The singular locus of $\mathbb{Z}_{p^2} \otimes M_G^{\text{loc}}$ is precisely where p and all the coordinates X_i , $1 \leq i \leq r-t$, vanish. When $t = 2$, the irregular points are precisely those points in the singular locus where one of Y or Z also vanishes.

Set

$$U = \begin{cases} \text{Spec } \frac{\mathbb{Z}_{p^2}[u_1, \dots, u_{r-1}]}{(\sum_i u_i^2 + p)} & \text{if } t = 1, \\ \text{Spec } \frac{\mathbb{Z}_{p^2}[u_1, \dots, u_{r-2}, v]}{(\sum_i u_i^2 + pv)} & \text{if } t = 2. \end{cases}$$

Every singular point of $\mathbb{Z}_{p^2} \otimes M_G^{\text{loc}}$ has a Zariski open neighborhood isomorphic to U .

It is now an easy observation that U is regular everywhere when $t = 1$. When $t = 2$, it is regular outside of points where all the coordinates u_1, \dots, u_{r-2}, v vanish: these are precisely the irregular points defined above.

When $t = 1$, the completion of U at its singular point is $\mathbb{Z}_{p^2}[[u_1, \dots, u_{r-2}]]/(\sum_i u_i^2 + p)$. Here, we assume that $r \geq 3$, since otherwise $\dim M_G^{\text{loc}} \leq 1$, and being healthy is a vacuous condition.

Similarly, when $t = 2$, the completion of U at any regular $\overline{\mathbb{F}}_p$ -valued point in the singular locus of the special fiber is isomorphic to $W(\overline{\mathbb{F}}_p)[[u_1, \dots, u_{r-2}, w]]/(\sum_i u_i^2 + pw + p)$. Here again, we assume that $r \geq 3$.

We also need to consider the completion of U at the generic point of the singular locus. If we complete instead at an algebraically closed point over this generic point, a quick computation shows that we obtain a ring isomorphic to $W(k)[[u_1, \dots, u_{r-2}]]/(\sum_i u_i^2 + pv)$. Here, k is an algebraically closed field containing $\mathbb{F}_{p^2}((v))$, and we view v as an element of $W(k)$ via the Teichmüller lift.

In all three cases, if \mathfrak{m} is the maximal ideal of the complete local ring, we see that $p \in \mathfrak{m}^2 \setminus \mathfrak{m}^3$. So, we can conclude using Theorem 2.14(ii) that the complete local rings of U at any regular, singular point are quasi-healthy. Since the complete local rings at the non-singular points are also quasi-healthy by Theorem 2.14, we see that the regular locus of U is locally healthy. This proves the proposition. \square

2.17 Assume that $t = 2$. We will now construct a regular, locally healthy resolution M_G^{ref} of M_G^{loc} . Fix any quadratic extension F/\mathbb{Q} in which p is inert. The two isotropic lines in $\text{disc}(L)_{\mathbb{F}_{p^2}}$, via the correspondence noted in the proof of (2.4), determine two self-dual lattices in L_F containing $L_{\mathcal{O}_{F,(p)}}$. Fix one of them and denote it by L^\diamond .

Given a $\mathbb{Z}_{(p)}$ -algebra R , we take $M_G^{\text{ref}}(R)$ to be the set of pairs $(F^1 L_R, F^1 L_R^\diamond)$, where the following hold.

- $F^1 L_R \subset L_R$ is an isotropic line.
- $F^1 L_R^\diamond \subset L_R^\diamond = L^\diamond \otimes_{\mathbb{Z}_{(p)}} R$ is an isotropic $\mathcal{O}_{F,(p)} \otimes_{\mathbb{Z}_{(p)}} R$ -sub-module that is locally a direct summand of rank 1.
- Under the natural map $\mathcal{O}_{F,(p)} \otimes_{\mathbb{Z}_{(p)}} L_R \rightarrow L_R^\diamond$, $\mathcal{O}_{F,(p)} \otimes F^1 L_R$ maps into $F^1 L_R^\diamond$.

PROPOSITION 2.18.

- (i) M_G^{ref} is represented by a regular, locally healthy projective $\mathbb{Z}_{(p)}$ -scheme of relative dimension $r - 2$.
- (ii) The natural map $q : M_G^{\text{ref}} \rightarrow M_G^{\text{loc}}$ is G -equivariant and an isomorphism over the regular locus of the target.

Proof. Let $Q(L^\diamond)$ be the smooth quadric over $\mathcal{O}_{F,(p)}$ attached to L^\diamond , and let M_G^{loc} be the Weil restriction of $Q(L^\diamond)$ from $\mathcal{O}_{F,(p)}$ to $\mathbb{Z}_{(p)}$. By definition, we have an inclusion of functors

$$M_G^{\text{ref}} \hookrightarrow M_G^{\text{loc}} \times_{\mathbb{Z}_{(p)}} M_G^{\text{loc}}.$$

This is easily seen to be a closed immersion, so that M_G^{ref} is represented by a projective $\mathbb{Z}_{(p)}$ -scheme. Since G acts trivially on $\text{disc}(L)$, its action on L_F preserves L^\diamond . This endows M_G^{ref} with a natural G -action compatible with its projection onto M_G^{loc} .

Fix an identification of the completion of $\mathcal{O}_{F,(p)}$ with \mathbb{Z}_{p^2} . Using the isomorphism of \mathbb{Z}_{p^2} -algebras $\mathbb{Z}_{p^2} \otimes_{\mathbb{Z}_{(p)}} \mathcal{O}_{F,(p)} \xrightarrow{\cong} \mathbb{Z}_{p^2} \times \mathbb{Z}_{p^2}$, we obtain maps

$$\mathbb{Z}_{p^2} \otimes_{\mathbb{Z}_{(p)}} L \rightarrow \mathbb{Z}_{p^2} \otimes_{\mathbb{Z}_{(p)}} L^\diamond = (\mathbb{Z}_{p^2} \otimes_{\mathcal{O}_{F,(p)}} L^\diamond) \oplus (\mathbb{Z}_{p^2} \otimes_{\mathcal{O}_{F,(p)}} L^\diamond). \tag{2.18.1}$$

As in the proof of (2.16), we can find compatible bases for $\mathbb{Z}_{p^2} \otimes_{\mathbb{Z}_{(p)}} L$ and $\mathbb{Z}_{p^2} \otimes_{\mathcal{O}_{F,(p)}} L^\diamond$ such that the map in (2.18.1) has the shape

$$\begin{aligned} \mathbb{Z}_{p^2} \otimes_{\mathbb{Z}_{(p)}} L &\rightarrow (\mathbb{Z}_{p^2} \otimes_{\mathcal{O}_{F,(p)}} L^\diamond) \oplus (\mathbb{Z}_{p^2} \otimes_{\mathcal{O}_{F,(p)}} L^\diamond), \\ (X_1, \dots, X_{r-2}, Y, Z) &\mapsto ((X_1, \dots, X_{r-2}, pY, Z), (X_1, \dots, X_{r-2}, Y, pZ)), \end{aligned}$$

and so that the quadratic forms on $\mathbb{Z}_{p^2} \otimes_{\mathbb{Z}_{(p)}} L$ and $\mathbb{Z}_{p^2} \otimes_{\mathcal{O}_{F,(p)}} L^\diamond$ are given by the formulas $\sum_i X_i^2 + pYZ$ and $\sum_i X_i^2 + YZ$, respectively.

Let $C \subset M_G^{\text{loc}}$ be the irregular locus: then $\mathbb{Z}_{p^2} \otimes C \subset \mathbb{Z}_{p^2} \otimes M_G^{\text{loc}}$ can be identified with the locus in the special fiber where the coordinates X_1, \dots, X_{r-2} , as well as (exactly) one of Y or Z , vanish. Set $V_1 = M_G^{\text{loc}} \setminus C$; then we see that, as open sub-schemes of $\mathbb{Z}_{p^2} \otimes M_G^{\text{loc}}$, we have

$$\mathbb{Z}_{p^2} \otimes V_1 = \bigcup_{i=1}^{r-1} \{X_i \neq 0\} \cup \{YZ \neq 0\}.$$

This immediately shows that, if R is an $\mathbb{Z}_{(p)}$ -algebra with $F^1 L_R \in V_1(R)$, the image of $\mathcal{O}_{F,(p)} \otimes_{\mathbb{Z}_{(p)}} F^1 L_R$ in L_R^\diamond is locally a direct summand as an $\mathcal{O}_{F,(p)} \otimes_{\mathbb{Z}_{(p)}} R$ -module. In other words, the map $q : q^{-1}(V_1) \rightarrow V_1$ is an isomorphism. This shows (ii).

Using (2.16), we see that $q^{-1}(V_1)$ is regular and locally healthy.

To show that M_G^{ref} is regular and locally healthy, we need to show that its complete local rings at points in characteristic p are quasi-healthy regular. For this, we can now work in the neighborhood of a singular point in $\mathbb{Z}_{p^2} \otimes M_G^{\text{ref}}$ where one of Y or Z vanishes.

Without loss of generality, we can assume that it is the coordinate Y that vanishes, so that $Z \neq 0$. If we set $x_i = X_i/Z$ and $y = Y/Z$, just as in the proof of *loc. cit.* we can work over a Zariski open affine neighborhood $V_2 \subset \mathbb{Z}_{p^2} \otimes M_G^{\text{loc}}$ of the form $\text{Spec } A$ with $A = \mathbb{Z}_{p^2}[x_1, \dots, x_{r-2}, y]/(\sum_i x_i^2 + py)$.

Now, for any \mathbb{Z}_{p^2} -algebra R , the set $q^{-1}(V_2)(R)$ can be identified with the space of pairs $(F^1L_R, F^1(R \otimes_{\mathcal{O}_{F,(p)}} L^\diamond))$, where:

- $F^1L_R \subset L_R$ is an isotropic line spanned by an element with coordinates $(x_1, \dots, x_{r-2}, y, 1)$ for $x_1, \dots, x_{r-2}, y \in R$;
- $F^1(R \otimes_{\mathcal{O}_{F,(p)}} L^\diamond) \subset R \otimes_{\mathcal{O}_{F,(p)}} L^\diamond$ is an isotropic line containing an element with coordinates $(x_1, \dots, x_{r-2}, y, p)$.

Therefore, we find that, as schemes over V_2 , we have

$$q^{-1}(V_2) \xrightarrow{\sim} \text{Proj } A[W_1, \dots, W_{r-2}, U, T]/I,$$

where I is the ideal generated by the elements $\sum_{k=1}^{r-2} W_k^2 + UT$, $x_jW_i - x_iW_j$, for $1 \leq i, j \leq r-2$, $pW_i - x_iU$, $yW_i - x_iT$, for $1 \leq i \leq r-2$, and $yU - pT$.

The scheme $q^{-1}(V_2)$ is covered by three kinds of affine open sub-schemes: one where $U \neq 0$, one where $T \neq 0$ and one where $W_i \neq 0$ for some i . We will consider each in turn.

Setting $w_i = W_i/U$ and $t = T/U$, we can easily see that $q^{-1}(V_2) \cap \{U \neq 0\}$ is isomorphic to

$$\text{Spec } \frac{\mathbb{Z}_{p^2}[w_1, \dots, w_{r-2}, t]}{(\sum_k w_k^2 + t)}.$$

This is clearly smooth over \mathbb{Z}_{p^2} and is therefore regular and locally healthy.

Similarly, setting $w_i = W_i/T$ and $u = U/T$ instead, we find that $q^{-1}(V_2) \cap \{T \neq 0\}$ is isomorphic to

$$\text{Spec } \frac{\mathbb{Z}_{p^2}[w_1, \dots, w_{r-2}, u, y]}{(\sum_k w_k^2 + u, uy - p)} = \text{Spec } \frac{\mathbb{Z}_{p^2}[w_1, \dots, w_{r-2}, y]}{(p + y(\sum_k w_k^2))}.$$

This is a regular scheme over \mathbb{Z}_{p^2} .

Consider the complete local ring of $q^{-1}(V_2) \cap \{T \neq 0\}$ at the point where p and all the coordinates y, w_1, \dots, w_{r-2} vanish. It is

$$R = \frac{\mathbb{Z}_{p^2}[[w_1, \dots, w_{r-2}, y]]}{(p + y(\sum_k w_k^2))}.$$

It admits a surjection to $\mathbb{Z}_{p^2}[[w_1, y]]/(p + yw_1^2)$ and so we can use Theorem 2.14(i) to conclude that R is quasi-healthy. One can check that the order of vanishing of p at all other points of $q^{-1}(V_2) \cap \{T \neq 0\}$ is at most 2 and so we can use assertion (ii) of Theorem 2.14 to conclude that all complete local rings of $q^{-1}(V_2) \cap \{T \neq 0\}$ are quasi-healthy. Hence, this affine open sub-scheme is also locally healthy.

Finally, setting $w_j = W_j/W_1$ for $j \neq 1$, $u = U/W_1$ and $t = T/W_1$, we find that $q^{-1}(V_2) \cap \{W_1 \neq 0\}$ is isomorphic to

$$\text{Spec } \frac{\mathbb{Z}_{p^2}[x_1, w_2, \dots, w_{r-2}, u, t]}{(ux_1 - p, 1 + \sum_{k=1}^{r-2} w_k^2 + ut)}.$$

Set $B' = \mathbb{Z}_{p^2}[x_1, u]/(ux_1 - p)$; then $q^{-1}(V_2) \cap \{W_1 \neq 0\}$ is smooth over $\text{Spec } B'$ and so it suffices to check the criterion of Theorem 2.14(ii) for the complete local rings of B' , which is easy. This shows that $q^{-1}(V_2) \cap \{W_1 \neq 0\}$ is locally healthy, and an identical proof shows that $q^{-1}(V_2) \cap \{W_i \neq 0\}$ is locally healthy for any i . □

Remark 2.19. The map $M_G^{\text{ref}} \rightarrow M_G^{\text{loc}}$ is, up to unique isomorphism, independent of the choice of both F and L^\diamond . In fact, it is simply the blow-up of the singular locus of M_G^{loc} .

3. GSpin Shimura varieties

Let (L, Q) be a quadratic space over $\mathbb{Z}_{(p)}$ of signature $(n, 2)$ with $n \geq 1$. By this, we mean that the largest positive definite sub-space of $L_{\mathbb{R}}$ has dimension n . Let G (respectively G_0) be the smooth $\mathbb{Z}_{(p)}$ -group scheme attached to L in §2.5, so that $G_{\mathbb{Q}} = \text{GSpin}(L_{\mathbb{Q}}, Q)$ (respectively $G_{0,\mathbb{Q}} = \text{SO}(L_{\mathbb{Q}}, Q)$).

3.1 Let X be the space of oriented negative definite 2-planes in $L_{\mathbb{R}}$. The points of X correspond to certain Hodge structures of weight 0 on the vector space $L_{\mathbb{Q}}$, polarized by Q : fix $\mathbf{h} \in X$, and suppose that $(e_{\mathbf{h}}, f_{\mathbf{h}})$ is an oriented, orthogonal basis for the oriented negative definite 2-plane attached to \mathbf{h} with $Q(e_{\mathbf{h}}) = Q(f_{\mathbf{h}}) = -1$. Also, fix a square root of -1 , $\sqrt{-1} \in \mathbb{C}$. Set

$$L_{\mathbf{h}}^{p,q} = \begin{cases} \langle e_{\mathbf{h}} + \sqrt{-1}f_{\mathbf{h}} \rangle \subset L_{\mathbb{C}} & \text{if } (p, q) = (-1, 1), \\ \langle e_{\mathbf{h}}, f_{\mathbf{h}} \rangle^{\perp} \subset L_{\mathbb{C}} & \text{if } (p, q) = (0, 0), \\ \langle e_{\mathbf{h}} - \sqrt{-1}f_{\mathbf{h}} \rangle \subset L_{\mathbb{C}} & \text{if } (p, q) = (1, -1), \\ 0 & \text{otherwise.} \end{cases}$$

Then $L_{\mathbf{h}}$ is a $\mathbb{Z}_{(p)}$ -Hodge structure of weight 0 with underlying $\mathbb{Z}_{(p)}$ -module L ; the associated \mathbb{Q} -Hodge structure is polarized by Q . In fact, each $\mathbf{h} \in X$ gives rise to a unique homomorphism $\mathbb{S} \rightarrow G_{\mathbb{R}}$ which induces the Hodge structure $L_{\mathbf{h}}$ on L and whose restriction to the diagonal subgroup $\mathbb{G}_{m,\mathbb{R}} \subset \mathbb{S}$ is the canonical central embedding $\mathbb{G}_{m,\mathbb{R}} \hookrightarrow G_{\mathbb{R}}$. Here, of course, $\mathbb{S} = \text{Res}_{\mathbb{C}/\mathbb{R}} \mathbb{G}_{m,\mathbb{R}}$ is the Deligne torus.

The map carrying \mathbf{h} to the line $L_{\mathbf{h}}^{1,-1}$ embeds X as an open sub-space of the quadric $\check{X} \subset \mathbb{P}(L_{\mathbb{C}})$ determined by Q . The two connected components of X are switched by complex conjugation on \check{X} . The pairs $(G_{\mathbb{Q}}, X)$ and $(G_{0,\mathbb{Q}}, X)$ are Shimura data, which, since $n \geq 1$, have reflex field \mathbb{Q} .

3.2 Fix a compact open sub-group $K \subset G(\mathbb{A}_f)$ with image $K_0 \subset G_0(\mathbb{A}_f)$, and let $\text{Sh}_K := \text{Sh}_K(G_{\mathbb{Q}}, X)$ and $\text{Sh}_{K_0} := \text{Sh}_{K_0}(G_{0,\mathbb{Q}}, X)$ be the associated Shimura varieties over \mathbb{Q} . We will assume that K is of the form $K_p K^p$, where $K_p = G(\mathbb{Z}_p) \subset G(\mathbb{Q}_p)$ and $K^p \subset G(\mathbb{A}_f^p)$. We will also assume that K^p is chosen to be small enough so that Sh_K is a smooth variety and not just an algebraic space. By weak approximation for $G_{\mathbb{Q}}$ (which can be deduced from weak approximation for its derived group, which is simply connected [PR94, Theorem 7.8]), we have identifications of complex analytic varieties:

$$\text{Sh}_{K,\mathbb{C}}^{\text{an}} = G(\mathbb{Z}_{(p)}) \backslash (X \times G(\mathbb{A}_f^p) / K^p), \tag{3.2.1}$$

$$\text{Sh}_{K_0,\mathbb{C}}^{\text{an}} = G_0(\mathbb{Z}_{(p)}) \backslash (X \times G_0(\mathbb{A}_f^p) / K_0^p). \tag{3.2.2}$$

From this description, we find that the map $\text{Sh}_K \rightarrow \text{Sh}_{K_0}$ is a finite (étale) Galois cover with Galois group

$$\Delta(K) := \mathbb{A}_f^{\times} / \mathbb{Q}^{\times} (K \cap \mathbb{A}_f^{\times}) = \mathbb{A}_f^{p,\times} / \mathbb{Z}_{(p)}^{\times} (K^p \cap \mathbb{A}_f^{p,\times}).$$

Here, we are viewing \mathbb{A}_f^{\times} as a central sub-group of $G(\mathbb{A}_f^{\times})$.

3.3 Let $R \subset \mathbb{R}$ be a $\mathbb{Z}_{(p)}$ -algebra. Recall that a variation of (pure) R -Hodge structures over a smooth complex algebraic variety S is a pair $(\mathbf{U}_B, F^{\bullet}(\mathbf{U}_B \otimes \mathcal{O}_{S^{\text{an}}}))$, where \mathbf{U}_B is a local system of finite free R -modules over S^{an} and $F^{\bullet}(\mathbf{U}_B \otimes \mathcal{O}_{S^{\text{an}}})$ is a descending filtration by sub-vector bundles over S^{an} such that, for every point $s \in S(\mathbb{C})$, the induced pair $(\mathbf{U}_{B,s}, F^{\bullet}(\mathbf{U}_{B,s} \otimes_R \mathbb{C}))$ is a pure R -Hodge structure. An algebraic variation of Hodge structures over S is a tuple $(\mathbf{U}_B, \mathbf{U}_{\text{dR}}, F^{\bullet} \mathbf{U}_{\text{dR}}, \iota)$, where \mathbf{U}_{dR} is a vector bundle over S equipped with an integrable connection

and a filtration $F^\bullet \mathbf{U}_{\text{dR}}$, and $\iota : \mathbf{U}_{\text{dR}}^{\text{an}} \xrightarrow{\cong} \mathbf{U}_B \otimes \mathcal{O}_{S^{\text{an}}}$ is a parallel isomorphism (the right-hand side being endowed with the trivial integrable connection) of vector bundles over S^{an} such that the pair $(\mathbf{U}_B, \iota(F^\bullet \mathbf{U}_{\text{dR}}^{\text{an}}))$ is a variation of R -Hodge structures.

There is a natural exact tensor functor from the category of algebraic R -representations of G (respectively G_0) to the category of algebraic variations of R -Hodge structures on $\text{Sh}_{K,\mathbb{C}}$ (respectively $\text{Sh}_{K_0,\mathbb{C}}$).⁶ This is reasonably well known, but, so as to fix notation, we will now briefly describe it for G and Sh_K ; the situation for G_0 and Sh_{K_0} is completely analogous.

Suppose that we are given an algebraic R -representation U of G . We can view the R -module U as a representation of the discrete group $G(\mathbb{Z}_{(p)})$. Using the uniformization in (3.2.1), we find that the constant local system $U \times X \times G(\mathbb{A}_f^p)/K^p$ (where we equip U with the discrete topology) over $X \times G(\mathbb{A}_f^p)/K^p$ descends to an R -local system \mathbf{U}_B over $\text{Sh}_{K,\mathbb{C}}^{\text{an}}$.

We also have the trivial vector bundle $(U \otimes_R \mathbb{C}) \times X$ over X , where we equip $U \otimes_R \mathbb{C}$ with the natural complex topology. This has a natural, $G(\mathbb{R})$ -equivariant descending filtration by vector sub-bundles $F^\bullet((U \otimes_R \mathbb{C}) \times X)$ such that, at any point $\mathbf{h} \in X$, the induced filtration $F^\bullet(U_{\mathbf{h}} \otimes_R \mathbb{C})$ is simply the Hodge filtration induced by the homomorphism $\mathbb{S} \rightarrow G_{\mathbb{R}}$ attached to \mathbf{h} . Being $G(\mathbb{R})$ -equivariant, and in particular $G(\mathbb{Z}_{(p)})$ -equivariant, the pair $((U \otimes_R \mathbb{C}) \times X, F^\bullet((U \otimes_R \mathbb{C}) \times X))$ descends to a filtered vector bundle over $\text{Sh}_{K,\mathbb{C}}^{\text{an}}$; we will denote this descent by $(\mathbf{U}_{\text{dR},\mathbb{C}}^{\text{an}}, F^\bullet \mathbf{U}_{\text{dR},\mathbb{C}}^{\text{an}})$. By construction, $\mathbf{U}_{\text{dR},\mathbb{C}}^{\text{an}}$ is canonically isomorphic to $\mathbf{U}_B \otimes_R \mathcal{O}_{\text{Sh}_{K,\mathbb{C}}^{\text{an}}}$, and we find that the pair $(\mathbf{U}_B, F^\bullet \mathbf{U}_{\text{dR},\mathbb{C}}^{\text{an}})$ is a variation of R -Hodge structures over $\text{Sh}_{K,\mathbb{C}}^{\text{an}}$.

Now, $\mathbf{U}_{\text{dR},\mathbb{C}}^{\text{an}}$ algebraizes to an algebraic vector bundle with integrable connection $\mathbf{U}_{\text{dR},\mathbb{C}}$. This is essentially due to Baily and Borel [BB66, Theorem 10.14]; cf. also [Har85, (3.1)]. Moreover, by the projectivity of Grassmannians, the filtration $F^\bullet \mathbf{U}_{\text{dR},\mathbb{C}}^{\text{an}}$ algebraizes to a filtration $F^\bullet \mathbf{U}_{\text{dR},\mathbb{C}}$. This finishes our construction of the algebraic variation of R -Hodge structures attached to the representation U : we will denote it by $\mathbb{V}_{\mathbb{C}}(U)$. One can check that $U \mapsto \mathbb{V}_{\mathbb{C}}(U)$ is functorial, exact and respects tensor operations.

As shown in [Del79, § 1.1], the variations of Hodge structures obtained in this fashion satisfy Griffiths’s transversality: the connection on $\mathbf{U}_{\text{dR},\mathbb{C}}$ carries $F^i \mathbf{U}_{\text{dR},\mathbb{C}}$ to $F^{i-1} \mathbf{U}_{\text{dR},\mathbb{C}} \otimes \Omega_{\text{Sh}_{K,\mathbb{C}}/\mathbb{C}}^1$.

3.4 Let $C = C(L, Q)$ be the Clifford algebra for (L, Q) . The above construction applied to the representation $H = C$, on which G acts via left multiplication, produces an algebraic variation of $\mathbb{Z}_{(p)}$ -Hodge structures $\mathbb{V}_{\mathbb{C}}(H) = (\mathbf{H}_B, \mathbf{H}_{\text{dR},\mathbb{C}}, F^\bullet \mathbf{H}_{\text{dR},\mathbb{C}})$ over $\text{Sh}_{K,\mathbb{C}}^{\text{an}}$. Since the right C -action and grading on H are G -equivariant, they are both naturally inherited by $\mathbb{V}_{\mathbb{C}}(H)$.

The tensor $\pi \in H_{\mathbb{Q}}^{\otimes(2,2)}$ is G -invariant, and we can view it as a map of G -representations $\pi : \mathbb{Q} \rightarrow H_{\mathbb{Q}}^{\otimes(2,2)}$, with G acting trivially on \mathbb{Q} . The functoriality of our construction now shows that π induces a map of algebraic variations of \mathbb{Q} -Hodge structures $\mathbb{V}_{\mathbb{C}}(\mathbb{Q}) \rightarrow \mathbb{V}_{\mathbb{C}}(H_{\mathbb{Q}}^{\otimes(2,2)}) = \mathbb{V}_{\mathbb{C}}(H_{\mathbb{Q}})^{\otimes(2,2)}$.

Explicitly, this means that we have a global section $\pi_B \in H^0(\text{Sh}_{K,\mathbb{C}}^{\text{an}}, \mathbf{H}_B^{\otimes(2,2)} \otimes \mathbb{Q})$ and a section $\pi_{\text{dR},\mathbb{C}} \in H^0(\text{Sh}_{K,\mathbb{C}}, F^0 \mathbf{H}_{\text{dR},\mathbb{C}}^{\otimes(2,2)})$ that is parallel for the connection on $\mathbf{H}_{\text{dR},\mathbb{C}}^{\otimes(2,2)}$. Moreover, π_B is carried to $\pi_{\text{dR},\mathbb{C}}$ under the comparison isomorphism

$$\mathbf{H}_B^{\otimes(2,2)} \otimes \mathcal{O}_{\text{Sh}_{K,\mathbb{C}}^{\text{an}}} \xrightarrow{\cong} \mathbf{H}_{\text{dR},\mathbb{C}}^{\otimes(2,2),\text{an}}.$$

In particular, for every point $s \in \text{Sh}_K(\mathbb{C})$, the fiber $\pi_{B,s} \in \mathbf{H}_{B,s}^{\otimes(2,2)} \otimes \mathbb{Q}$ is a Hodge tensor.

⁶ Such a result is true for any Shimura variety.

We can view π as an idempotent endomorphism of $\mathbb{V}_{\mathbb{C}}(H_{\mathbb{Q}}^{\otimes(1,1)})$ and, by construction, its image is precisely $\mathbb{V}_{\mathbb{C}}(L_{\mathbb{Q}}) = (\mathbf{L}_{B,\mathbb{Q}}, \mathbf{L}_{\mathrm{dR},\mathbb{C}}, F^{\bullet}\mathbf{L}_{\mathrm{dR},\mathbb{C}})$, the algebraic variation of \mathbb{Q} -Hodge structures attached to the representation $L_{\mathbb{Q}}$.

In fact, it follows from (1.5) that the image under π of $\mathbb{V}_{\mathbb{C}}(H^{\otimes(1,1)}) \subset \mathbb{V}_{\mathbb{C}}(H_{\mathbb{Q}}^{\otimes(1,1)})$ is exactly the variation of $\mathbb{Z}_{(p)}$ -Hodge structures $\mathbb{V}_{\mathbb{C}}(L^{\vee})$.

Moreover, the G -equivariant map $L_{\mathbb{Q}} \rightarrow L_{\mathbb{Q}}^{\vee}$ induced by the pairing gives rise to an isomorphism $\mathbb{V}_{\mathbb{C}}(L_{\mathbb{Q}}) \xrightarrow{\sim} \mathbb{V}_{\mathbb{C}}(L_{\mathbb{Q}}^{\vee})$. This corresponds to a pairing $\mathbb{V}_{\mathbb{C}}(L_{\mathbb{Q}}) \otimes \mathbb{V}_{\mathbb{C}}(L_{\mathbb{Q}}) \rightarrow \mathbb{V}_{\mathbb{C}}(\mathbb{Q})$, which is in fact a polarization of variations of Hodge structures. $\mathbb{V}_{\mathbb{C}}(L)$ is precisely the pre-image of $\mathbb{V}_{\mathbb{C}}(L^{\vee})$ under this isomorphism.

In sum, we find that we can recover the tuple $\mathbb{V}_{\mathbb{C}}(L) = (\mathbf{L}_B, \mathbf{L}_{\mathrm{dR},\mathbb{C}}, F^{\bullet}\mathbf{L}_{\mathrm{dR},\mathbb{C}})$ from the data of $\mathbb{V}_{\mathbb{C}}(H)$ and the idempotent operator $\pi \in \mathbb{V}_{\mathbb{C}}(H_{\mathbb{Q}}^{\otimes(2,2)})$.

3.5 For $\delta \in C \cap C_{\mathbb{Q}}^{\times}$ satisfying $\delta^* = -\delta$, set $\mathcal{G}_{\delta,\mathbb{Q}} = \mathrm{GSp}(H_{\mathbb{Q}}, \psi_{\delta})$, so that we have an embedding $G_{\mathbb{Q}} \hookrightarrow \mathcal{G}_{\delta,\mathbb{Q}}$. Let \mathcal{X} be the space of lagrangian sub-spaces $W \subset H_{\mathbb{C}}$ (with respect to the form ψ_{δ}) such that the hermitian form $\sqrt{-1}\psi_{\delta}(w_1, \bar{w}_2)$ restricts to a (positive or negative) definite form on W : this is simply the union of the Siegel half-spaces attached to (H, ψ_{δ}) .

LEMMA 3.6. *One can choose δ so that the embedding $G_{\mathbb{Q}} \hookrightarrow \mathcal{G}_{\delta,\mathbb{Q}}$ induces an embedding of Shimura data $(G_{\mathbb{Q}}, X) \hookrightarrow (\mathcal{G}_{\delta,\mathbb{Q}}, \mathcal{X})$. \square*

Let $\mathcal{K}_p \subset \mathcal{G}_{\delta,\mathbb{Q}}(\mathbb{Q}_p)$ be the stabilizer of $H_{\mathbb{Z}_p}$. Then we have $K_p \subset \mathcal{K}_p \cap G(\mathbb{Q}_p)$. Let $K = K_p K^p \subset G(\mathbb{A}_f)$ be as above; then, for any compact open $\mathcal{K}^p \subset \mathcal{G}_{\delta}(\mathbb{A}_f^p)$ containing K_p , we have a finite, unramified⁷ map of canonical models of Shimura varieties over \mathbb{Q} :

$$\mathrm{Sh}_K \rightarrow \mathrm{Sh}_{\mathcal{K}}(\mathcal{G}_{\delta,\mathbb{Q}}, \mathcal{X}).$$

Here, $\mathcal{K} = \mathcal{K}_p \mathcal{K}^p$. We will call this a *Kuga-Satake map*.

3.7 The scheme $\mathrm{Sh}_{\mathcal{K}} := \mathrm{Sh}_{\mathcal{K}}(\mathcal{G}_{\delta,\mathbb{Q}}, \mathcal{X})$ has a natural moduli description. To describe this, we will work with abelian schemes up to prime-to- p isogeny. More precisely, given a scheme T , the category $AV_{(p)}(T)$ of abelian schemes up to prime-to- p isogeny has for its objects abelian schemes A over T , where, for two abelian schemes A and B over T , the space of morphisms from A to B is the $\mathbb{Z}_{(p)}$ -module

$$\mathrm{Hom}(A, B)_{(p)} = \mathrm{Hom}(A, B) \otimes \mathbb{Z}_{(p)}.$$

Given an abelian scheme A over T , a *quasi-polarization* (or simply *polarization*) of A in $AV_{(p)}(T)$ will be an element $\lambda \in \mathrm{Hom}(A, A^{\vee})_{(p)}$ that is a positive multiple of a polarization $\lambda' : A \rightarrow A^{\vee}$.

Given any scheme T , a prime $\ell \neq p$ invertible in T and an abelian scheme $f : A \rightarrow T$, we can consider the associated relative first ℓ -adic cohomology sheaf $R^1 f_* \underline{\mathbb{Q}}_{\ell}$. If p is invertible in T , we can also consider the p -adic sheaf $R^1 f_* \underline{\mathbb{Z}}_p$. Both of these constructions are invariants of the prime-to- p isogeny class of A . Let $\underline{\mathbb{Q}}_{\ell}(-1)$ be the Tate twist: it is the relative first ℓ -adic cohomology of $\mathbb{G}_{m,T}$ over T .

Given an abelian scheme $f : A \rightarrow T$ over a $\mathbb{Z}_{(p)}$ -scheme T , the following constructions are invariants of its prime-to- p isogeny class: the p -divisible group $A[p^{\infty}]$; the p -adic étale cohomology $R^1 f_* \underline{\mathbb{Z}}_p$; for $\ell \neq p$, the rational ℓ -adic cohomology $R^1 f_* \underline{\mathbb{Q}}_{\ell}$; and the \mathbb{A}_f^p -adic cohomology $R^1 f_* \underline{\mathbb{A}}_f^p$. If T is an \mathbb{F}_p -scheme, we also have the degree 1 crystalline cohomology of A over T : for instance, this can be viewed as the Dieudonné crystal associated with the p -divisible group $A[p^{\infty}]$.

⁷ Note that, over \mathbb{C} and locally in the complex analytic topology, this map is isomorphic to the closed immersion $X \hookrightarrow \mathcal{X}$.

3.8 For some purposes, it is useful to embed $AV_{(p)}(T)$ in the category of group schemes over T . We will follow [Har, § 1] for this. Given an abelian scheme $A \rightarrow T$, let $\mathcal{T}^p(A)$ be the inverse system consisting of prime-to- p isogenies⁸ $B \rightarrow A$. Then $\mathcal{T}^p(A)$ has a co-final system consisting of the finite étale multiplication-by- n endomorphisms $[n] : A \rightarrow A$ for $n \in \mathbb{Z}$ with $p \nmid n$. Therefore, the inverse limit

$$\hat{A}^{(p)} = \varprojlim_{(B \rightarrow A) \in \mathcal{T}^p(A)} B$$

exists in the category of group schemes over T . One sees that $A \mapsto \hat{A}^{(p)}$ is a fully faithful functor from $AV^{(p)}(T)$ to the category of T -group schemes.

The main purpose of this construction is the following abuse of terminology: given $A \in AV^{(p)}(T)$ and a T -scheme T' , we will write $H^0(T', A)$ for the $\mathbb{Z}_{(p)}$ -module $H^0(T', \hat{A}^{(p)})$. This allows us to speak of ‘sections of A ’.

From now on we will suppress the qualifying phrase ‘up to prime-to- p isogeny’: *all abelian schemes will only be considered in the prime-to- p isogeny category*. The cohomological constructions we are concerned with will all be invariants of the prime-to- p isogeny class.

3.9 Suppose that T is a $\mathbb{Z}_{(p)}$ -scheme and $f : A \rightarrow T$ is an abelian scheme. Given a polarization $\lambda : A \rightarrow A^\vee$, we get an induced non-degenerate Poincaré pairing of \mathbb{A}_f^p -sheaves

$$\psi_\lambda : R^1 f_* \mathbb{A}_f^p \otimes R^1 f_* \mathbb{A}_f^p \rightarrow \mathbb{A}_f^p(-1).$$

Suppose that we are also given an isomorphism

$$\eta : H \otimes \mathbb{A}_f^p \xrightarrow{\sim} R^1 f_* \mathbb{A}_f^p$$

of \mathbb{A}_f^p -sheaves over T . Then we obtain two different non-degenerate pairings on $H \otimes \mathbb{A}_f^p$: the first is the constant pairing into \mathbb{A}_f^p arising from ψ_δ , which we will again call ψ_δ ; and the second is the pairing $\eta^* \psi_\lambda$ into $\mathbb{A}_f^p(-1)$ obtained by pulling back ψ_λ along η . In particular, the existence of η implies that $\mathbb{A}_f^p(-1)$ is trivializable over T . We say that η *preserves polarizations* if, for some choice of isomorphism $\mathbb{A}_f^p(-1) \xrightarrow{\sim} \mathbb{A}_f^p$, the pairings $\eta^* \psi_\lambda$ and ψ_δ agree.

In the above situation, we will consider the étale sheaf $I^p(A, \lambda)$ over T , whose sections are polarization preserving isomorphisms of \mathbb{A}_f^p -sheaves $\eta : H \otimes \mathbb{A}_f^p \xrightarrow{\sim} R^1 f_* \mathbb{A}_f^p$. Note that $I^p(A, \lambda)$ is a pseudo-torsor under $\mathcal{G}_\delta(\mathbb{A}_f^p)$ via its action through pre-composition.

For any $\mathbb{Z}_{(p)}$ -scheme T , let $\mathcal{S}_\mathcal{K}(T)$ be the set of isomorphism classes of tuples $(A, \lambda, [\eta])$, where:

- (A, λ) is a polarized abelian scheme over T ;
- $[\eta]$ is a \mathcal{K}^p -level structure: it is a section of the quotient sheaf $I^p(A, \lambda)/\mathcal{K}^p$.

For \mathcal{K} sufficiently small, the functor $\mathcal{S}_\mathcal{K}$ is (represented by) a quasi-projective scheme over $\mathbb{Z}_{(p)}$, whose generic fiber is canonically identified with $\text{Sh}_\mathcal{K} := \text{Sh}_\mathcal{K}(\mathcal{G}_{\delta, \mathbb{Q}}, \mathcal{X})$.

3.10 Over $\mathcal{S}_\mathcal{K}$, we have the tautological tuple $(A, \lambda, [\eta])$. Let $(A_{\text{Sh}_\mathcal{K}}^{\text{KS}}, \lambda_{\text{Sh}_\mathcal{K}}^{\text{KS}}, [\eta^{\text{KS}}])$ be the induced tuple over $\text{Sh}_\mathcal{K}$. We will refer to $A_{\text{Sh}_\mathcal{K}}^{\text{KS}}$ as the *Kuga–Satake abelian scheme* over $\text{Sh}_\mathcal{K}$.

The identification of $\text{Sh}_\mathcal{K}$ with the generic fiber of the moduli scheme $\mathcal{S}_\mathcal{K}$ has the following property: over $\text{Sh}_{K, \mathbb{C}}^{\text{an}}$, the algebraic variation of $\mathbb{Z}_{(p)}$ -Hodge structures obtained from the degree 1 cohomology of $A_{\text{Sh}_{K, \mathbb{C}}}^{\text{KS}}$ is canonically identified with $\mathbb{V}_\mathbb{C}(H)$.

⁸ These are finite, flat homomorphisms, whose kernel is an étale group scheme of order not divisible by p .

Since $\mathbb{V}_{\mathbb{C}}(H)$ carries a $\mathbb{Z}/2\mathbb{Z}$ -grading and a right C -action, we conclude that $A_{\text{Sh}_{K,\mathbb{C}}}^{\text{KS}}$, and hence $A_{\text{Sh}_{K,\mathbb{C}}}^{\text{KS}}$, admits a canonical $\mathbb{Z}/2\mathbb{Z}$ -grading and a (left) C -action.⁹ Here, we are using the (anti-)equivalence of categories between abelian schemes over $\text{Sh}_{K,\mathbb{C}}$ and polarizable variations of $\mathbb{Z}_{(p)}$ -Hodge structures over $\text{Sh}_{K,\mathbb{C}}$ of weight 1.

The above implies that the degree 1 relative Betti cohomology of $A_{\text{Sh}_{K,\mathbb{C}}}^{\text{KS}}$ with coefficients in $\mathbb{Z}_{(p)}$ can be identified with \mathbf{H}_B as a $\mathbb{Z}_{(p)}$ -local system. Similarly, the degree 1 relative de Rham cohomology of $A_{\text{Sh}_{K,\mathbb{C}}}^{\text{KS}}$ can be identified with $(\mathbf{H}_{\text{dR},\mathbb{C}}, F^\bullet \mathbf{H}_{\text{dR},\mathbb{C}})$ as a filtered vector bundle with integrable connection.

Let \mathbf{H}_ℓ (respectively \mathbf{H}_p) be the relative degree 1 étale cohomology of $A_{\text{Sh}_K}^{\text{KS}}$ over Sh_K with coefficients in \mathbb{Q}_ℓ for $\ell \neq p$ (respectively \mathbb{Z}_p). Then, over $\text{Sh}_{K,\mathbb{C}}^{\text{an}}$, Artin’s comparison theorem gives us canonical isomorphisms of local systems:

$$\alpha_\ell : \mathbf{H}_B \otimes \mathbb{Q}_\ell \xrightarrow{\cong} \mathbf{H}_\ell|_{\text{Sh}_{K,\mathbb{C}}^{\text{an}}} \quad \text{for } \ell \neq p, \tag{3.10.1}$$

$$\alpha_p : \mathbf{H}_B \otimes \mathbb{Z}_p \xrightarrow{\cong} \mathbf{H}_p|_{\text{Sh}_{K,\mathbb{C}}^{\text{an}}}. \tag{3.10.2}$$

Let $(\mathbf{H}_{\text{dR},\mathbb{Q}}, F^\bullet \mathbf{H}_{\text{dR},\mathbb{Q}})$ be the relative degree 1 de Rham cohomology of $A_{\text{Sh}_K}^{\text{KS}}$ over Sh_K : this is equipped with the Hodge filtration and the Gauss–Manin connection. Over $\text{Sh}_{K,\mathbb{C}}^{\text{an}}$, there now exists a canonical de Rham comparison isomorphism, parallel for the trivial connection on the left-hand side:

$$\alpha_{\text{dR}} : \mathbf{H}_B \otimes \mathcal{O}_{\text{Sh}_{K,\mathbb{C}}^{\text{an}}} \xrightarrow{\cong} \mathbf{H}_{\text{dR},\mathbb{C}}^{\text{an}} \xrightarrow{\cong} \mathbf{H}_{\text{dR},\mathbb{Q}}|_{\text{Sh}_{K,\mathbb{C}}^{\text{an}}}. \tag{3.10.3}$$

Note that the first isomorphism in this composition carries $\pi_B \otimes 1$ to $\pi_{\text{dR},\mathbb{C}}$.

PROPOSITION 3.11.

- (i) The structures of the $\mathbb{Z}/2\mathbb{Z}$ -grading and the C -action on $A_{\text{Sh}_{K,\mathbb{C}}}^{\text{KS}}$ descend (necessarily uniquely) to $A_{\text{Sh}_K}^{\text{KS}}$.
- (ii) For any prime ℓ , the global section $\alpha_\ell(\pi_B \otimes 1)$ of $\mathbf{H}_\ell^{\otimes(2,2)} \otimes \mathbb{Q}$ over $\text{Sh}_{K,\mathbb{C}}^{\text{an}}$ arises from a (necessarily unique) section

$$\pi_\ell \in H^0(\text{Sh}_K, \mathbf{H}_\ell^{\otimes(2,2)} \otimes \mathbb{Q}).$$

- (iii) The section $\pi_{\text{dR},\mathbb{C}}$ of $\mathbf{H}_{\text{dR},\mathbb{C}}^{\otimes(2,2)}$ descends (necessarily uniquely) to a parallel section

$$\pi_{\text{dR},\mathbb{Q}} \in H^0(\text{Sh}_K, F^0 \mathbf{H}_{\text{dR},\mathbb{Q}}^{\otimes(2,2)}).$$

Proof. This can be extracted from [Kis10, § 2.2]. We sketch the proof.

There is a canonical pro-finite Galois cover $\text{Sh}_{K^p} \rightarrow \text{Sh}_K$ with Galois group K_p :

$$\text{Sh}_{K^p} = \varprojlim_{K'_p \subset K_p} \text{Sh}_{K'_p K^p}.$$

Here, K'_p runs over the compact open sub-groups of K_p .

If we fix a connected component $S \subset \text{Sh}_K$ and a geometric point $\bar{s} \rightarrow S$, pulling the pro-finite cover Sh_{K^p} back over S gives us a K_p -torsor over S , which in turn corresponds to a map $\pi_1(S, \bar{s}) \rightarrow K_p$. The restriction of \mathbf{H}_p to S is precisely the p -adic sheaf attached to the composite representation

$$\pi_1(S, \bar{s}) \rightarrow K_p = G(\mathbb{Z}_p) \subset \text{Aut}(H \otimes \mathbb{Z}_p).$$

⁹ The right action is converted to a left, because cohomology is a contravariant functor.

Since the $\mathbb{Z}/2\mathbb{Z}$ -grading and the C -action on $H \otimes \mathbb{Z}_p$ are K_p -invariant, and hence $\pi_1(S, \bar{s})$ -invariant, we find that the corresponding structures on $\mathbf{H}_B \otimes \mathbb{Z}_p$ over $S_{\mathbb{C}}^{\text{an}}$ must in fact descend to structures on \mathbf{H}_p over S . From this, one can deduce (i). The essential point is the following simple consequence of Galois descent: suppose that A and B are abelian varieties over a characteristic 0 field κ , and $f : A_{\bar{\kappa}} \rightarrow B_{\bar{\kappa}}$ is a map of abelian varieties over an algebraic closure $\bar{\kappa}/\kappa$ such that the induced map on p -adic Tate modules $T_p(f) : T_p(A_{\bar{\kappa}}) \rightarrow T_p(B_{\bar{\kappa}})$ is equivariant for the action of the absolute Galois group $\text{Gal}(\bar{\kappa}/\kappa)$. Then f is defined over κ .

Similarly, since the tensor $\pi \in H^{\otimes(2,2)} \otimes \mathbb{Q}_p$ is also K_p -invariant, $\alpha_p(\pi_B \otimes 1)$ must descend to a section $\pi_p \in H^0(\text{Sh}_K, \mathbf{H}_p^{\otimes(2,2)} \otimes \mathbb{Q}_p)$. This shows the case $\ell = p$ of (ii). The de Rham case (iii) and the $\ell \neq p$ case of (ii) follow from this and the fact that all Hodge cycles on abelian varieties are absolutely Hodge [DMOS82, Ch. I]; cf. the proof of [Kis10, 2.2.2]. There are two main points [DMOS82, §I.2].

- Given a projective variety X over κ and an algebraically closed overfield $K \supset \bar{\kappa}$, any absolute Hodge cycle β over X_K is already defined over $X_{\bar{\kappa}}$; that is, the de Rham realization β_{dR} of β lies in $H_{\text{dR}}^{\bullet}(X_{\bar{\kappa}}/\bar{\kappa})^{\otimes}$.
- If one cohomological realization of β is fixed by $\text{Gal}(\bar{\kappa}/\kappa)$, then all of its realizations are.

We apply this to the situation where κ is the function field of an irreducible component S as above, K is the function field of an irreducible component of $S_{\mathbb{C}}$, X is the fiber of $A_{\text{Sh}_K}^{\text{KS}}$ over κ and β is the Hodge (hence absolutely Hodge) cycle over X_K obtained from the fiber of the Betti realization π_B . □

3.12 The descent of the realizations of π proven above now allows us to descend $\mathbb{V}_{\mathbb{C}}(L)$ over Sh_K .

For a prime $\ell \neq p$, let $\mathbf{L}_{\ell} \subset \mathbf{H}_{\ell}^{\otimes(1,1)}$ be the image of the idempotent operator π_{ℓ} . Then \mathbf{L}_{ℓ} is a \mathbb{Q}_{ℓ} -local system over Sh_K equipped with a non-degenerate pairing $\mathbf{L}_{\ell} \times \mathbf{L}_{\ell} \rightarrow \underline{\mathbb{Q}}_{\ell}$. Over $\text{Sh}_{K,\mathbb{C}}^{\text{an}}$, we have a canonical comparison isomorphism (respecting pairings)

$$\mathbf{L}_B \otimes \mathbb{Q}_{\ell} \xrightarrow{\cong} \mathbf{L}_{\ell}|_{\text{Sh}_{K,\mathbb{C}}^{\text{an}}}.$$

For $\ell = p$, as in §3.4, we can show that the image of $\mathbf{H}_p^{\otimes(1,1)}$ under π_p is a descent \mathbf{L}_p^{\vee} of the \mathbb{Z}_p -local system $\mathbf{L}_B^{\vee} \otimes \mathbb{Z}_p$ over $\text{Sh}_{K,\mathbb{C}}^{\text{an}}$. It is equipped with a quadratic form with values in $\underline{\mathbb{Q}}_p$, and $\mathbf{L}_p \subset \mathbf{L}_p^{\vee}$ is recovered as the largest sub-local system whose pairing with \mathbf{L}_p^{\vee} takes values in $\underline{\mathbb{Z}}_p$. Again, we have a canonical isometric comparison isomorphism

$$\mathbf{L}_B \otimes \mathbb{Z}_p \xrightarrow{\cong} \mathbf{L}_p|_{\text{Sh}_{K,\mathbb{C}}^{\text{an}}}.$$

Similarly, let $\mathbf{L}_{\text{dR},\mathbb{Q}} \subset \mathbf{H}_{\text{dR},\mathbb{Q}}^{\otimes(1,1)}$ be the image of the idempotent operator $\pi_{\text{dR},\mathbb{Q}}$. By construction, $\pi_{\text{dR},\mathbb{Q}}$ respects the Hodge filtration and is parallel for the Gauss–Manin connection. Therefore, $\mathbf{L}_{\text{dR},\mathbb{Q}}$ inherits the connection as well as a filtration $F^{\bullet} \mathbf{L}_{\text{dR},\mathbb{Q}}$, and $(\mathbf{L}_{\text{dR},\mathbb{Q}}, F^{\bullet} \mathbf{L}_{\text{dR},\mathbb{Q}})$ is a descent of $(\mathbf{L}_{\text{dR},\mathbb{C}}, F^{\bullet} \mathbf{L}_{\text{dR},\mathbb{C}})$ as a filtered vector bundle with flat connection.

3.13 Let $\mathbf{H}_{\mathbb{A}_f^p}$ be the \mathbb{A}_f^p -valued degree 1 étale cohomology of $A_{\text{Sh}_K}^{\text{KS}}$: it can be viewed as a descent of the analytic local system $\mathbf{H}_B \otimes_{\mathbb{Z}(p)} \mathbb{A}_f^p$. There is a unique idempotent operator $\pi_{\mathbb{A}_f^p}$ on $\mathbf{H}_{\mathbb{A}_f^p}^{\otimes(1,1)}$ such that, for any prime $\ell \neq p$, the ℓ -adic component of $\pi_{\mathbb{A}_f^p}$ is π_{ℓ} . Let $\mathbf{L}_{\mathbb{A}_f^p} \subset \mathbf{H}_{\mathbb{A}_f^p}^{\otimes(1,1)}$ be the image of $\pi_{\mathbb{A}_f^p}$: its ℓ -adic component for any $\ell \neq p$ is simply \mathbf{L}_{ℓ} .

Let I_G^p be the sub-sheaf of $I^p(A_{\text{Sh}_K}^{\text{KS}}, \lambda_{\text{Sh}_K}^{\text{KS}})$ consisting of C -equivariant, $\mathbb{Z}/2\mathbb{Z}$ -graded isomorphisms

$$\eta : H \otimes \underline{\mathbb{A}}_f^p \xrightarrow{\cong} \mathbf{H}_{\mathbb{A}_f^p}$$

that carry $L \otimes \underline{\mathbb{A}}_f^p \subset H^{\otimes(1,1)} \otimes \underline{\mathbb{A}}_f^p$ onto $\mathbf{L}_{\mathbb{A}_f^p} \subset \mathbf{H}_{\mathbb{A}_f^p}^{\otimes(1,1)}$. Note that $G(\mathbb{A}_f^p)$ naturally acts on the right on I_G^p via pre-composition, making it a torsor under $G(\mathbb{A}_f^p)$. We have a natural map of quotient sheaves

$$I_G^p/K^p \rightarrow I^p(A_{\text{Sh}_K}^{\text{KS}}, \lambda_{\mathbb{Q}}^{\text{KS}})/K^p.$$

A section of I_G^p/K^p will be called a K^p -level structure.

PROPOSITION 3.14. *There is a canonical K^p -level structure $[\eta_G] \in H^0(\text{Sh}_K, I_G^p/K^p)$ such that $[\eta^{\text{KS}}]$ is its image in $I^p(A_{\text{Sh}_K}^{\text{KS}}, \lambda_{\text{Sh}_K}^{\text{KS}})/K^p$.*

Proof. Consider the pro-étale cover $\mathcal{S}_{K^p} \rightarrow \mathcal{S}_K$. Here,

$$\mathcal{S}_{K^p} = \varprojlim_{K'^p \subset K^p} \mathcal{S}_{K^p K'^p},$$

where K'^p varies over the compact open sub-groups of K^p . Then there is a canonical isomorphism of $\mathcal{G}_\delta(\mathbb{A}_f^p)$ -torsors:

$$I^p(A, \lambda) \xrightarrow{\cong} \mathcal{S}_{K^p} \times^{K^p} \mathcal{G}_\delta(\mathbb{A}_f^p) := (\mathcal{S}_{K^p} \times \mathcal{G}_\delta(\mathbb{A}_f^p))/K^p.$$

The canonical K -level structure $[\eta]$ over \mathcal{S}_K is now obtained from the tautological section $\eta \in H^0(\mathcal{S}_{K^p}, I^p(A, \lambda))$.

Similarly, the $G(\mathbb{A}_f^p)$ -torsor I_G^p over Sh_K has a canonical reduction of structure group to a K^p -torsor pro-represented by the pro-finite cover $\text{Sh}_{K^p} \rightarrow \text{Sh}_K$, where

$$\text{Sh}_{K^p} = \varprojlim_{K'^p \subset K^p} \text{Sh}_{K^p K'^p}.$$

Here, K'^p varies over the compact open sub-groups of K^p .

Therefore, the image $[\eta_G] \in H^0(\text{Sh}_K, I_G^p/K^p)$ of the tautological section $\eta_G \in H^0(\text{Sh}_{K^p}, I_G^p)$ is the K^p -level structure we seek. \square

3.15 Notice that, for any algebraic representation U of G , the $G(\mathbb{A}_f^p)$ -torsor I_G^p allows us to functorially descend the \mathbb{A}_f^p -adic local system $\mathbf{U}_B \otimes \mathbb{A}_f^p$ over Sh_K . We take it to be the contraction product:

$$\mathbf{U}_{\mathbb{A}_f^p} := I_G^p \times^{G(\mathbb{A}_f^p)} \mathbf{U}(\mathbb{A}_f^p).$$

Similarly, the K_p -torsor $I_{p,G} := \text{Sh}_{K^p} \rightarrow \text{Sh}_K$ used in the proof of (3.11) allows us to functorially descend the p -adic sheaf $\mathbf{U}_B \otimes \mathbb{Z}_p$:

$$\mathbf{U}_p := I_{p,G} \times^{G(\mathbb{Z}_p)} \mathbf{U}(\mathbb{Z}_p).$$

We can also descend the filtered vector bundle with connection $(\mathbf{U}_{\text{dR},\mathbb{C}}, F^\bullet \mathbf{U}_{\text{dR},\mathbb{C}})$ canonically to a pair $(\mathbf{U}_{\text{dR},\mathbb{Q}}, F^\bullet \mathbf{U}_{\text{dR},\mathbb{Q}})$ over Sh_K . To do this, consider the functor $\mathcal{P}_{\text{dR},\mathbb{Q}}$ on Sh_K -schemes that assigns to any Sh_K -scheme T the set

$$\mathcal{P}_{\text{dR},\mathbb{Q}}(T) = \left(\begin{array}{l} C\text{-equivariant } \mathbb{Z}/2\mathbb{Z}\text{-graded } \mathcal{O}_T\text{-module isomorphisms} \\ \xi : H \otimes_{\mathbb{Z}(p)} \mathcal{O}_T \xrightarrow{\cong} \mathbf{H}_{\text{dR},\mathbb{Q},T} \\ \text{carrying } L \otimes \mathcal{O}_T \subset \mathbf{H}^{\otimes(2,2)} \otimes \mathcal{O}_T \text{ onto } \mathbf{L}_{\text{dR},\mathbb{Q},T} \subset \mathbf{H}_{\text{dR},\mathbb{Q},T}^{\otimes(2,2)} \end{array} \right).$$

The algebraic group $G_{\mathbb{Q}}$ acts on $\mathcal{P}_{\mathrm{dR},\mathbb{Q}}$ by pre-composition. By working over $\mathrm{Sh}_{K,\mathbb{C}}^{\mathrm{an}}$, we can show that $\mathcal{P}_{\mathrm{dR},\mathbb{Q}}$ is a $G_{\mathbb{Q}}$ -torsor over Sh_K . Observe that the connection on $\mathbf{H}_{\mathrm{dR},\mathbb{Q}}$ equips $\mathcal{P}_{\mathrm{dR},\mathbb{Q}}$ with an integrable connection. For any representation U as above, we can now take $\mathbf{U}_{\mathrm{dR},\mathbb{Q}}$ to be the contraction product:

$$\mathbf{U}_{\mathrm{dR},\mathbb{Q}} = \mathcal{P}_{\mathrm{dR},\mathbb{Q}} \times^{G_{\mathbb{Q}}} U := (\mathcal{P}_{\mathrm{dR},\mathbb{Q}} \times_{\mathrm{Spec} \mathbb{Q}} U) / G_{\mathbb{Q}}.$$

Here, we are viewing U as a vector bundle over $\mathrm{Spec} \mathbb{Q}$, and $G_{\mathbb{Q}}$ acts diagonally on the product $\mathcal{P}_{\mathrm{dR},\mathbb{Q}} \times U$.

To construct the filtration $F^{\bullet} \mathbf{U}_{\mathrm{dR},\mathbb{Q}}$, we fix an isotropic line $F^1 L \subset L$. The stabilizer of this line is a parabolic sub-group $P_{\mathbb{Q}} \subset G_{\mathbb{Q}}$. Now, consider the sub-functor $\mathcal{P}_{\mathrm{dR},P_{\mathbb{Q}}} \subset \mathcal{P}_{\mathrm{dR},\mathbb{Q}}$ given by

$$\mathcal{P}_{\mathrm{dR},P_{\mathbb{Q}}}(T) = (\xi \in \mathcal{P}_{\mathrm{dR},\mathbb{Q}}(T) : \xi(F^1 L \otimes \mathcal{O}_T) = F^1 \mathbf{L}_{\mathrm{dR},\mathbb{Q},T}).$$

This is a $P_{\mathbb{Q}}$ -torsor over Sh_K .

Choose any co-character $\mu : \mathbb{G}_{m,\mathbb{Q}} \rightarrow P_{\mathbb{Q}}$ splitting the two-step filtration $0 \subset F^1 H_{\mathbb{Q}} \subset H_{\mathbb{Q}}$ with $F^1 H_{\mathbb{Q}} = \ker(F^1 L_{\mathbb{Q}})$ (cf. 1.9). On all $G_{\mathbb{Q}}$ -representations U , the action of $\mu(\mathbb{G}_m)$ produces a grading and hence a filtration $F^{\bullet} U_{\mathbb{Q}}$ that is stabilized by $P_{\mathbb{Q}}$, and is independent of the choice of μ . We now set

$$F^{\bullet} \mathbf{U}_{\mathrm{dR},\mathbb{Q}} := \mathcal{P}_{\mathrm{dR},P_{\mathbb{Q}}} \times^{P_{\mathbb{Q}}} F^{\bullet} U.$$

The pair $(\mathbf{U}_{\mathrm{dR},\mathbb{Q}}, F^{\bullet} \mathbf{U}_{\mathrm{dR},\mathbb{Q}})$ is the descent that we seek.

It is easy to check that when $U = H$ (for which it is essentially tautological) or $U = L$, these constructions agree with the ones already given above.

In sum, we have defined a functor $\mathbb{V}_{\mathbb{Q}}$ from the category of algebraic representations U of G to the category of tuples

$$(\mathbf{U}_B, (\mathbf{U}_p, \alpha_p), (\mathbf{U}_{\mathbb{A}_f^p}, \alpha_{\mathbb{A}_f^p}), (\mathbf{U}_{\mathrm{dR},\mathbb{Q}}, F^{\bullet} \mathbf{U}_{\mathrm{dR},\mathbb{Q}}, \alpha_{\mathrm{dR}})),$$

where \mathbf{U}_B is a local system over $\mathrm{Sh}_{K,\mathbb{C}}^{\mathrm{an}}$; \mathbf{U}_p (respectively $\mathbf{U}_{\mathbb{A}_f^p}$) are locally constant sheaves over Sh_K equipped with comparison isomorphisms:

$$\begin{aligned} \alpha_p &: \mathbf{U}_B \otimes \mathbb{Z}_p \xrightarrow{\cong} \mathbf{U}_p|_{\mathrm{Sh}_{K,\mathbb{C}}^{\mathrm{an}}}, \\ \alpha_{\mathbb{A}_f^p} &: \mathbf{U}_B \otimes \mathbb{A}_f^p \xrightarrow{\cong} \mathbf{U}_{\mathbb{A}_f^p}|_{\mathrm{Sh}_{K,\mathbb{C}}^{\mathrm{an}}}. \end{aligned}$$

Moreover, $(\mathbf{U}_{\mathrm{dR},\mathbb{Q}}, F^{\bullet} \mathbf{U}_{\mathrm{dR},\mathbb{Q}})$ is a filtered vector bundle over Sh_K with integrable connection, equipped with a parallel comparison isomorphism:

$$\alpha_{\mathrm{dR}} : \mathbf{U}_B \otimes \mathcal{O}_{\mathrm{Sh}_{K,\mathbb{C}}^{\mathrm{an}}} \xrightarrow{\cong} \mathbf{U}_{\mathrm{dR},\mathbb{Q}}|_{\mathrm{Sh}_{K,\mathbb{C}}^{\mathrm{an}}}.$$

Finally, we require that $(\mathbf{U}_B, \alpha_{\mathrm{dR}}^{-1}(F^{\bullet} \mathbf{U}_{\mathrm{dR},\mathbb{Q}}|_{\mathrm{Sh}_{K,\mathbb{C}}^{\mathrm{an}}}))$ is a variation of Hodge structures over $\mathrm{Sh}_{K,\mathbb{C}}^{\mathrm{an}}$.

The functor $\mathbb{V}_{\mathbb{C}}$ from § 3.3 factors through $\mathbb{V}_{\mathbb{Q}}$ in the obvious way.

4. Integral canonical models I: the self-dual case

Throughout this section, we will assume that (L, Q) is a self-dual quadratic space over $\mathbb{Z}_{(p)}$.

4.1 We return now to the Kuga–Satake map $\mathrm{Sh}_K \rightarrow \mathrm{Sh}_{\mathcal{K}}$. We will always assume that K^p and \mathcal{K}^p are chosen such that $\mathrm{Sh}_{\mathcal{K}}$ admits the above description as a fine moduli scheme over \mathbb{Q} with integral model $\mathcal{S}_{\mathcal{K}}$ over $\mathbb{Z}_{(p)}$.

DEFINITION 4.2. A pro-scheme X over $\mathbb{Z}_{(p)}$ satisfies the *extension property* if, for any regular, locally healthy $\mathbb{Z}_{(p)}$ -scheme S , any map $S \otimes \mathbb{Q} \rightarrow X$ extends to a map $S \rightarrow X$.

Let Sh_{K_p} (respectively $\text{Sh}_{K_{0,p}}$) be the pro-variety attached to the inverse system $\{\text{Sh}_{K_p K^p}\}$ (respectively $\{\text{Sh}_{K_{0,p} K_0^p}\}$); here, K^p varies over the compact open sub-groups of $G(\mathbb{A}_f^p)$.

DEFINITION 4.3. A model \mathcal{S}_{K_p} (respectively $\mathcal{S}_{K_{0,p}}$) for Sh_{K_p} (respectively $\text{Sh}_{K_{0,p}}$) over $\mathbb{Z}_{(p)}$ is an *integral canonical model* if it is regular, locally healthy and has the extension property. If \mathcal{S}_{K_p} is an integral canonical model for Sh_{K_p} , and if $K = K_p K^p$ is a compact open sub-group in $G(\mathbb{A}_f)$, then we will call $\mathcal{S}_K := \mathcal{S}_{K_p} / K^p$ the *integral canonical model* for Sh_K . A similar convention will hold for quotients $\mathcal{S}_{K_0} := \mathcal{S}_{K_{0,p}} / K_0^p$ by compact open sub-groups $K_0^p \subset G_0(\mathbb{A}_f^p)$.

For any compact open sub-group $\mathcal{K}' \subset \mathcal{K}$, with $\mathcal{K}'_p = \mathcal{K}_p$, set $K' = \mathcal{K}' \cap G(\mathbb{A}_f)$. Let $\mathcal{S}_{K'}$ be the normalization of the Zariski closure of $\text{Sh}_{K'}$ in $\mathcal{S}_{K'}$. Consider the pro-scheme

$$\mathcal{S}_{K_p} = \varprojlim_{K' \subset K} \mathcal{S}_{K'}$$

Here, K' varies over compact open sub-groups of K with $K'_p = K_p$.

We have the following result, due (independently) to Kisin [Kis10, 2.3.8 and 3.4.14] and Vasiu [Vas99].

THEOREM 4.4. *The scheme \mathcal{S}_{K_p} is a smooth integral canonical model for Sh_{K_p} over $\mathbb{Z}_{(p)}$. Moreover, the finite Galois cover $\text{Sh}_{K_p} \rightarrow \text{Sh}_{K_{0,p}}$ extends to a pro-finite Galois cover $\mathcal{S}_{K_p} \rightarrow \mathcal{S}_{K_{0,p}}$, where $\mathcal{S}_{K_{0,p}}$ is a smooth integral canonical model for $\text{Sh}_{K_{0,p}}$ over $\mathbb{Z}_{(p)}$. \square*

Though we will not present the proof in its entirety, we will need some of its ingredients, which we describe now. They are extracted mainly from [Kis10].

4.5 By construction, the polarized abelian scheme $(A_{\text{Sh}_K}^{\text{KS}}, \lambda_{\text{Sh}_K}^{\text{KS}})$ extends to a polarized abelian scheme $(A^{\text{KS}}, \lambda^{\text{KS}})$ over \mathcal{S}_K . By the theory of Nerón models (cf. [FC90, I.2.7]), A^{KS} has a unique $\mathbb{Z}/2\mathbb{Z}$ -grading and a C -action extending those on $A_{\text{Sh}_K}^{\text{KS}}$.

For any \mathbb{F}_p -scheme S , let $(S/\mathbb{Z}_p)_{\text{cris}}$ be the big crystalline site for S over $\text{Spec } \mathbb{Z}_p$ (cf. [BM90, p. 178]) and let $\mathcal{O}_S^{\text{cris}}$ be the structure sheaf of $(S/\mathbb{Z}_p)_{\text{cris}}$. Recall that an object in $(S/\mathbb{Z}_p)_{\text{cris}}$ is a triple (U, T, γ) , where U is an S -scheme, $U \hookrightarrow T$ is a nilpotent thickening of \mathbb{Z}_p -schemes with ideal of definition $\mathcal{J}_{(U \hookrightarrow T)}$ and γ is a divided power structure on $\mathcal{J}_{(U \hookrightarrow T)}$ that is compatible with the natural divided power structure on the ideal $p\mathcal{O}_T$. For any sheaf G over $(S/\mathbb{Z}_p)_{\text{cris}}$, and any object (U, T, γ) in $(S/\mathbb{Z}_p)_{\text{cris}}$, we denote by G_T the restriction of G to the fppf site over T .

Let $A_{\mathbb{F}_p}^{\text{KS}}$ be the fiber of A^{KS} over $\mathcal{S}_{K, \mathbb{F}_p}$. Let \mathbf{H}_{cris} be the first crystalline cohomology of $A_{\mathbb{F}_p}^{\text{KS}}$ over $\mathcal{S}_{K, \mathbb{F}_p}$. This is a crystal of locally free $\mathcal{O}_{\mathcal{S}_{K, \mathbb{F}_p}}^{\text{cris}}$ -modules over $(\mathcal{S}_{K, \mathbb{F}_p} / \mathbb{Z}_p)_{\text{cris}}$. Let $\hat{\mathcal{S}}_{K, \mathbb{Z}_p}$ be the completion of \mathcal{S}_K along its special fiber. We then have a natural identification

$$\mathbf{H}_{\text{dR}}|_{\hat{\mathcal{S}}_{K, \mathbb{Z}_p}} = \varprojlim_n \mathbf{H}_{\text{cris}, \mathcal{S}_{K, \mathbb{Z}/p^n}}$$

of coherent sheaves over $\hat{\mathcal{S}}_{K, \mathbb{Z}_p}$. In other words, for any \mathcal{S}_K -scheme T in which p is nilpotent, we have a canonical identification of coherent sheaves

$$\mathbf{H}_{\text{dR}}|_T = \mathbf{H}_{\text{cris}, T}$$

The crystal \mathbf{H}_{cris} has more structure: it is an F -crystal. More precisely, let Fr be the absolute Frobenius endomorphism on $\mathcal{S}_{K, \mathbb{F}_p}$. Then $\text{Fr}^* \mathbf{H}_{\text{cris}}$ is identified with the relative crystalline

cohomology of the Frobenius pull-back $\text{Fr}^* A_{\mathbb{F}_p}^{\text{KS}}$, and the relative Frobenius map $A_{\mathbb{F}_p}^{\text{KS}} \rightarrow \text{Fr}^* A_{\mathbb{F}_p}^{\text{KS}}$ induces a map of crystals

$$F : \text{Fr}^* \mathbf{H}_{\text{cris}} \rightarrow \mathbf{H}_{\text{cris}}.$$

If T is any $\mathcal{S}_{K, \mathbb{F}_p}$ -scheme, we get an induced map of coherent sheaves

$$F : \text{Fr}^* \mathbf{H}_{\text{dR}}|_T \rightarrow \mathbf{H}_{\text{dR}}|_T.$$

The kernel of this map is precisely the Hodge filtration $\text{Fr}^* F^1 \mathbf{H}_{\text{dR}}|_T \subset \text{Fr}^* \mathbf{H}_{\text{dR}}|_T$.

4.6 Suppose that we have a point $s \in \mathcal{S}_K(k)$, where k is a perfect field of characteristic p . Set $W = W(k)$; then the restriction \mathbf{H}_{cris} (respectively $\mathbf{H}_{\text{cris}}^{\otimes(1,1)}$) to $(\text{Spec } k/\mathbb{Z}_p)_{\text{cris}}$ corresponds to the W -module $H_{\text{cris}}^1(A_s^{\text{KS}}/W)$ (respectively $\text{End}(H_{\text{cris}}^1(A_s^{\text{KS}}/W))$). If σ is the canonical Frobenius lift on W , F induces a map

$$F_s : \sigma^* \mathbf{H}_{\text{cris},s} \rightarrow \mathbf{H}_{\text{cris},s},$$

giving $\mathbf{H}_{\text{cris},s}$ the structure of an F -crystal over W . Conjugation by F_s induces an F -isocrystal structure on $\mathbf{H}_{\text{cris},s}[p^{-1}]^{\otimes(1,1)}$.

PROPOSITION 4.7. Assume that W admits an embedding into \mathbb{C} . Then there exists a canonical F -invariant tensor $\pi_{\text{cris},s} \in \mathbf{H}_{\text{cris},s}^{\otimes(2,2)}$ such that, given any finite extension $E/W_{\mathbb{Q}}$, an algebraic closure \bar{E}/E of E and any lift $\tilde{s} \in \mathcal{S}_K(\mathcal{O}_E)$ of s , the following properties hold.

(i) Let $\tilde{s}_E \in \text{Sh}_K(E)$ be the attached E -valued point. Under the Berthelot–Ogus comparison isomorphism

$$\mathbf{H}_{\text{dR},\tilde{s}_E} \xrightarrow{\cong} \mathbf{H}_{\text{cris},s} \otimes_W E,$$

$\pi_{\text{dR},\mathbb{Q},\tilde{s}_E} \otimes 1$ is carried to $\pi_{\text{cris},s} \in \mathbf{H}_{\text{cris},s}^{\otimes(2,2)}$.

(ii) Let $\tilde{s}_{\bar{E}} \in \text{Sh}_K(\bar{E})$ be the attached \bar{E} -valued point. The crystalline comparison isomorphism

$$\mathbf{H}_{p,\tilde{s}_{\bar{E}}} \otimes_{\mathbb{Z}_p} B_{\text{cris}} \xrightarrow{\cong} \mathbf{H}_{\text{cris},s} \otimes_W B_{\text{cris}}$$

respects grading, is C -equivariant and carries $\pi_{p,\tilde{s}_{\bar{E}}} \otimes 1$ to $\pi_{\text{cris},s} \otimes 1$.

(iii) There exists a C -equivariant isomorphism of $\mathbb{Z}/2\mathbb{Z}$ -graded W -modules

$$H \otimes_{\mathbb{Z}(p)} W \xrightarrow{\cong} \mathbf{H}_{\text{cris},s}$$

carrying π to $\pi_{\text{cris},s}$.

(iv) $\pi_{\text{cris},s}$ is an idempotent projector on $\mathbf{H}_{\text{cris},s}^{\otimes(1,1)}$. Set

$$\mathbf{L}_{\text{cris},s} = \text{im } \pi_{\text{cris},s} \subset \mathbf{H}_{\text{cris},s}^{\otimes(1,1)}.$$

Then $\mathbf{L}_{\text{cris},s}$ is a self-dual quadratic space over W that is isometric to $L \otimes W$.

(v) Set $\mathbf{L}_{\text{dR},s} = \mathbf{L}_{\text{cris},s} \otimes k$; then the Hodge filtration $F^1 \mathbf{H}_{\text{dR},s} \subset \mathbf{H}_{\text{dR},s}$ is $\text{GSpin}(\mathbf{L}_{\text{dR},s})$ -split. More precisely, there exists a canonical isotropic line $F^1 \mathbf{L}_{\text{dR},s} \subset \mathbf{L}_{\text{dR},s}$ such that

$$F^1 \mathbf{H}_{\text{dR},s} = \ker(F^1 \mathbf{L}_{\text{dR},s}) = \text{im}(F^1 \mathbf{L}_{\text{dR},s}).$$

Proof. Fix E/\mathbb{Q}_p and \tilde{s} as in the proposition. Then π_{p,\tilde{s}_E} is a Galois-invariant tensor in $\mathbf{H}_{p,\tilde{s}_E}^{\otimes(2,2)}$. Therefore, under the crystalline comparison isomorphism, it is carried to an F -invariant tensor $\pi_{\text{cris},s} \in \mathbf{H}_{\text{cris},s}^{\otimes(2,2)}[p^{-1}]$. In fact, $\pi_{\text{cris},s}$ belongs to $\mathbf{H}_{\text{cris},s}^{\otimes(2,2)}$ [Kis10, 1.3.6(1) and 1.4.3(1)]. We have to show that it has the desired properties.

The comparison isomorphisms are functorial and are therefore C -equivariant and respect gradings. The de Rham comparison isomorphism

$$\mathbf{H}_{p,\tilde{s}_E} \otimes_{\mathbb{Z}_p} B_{\text{dR}} \xrightarrow{\sim} \mathbf{H}_{\text{dR},\tilde{s}_E} \otimes_E B_{\text{dR}}$$

respects $\mathbb{Z}/2\mathbb{Z}$ -grading, is C -equivariant and carries $\pi_{p,\tilde{s}_E} \otimes 1$ to $\pi_{\text{dR},\mathbb{Q},\tilde{s}_E} \otimes 1$. This is a result of Blasius and Wintenberger [Bla94], when \tilde{s}_E arises from a point valued in a number field. For the generality we need, cf. [Moo98, 5.6.3].

The crystalline and de Rham comparison isomorphisms are compatible with the Berthelot–Ogus isomorphism. This implies that the isomorphism in (i) carries $\pi_{\text{dR},\mathbb{Q},\tilde{s}_E}$ to $\pi_{\text{cris},s}$. In particular, (i) holds for all lifts \tilde{s} if and only if (ii) does. That (i) is true for all lifts follows from a parallel transport argument; cf. the proof of [Kis10, 2.3.5].

Now, [Kis10, 1.4.3(3)] shows that there exists a C -equivariant $\mathbb{Z}/2\mathbb{Z}$ -graded isomorphism

$$\mathbf{H}_{p,\tilde{s}_E} \otimes_{\mathbb{Z}_p} W \xrightarrow{\sim} \mathbf{H}_{\text{cris},s}$$

carrying π_{p,\tilde{s}_E} to $\pi_{\text{cris},s}$. The main input is (1.4), which shows that $\text{GSpin}(L)$ is the point-wise stabilizer in $\text{GL}_C^+(H)$ of π .

So, to show (iii), we only have to observe that there exists a C -equivariant graded isomorphism

$$H \otimes_{\mathbb{Z}(p)} \mathbb{Z}_p \xrightarrow{\sim} \mathbf{H}_{p,\tilde{s}_E}$$

carrying π to π_{p,\tilde{s}_E} . Indeed, we can fix an embedding $\sigma : \bar{E} \hookrightarrow \mathbb{C}$ and identify $\mathbf{H}_{p,\tilde{s}_E}$ with the Betti cohomology $\mathbf{H}_{B,\sigma(\tilde{s}_E)} \otimes \mathbb{Z}_p$. Now the assertion is clear from the construction of the sheaf \mathbf{H}_B in § 3.3.

Part (iv) is immediate from (iii), and the first assertion of (v) follows from [Kis10, 1.4.3(4)]. The second assertion of (v) follows from the discussion in § 1.9. Note that we are viewing $\mathbf{L}_{\text{dR},s}$ as a sub-space of $\mathbf{H}_{\text{dR},s}^{\otimes(1,1)}$ via the canonical identification $\mathbf{H}_{\text{cris},s} \otimes k = \mathbf{H}_{\text{dR},s}$, and we are identifying $\text{GSpin}(\mathbf{L}_{\text{dR},s})$ with the sub-group of C -equivariant, graded isomorphisms of $\mathbf{H}_{\text{dR},s}$, which preserve $\mathbf{L}_{\text{dR},s}$ under conjugation. \square

4.8 Suppose that we are given a lift $\tilde{s} \in \mathcal{S}_K(W)$. This equips $\mathbf{H}_{\text{cris},s}$ with a filtration $F^1 \mathbf{H}_{\text{cris},s}$: it is the pull-back of the Hodge filtration $F^1 \mathbf{H}_{\text{dR},\tilde{s}} \subset \mathbf{H}_{\text{dR},\tilde{s}}$. This filtration is *strongly divisible*. That is, we have

$$F_s(\sigma^*(p^{-1}F^1 \mathbf{H}_{\text{cris},s} + \mathbf{H}_{\text{cris},s})) = \mathbf{H}_{\text{cris},s}.$$

This is equivalent to the assertion that $\text{Fr}^* F^1 \mathbf{H}_{\text{dR},s}$ is the kernel of the mod- p Frobenius $\text{Fr}^* \mathbf{H}_{\text{dR},s} \rightarrow \mathbf{H}_{\text{dR},s}$.

If we now endow $\mathbf{H}_{\text{cris},s}^{\otimes(1,1)}$ with its induced filtration and $\mathbf{H}_{\text{cris},s}^{\otimes(1,1)}[p^{-1}]$ with the conjugation action of F_s , then we again obtain a strongly divisible module, in the sense that the following identity holds (cf. [Laf80, 4.2]):

$$F_s(\sigma^*(p^{-1}F^1 \mathbf{H}_{\text{cris},s}^{\otimes(1,1)} + F^0 \mathbf{H}_{\text{cris},s}^{\otimes(1,1)} + p\mathbf{H}_{\text{cris},s}^{\otimes(1,1)})) = \mathbf{H}_{\text{cris},s}^{\otimes(1,1)}. \tag{4.8.1}$$

Applying the F -equivariant, filtration preserving projector $\pi_{\text{cris},s}$ to (4.8.1) now gives us

$$F_s(\sigma^*(p^{-1}F^1 \mathbf{L}_{\text{cris},s} + F^0 \mathbf{L}_{\text{cris},s} + p\mathbf{L}_{\text{cris},s})) = \mathbf{L}_{\text{cris},s}. \tag{4.8.2}$$

4.9 Choose any co-character $\mu_0 : \mathbb{G}_m \otimes k \rightarrow \mathrm{GSpin}(\mathbf{L}_{\mathrm{dR},s})$ splitting the Hodge filtration and let $\mu : \mathbb{G}_m \otimes W \rightarrow \mathrm{GSpin}(\mathbf{L}_{\mathrm{cris},s})$ be any lift of μ_0 . It determines a lift $F^1 \mathbf{H}_{\mathrm{cris},s} \subset \mathbf{H}_{\mathrm{cris},s}$ of the Hodge filtration as well as a splitting $\mathbf{H}_{\mathrm{cris},s} = F^1 \mathbf{H}_{\mathrm{cris},s} \oplus \overline{F}^1 \mathbf{H}_{\mathrm{cris},s}$. Since μ factors through $\mathrm{GSpin}(\mathbf{L}_{\mathrm{cris},s})$, it induces a splitting

$$\mathbf{L}_{\mathrm{cris},s} = F^1 \mathbf{L}_{\mathrm{cris},s} \oplus \mathbf{L}_{\mathrm{cris},s}^0 \oplus \overline{F}^1 \mathbf{L}_{\mathrm{cris},s},$$

where $F^1 \mathbf{L}_{\mathrm{cris},s} \subset \mathbf{L}_{\mathrm{cris},s}$ is an isotropic line lifting $F^1 \mathbf{L}_{\mathrm{dR},s}$. We again have

$$F^1 \mathbf{H}_{\mathrm{cris},s} = \ker(F^1 \mathbf{L}_{\mathrm{cris},s}) = \mathrm{im}(F^1 \mathbf{L}_{\mathrm{cris},s}).$$

As usual, we are viewing $\mathbf{L}_{\mathrm{cris},s}$ as a space of endomorphisms of $\mathbf{H}_{\mathrm{cris},s}$.

Let $U \subset \mathrm{GL}(\mathbf{H}_{\mathrm{cris},s})$ be the opposite unipotent attached to this splitting: namely, it is the unipotent radical of the parabolic sub-group attached to the filtration $\overline{F}^1 \mathbf{H}_{\mathrm{cris},s}$. Its intersection U_G with $\mathrm{GSpin}(\mathbf{L}_{\mathrm{cris},s})$ is again the unipotent radical of a parabolic sub-group of $\mathrm{GSpin}(\mathbf{L}_{\mathrm{cris},s})$.

Let \widehat{U} (respectively \widehat{U}_G) be the completion of U (respectively U_G) along the identity section. Let R (respectively R_G) be the ring of formal functions on \widehat{U} (respectively \widehat{U}_G); then we have a surjection $R \twoheadrightarrow R_G$ of formally smooth W -algebras.

Set $\mathbf{H}_R = \mathbf{H}_{\mathrm{cris},s} \otimes_W R$, and equip it with the constant filtration arising from the filtration $F^1 \mathbf{H}_{\mathrm{cris},s}$. Choose compatible isomorphisms

$$R \xrightarrow{\cong} W[[t_1, \dots, t_r]], \quad R_G \xrightarrow{\cong} W[[t_1, \dots, t_d]]$$

such that the identity sections are identified with the maps $t_i \mapsto 0$. Equip R with the Frobenius lift $\varphi : t_i \mapsto t_i^p$. Equip \mathbf{H}_R with the map

$$\mathbf{F}_R : \varphi^* \mathbf{H}_R = \sigma^* \mathbf{H}_{\mathrm{cris},s} \otimes_W R \xrightarrow{\mathbf{F}_s \otimes 1} \mathbf{H}_{\mathrm{cris},s} \otimes_W R = \mathbf{H}_R \xrightarrow{g} \mathbf{H}_R,$$

where $g \in \widehat{U}(R)$ is the tautological element.

By [Moo98, Theorem 4.4], there is a unique topologically quasi-nilpotent integrable connection $\nabla_R : \mathbf{H}_R \rightarrow \mathbf{H}_R \otimes_R \widehat{\Omega}_{R/W}^1$ for which \mathbf{F}_R is parallel. The tuple $(\mathbf{H}_R, \mathbf{F}_R, \nabla_R)$ determines, and is determined by, a unique F -crystal over $(\mathrm{Spec}(R \otimes \mathbb{F}_p)/\mathbb{Z}_p)_{\mathrm{cris}}$, which we will denote by \mathbf{H} . The evaluation of \mathbf{H} on the pro-nilpotent divided power thickening $\mathrm{Spec}(R \otimes \mathbb{F}_p) \hookrightarrow \mathrm{Spf} R$ is identified with \mathbf{H}_R . Similarly, a change of scalars along $R \rightarrow R_G$ gives us a tuple $(\mathbf{H}_{R_G}, \mathbf{F}_{R_G}, \nabla_{R_G}, F^1 \mathbf{H}_{R_G})$ over R_G . This corresponds to a unique F -crystal \mathbf{H}_G over $(\mathrm{Spec}(R_G \otimes \mathbb{F}_p)/\mathbb{Z}_p)_{\mathrm{cris}}$.

Observe that $\mathbf{H}_{R_G} = \mathbf{H}_{\mathrm{cris},s} \otimes_W R_G$ is equipped with the constant tensor $\pi_{R_G} = \pi_{\mathrm{cris},s} \otimes 1 \in \mathbf{H}_{R_G}^{\otimes(2,2)}$. We can view π_{R_G} as an idempotent operator on $\mathbf{H}_{R_G}^{\otimes(1,1)}$. Write \mathbf{L}_{R_G} for its image: this is a direct summand of $\mathbf{H}_{R_G}^{\otimes(1,1)}$, which can be identified with $\mathbf{L}_{\mathrm{cris},s} \otimes R_G$.

PROPOSITION 4.10.

- (i) The tensor $\pi_{R_G} \in \mathbf{H}_{R_G}^{\otimes(2,2)}$ is parallel, lies in $F^0 \mathbf{H}_{R_G}^{\otimes(2,2)}$ and is F -invariant in $\mathbf{H}_{R_G}^{\otimes(2,2)}[p^{-1}]$.
- (ii) The direct summand $\mathbf{L}_{R_G} \subset \mathbf{H}_{R_G}^{\otimes(1,1)}$ is stable under the connection ∇_{R_G} and $\mathbf{L}_{R_G}[p^{-1}]$ is stable under the conjugation action of \mathbf{F}_{R_G} . Furthermore, \mathbf{L}_{R_G} with its induced filtration $F^\bullet \mathbf{L}_{R_G}$ is strongly divisible:

$$\mathbf{F}_{R_G}(\varphi^*(p^{-1}F^1 \mathbf{L}_{R_G} + F^0 \mathbf{L}_{R_G} + p \mathbf{L}_{R_G})) = \mathbf{L}_{R_G}.$$

Proof. Part (i) is essentially by construction; cf. [Moo98, 4.8]. Part (ii) also follows by construction: we can identify $(\mathbf{L}_{R_G}, F^\bullet \mathbf{L}_{R_G})$ with $(\mathbf{L}_{\text{cris},s}, F^\bullet \mathbf{L}_{\text{cris},s}) \otimes_W R_G$, so that \mathbf{F}_{R_G} is identified with $g \circ (\mathbf{F}_s \otimes 1)$. Here, $g \in U_G(R_G)$ is the tautological element. The result now is a consequence of (4.8.2). \square

4.11 Now assume that $k = \overline{\mathbb{F}}_p$, so that s is a closed point. Set $R_s = \widehat{\mathcal{O}}_{\mathcal{S}_K,s}$ and let \widehat{U}_s be the completion of \mathcal{S}_K at s . Let $\widehat{U}' = \text{Spf } R'$ be the universal deformation space for the abelian variety A_s^{KS} ; then \widehat{U}_s is naturally identified with a closed formal sub-scheme of \widehat{U}' . Restricting \mathbf{H}_{cris} to $\text{Spec}(R' \otimes \mathbb{F}_p)$ gives rise to an F -crystal \mathbf{H}' over $(\text{Spec}(R' \otimes \mathbb{F}_p)/\mathbb{Z}_p)_{\text{cris}}$. Evaluating \mathbf{H}' along the pro-nilpotent divided power thickening $\text{Spec}(R' \otimes \mathbb{F}_p) \hookrightarrow \text{Spf}(R')$ gives us a finite free R' -module $\mathbf{H}_{R'}$, which can be identified with the evaluation of \mathbf{H}_{dR} on $\text{Spec } R'$. The Hodge filtration on \mathbf{H}_{dR} equips $\mathbf{H}_{R'}$ with a direct summand $F^1 \mathbf{H}_{R'} \subset \mathbf{H}_{R'}$.

The W -algebra R admits the identity section $j : R \rightarrow W$ with $t_i \mapsto 0$ for all i . Also, by Grothendieck–Messing theory [Mes72, V.1.6], the lift $F^1 \mathbf{H}_{\text{cris},s} \subset \mathbf{H}_{\text{cris},s}$ corresponds to a lift $\tilde{s} : \text{Spf } W \rightarrow \text{Spf}(R')$ of s . In fact, we have a canonical isomorphism of W -modules $\mathbf{H}_{\text{dR},\tilde{s}} \xrightarrow{\cong} \mathbf{H}_{\text{cris},s}$ that identifies $F^1 \mathbf{H}_{\text{dR},\tilde{s}}$ with $F^1 \mathbf{H}_{\text{cris},s}$. Write $j' : R' \rightarrow W$ for the section corresponding to the lift \tilde{s} .

By construction, we now have canonical isomorphisms $\iota_0 : \mathbf{H}_R \otimes_{R,j} W \xrightarrow{\cong} \mathbf{H}_{\text{cris},s}$ and $\iota'_0 : \mathbf{H}_{R'} \otimes_{R',j'} W \xrightarrow{\cong} \mathbf{H}_{\text{cris},s}$.

THEOREM 4.12.

- (i) *There exists an isomorphism (necessarily unique) of augmented W -algebras $f : (R, j) \xrightarrow{\cong} (R', j')$ with the following property: let $f^* : (\text{Spec}(R \otimes \mathbb{F}_p)/\mathbb{Z}_p)_{\text{cris}} \rightarrow (\text{Spec}(R' \otimes \mathbb{F}_p)/\mathbb{Z}_p)_{\text{cris}}$ be the induced map on crystalline sites. Then there exists an isomorphism (necessarily unique) of F -crystals $\alpha : f^* \mathbf{H} \xrightarrow{\cong} \mathbf{H}'$ such that:

 - (a) *the induced isomorphism $\alpha_R : \mathbf{H}_R \otimes_{R,f} R' \xrightarrow{\cong} \mathbf{H}_{R'}$ carries $F^1 \mathbf{H}_R \otimes_{R,f} R'$ onto $F^1 \mathbf{H}_{R'}$;*
 - (b) *the induced isomorphism $j^* \alpha_R : \mathbf{H}_R \otimes_{R,j} W \xrightarrow{\cong} \mathbf{H}_{R'} \otimes_{R',j'} W$ is equal to $(\iota'_0)^{-1} \circ \iota_0$.**
- (ii) *The induced isomorphism $\text{Spf}(f) : \widehat{U}' \xrightarrow{\cong} \widehat{U}$ carries \widehat{U}_s onto \widehat{U}_G .*
- (iii) *The restriction of the tensor $\pi_{\text{dR},\mathbb{Q}}$ to $\text{Spec}(R_s \otimes \mathbb{Q})$ extends to a parallel section*

$$\pi_{\text{dR},R_s} \in H^0(\text{Spec } R_s, F^0 \mathbf{H}_{\text{dR}}^{\otimes(2,2)})^{\nabla=0}.$$

π_{dR,R_s} is an idempotent operator on $\mathbf{H}_{\text{dR}}^{\otimes(1,1)}|_{\text{Spec } R_s}$.

Proof. Part (i) is due to Faltings; cf. [Fal99, § 7]. Part (ii) is shown during the course of the proof of [Kis10, 2.3.5].

Part (iii) is a special case of [Kis10, 2.3.9]. We sketch the proof: from f and α_R , we obtain isomorphisms $f_G : R_G \xrightarrow{\cong} R_s$ and $\alpha_{R_G} : \mathbf{H}_{R_G} \otimes_{R_G,f_G} R_s \xrightarrow{\cong} \mathbf{H}_{\text{dR},R_s}$. From (4.10), we then obtain a parallel tensor:

$$\alpha_{R_G}(\pi_{R_G}) \in H^0(\text{Spec } R_s, F^0 \mathbf{H}_{\text{dR}}^{\otimes(2,2)})^{\nabla=0}.$$

By construction, the evaluation of this tensor along the map $R_s \xrightarrow{\tilde{s}} W \hookrightarrow W_{\mathbb{Q}}$ agrees with that of $\pi_{\text{dR},\mathbb{Q}}$. Therefore, $\alpha_{R_G}(\pi_{R_G})|_{\text{Spec}(R_s \otimes \mathbb{Q})}$ and $\pi_{\text{dR},\mathbb{Q}}|_{\text{Spec}(R_s \otimes \mathbb{Q})}$ are both parallel tensors in $\mathbf{H}_{\text{dR}}^{\otimes(2,2)}|_{\text{Spec}(R_s \otimes \mathbb{Q})}$ that agree at a point. They must therefore agree everywhere. Thus, $\pi_{\text{dR},R_s} := \alpha_{R_G}(\pi_{R_G})$ is the extension we seek in (iii). \square

COROLLARY 4.13.

- (i) $\pi_{\mathrm{dR},\mathbb{Q}}$ extends to a parallel section π_{dR} of $F^0\mathbf{H}_{\mathrm{dR}}^{\otimes(2,2)}$ over \mathcal{S}_K . Moreover, π_{dR} is an idempotent projector on $\mathbf{H}_{\mathrm{dR}}^{\otimes(1,1)}$, whose image is a vector sub-bundle \mathbf{L}_{dR} extending $\mathbf{L}_{\mathrm{dR},\mathbb{Q}}$.
- (ii) Consider the functor $\mathcal{P}_{\mathrm{dR}}$ on \mathcal{S}_K -schemes given by

$$\mathcal{P}_{\mathrm{dR}}(T) = \left(\begin{array}{l} C\text{-equivariant } \mathbb{Z}/2\mathbb{Z}\text{-graded } \mathcal{O}_T\text{-module isomorphisms} \\ \xi : H \otimes_{\mathbb{Z}(p)} \mathcal{O}_T \xrightarrow{\cong} \mathbf{H}_{\mathrm{dR},T} \\ \text{carrying } L \otimes \mathcal{O}_T \subset H^{\otimes(1,1)} \otimes \mathcal{O}_T \text{ onto } \mathbf{L}_{\mathrm{dR}} \subset \mathbf{H}_{\mathrm{dR}}^{\otimes(1,1)} \end{array} \right).$$

Then $\mathcal{P}_{\mathrm{dR}}$ is a G -torsor over \mathcal{S}_K with generic fiber $\mathcal{P}_{\mathrm{dR},\mathbb{Q}}$.

Proof. We only need to prove these properties over the formal completion \widehat{U}_s at a point $s \in \mathcal{S}_K(\overline{\mathbb{F}}_p)$. Part (i) is immediate from (4.12)(iii).

To prove (ii), it is enough to show that, if we view $\mathrm{Spec} R_G$ as an \mathcal{S}_K -scheme via the maps $\mathrm{Spec} R_G \xrightarrow{\cong} \mathrm{Spec} R_s \rightarrow \mathcal{S}_K$ of (4.12), the set $\mathcal{P}_{\mathrm{dR}}(\mathrm{Spec} R_G)$ is non-empty.

In other words, we have to show that there exists a C -equivariant, graded isomorphism of R_G -modules $H \otimes_{\mathbb{Z}(p)} R_G \xrightarrow{\cong} \mathbf{H}_{R_G}$ carrying $L \otimes R_G$ onto $\mathbf{L}_{R_G} := \mathrm{im}(\pi_{R_G})$. This is immediate from (4.7)(iii) and the very construction of \mathbf{H}_{R_G} . \square

4.14 Since $\mathcal{S}_{K,\mathbb{Z}_p}$ is smooth over \mathbb{Z}_p , as a crystal, $\mathbf{H}_{\mathrm{cris}}$ is determined by $\mathbf{H}_{\mathrm{dR}}|_{\widehat{\mathcal{S}}_{K,\mathbb{Z}_p}}$ equipped with its Gauss–Manin connection. In particular, evaluation along the formal PD thickening $\mathcal{S}_{K,\mathbb{F}_p} \hookrightarrow \widehat{\mathcal{S}}_{K,\mathbb{Z}_p}$ gives us a canonical isomorphism

$$H^0((\mathcal{S}_{K,\mathbb{F}_p}/\mathbb{Z}_p)_{\mathrm{cris}}, \mathbf{H}_{\mathrm{cris}}^{\otimes}) \xrightarrow{\cong} H^0(\widehat{\mathcal{S}}_{K,\mathbb{Z}_p}, \mathbf{H}_{\mathrm{dR}}^{\otimes})^{\nabla=0}.$$

Therefore, there exists a unique global section

$$\pi_{\mathrm{cris}} \in H^0((\mathcal{S}_{K,\mathbb{F}_p}/\mathbb{Z}_p)_{\mathrm{cris}}, \mathbf{H}_{\mathrm{cris}}^{\otimes(2,2)}),$$

whose evaluation at $\mathcal{S}_{K,\mathbb{F}_p} \hookrightarrow \widehat{\mathcal{S}}_{K,\mathbb{Z}_p}$ is the restriction of the parallel section $\pi_{\mathrm{dR}} \in H^0(\mathcal{S}_K, \mathbf{H}_{\mathrm{dR}}^{\otimes(2,2)})$.

Again, we can view π_{cris} as an idempotent endomorphism of the crystal $\mathbf{H}_{\mathrm{cris}}^{\otimes(1,1)}$, and we denote by $\mathbf{L}_{\mathrm{cris}}$ the image of π_{cris} in $\mathbf{H}_{\mathrm{cris}}^{\otimes(1,1)}$. If $s \rightarrow \mathcal{S}_{K,\mathbb{F}_p}$ is a point valued in an algebraically closed field $k(s)$, then the restriction of π_{cris} to $(\mathrm{Spec} k(s)/\mathbb{Z}_p)_{\mathrm{cris}}$ determines and is determined by its evaluation along the formal thickening $\mathrm{Spec} k(s) \hookrightarrow \mathrm{Spf} W(k(s))$.

Suppose that we have a lift $\tilde{s} \in \mathcal{S}_K(W)$ of s . Then there is a natural isomorphism $\mathbf{H}_{\mathrm{cris},s} \xrightarrow{\cong} \mathbf{H}_{\mathrm{dR},\tilde{s}}$. By the definition of π_{cris} , its evaluation over $\mathrm{Spec} k(s) \hookrightarrow \mathrm{Spf} W(k(s))$ must map to the tensor $\pi_{\mathrm{dR},\tilde{s}} \in \mathbf{H}_{\mathrm{dR},\tilde{s}}^{\otimes(2,2)}$ under this isomorphism. In particular, we find that this evaluation is exactly the tensor $\pi_{\mathrm{cris},s}$ defined in (4.7). Similarly, the restriction of $\mathbf{L}_{\mathrm{cris}}$ to $(\mathrm{Spec} k(s)/\mathbb{Z}_p)_{\mathrm{cris}}$ determines and is determined by the F -isocrystal $\mathbf{L}_{\mathrm{cris},s}$ seen in *loc. cit.*

4.15 The Gauss–Manin connection on $\mathbf{H}_{\mathrm{dR}}^{\otimes(2,2)}$ restricts to a connection on \mathbf{L}_{dR} . The filtration $F^\bullet \mathbf{L}_{\mathrm{dR}}$ on \mathbf{L}_{dR} satisfies Griffith’s transversality. That is, for any integer i , we have

$$\nabla(F^i \mathbf{L}_{\mathrm{dR}}) \subset F^{i-1} \mathbf{L}_{\mathrm{dR}} \otimes \Omega^1_{\mathcal{S}_K/\mathbb{Z}(p)}.$$

Indeed, this can be checked over \mathbb{C} , where it holds by construction. Thus, we obtain $\mathcal{O}_{\mathcal{S}_K}$ -linear Kodaira–Spencer maps:

$$\mathrm{gr}_F^0 \mathbf{L}_{\mathrm{dR}} \rightarrow \mathrm{gr}_F^{-1} \mathbf{L}_{\mathrm{dR}} \otimes \Omega^1_{\mathcal{S}_K/\mathbb{Z}_{(p)}}, \tag{4.15.1}$$

$$F^1 \mathbf{L}_{\mathrm{dR}} \rightarrow \mathrm{gr}_F^0 \mathbf{L}_{\mathrm{dR}} \otimes \Omega^1_{\mathcal{S}_K/\mathbb{Z}_{(p)}}. \tag{4.15.2}$$

PROPOSITION 4.16. *The map in (4.15.1) is an isomorphism and the map in (4.15.2) is injective with image a local direct summand of the right-hand side.*

Proof. Both statements can be studied over the complete local ring R_s at a point $s \in \mathcal{S}_K(\overline{\mathbb{F}}_p)$. As in the proof of (4.13), we can work instead over the ring R_G , and study the Kodaira–Spencer maps induced by ∇_{R_G} :

$$\mathrm{gr}_F^0 \mathbf{L}_{R_G} \rightarrow \mathrm{gr}_F^{-1} \mathbf{L}_{R_G} \otimes \hat{\Omega}^1_{R_G/W}, \quad F^1 \mathbf{L}_{R_G} \rightarrow \mathrm{gr}_F^0 \mathbf{L}_{R_G} \otimes \hat{\Omega}^1_{R_G/W}.$$

We can identify $\hat{\Omega}^1_{R_G/W}$ canonically with $\mathrm{Hom}(\mathrm{Lie} U_G, R_G)$, and so we can rewrite these maps equivalently as

$$\mathrm{Lie} U_G \otimes R_G \rightarrow \mathrm{Hom}(\mathrm{gr}_F^0 \mathbf{L}_{R_G}, \mathrm{gr}_F^{-1} \mathbf{L}_{R_G}), \quad \mathrm{Lie} U_G \otimes R_G \rightarrow \mathrm{Hom}(F^1 \mathbf{L}_{R_G}, \mathrm{gr}_F^0 \mathbf{L}_{R_G}).$$

The proposition now amounts to showing that both of these maps are isomorphisms. By Nakayama’s lemma, it suffices to do so after a change of scalars along the augmentation map $j : R_G \rightarrow W$. But then we obtain maps:

$$\mathrm{Lie} U_G \rightarrow \mathrm{Hom}(\mathrm{gr}_F^0 \mathbf{L}_{\mathrm{cris},s}, \mathrm{gr}_F^{-1} \mathbf{L}_{\mathrm{cris},s}), \quad \mathrm{Lie} U_G \rightarrow \mathrm{Hom}(F^1 \mathbf{L}_{\mathrm{cris},s}, \mathrm{gr}_F^0 \mathbf{L}_{\mathrm{cris},s}).$$

We claim that the second of these maps is (up to sign) the natural map obtained by the action of U_G on $\mathbf{L}_{\mathrm{cris},s}$, and that the first is obtained from the second via the self-duality of $\mathbf{L}_{\mathrm{cris},s}$. The proposition will follow immediately from this claim.

Consider instead the Kodaira–Spencer map on \mathbf{H}_{R_G} : $F^1 \mathbf{H}_{R_G} \rightarrow \mathrm{gr}_F^1 \mathbf{H}_{R_G} \otimes \hat{\Omega}^1_{R_G/W}$, which we can view as a map:

$$\mathrm{Lie} U_G \otimes R_G \rightarrow \mathrm{Hom}(F^1 \mathbf{H}_{R_G}, \mathrm{gr}_F^1 \mathbf{H}_{R_G}). \tag{4.16.1}$$

Tensoring (4.16.1) along j gives us

$$\mathrm{Lie} U_G \rightarrow \mathrm{Hom}(F^1 \mathbf{H}_{\mathrm{cris},s}, \mathrm{gr}_F^1 \mathbf{H}_{\mathrm{cris},s}).$$

To prove our claim, it is enough to show that, up to sign, this map is identified with the canonical inclusion $\mathrm{Lie} U_G \hookrightarrow \mathrm{Hom}(F^1 \mathbf{H}_{\mathrm{cris},s}, \mathrm{gr}_F^1 \mathbf{H}_{\mathrm{cris},s})$ induced by the action of U_G on $\mathbf{H}_{\mathrm{cris},s}$. This last assertion can be extracted from [Moo98, 4.5]; cf. also [MaS11, 1.4.2.2(3)]. \square

4.17 Over \mathcal{S}_K , we now have two tautological line bundles: first, we have the Hodge or canonical bundle ω^{KS} attached to the top exterior power of the sheaf of differentials $\Omega^1_{\mathrm{AKS}/\mathcal{S}_K}$. Second, we have the line bundle $F^1 \mathbf{L}_{\mathrm{dR}}$. These are closely related, as the next result shows.

We will need a little preparation. Fix an isotropic line $F^1 L \subset L$, and let $P \subset G$ be the parabolic sub-group stabilizing it. Let $F^1 H = \ker(F^1 L) \subset H$ be the corresponding isotropic sub-space of H . The G -torsor $\mathcal{P}_{\mathrm{dR}}$ introduced in (4.13)(ii) has a natural reduction of structure group to a P -torsor $\mathcal{P}_{\mathrm{dR},P}$. Indeed, we can take $\mathcal{P}_{\mathrm{dR},P}$ to be the sub-functor of $\mathcal{P}_{\mathrm{dR}}$ such that, for any \mathcal{S}_K -scheme T , we have

$$\mathcal{P}_{\mathrm{dR},P}(T) = \{\xi \in \mathcal{P}_{\mathrm{dR}}(T) : \xi(F^1 H \otimes \mathcal{O}_T) = F^1 \mathbf{H}_{\mathrm{dR},T}\}.$$

The proof of *loc. cit.* shows that this is indeed a P -torsor. Given such a torsor, one immediately gets a functor from $\mathbb{Z}_{(p)}$ -representations of the group scheme P to vector bundles over \mathcal{S}_K . More precisely, given a $\mathbb{Z}_{(p)}$ -representation U of P , we can view it as a trivial vector bundle over $\text{Spec } \mathbb{Z}_{(p)}$ with a P -action, and then take the corresponding \mathcal{S}_K -vector bundle to be the quotient $(\mathcal{P}_{\text{dR},P} \times_{\text{Spec } \mathbb{Z}_{(p)}} U)/P$, where P acts diagonally.

PROPOSITION 4.18. *There exists a canonical isomorphism of line bundles:*

$$\omega^{\text{KS},\otimes 2} \xrightarrow{\cong} ((F^1 \mathbf{L}_{\text{dR}})(-1))^{\otimes 2^{n+1}}.$$

Here, the (-1) denotes the twist by the (trivial) line bundle attached to the spinor character $\nu : P \rightarrow \mathbb{G}_m$. In particular, $F^1 \mathbf{L}_{\text{dR}}$ is a relatively ample line bundle for \mathcal{S}_K over $\mathbb{Z}_{(p)}$.

Proof. This follows from an argument of Maulik [Mau14, §5]. The main point is that both bundles involved are canonical extensions over the integral canonical model of automorphic line bundles.

The line bundle ω^{KS} is attached via $\mathcal{P}_{\text{dR},P}$ to the P -representation $\det(F^1 H)$, and $F^1 \mathbf{L}_{\text{dR}}(-1)$ is the line bundle attached to the representation $F^1 L(\nu)$.

The left multiplication map $L \otimes H \rightarrow H$ induces an isomorphism of P -representations

$$\text{gr}_F^{-1} L \otimes F^1 H \xrightarrow{\cong} \text{gr}_F^0 H.$$

Therefore, we have a canonical isomorphism of P -representations

$$\det(H) \xrightarrow{\cong} \det(F^1 H) \otimes \det(\text{gr}_F^0 H) \xrightarrow{\cong} \det(F^1 H)^{\otimes 2} \otimes (\text{gr}_F^{-1} L)^{\otimes 2^{n+1}}.$$

Since $(\text{gr}_F^{-1} L)^\vee \xrightarrow{\cong} F^1 L$, this gives us a canonical isomorphism of P -representations

$$\det(F^1 H)^{\otimes 2} \xrightarrow{\cong} (F^1 L)^{\otimes 2^{n+1}} \otimes \det(H).$$

On the other hand, the symplectic form ψ_δ on H induces a canonical isomorphism of P -representations

$$\text{gr}_0^F H \xrightarrow{\cong} (F^1 H)^\vee(\nu).$$

This shows that we have

$$\det(H) \xrightarrow{\cong} \mathbb{Z}_{(p)}(\nu^{2^{n+1}}),$$

completing the proof of the claimed isomorphism.

The last statement of the lemma follows, since it is known that the bundle $\tilde{\omega}^{\text{KS}}$ is relatively ample; cf. for example [Lan08, 7.2.4.1(2)]. □

4.19 Fix a perfect field k of characteristic p as usual, and set $W = W(k)$. Fix a point $s \in \mathcal{S}_K(k)$, and consider the Frobenius map $\mathbf{F} : \text{Fr}^* \mathbf{H}_{\text{dR},s} \rightarrow \mathbf{H}_{\text{dR},s}$: its image is a sub-space $\mathbf{C}_s^1 \subset \mathbf{H}_{\text{dR},s}$. We claim that there exists an isotropic line $\mathbf{N}_s^1 \subset \mathbf{L}_{\text{dR},s}$ such that, in the notation of § 1.9, $\mathbf{C}_s^1 = \ker(\mathbf{N}_s^1)$.

Indeed, fix a lift $\tilde{s} \in \mathcal{S}_K(W)$, and let μ be a co-character of $\text{GSpin}(\mathbf{L}_{\text{dR},\tilde{s}})$ that splits the Hodge filtration on $\mathbf{H}_{\text{dR},\tilde{s}}$. We obtain a decomposition into weight spaces for μ :

$$\begin{aligned} \mathbf{H}_{\text{dR},\tilde{s}} &= F^1 \mathbf{H}_{\text{dR},\tilde{s}} \oplus \mathbf{H}_{\text{dR},\tilde{s}}^0, \\ \mathbf{L}_{\text{dR},\tilde{s}} &= F^1 \mathbf{L}_{\text{dR},\tilde{s}} \oplus \mathbf{L}_{\text{dR},\tilde{s}}^0 \oplus \mathbf{L}_{\text{dR},\tilde{s}}^{-1}. \end{aligned}$$

By strong divisibility §4.8, C_s^1 is simply the image in $L_{dR,s}$ of $F_s(\sigma^*H_{dR,s}^0)$. Now, $L_{dR,\bar{s}}^{-1}$ is an isotropic line satisfying

$$\ker(L_{dR,\bar{s}}^{-1}) = H_{dR,\bar{s}}^0.$$

Moreover, by (4.8.2), $F_s(p\sigma^*L_{dR,\bar{s}}^{-1})$ is an isotropic line in $L_{dR,\bar{s}}$. One can now easily check that the image of this line in $L_{dR,s}$ is our desired N_s^1 .

LEMMA 4.20. *With the notation as above, let $\Lambda \subset L_{dR,\bar{s}}$ be a direct summand satisfying $F_s(\Lambda) = \Lambda$. Let $\bar{\Lambda}$ be the image of Λ in $L_{dR,s}$. Then N_s^1 is contained in $\bar{\Lambda}$ only if $F^1L_{dR,s}$ is also contained in $\bar{\Lambda}$.*

Proof. Choose a generator w for $L_{dR,\bar{s}}^{-1}$; then, by the construction above, the image of $F_s(pw)$ in $L_{dR,s}$ is a generator for N_s^1 . Therefore, if N_s^1 is contained in $\bar{\Lambda}$, then we can find $v \in \Lambda$ such that $F_s(pw) - v \in pL_{dR,\bar{s}}$.

But, by our hypothesis, $v = F_s(v')$ for $v' \in \Lambda$. Then, by (4.8.2), we must have

$$pw - v' \in F^1L_{dR,\bar{s}} + pL_{dR,\bar{s}}.$$

This implies that $F^1L_{dR,s}$ is generated by the image of $v' \in \Lambda$. □

5. Special endomorphisms

We will now drop the self-duality assumption on (L, Q) until further notice.

5.1 Let T be an $\text{Sh}_{K,\mathbb{C}}$ -scheme; then functoriality of cohomology gives us a natural map

$$\text{End}(A_T^{\text{KS}})_{(p)} \rightarrow H^0(T^{\text{an}}, \mathbf{H}_B^{\otimes(1,1)}).$$

DEFINITION 5.2. An endomorphism $f \in \text{End}(A_T^{\text{KS}})_{(p)}$ is *special* if it gives rise to a section of $L_B \subset \mathbf{H}_B^{\otimes(1,1)}$ under the above map. It follows from the definition that f is special if and only if its fiber f_s at every point $s \rightarrow T$ is special. In fact, it is enough to require this for one point s in each connected component of T^{an} .

Let T be any Sh_K -scheme; then, for any prime ℓ , we have a natural map

$$\text{End}(A_T^{\text{KS}})_{(p)} \rightarrow H^0(T, \mathbf{H}_\ell^{\otimes(1,1)}).$$

DEFINITION 5.3. Fix a prime $\ell \neq p$. An endomorphism $f \in \text{End}(A_T^{\text{KS}})_{(p)}$ is ℓ -special if it gives rise to a section of $L_\ell \subset \mathbf{H}_\ell^{\otimes(1,1)}$ under the above map. We say that f is p -special if it gives rise to a section of $L_p \subset \mathbf{H}_p^{\otimes(1,1)}$ under the corresponding p -adic realization map.

For any prime ℓ , we denote the space of ℓ -special endomorphisms by $L_\ell(A_T^{\text{KS}})$.

One immediately sees that f is ℓ -special if and only if, in every connected component of T , there exists a point s such that the fiber f_s at s is ℓ -special. In particular, ℓ -specialness is a condition that can be checked at geometric points.

LEMMA 5.4. *Suppose that T is an Sh_K -scheme and that $f \in \text{End}(A_T^{\text{KS}})_{(p)}$. Then the following are equivalent:*

- (i) f is ℓ -special for all primes ℓ ;
- (ii) f is ℓ -special for some prime ℓ ;
- (iii) the restriction of f over $T \otimes_{\mathbb{Q}} \mathbb{C}$ is special.

Proof. Let $s \rightarrow T$ be any \mathbb{C} -valued point. Then it is clear from the definitions that f_s is special if and only if it is ℓ -special for some (and hence any) prime ℓ . Using this, the lemma easily follows. \square

DEFINITION 5.5. Let T and f be as above. Then we say that f is *special* if it satisfies any of the equivalent conditions of (5.4). We denote the space of special endomorphisms of A_T^{KS} by $L(A_T^{\text{KS}})$.

Over \mathbb{C} , $L(A_s^{\text{KS}})$ can be described purely Hodge theoretically, as follows.

PROPOSITION 5.6. *If $s \in \text{Sh}_K(\mathbb{C})$, then*

$$L(A_s^{\text{KS}}) = \mathbf{L}_{B,s} \cap (\mathbf{L}_{B,s} \otimes \mathbb{C})^{0,0}.$$

In particular, for any Sh_K -scheme T , $\text{rk}_{\mathbb{Z}(p)} L(A_T^{\text{KS}}) \leq n$.

Proof. This is clear from the definitions. \square

5.7 We now assume that (L, Q) is self-dual, and turn to the investigation of specialness over the integral canonical model \mathcal{S}_K . For any \mathcal{S}_K -scheme T and any $\ell \neq p$, the definition of an ℓ -special endomorphism carries over directly from (5.3). We will now develop a version of p -specialness that works also over the integral model.

DEFINITION 5.8. Suppose that $s \rightarrow \mathcal{S}_{K, \mathbb{F}_p}$ is a k -valued point. Then we obtain a map

$$\text{End}(A_s^{\text{KS}})_{(p)} \rightarrow \text{End}(\mathbf{H}_{\text{cris},s}).$$

An endomorphism $f \in \text{End}(A_s^{\text{KS}})_{(p)}$ is *p -special* if it gives rise to an element of $\mathbf{L}_{\text{cris},s}$ under the above map.

LEMMA 5.9. *Let T be a \mathcal{S}_K -scheme in which p is locally nilpotent, and suppose that we have $f \in \text{End}(A_T^{\text{KS}})_{(p)}$. Then the following are equivalent.*

- (i) *For every point $s \rightarrow T$ valued in a perfect field, the fiber $f_s \in \text{End}(A_s^{\text{KS}})_{(p)}$ is p -special.*
- (ii) *In every connected component of T , there exists a point s valued in a perfect field such that the fiber f_s is p -special.*

Proof. This is an immediate consequence of the definition, the fact that the endomorphism scheme $\text{End}(A^{\text{KS}})_{(p)}$ of A^{KS} is locally noetherian and (5.10) below, applied to the crystal $\mathbf{H}_{\text{cris}}^{\otimes(1,1)} / \mathbf{L}_{\text{cris}}$. \square

LEMMA 5.10. *Suppose that T is a connected, locally noetherian \mathbb{F}_p -scheme, and that \mathbf{M} is a crystal of vector bundles over T equipped with a global section $e \in \Gamma((T/\mathbb{Z}_p)_{\text{cris}}, \mathbf{M})$. Suppose that, for some point $x \rightarrow T$, the induced global section*

$$e_x \in \Gamma((\text{Spec } k(x)/\mathbb{Z}_p)_{\text{cris}}, \mathbf{M}|_x)$$

vanishes. Then, for every point $y \rightarrow T$, the induced global section

$$e_y \in \Gamma((\text{Spec } k(y)/\mathbb{Z}_p)_{\text{cris}}, \mathbf{M}|_y)$$

also vanishes.

Proof. Since T is connected and locally noetherian, the lemma will follow if we can prove the following claim.

CLAIM. Suppose that x and y are points of T such that x is a specialization of y , and such that the prime ideal corresponding to y has height 1 in $\mathcal{O}_{T,x}$; then e_x vanishes if and only if e_y vanishes.

Let x^{perf} be the point attached to a perfect closure of $k(x)$. By [BM90, 1.3.5], e_x vanishes if and only if $e_{x^{\text{perf}}}$ vanishes. So, we can assume that $k = k(x)$ is perfect.

Let $\mathfrak{p}_y \subset \mathcal{O}_{T,x}$ be the prime ideal corresponding to y , and let R be the completion of the normalization of $\mathcal{O}_{T,x}/\mathfrak{p}_y$. By our hypotheses, R is an equicharacteristic complete DVR with residue field k , and so is isomorphic to $k[[t]]$. By pulling \mathbf{M} back to $\text{Spec } R$, we are further reduced to the situation where $T = \text{Spec } k[[t]]$, $x = \text{Spec } k$ and $y = \text{Spec } k((t))$.

Let M be the evaluation of \mathbf{M} along the formal divided power thickening $\text{Spec } k[[t]] \hookrightarrow \text{Spf } W[[t]]$: it is equipped with a flat, topologically quasi-nilpotent connection $\nabla : M \rightarrow M \otimes \widehat{\Omega}_{W(k)[[t]]}^1$. In other words, we have a derivation $D : M \rightarrow M$ over d/dt such that a sufficiently large iteration of D carries M into pM . The crystal \mathbf{M} is determined by the finite free $W(k)[[t]]$ -module M and the connection ∇ . In particular, the global sections of \mathbf{M} are identified with the module of horizontal elements $M^{\nabla=0}$.

The corresponding crystal over x is the one attached to the $W(k)$ -module $M_0 = M/tM$. The restriction from global sections over T to global sections over x is just the reduction-mod- t map $M^{\nabla=0} \rightarrow M_0$. It is easy to see that this map is injective: indeed, suppose that we have $m \in tM$ such that $D(m) = 0$, and suppose that $n \in \mathbb{Z}_{>0}$ is the largest integer such that $m \in t^n M$ (such an n exists if and only if m is non-zero). Write $m = t^n m'$ for some $m' \notin tM$. We then have

$$0 = D(m) = D(t^n m') = nt^{n-1} m' + t^n D(m').$$

Dividing by t^{n-1} , this gives us $nm' \in tM$, which implies that $m' \in tM$, contradicting our assumption that m is non-zero. So, we find that a global section of a crystal over T is 0 precisely if it restricts to 0 over x .

By [BM90, 1.3.5] again, restriction from global sections over T to global sections over y is injective. Therefore, a global section of a crystal over T is 0 precisely when it restricts to 0 over y . This proves the claim and the lemma. \square

DEFINITION 5.11. Let T be an \mathcal{S}_K -scheme and let $f \in \text{End}(A_T^{\text{KS}})_{(p)}$. If $p\mathcal{O}_T = 0$, we will say that f is p -special if it satisfies the equivalent conditions of (5.9). In general, we will say that f is p -special if its restrictions to $T \otimes \mathbb{F}_p$ and $T \otimes \mathbb{Q}$ are both p -special.

We will say that f is special if it is ℓ -special for every prime ℓ .

Given any prime ℓ , write $L_\ell(A_T^{\text{KS}})$ for the space of ℓ -special endomorphisms, and $L(A_T^{\text{KS}})$ for the space of special endomorphisms. By definition, we have

$$L(A_T^{\text{KS}}) = \bigcap_{\ell \text{ prime}} L_\ell(A_T^{\text{KS}}).$$

LEMMA 5.12. The space $L(A_T^{\text{KS}}) \subset \text{End}(A_T^{\text{KS}})_{(p)}$ is point-wise fixed by the Rosati involution induced from the polarization λ^{KS} . In particular, $f \mapsto f \circ f$ defines a positive definite $\mathbb{Z}_{(p)}$ -quadratic form on $L(A_T^{\text{KS}})$.

Proof. Consider the G -equivariant symplectic pairing $\psi_\delta : H_{\mathbb{Q}} \otimes H_{\mathbb{Q}} \rightarrow \mathbb{Q}(\nu)$: by construction, for any prime $\ell \neq p$, the corresponding pairing of ℓ -adic sheaves $\mathbf{H}_\ell \otimes \mathbf{H}_\ell \rightarrow \underline{\mathbb{Q}}_\ell(-1)$ is, up to scalars, identified with the polarization pairing induced from λ^{KS} .

The character ψ_δ gives rise to an isomorphism of G -representations $\lambda(\psi_\delta) : H_{\mathbb{Q}} \xrightarrow{\cong} H_{\mathbb{Q}}^\vee(\nu)$. If we identify $H^{\otimes(1,1)}$ with $\text{End}(H)$, we now obtain a G -equivariant involution:

$$i_\delta : H_{\mathbb{Q}}^{\otimes(1,1)} \rightarrow H_{\mathbb{Q}}^{\otimes(1,1)},$$

$$f \mapsto \lambda(\psi_\delta)^{-1} \circ f^\vee(\nu) \circ \lambda(\psi_\delta).$$

Further, for every $v \in L_{\mathbb{Q}}$, $v^* = v \in C_{\mathbb{Q}}$. Therefore, for $z_1, z_2 \in H_{\mathbb{Q}}$, we have

$$\psi_\delta(vz_1, z_2) = \text{Trd}(vz_1\delta z_2^*) = \text{Trd}(z_1\delta z_2^*v) = \text{Trd}(z_1\delta(vz_2)^*) = \psi_\delta(z_1, vz_2).$$

This shows that $i_\delta(v) = v$ for all $v \in L_{\mathbb{Q}}$.

For any prime $\ell \neq p$, the G -equivariant map i_δ induces a map of sheaves over Sh_K (and hence \mathcal{S}_K): $i_\delta : \mathbf{H}_\ell^{\otimes(1,1)} \rightarrow \mathbf{H}_\ell^{\otimes(1,1)}$. By construction, $i_\delta|_{L_\ell} \equiv 1$.

Now, the restriction of i_δ to the ℓ -adic realization of any endomorphism of A^{KS} is exactly the Rosati involution arising from λ^{KS} . This gives us the first assertion. The second follows from the positivity of the Rosati involution. □

We have the following result.

LEMMA 5.13. *Let T be an \mathcal{S}_K -scheme such that every generic point of $T \otimes \mathbb{F}_p$ is the specialization of a point in $T \otimes \mathbb{Q}$. Then an endomorphism $f \in \text{End}(A_T^{\text{KS}})_{(p)}$ is p -special over $T \otimes \mathbb{Q}$ if and only if it is p -special over $T \otimes \mathbb{F}_p$. In particular, in this situation, if f is p -special, then it is in fact special.*

Proof. First, assume that $T = \text{Spec } \mathcal{O}_E$, for some complete discrete valuation ring \mathcal{O}_E with characteristic 0 fraction field E and characteristic p perfect residue field k . Let $s = \text{Spec } k$, and let $\bar{\eta} = \text{Spec } \bar{E}$, for some algebraic closure \bar{E}/E . Then the result holds because the p -adic comparison isomorphism for A_T^{KS} carries $L_{p,\bar{\eta}} \otimes B_{\text{cris}}$ into $L_{\text{cris},s} \otimes B_{\text{cris}}$ (cf. (4.7)(ii)).

For general T , in every connected component of $T \otimes \mathbb{F}_p$, we can find a point s valued in an algebraically closed field that is the specialization of a point valued in a complete discrete valuation field of mixed characteristic $(0, p)$. Now we can apply the result of the previous paragraph to s .

The last assertion follows because ℓ -specialness is independent of the prime ℓ in characteristic 0. □

5.14 Fix a point $x_0 \in \mathcal{S}_K(k)$, where, as always, k is a perfect field of characteristic p . Set $W = W(k)$, and let R_{x_0} be the complete local ring of $\mathcal{S}_{K,W}$ at x_0 ; let $\mathfrak{m} \subset R_{x_0}$ be its maximal ideal. Also, let $\widehat{U} = \text{Spf } R_{x_0}$ be the corresponding formal scheme over W .

Let Λ be a quadratic space over $\mathbb{Z}_{(p)}$; we assume that Λ is finite free over $\mathbb{Z}_{(p)}$. Suppose that we have an isometric map $\iota_0 : \Lambda_0 \rightarrow L(A_{x_0}^{\text{KS}})$. We will consider the deformation functor $\text{Def}_{(x_0, \iota_0)}$, defined as follows: for any $B \in \text{Art}_W$, we have

$$\text{Def}_{(x_0, \iota_0)}(B) = \{(x, \iota) : x \in \widehat{U}(B), \iota : \Lambda_0 \rightarrow L(A_x^{\text{KS}}) \text{ isometric map lifting } \iota_0\}.$$

Here, Art_W is the category of local artinian W -algebras with residue field k .

If $\Lambda = \{v\}$ consists of a single element, and if $\iota_0(v) = f_0$, we will write $\text{Def}_{(x_0, f_0)}$ for the corresponding deformation functor.

The functor $\text{Def}_{(x_0, \iota_0)}$ is represented by a closed formal sub-scheme $\widehat{U}_{\iota_0} \subset \widehat{U}$. This can be seen from the fact that the endomorphism scheme of an abelian scheme is representable and unramified over the base. Again, if $\Lambda = \{v\}$ is a singleton with $\iota_0(v) = f_0$, we will write \widehat{U}_{f_0} for the corresponding formal sub-scheme.

5.15 Suppose that we have a surjection $\mathcal{O} \rightarrow \overline{\mathcal{O}}$ in Art_W , whose kernel I admits nilpotent divided powers. Suppose also that we have $(\overline{x}, \overline{\iota}) \in \widehat{U}_{\iota_0}(\overline{\mathcal{O}})$ giving rise to an abelian scheme $A_{\overline{x}}^{\text{KS}}$ over $\overline{\mathcal{O}}$ equipped with an isometric map $\overline{\iota} : \Lambda \rightarrow L(A_{\overline{x}}^{\text{KS}})$ lifting ι_0 .

Let $\mathbf{H}_{\mathcal{O}}$ be the \mathcal{O} -module obtained by evaluating $\mathbf{H}_{\text{cris,Spec } \overline{\mathcal{O}}}$ on $\text{Spec } \mathcal{O}$, and let $\mathbf{L}_{\mathcal{O}} \subset \text{End}(\mathbf{H}_{\mathcal{O}})$ be the corresponding quadratic space. Denote by $\mathbf{H}_{\overline{\mathcal{O}}}$ and $\mathbf{L}_{\overline{\mathcal{O}}}$ the induced modules over $\overline{\mathcal{O}}$; then $\mathbf{H}_{\overline{\mathcal{O}}} = \mathbf{H}_{\text{dR}, \overline{x}}$ is equipped with its Hodge filtration $F^1 \mathbf{H}_{\overline{\mathcal{O}}}$.

The crystalline realization $L(A_{\overline{x}}^{\text{KS}}) \rightarrow \mathbf{L}_{\overline{\mathcal{O}}}$ composed with $\overline{\iota}$ gives us a map $\iota_{\overline{\mathcal{O}}} : \Lambda \rightarrow \mathbf{L}_{\overline{\mathcal{O}}}$. The image of $\iota_{\overline{\mathcal{O}}}(\Lambda)$ in $\mathbf{L}_{\overline{\mathcal{O}}}$ preserves the Hodge filtration $F^1 \mathbf{H}_{\overline{\mathcal{O}}}$ and so lands in $F^0 \mathbf{L}_{\overline{\mathcal{O}}}$.

By Serre–Tate (cf. [Kat81, 1.2.1]) and Grothendieck–Messing (cf. [Mes72, V.1.6]), we have a natural bijection:

$$\left(\begin{array}{c} \text{Isomorphism classes of} \\ \text{abelian schemes over } \mathcal{O} \text{ lifting } A_{\overline{x}}^{\text{KS}} \end{array} \right) \xrightarrow{\cong} \left(\begin{array}{c} \text{Direct summands } F^1 \mathbf{H}_{\mathcal{O}} \subset \mathbf{H}_{\mathcal{O}} \\ \text{lifting } F^1 \mathbf{H}_{\overline{\mathcal{O}}} \end{array} \right).$$

This works as follows: for any lift A_x of $A_{\overline{x}}^{\text{KS}}$, we have a canonical identification $H_{\text{dR}}^1(A_x/\mathcal{O}) = \mathbf{H}_{\mathcal{O}}$ that carries the Hodge filtration on $H_{\text{dR}}^1(A_x/\mathcal{O})$ to the corresponding summand $F_x^1 \mathbf{H}_{\mathcal{O}} \subset \mathbf{H}_{\mathcal{O}}$.

PROPOSITION 5.16. *The bijection above induces further bijections:*

$$\begin{aligned} (\text{Lifts } x \in \widehat{U}(\mathcal{O}) \text{ of } \overline{x}) &\xrightarrow{\cong} (\text{Isotropic lines } F^1 \mathbf{L}_{\mathcal{O}} \subset \mathbf{L}_{\mathcal{O}} \text{ lifting } F^1 \mathbf{L}_{\overline{\mathcal{O}}}), \\ (\text{Lifts } (x, \iota) \in \widehat{U}_{\iota_0}(\mathcal{O}) \text{ of } (\overline{x}, \overline{\iota})) &\xrightarrow{\cong} \left(\begin{array}{c} \text{Isotropic lines } F^1 \mathbf{L}_{\mathcal{O}} \subset \mathbf{L}_{\mathcal{O}} \text{ lifting } F^1 \mathbf{L}_{\overline{\mathcal{O}}} \\ \text{and orthogonal to the sub-space } \iota_{\overline{\mathcal{O}}}(\Lambda) \end{array} \right). \end{aligned}$$

In particular, both of the sets in the latter bijection are empty, unless $F^1 \mathbf{L}_{\overline{\mathcal{O}}}$ lies in the image in $\mathbf{L}_{\overline{\mathcal{O}}}$ of the sub-module $\iota_{\overline{\mathcal{O}}}(\Lambda)^\perp \subset \mathbf{L}_{\overline{\mathcal{O}}}$.

Proof. In the first of the claimed bijections, there is a natural map in one direction: given a lift $x \in \widehat{U}(\mathcal{O})$ and the identification $\mathbf{L}_{\text{dR}, x} = \mathbf{L}_{\mathcal{O}}$, the Hodge filtration $F^1 \mathbf{L}_{\text{dR}, x}$ gives us an isotropic line $F^1 \mathbf{L}_{\mathcal{O}}$ lifting $F^1 \mathbf{L}_{\overline{\mathcal{O}}}$. Further, an endomorphism $f \in L(A_x^{\text{KS}})$ lifts to an endomorphism of A_x^{KS} if and only if its crystalline realization $f_{\mathcal{O}} \in \mathbf{L}_{\mathcal{O}}$ preserves the Hodge filtration $F^1 \mathbf{H}_{\mathcal{O}}$. Since $F^1 \mathbf{H}_{\mathcal{O}}$ is the kernel of any generator of $F^1 \mathbf{L}_{\mathcal{O}}$, it is easy to see that $f_{\mathcal{O}}$ preserves $F^1 \mathbf{H}_{\mathcal{O}}$ if and only if it is orthogonal to $F^1 \mathbf{L}_{\mathcal{O}}$.

So, it is enough to show that the first map is a bijection. For this, we can work successively with the thickenings $\mathcal{O}/I^{[r-1]} \rightarrow \mathcal{O}/I^{[r]}$ (where $I^{[r]}$ denotes the r th-divided power of I), and assume that $I^2 = 0$. If $\mathfrak{m}_{\mathcal{O}} \subset \mathcal{O}$ is the maximal ideal, we can even work successively with the thickenings $\mathcal{O}/\mathfrak{m}_{\mathcal{O}}^{r-1} I \rightarrow \mathcal{O}/\mathfrak{m}_{\mathcal{O}}^r I$, and further assume that $\mathfrak{m}_{\mathcal{O}} I = 0$. In this case, we find that both sides of the map in question are vector spaces over k of the same dimension, namely $n \cdot \dim_k I$, and that the map is a map of k -vector spaces. Since it is clearly injective, we see that the map must in fact be a bijection.

As for the final assertion, we have only included it to highlight the fact that, when $\iota_{\overline{\mathcal{O}}}(\Lambda)$ is not a direct summand of $\mathbf{L}_{\overline{\mathcal{O}}}$, the formation of its orthogonal complement is not well behaved with respect to arbitrary base change. So, even though $F^1 \mathbf{L}_{\overline{\mathcal{O}}}$ is orthogonal to $\iota_{\overline{\mathcal{O}}}(\Lambda)$, it need not be in the image of the orthogonal complement of $\iota_{\overline{\mathcal{O}}}(\Lambda)$. \square

COROLLARY 5.17. *Let the notation be as above. Suppose that we have a lift $x \in \widehat{U}(\mathcal{O})$ of \overline{x} corresponding to an isotropic line $F^1 \mathbf{L}_{\mathcal{O}} \subset \mathbf{L}_{\mathcal{O}}$. Let $J \subset \mathcal{O}$ be the smallest ideal such that $\overline{\iota}$ lifts to an isometric map $\Lambda \rightarrow L(A_x^{\text{KS}} \otimes_{\mathcal{O}} (\mathcal{O}/J))$. Then J is generated by the elements $[\overline{\iota}(v), w]$, where v varies over the elements of Λ and w is any basis element of $F^1 \mathbf{L}_{\mathcal{O}}$. In particular, $\widehat{U}_{\iota_0} \subset \widehat{U}$ is cut out by $r = \text{rank } \iota_0(\Lambda)$ equations.*

Proof. The assertion about J is immediate.

Set $R = \widehat{\mathcal{O}}_{\mathcal{S}_K, x_0}$. Let $I \subset R$ be the ideal defining \widehat{U}_{ι_0} . Applying the first assertion with $\mathcal{O} = R/\mathfrak{m}I$ and $\overline{\mathcal{O}} = k$ shows that $I/\mathfrak{m}I$ is generated by r elements. Now the last assertion follows from Nakayama’s lemma. \square

5.18 Assume that ι_0 maps Λ injectively onto a direct summand of $L(A_{s_0}^{\text{KS}})$. Let $\iota_{\text{dR}} : \Lambda \otimes R_{\iota_0} \rightarrow \mathbf{L}_{\text{dR}, R_{\iota_0}}$ be the de Rham realization of the universal isometry $\iota : \Lambda \rightarrow L(A^{\text{KS}}|_{\widehat{U}_{\iota_0}})$: it factors through $F^0 \mathbf{L}_{\text{dR}, R_{\iota_0}}$. Write $\mathbf{\Lambda}_{\text{dR}}$ for the image of this map, and let $\overline{\mathbf{\Lambda}}_{\text{dR}}$ be the image of $\mathbf{\Lambda}_{\text{dR}}$ in $\text{gr}_F^0 \mathbf{L}_{\text{dR}, R_{\iota_0}}$.

The Kodaira–Spencer map over R restricts to a map

$$\text{gr}_F^0 \mathbf{L}_{\text{dR}, R_{\iota_0}} \rightarrow \text{gr}_F^{-1} \mathbf{L}_{\text{dR}, R_{\iota_0}} \otimes \widehat{\Omega}_{R_{\iota_0}/W}^1.$$

Since $\mathbf{\Lambda}_{\text{dR}}$ is generated by sections that are parallel for the connection, this map factors as

$$\frac{\text{gr}_F^0 \mathbf{L}_{\text{dR}, R_{\iota_0}}}{\overline{\mathbf{\Lambda}}_{\text{dR}}} \rightarrow \text{gr}_F^{-1} \mathbf{L}_{\text{dR}, R_{\iota_0}} \otimes \widehat{\Omega}_{R_{\iota_0}/W}^1. \tag{5.18.1}$$

As a direct consequence of (5.16) (use $\mathcal{O} = k[\epsilon]$ and $\overline{\mathcal{O}} = k$), we have the following result.

COROLLARY 5.19. *The formal scheme \widehat{U}_{ι_0} is formally smooth over W if and only if $\mathbf{\Lambda}_{\text{dR}}$ maps isomorphically onto $\overline{\mathbf{\Lambda}}_{\text{dR}}$. In this case, the Kodaira–Spencer map in (5.18.1) is an isomorphism. \square*

5.20 Assume that $\Lambda = \{v\}$ is a singleton with $\iota_0(v) = f_0 \neq 0$. For simplicity, set $R = R_{x_0}$, let $I_{f_0} \subset R$ be the ideal defining $\widehat{U}_{f_0} \subset \widehat{U}$ and set $R_{f_0} = R/I_{f_0}$, so that $\widehat{U}_{f_0} = \text{Spf } R_{f_0}$. It follows from (5.17) that I_{f_0} is principal, generated by a single element a_{f_0} . Let $f_{0, \text{dR}} \in \mathbf{L}_{\text{dR}, x_0}$ be the de Rham realization of f_0 . Most of the following result is essentially a retread of [Del81, Proposition 1.5].

PROPOSITION 5.21.

- (i) \widehat{U}_{f_0} is flat over \mathbb{Z}_p ; that is, $p \nmid a_{f_0}$;
- (ii) if $f_{0, \text{dR}} \neq 0$ and $\nu_p(f_0 \circ f_0) \neq 1$, then \widehat{U}_{f_0} is formally smooth.

Proof. For the first statement, we use the argument from the proof of [Del81, 1.6]. As in *loc. cit.*, we reduce immediately to the following assertion: f_0 does not propagate to a special endomorphism of $\widetilde{A}_{R \otimes \mathbb{F}_p}^{\text{KS}}$. Suppose that such a propagation did exist; then we can consider its crystalline realization $f_R \in \mathbf{L}_{\text{dR}, R}$. Choose $k \in \mathbb{Z}_{\geq 0}$ minimal with respect to the condition that $p^{-k} f_R$ belongs to $\mathbf{L}_{\text{dR}, R}$.

Fix a Frobenius lift $\varphi : R \rightarrow R$; this, combined with the F -crystal structure on $\mathbf{H}_{\text{dR}, R}$, endows $\mathbf{L}_{\text{dR}, R}[p^{-1}]$ with a φ -semi-linear Frobenius endomorphism F . Now, $F(f_R) = f_R$, which implies that $F(p^{-k} f_R) = p^{-k} f_R$. By strong divisibility (4.10)(ii), $p^{-k} f_R$ lies in $F^0 \mathbf{L}_{\text{dR}, R} + p \mathbf{L}_{\text{dR}, R}$. In particular, the image \bar{f}_R of $p^{-k} f_R$ in $\mathbf{L}_{\text{dR}, R \otimes \mathbb{F}_p}$ is a non-zero horizontal element that lies in $F^0 \mathbf{L}_{\text{dR}, R \otimes \mathbb{F}_p}$.

But (4.16) shows that the connection on $\mathbf{L}_{\text{dR}, R \otimes \mathbb{F}_p}$ induces an R -linear Kodaira–Spencer isomorphism

$$\text{gr}_F^0 \mathbf{L}_{\text{dR}, R \otimes \mathbb{F}_p} \xrightarrow{\cong} \text{gr}_F^{-1} \mathbf{L}_{\text{dR}, R \otimes \mathbb{F}_p} \otimes \widehat{\Omega}_{R \otimes \mathbb{F}_p/k}^1.$$

This shows that \bar{f}_R must actually lie in $F^1\mathbf{L}_{\mathrm{dR},R\otimes\mathbb{F}_p}$. But once again the connection on $\mathbf{L}_{\mathrm{dR},R\otimes\mathbb{F}_p}$ sets up an R -linear embedding:

$$F^1\mathbf{L}_{\mathrm{dR},R\otimes\mathbb{F}_p} \hookrightarrow \mathrm{gr}_F^0\mathbf{L}_{\mathrm{dR},R\otimes\mathbb{F}_p} \otimes \widehat{\Omega}_{R\otimes\mathbb{F}_p/k}^1.$$

This shows that $\bar{f}_R = 0$, which is a contradiction.

We move on to (ii): since it is defined by a single equation within the formally smooth formal scheme \widehat{U} , \widehat{U}_{f_0} is formally smooth precisely when the map on tangent spaces

$$\widehat{U}_{f_0}(k[\epsilon]) \rightarrow \widehat{U}(k[\epsilon])$$

is not bijective.

Therefore, by applying (5.16) with $\mathcal{O} = k[\epsilon]$, $\bar{\mathcal{O}} = k$ and $\Lambda = \{v\}$, with $\iota_0(v) = f_0$, we see that \widehat{U}_{f_0} fails to be formally smooth precisely when every isotropic lift $F^1(\mathbf{L}_{\mathrm{dR},x_0} \otimes k[\epsilon])$ of $F^1\mathbf{L}_{\mathrm{dR},x_0}$ is orthogonal to $f_{0,\mathrm{dR}}$. It is easy to check that this can happen if and only if $F^1\mathbf{L}_{\mathrm{dR},x_0}$ contains $f_{0,\mathrm{dR}}$.

Suppose therefore that $f_{0,\mathrm{dR}}$ is contained in $F^1\mathbf{L}_{\mathrm{dR},x_0}$. Choose any isotropic line $F^1\mathbf{L}_{\mathrm{cris},x_0} \subset \mathbf{L}_{\mathrm{cris},x_0}$ lifting $F^1\mathbf{L}_{\mathrm{dR},x_0}$. If $f_{0,\mathrm{cris}} \in \mathbf{L}_{\mathrm{cris},x_0}$ is the crystalline realization of f_0 , then $F^1\mathbf{L}_{\mathrm{cris},x_0}$ is generated by an element w of the form $f_{0,\mathrm{cris}} + pv$ for some $v \in \mathbf{L}_{\mathrm{cris},x_0}$. We have

$$w - pv = f_{0,\mathrm{cris}} = F(f_{0,\mathrm{cris}}) = F(w) - pF(v). \tag{5.21.1}$$

By strong admissibility of $\mathbf{L}_{\mathrm{cris},x_0}$, we have $F(w) \in p\mathbf{L}_{\mathrm{cris},x_0}$. This shows that

$$F(v) = \frac{1}{p}F(w) + v - \frac{1}{p}w \in p^{-1}\mathbf{L}_{\mathrm{cris},x_0} \setminus \mathbf{L}_{\mathrm{cris},x_0}. \tag{5.21.2}$$

Applying strong admissibility once again, we conclude that $v \notin F^0\mathbf{L}_{\mathrm{cris},x_0} + p\mathbf{L}_{\mathrm{cris},x_0}$; here, $F^0\mathbf{L}_{\mathrm{cris},x_0} = (F^1\mathbf{L}_{\mathrm{cris},x_0})^\perp$. In particular, $[f_{0,\mathrm{cris}}, v]$ belongs to W^\times .

We now have

$$0 = [f_{0,\mathrm{cris}} + pv, f_{0,\mathrm{cris}} + pv] = [f_{0,\mathrm{cris}}, f_{0,\mathrm{cris}}] + 2p[f_{0,\mathrm{cris}}, v] + p^2[v, v]. \tag{5.21.3}$$

Since $[f_{0,\mathrm{cris}}, v]$ is a unit, this implies that $\nu_p(f_0 \circ f_0) = 1$. □

COROLLARY 5.22. *For every \mathcal{S}_K -scheme T and every prime ℓ , we have*

$$L_p(A_T^{\mathrm{KS}}) \hookrightarrow L_\ell(A_T^{\mathrm{KS}}).$$

In particular, $L(A_T^{\mathrm{KS}}) = L_p(A_T^{\mathrm{KS}})$.

Proof. It follows from the definitions that it suffices to prove the corollary when T is a point $x_0 : \mathrm{Spec} k \rightarrow \mathcal{S}_K$. If $f_0 \in L_p(A_{x_0}^{\mathrm{KS}})$, then, by (5.21), \widehat{U}_{f_0} is flat. This implies that there exists a finite extension $E/W_\mathbb{Q}$ and a lift $x : \mathrm{Spec} \mathcal{O}_E \rightarrow \mathcal{S}_K$ of x_0 such that f_0 lifts to a special endomorphism f of A_x^{KS} . Now (5.13) shows that f , and hence f_0 , is ℓ -special for every ℓ . □

Remark 5.23. From now on, we will refer to p -special endomorphisms simply as special endomorphisms.

5.24 The special endomorphisms are, in a precise sense, objects defined over the integral model \mathcal{S}_{K_0} of the orthogonal Shimura variety Sh_{K_0} . Let $\Delta(K)$ be as in §3.2: it is the Galois group of the finite étale cover $\mathcal{S}_K \rightarrow \mathcal{S}_{K_0}$. Given $[z] \in \Delta(K)$ attached to an element $z \in \mathbb{A}_f^{p,\times}$, its action on \mathcal{S}_K carries $(A^{\text{KS}}, [\eta_G^{\text{KS}}])$ to $(A^{\text{KS}}, [\eta_G^{\text{KS}} \circ z])$. Here, $[\eta_G^{\text{KS}}]$ is the canonical K^p -level structure on A^{KS} (cf. 3.14).

On the other hand, viewing z as a tuple $(z_\ell)_{\ell \neq p}$, we can construct the element $n(z) = \prod_{\ell \neq p} \ell^{-\nu_\ell(z)} \in \mathbb{Z}_{(p)}^\times$. This acts on A^{KS} by multiplication and induces an isomorphism of pairs

$$(A^{\text{KS}}, [\eta_G^{\text{KS}} \circ z]) \xrightarrow{\cong} (A^{\text{KS}}, [\eta_G^{\text{KS}}]).$$

Conjugation by $n(z)$ now produces a canonical identification between the endomorphism sheaves $\underline{\text{End}}([z]^* A^{\text{KS}})_{(p)}$ and $\underline{\text{End}}(A^{\text{KS}})_{(p)}$. This gives us a canonical descent datum that allows us to descend the sheaf $\underline{\text{End}}(A^{\text{KS}})_{(p)}$ to a sheaf \mathbf{E} of $\mathbb{Z}_{(p)}$ -algebras over \mathcal{S}_{K_0} .

It follows now that, given a map $T \rightarrow \mathcal{S}_{K_0}$, we can always attach to it the ‘endomorphism’ algebra $\mathbf{E}(T)$. Moreover, \mathbf{E} comes equipped with canonical realization functors into $\mathbf{H}_p^{\otimes(1,1)}$ over the generic fiber, and into $\mathbf{H}_{\text{cris}}^{\otimes(1,1)}$ over the special fiber. In particular, we can speak of the space of ‘special endomorphisms’ $L(T) \subset \mathbf{E}(T)$, whose realizations at every closed point land in $L_{?} \subset \mathbf{H}_{?}^{\otimes(1,1)}$, where $? = p$, for a point in the generic fiber, and $? = \text{cris}$, for a point in the special fiber. If T is in fact an \mathcal{S}_K -scheme, then we will have $L(T) = L(A_T^{\text{KS}})$.

6. Cycles defined by special endomorphisms

In this section, (L, Q) will be a quadratic space over $\mathbb{Z}_{(p)}$ of signature $(n, 2)$. We will *not* assume it to be self-dual.

6.1 Suppose that (\tilde{L}, \tilde{Q}) is another $\mathbb{Z}_{(p)}$ -quadratic space of signature $(n + d, 2)$ equipped with an embedding

$$(L, Q) \hookrightarrow (\tilde{L}, \tilde{Q}),$$

so that L is a direct summand of \tilde{L} . Let $\Lambda = L^\perp \subset \tilde{L}$; by our assumptions, it is positive definite over \mathbb{R} . Consider the Shimura datum $(\tilde{G}_{\mathbb{Q}}, \tilde{X})$, where \tilde{G} is the smooth $\mathbb{Z}_{(p)}$ -group scheme attached to \tilde{L} : we have an embedding

$$(G_{\mathbb{Q}}, X) \hookrightarrow (\tilde{G}_{\mathbb{Q}}, \tilde{X}).$$

Set $\tilde{K}_p = \tilde{G}(\mathbb{Z}_p)$, and let $\tilde{K}^p \subset \tilde{G}(\mathbb{A}_f^p)$ be a compact open sub-group with $K^p \subset \tilde{K}^p$. We then get a map

$$\text{Sh}_K \rightarrow \text{Sh}_{\tilde{K}}(\tilde{G}_{\mathbb{Q}}, \tilde{X})$$

of Shimura varieties over \mathbb{Q} , which is finite and unramified. Here, as usual, $\tilde{K} = \tilde{K}_p \tilde{K}^p$. For simplicity, write $\text{Sh}_{\tilde{K}}$ for the second Shimura variety. We will also assume that \tilde{K}^p is small enough, so that $\text{Sh}_{\tilde{K}}$ is an algebraic variety, and so that we have the polarized Kuga–Satake abelian scheme $(\tilde{A}_{\text{Sh}_{\tilde{K}}}^{\text{KS}}, \tilde{\lambda}^{\text{KS}})$ over it.

6.2 We will need to briefly review the Serre tensor construction. Let B be a semi-simple associative algebra over \mathbb{Q} , and let $\mathcal{O} \subset B$ be a $\mathbb{Z}_{(p)}$ -order. Let M be a finitely generated, projective \mathcal{O} -module. Suppose that we are given a $\mathbb{Z}_{(p)}$ -scheme S and an abelian scheme $A \rightarrow S$

equipped with an action of \mathcal{O} . Then there is a canonical abelian scheme $\underline{\mathrm{Hom}}_{\mathcal{O}}(M, A)$ over S with the property that, for any S -scheme T , we have an identification of $\mathbb{Z}_{(p)}$ -modules (cf. 3.8):

$$H^0(T, \underline{\mathrm{Hom}}_{\mathcal{O}}(M, A)) = \mathrm{Hom}_{\mathcal{O}}(M, H^0(T, A)).$$

This essentially follows from [Lan08, 5.2.3.9]. We will only need the construction when M is in fact a free \mathcal{O} -module, in which case $\underline{\mathrm{Hom}}_{\mathcal{O}}(M, A)$ can be constructed as a product of $\mathrm{rank}_{\mathcal{O}} M$ copies of A .

Let $\mathbf{H}(A)$ be a degree-1 (de Rham, p -adic, ℓ -adic or crystalline) cohomology sheaf over S attached to A : it has a right action by \mathcal{O} . Then we have a canonical isomorphism

$$\mathbf{H}(A) \otimes_{\mathcal{O}} M \xrightarrow{\sim} \mathbf{H}(\underline{\mathrm{Hom}}_{\mathcal{O}}(M, A)).$$

6.3 Now, $\tilde{C} = C(\tilde{L})$ is a free module over C via the left multiplication action (this follows for instance from [Bou07, §9, Corollaire 3, $n^\circ 3$]). So, we can apply the Serre tensor construction to obtain the abelian scheme $\underline{\mathrm{Hom}}_C(\tilde{C}, A_{\mathrm{Sh}_K}^{\mathrm{KS}})$.

LEMMA 6.4. *The sheaf $\underline{\mathrm{Hom}}_C(\tilde{C}, A_{\mathrm{Sh}_K}^{\mathrm{KS}})$ has a natural $\mathbb{Z}/2\mathbb{Z}$ -grading, as well as a \tilde{C} action compatible with the grading. Moreover, there exists a canonical \tilde{C} -equivariant isomorphism of $\mathbb{Z}/2\mathbb{Z}$ -graded abelian schemes over Sh_K :*

$$i : \tilde{A}_{\mathrm{Sh}_K}^{\mathrm{KS}} \xrightarrow{\sim} \underline{\mathrm{Hom}}_C(\tilde{C}, A_{\mathrm{Sh}_K}^{\mathrm{KS}}).$$

Proof. The $\mathbb{Z}/2\mathbb{Z}$ -grading is simply the diagonal grading, and the action of \tilde{C} is via pre-composition by right multiplication.

The proof is now quite standard, and essentially comes down to the existence of the \tilde{C} -equivariant isomorphism of $\mathbb{Z}/2\mathbb{Z}$ -graded G -representations:

$$H \otimes_C \tilde{C} \rightarrow \tilde{H}, \tag{6.4.1}$$

$$w \otimes z \mapsto w \cdot z. \tag{6.4.2}$$

It gives rise to an isomorphism of tuples $\mathbb{V}_{\mathbb{Q}}(H \otimes_C \tilde{C}) \xrightarrow{\sim} \mathbb{V}_{\mathbb{Q}}(\tilde{H})$. Over $\mathrm{Sh}_{K, \mathbb{C}}$, $\mathbb{V}_{\mathbb{C}}(H \otimes_C \tilde{C})$ (respectively $\mathbb{V}_{\mathbb{C}}(\tilde{H})$) is the $\mathbb{Z}_{(p)}$ -variation of Hodge structures obtained from the cohomology of $\underline{\mathrm{Hom}}_C(\tilde{C}, A_{\mathrm{Sh}_{K, \mathbb{C}}}^{\mathrm{KS}})$ (respectively $\tilde{A}_{\mathrm{Sh}_{K, \mathbb{C}}}^{\mathrm{KS}}$). So, the induced isomorphism $\mathbb{V}_{\mathbb{C}}(H \otimes_C \tilde{C}) \xrightarrow{\sim} \mathbb{V}_{\mathbb{C}}(\tilde{H})$ gives us an isomorphism of abelian schemes

$$i : \tilde{A}_{\mathrm{Sh}_{K, \mathbb{C}}}^{\mathrm{KS}} \xrightarrow{\sim} \underline{\mathrm{Hom}}_C(\tilde{C}, A_{\mathrm{Sh}_{K, \mathbb{C}}}^{\mathrm{KS}}).$$

But, by the functoriality of $\mathbb{V}_{\mathbb{Q}}$, the p -adic realization of i is defined over Sh_K . From this, as in the proof of (3.11), we conclude that i must be defined over Sh_K . \square

PROPOSITION 6.5. *There exists a canonical isometric embedding $\Lambda \hookrightarrow L(\tilde{A}_{\mathrm{Sh}_K}^{\mathrm{KS}})$ mapping onto a direct summand. Furthermore, for any Sh_K -scheme T , there exists a canonical embedding $L(A_T^{\mathrm{KS}}) \hookrightarrow L(\tilde{A}_T^{\mathrm{KS}})$ also mapping onto the direct summand. Under the canonical bilinear pairing $f \mapsto f \circ f$ on $L(\tilde{A}_T^{\mathrm{KS}})$, $L(A_T^{\mathrm{KS}})$ is identified with the orthogonal complement of Λ .*

Proof. Viewing Λ as a trivial representation of G , the natural embedding $\Lambda \hookrightarrow \tilde{L}$ is a map of G -representations. Applying the functor $\mathbb{V}_{\mathbb{Q}}$, we obtain a map $\mathbb{V}_{\mathbb{Q}}(\Lambda) \hookrightarrow \mathbb{V}_{\mathbb{Q}}(\tilde{L})$ and thus an embedding:

$$\Lambda \hookrightarrow \mathrm{Hom}(\mathbb{V}_{\mathbb{Q}}(\mathbb{Z}_{(p)}), \mathbb{V}_{\mathbb{Q}}(\tilde{L})) = L(\tilde{A}_{\mathrm{Sh}_K}^{\mathrm{KS}}).$$

Given an Sh_K -scheme T , we obtain an embedding of $\mathbb{Z}_{(p)}$ -algebras:

$$\text{End}(A_T^{\text{KS}})_{(p)} \hookrightarrow \text{End}(\tilde{A}_T^{\text{KS}})_{(p)}.$$

This is defined as follows: given an endomorphism f of A_T^{KS} , we obtain an endomorphism of \tilde{A}_T^{KS} carrying a map $\varphi : \tilde{C} \rightarrow H^0(T, A^{\text{KS}})$ to the map $f \circ \varphi$. Here, we are using (6.4) to identify \tilde{A}_T^{KS} with $\underline{\text{Hom}}_C(\tilde{C}, A_T^{\text{KS}})$. This embedding is compatible under cohomological realizations with the map $\mathbb{V}_{\mathbb{Q}}(H^{\otimes(1,1)}) \hookrightarrow \mathbb{V}_{\mathbb{Q}}(\tilde{H}^{\otimes(1,1)})$ obtained from the inclusion of G -representations $H^{\otimes(1,1)} \hookrightarrow H^{\otimes(1,1)} \otimes_C \tilde{C} \xrightarrow{\sim} \tilde{H}^{\otimes(1,1)}$.

In fact, this embedding carries $L(A_T^{\text{KS}})$ onto $\Lambda^\perp \subset \tilde{L}(A_T^{\text{KS}})$. To see this, observe that the embedding $L \hookrightarrow \tilde{L}$ of G -representations gives us a canonical map of p -adic sheaves $\mathbf{L}_p \rightarrow \tilde{\mathbf{L}}_p|_{\text{Sh}_K}$. We can now identify both $L(A_T^{\text{KS}})$ and Λ^\perp with those elements of $\tilde{L}(A_T^{\text{KS}})$ whose p -adic realization is a section of $\mathbf{L}_p \subset \tilde{\mathbf{L}}_p|_{\text{Sh}_K}$. \square

6.6 Recall from § 3.13 that, over $\text{Sh}_{\tilde{K}}$ (respectively Sh_K), we have the canonical $\tilde{G}(\mathbb{A}_f^p)$ -torsor $I_{\tilde{G}}^p$ (respectively $G(\mathbb{A}_f^p)$ -torsor I_G^p). There is a canonical $G(\mathbb{A}_f^p)$ -equivariant map $I_G^p \rightarrow I_{\tilde{G}}^p|_{\text{Sh}_K}$, which we can describe explicitly. Recall that I_G^p parameterizes C -equivariant graded isomorphisms

$$\eta : H \otimes \underline{\mathbb{A}}_f^p \xrightarrow{\sim} \mathbf{H}_{\mathbb{A}_f^p}$$

carrying $L \otimes \underline{\mathbb{A}}_f^p$ onto $\mathbf{L}_{\mathbb{A}_f^p}$.

Tensoring both sides of such an isomorphism with \tilde{C} over C and using the isomorphisms $H \otimes_C \tilde{C} \xrightarrow{\sim} \tilde{H}$ and $\mathbf{H}_{\mathbb{A}_f^p} \otimes_C \tilde{C} \xrightarrow{\sim} \tilde{\mathbf{H}}_{\mathbb{A}_f^p}$, we obtain a \tilde{C} -equivariant graded isomorphism

$$\tilde{\eta} : \tilde{H} \otimes \underline{\mathbb{A}}_f^p \xrightarrow{\sim} \tilde{\mathbf{H}}_{\mathbb{A}_f^p}.$$

It carries $L \otimes \underline{\mathbb{A}}_f^p \subset \tilde{H}^{\otimes(1,1)} \otimes \underline{\mathbb{A}}_f^p$ onto $\mathbf{L}_{\mathbb{A}_f^p} \subset \tilde{\mathbf{H}}_{\mathbb{A}_f^p}^{\otimes(1,1)}$. Moreover, if $\iota_0 : \Lambda \hookrightarrow \tilde{L}$ is the natural inclusion, $\tilde{\eta}$ carries $\iota_0 \otimes 1$ to the isometry $\iota_{\mathbb{A}_f^p} : \Lambda \otimes \underline{\mathbb{A}}_f^p \hookrightarrow \tilde{L} \otimes \underline{\mathbb{A}}_f^p$ induced by the inclusion $\Lambda \hookrightarrow L(\tilde{A}_{\text{Sh}_K}^{\text{KS}})$. This essentially shows the following result.

PROPOSITION 6.7. *There is a canonical map $I_G^p \rightarrow I_{\tilde{G}}^p|_{\text{Sh}_K}$. It identifies I_G^p with the sub-sheaf $I_l^p \subset I_{\tilde{G}}^p|_{\text{Sh}_K}$ consisting of the trivializations*

$$\tilde{\eta} : \tilde{H} \otimes \underline{\mathbb{A}}_f^p \xrightarrow{\sim} \tilde{\mathbf{H}}_{\mathbb{A}_f^p},$$

which carry $\iota_0 \otimes 1$ to $\iota_{\mathbb{A}_f^p}$.

Proof. We will only need the fact that I_G^p maps naturally to I_l^p , which we have already seen. The remainder of the proof is left to the reader. \square

LEMMA 6.8. *There exists a self-dual quadratic lattice (\tilde{L}, \tilde{Q}) over $\mathbb{Z}_{(p)}$ and an embedding*

$$(L, Q) \hookrightarrow (\tilde{L}, \tilde{Q})$$

of quadratic lattices carrying L onto a direct summand of \tilde{L} such that $\Lambda = L^\perp$ is positive definite. If $\text{disc}(L)$ is a cyclic abelian group, we can choose \tilde{L} such that $\text{rank } \tilde{L} \leq \text{rank } L + 1$.

Proof. For any $a \in \mathbb{Z}_{(p)}$, let $\langle a \rangle$ be the rank-1 quadratic $\mathbb{Z}_{(p)}$ -module $\mathbb{Z}_{(p)}$ equipped with the quadratic form with value a on 1.

Let $v \in L$ be such that $\nu_p(Q(v))$ is minimal. Then one easily checks that $\nu_p([w_1, w_2]_Q) \geq \nu_p(Q(v))$ for all $w_1, w_2 \in L$. In particular, for any $w \in L$, the projection $w - (2[w, v]_Q/[v, v]_Q)v$ onto $\langle v \rangle^\perp \subset L$ is well defined, and we obtain an orthogonal decomposition

$$L = \langle v \rangle \oplus \langle v \rangle^\perp.$$

Applying this iteratively, we find that (L, Q) can be diagonalized, so that it is isometric to a lattice of the form

$$\left(\bigoplus_{i=1}^n \langle a_i \rangle \right) \oplus \left(\bigoplus_{j=1}^2 \langle -b_j \rangle \right),$$

where a_i and b_j positive integers for varying i and j .

For any $b \in \mathbb{Z}_{(p)}$, the quadratic space $\langle b \rangle$ embeds as a direct summand spanned by the element $e + \frac{1}{2}b \cdot f$ within the hyperbolic plane $U = \mathbb{Z}_{(p)}e \oplus \mathbb{Z}_{(p)}f$ with $e^2 = f^2 = 0$ and $[e, f] = 1$. Also, given any positive element $a \in \mathbb{Z}_{(p)}^{>0}$, we can find a self-dual positive definite quadratic space $E_+(a)$ over $\mathbb{Z}_{(p)}$ such that $\langle a \rangle$ embeds as a direct summand of $E_+(a)$. In other words, we need a binary quadratic form over $\mathbb{Z}_{(p)}$, which primitively represents a and whose discriminant is a negative element of $\mathbb{Z}_{(p)}^\times$. Take the form to be $ax^2 + xy + cy^2$ with $c = pc'$ for $c' \in \mathbb{Z}_{(p)}^{>0}$ satisfying $1 - 4ac' < 0$. The discriminant is then $1 - 4pac'$, which is negative and a unit in $\mathbb{Z}_{(p)}$.

We now find that L embeds isometrically as a direct summand of $\tilde{L} = (\bigoplus_{i=1}^n E_+(a_i)) \oplus H^{\oplus 2}$.

Suppose now that $\text{disc}(L)$ is cyclic. This implies that at most one element in the set $\{a_i : 1 \leq i \leq n\} \cup \{b_1, b_2\}$ is a non-unit. If every element is a unit, then we can take $\tilde{L} = L$. Otherwise, without loss of generality, we can assume that exactly one of a_1 or b_1 is a non-unit. In the former case, we set $\tilde{L} = E_+(a_1) \oplus (\bigoplus_{i=2}^n \langle a_i \rangle) \oplus (\bigoplus_{j=1}^2 \langle -b_j \rangle)$. In the latter, we can take $\tilde{L} = (\bigoplus_{i=1}^n \langle a_i \rangle) \oplus \langle -b_2 \rangle \oplus U$. \square

6.9 We will now fix an embedding $(L, Q) \hookrightarrow (\tilde{L}, \tilde{Q})$ as in (6.8). In the case where $\text{disc}(L)$ is cyclic, we will assume in addition that \tilde{L} has been chosen so that $\text{rank } \Lambda \leq 1$.

Let $\mathcal{S}_{\tilde{K}}$ (respectively $\mathcal{S}_{\tilde{K}_0}$) be the integral canonical model for $\text{Sh}_{\tilde{K}}$ (respectively $\text{Sh}_{\tilde{K}_0}$) over $\mathbb{Z}_{(p)}$ (cf. 4.1). Assume that \tilde{K} is small enough. Then, over $\mathcal{S}_{\tilde{K}}$, we have the Kuga–Satake abelian scheme $(\tilde{A}^{\text{KS}}, \tilde{\lambda}^{\text{KS}}, [\tilde{\eta}^{\text{KS}}])$. Since $\mathcal{S}_{\tilde{K}}$ is smooth, and in particular normal, both the $\tilde{G}(\mathbb{A}_f^p)$ -torsor $I_{\tilde{G}}^p$ and the canonical K^p -level structure $[\tilde{\eta}_{\tilde{G}}] \in H^0(\text{Sh}_{\tilde{K}}, I_{\tilde{G}}^p/\tilde{K}^p)$ (cf. 3.13) extend over $\mathcal{S}_{\tilde{K}}$: we will denote these extensions by the same symbols.

The $G_0(\mathbb{A}_f^p)$ -torsor $I_{G_0}^p/\mathbb{G}_m(\mathbb{A}_f^p)$ has a canonical descent $I_{G_0}^p$ over $\mathcal{S}_{\tilde{K}_0}$: it parameterizes certain orientation preserving isometries

$$\tilde{\eta}_0 : \tilde{L} \otimes \mathbb{A}_f^p \xrightarrow{\sim} \tilde{L}_{\mathbb{A}_f^p}.$$

Again, we have a canonical section $[\tilde{\eta}_{G_0}^p]$ of $I_{G_0}^p/\tilde{K}_0^p$ over $\mathcal{S}_{\tilde{K}_0}$.

6.10 For any $\mathcal{S}_{\tilde{K}_0}$ -scheme T , write $\tilde{L}(T)$ for the group of special endomorphisms defined as in § 5.24; in particular, if T is actually a scheme over $\mathcal{S}_{\tilde{K}}$, we will have $\tilde{L}(T) = L(\tilde{A}_T^{\text{KS}})$.

Let $\iota_0 : \Lambda \hookrightarrow \tilde{L}$ be the natural embedding. Suppose that we are given an isometric map $\iota : \Lambda \rightarrow L(T)$; since Λ is positive definite, any such map has to be an embedding. Let $I_{\iota,0}^p \subset I_{G_0}^p|_T$

be the sub-sheaf of isomorphisms $\tilde{\eta}_0$ that carry ι_0 to ι ; then, by Witt's extension theorem, $I_{\iota,0}^p$ is a torsor over T under $G_0(\mathbb{A}_f^p)$.

If the map $T \rightarrow \mathcal{S}_{\tilde{K}_0}$ arises from a lift $T \rightarrow \mathcal{S}_{\tilde{K}}$, then one can also define a sub-sheaf $I_{\iota}^p \subset I_{\tilde{G}}^p|_T$: this is the pre-image of $I_{\iota,0}^p$ under the natural quotient map $I_{\tilde{G}}^p|_T \rightarrow I_{\tilde{G}_0}^p|_T$, and is thus a $G(\mathbb{A}_f^p)$ -torsor over T .

DEFINITION 6.11. For any $\mathcal{S}_{\tilde{K}_0}$ -scheme T , a Λ -structure for T is an isometric map $\iota : \Lambda \rightarrow L(T)$. Given an $\mathcal{S}_{\tilde{K}}$ -scheme T and a Λ -structure ι for T , a K^p -level structure on (T, ι) is a section $[\eta_{\iota}]$ of I_{ι}^p/K^p over T mapping to $[\tilde{\eta}_{\tilde{G}}]$ under the obvious map

$$I_{\iota}^p/K^p \rightarrow (I_{\tilde{G}}^p/\tilde{K}^p)|_T.$$

Completely analogously, supposing only that T is a $\mathcal{S}_{\tilde{K}_0}$ -scheme, one can also define the notion of a K_0^p -level structure on (T, ι) as a section $[\eta_{\iota,0}]$ of $I_{\iota,0}^p/K_0^p$ mapping to $[\tilde{\eta}_{\tilde{G}_0}]$ in $H^0(T, I_{\tilde{G}_0}^p/\tilde{K}_0^p)$.

6.12 Let $Z_{K^p}(\Lambda)$ be the functor on $\mathcal{S}_{\tilde{K}}$ -schemes whose value on any $\mathcal{S}_{\tilde{K}}$ -scheme T is given by

$$Z_{K^p}(\Lambda)(T) = \{(\iota, [\eta_{\iota}]) : \iota \text{ a } \Lambda\text{-structure for } T; [\eta_{\iota}] \text{ a } K^p\text{-level structure for } (T, \iota)\}.$$

Similarly, one defines a functor $Z_{K_0^p}(\Lambda)$ on $\mathcal{S}_{\tilde{K}_0}$ -schemes employing K_0^p -level structures.

PROPOSITION 6.13. The functor $Z_{K^p}(\Lambda)$ (respectively $Z_{K_0^p}(\Lambda)$) is represented by a scheme finite and unramified over $\mathcal{S}_{\tilde{K}}$ (respectively over $\mathcal{S}_{\tilde{K}_0}$). Moreover, the natural map $Z_{K^p}(\Lambda) \rightarrow Z_{K_0^p}(\Lambda)$ is finite étale.

Proof. It is clear from the definitions that the map $Z_{K^p}(\Lambda) \rightarrow Z_{K_0^p}(\Lambda)$ is finite étale, so it is enough to prove the remaining assertions for $Z_{K^p}(\Lambda)$. To show representability, we first note that $\underline{\text{End}}(\tilde{A}^{\text{KS}})_{(p)}$ is representable over $\mathcal{S}_{\tilde{K}}$ by an inductive limit of schemes that are locally of finite type. Indeed, if we fix a representative \tilde{A} in the prime-to- p isogeny class of \tilde{A}^{KS} , then the endomorphism scheme $\underline{\text{End}}(\tilde{A})$ is known to be representable by a scheme locally of finite type over $\mathcal{S}_{\tilde{K}}$, and we can identify $\underline{\text{End}}(\tilde{A}^{\text{KS}})_{(p)}$ with the inductive limit of the system $\{\underline{\text{End}}(\tilde{A})\}_{m \in \mathbb{Z} \setminus p\mathbb{Z}}$, where, for $m_1 \mid m_2$, the transition map from the copy of $\underline{\text{End}}(\tilde{A})$ in the m_1 th position to that in the m_2 th position is given by multiplication by m_2/m_1 . Since the property of being a special endomorphism is a closed condition on the base, we see that $Z_{K^p}(\Lambda)$ is also represented by an inductive scheme. To show that it is in fact represented by a scheme, it is enough to show that it is finite and unramified over $\mathcal{S}_{\tilde{K}}$.

The unramifiedness is a consequence of the fact that endomorphisms of abelian schemes lift uniquely (if at all) over nilpotent thickenings. The néronian property of abelian schemes over discrete valuation rings combines with the valuative criterion to show that $Z_{K^p}(\Lambda)$ is proper.

Thus, it only remains to show quasi-finiteness. For this, take any geometric point $s \rightarrow \mathcal{S}_{\tilde{K}}$. We view $[\tilde{\eta}_{\tilde{G}_0,s}]$ as a \tilde{K}_0^p -orbit of isometries

$$\tilde{\eta}_{0,s} : \tilde{L} \otimes \mathbb{A}_f^p \xrightarrow{\cong} \tilde{L}_{\mathbb{A}_f^p,s}.$$

If ι is a Λ -structure for s such that (s, ι) admits a K_0^p -level structure, then, for any $v \in \Lambda$, the pre-image of the \mathbb{A}_f^p -realization of $\iota(v)$ under $\tilde{\eta}_{0,s}$ must lie within the set $\tilde{L} \cap K_0^p \cdot \iota_0(v)$. Since \tilde{L} is a discrete sub-group of $\tilde{L}_{\mathbb{A}_f^p,s}$, this set is finite. So, we see that the possible Λ -structures for s (admitting a K_0^p -level structure) are finite in number. From this, it follows easily that the fiber of $Z_{K^p}(\Lambda)$ over s is finite. \square

The following result is presumably well known, but we include it for lack of a reference.

LEMMA 6.14. *Let $W = W(k)$ and let $E/W_{\mathbb{Q}}$ be a finite extension of ramification index $e \leq p - 1$. Let A be an abelian scheme over \mathcal{O}_E with special fiber A_0 . Then $\text{End}(A_E) \otimes W$ embeds in $\text{End}_W(H_{\text{cris}}^1(A_0/W))$ as a direct summand. In particular, $\text{End}(A_E) \otimes k$ maps injectively into $\text{End}_k(H_{\text{dR}}^1(A_0/k))$.*

Proof. Without loss of generality, we can assume that k is algebraically closed. There is a canonical comparison isomorphism

$$H_{\text{cris}}^1(A_0/W) \otimes_W \mathcal{O}_E \xrightarrow{\cong} H_{\text{dR}}^1(A/\mathcal{O}_E). \tag{6.14.1}$$

Note that this is where we need the hypothesis $e \leq p - 1$, which ensures that the map $\mathcal{O}_E \rightarrow k$ is a divided power thickening.

Set $M = \text{End}_W(H_{\text{cris}}^1(A_0/W))$: we have a direct summand $M^0 \subset M$ consisting of endomorphisms that preserve the Hodge filtration $F^1 H_{\text{dR}}^1(A/\mathcal{O}_E)$. Here, we use the canonical isomorphism (6.14.1) to view M as a group of endomorphisms of $H_{\text{dR}}^1(A/\mathcal{O}_E)$. The conjugation action of the semi-linear Frobenius on $H_{\text{cris}}^1(A_0/W)$ induces an isomorphism $F : \sigma^* M[1/p] \xrightarrow{\cong} M[1/p]$ such that $F(\sigma^* M^0) \subset M$. The last condition holds because the image of $F^1 H_{\text{dR}}^1(A/\mathcal{O}_E)$ in $H_{\text{dR}}^1(A_0/k)$ has the following property: its pull-back via σ is precisely the kernel of the map $F : \text{Fr}^* H_{\text{dR}}^1(A_0/k) \rightarrow H_{\text{dR}}^1(A_0/k)$.

One can deduce from classical Dieudonné theory and Grothendieck–Messing theory that there is a canonical isomorphisms of \mathbb{Z}_p -modules:

$$\text{End}(A[p^\infty]) \xrightarrow{\cong} (M^0)^{F=1}.$$

Moreover, by the nerónian property, $\text{End}(A_E) = \text{End}(A)$. Also, $\text{End}(A) \otimes \mathbb{Z}_p$ is a direct summand of $\text{End}(A[p^\infty])$, since any element of $\text{End}(A)$ that kills the p -torsion $A[p]$ has to be divisible by p .

So, to finish the proof, we have to show that $(M^0)^{F=1} \otimes W$ maps onto a direct summand of M . For this, set

$$M' = \{m \in M^0 : F^i(m) \in M^0, \text{ for all } i \in \mathbb{Z}_{\geq 0}\}.$$

We need to explain what we mean by $F^i(m)$. One defines this inductively: we set $F(m) = F(\sigma^* m)$ and $F^i(m) = F(\sigma^* F^{i-1}(m))$, where we are using the assumption that $F^{i-1}(m) \in M^0$ in each inductive step.

Now, F restricts to a (necessarily injective) map $\sigma^* M' \rightarrow M'$. Moreover, M' is a direct summand of M , and $(M^0)^{F=1} \otimes W$ clearly maps into M' . It follows from the Dieudonné–Manin classification of F -crystals over W [Man62] that there exists a largest F -stable direct summand $M^{\text{ét}} \subset M'$ such that F induces an isomorphism $\sigma^* M^{\text{ét}} \xrightarrow{\cong} M^{\text{ét}}$. Moreover, it is known [Kat73, 4.1.1] that the map

$$(M^{\text{ét}})^{F=1} \otimes_{\mathbb{Z}_p} W \rightarrow M^{\text{ét}}$$

is an isomorphism. It follows *a fortiori* that the map $(M^0)^{F=1} \otimes W \rightarrow M$ identifies its source with the direct summand $M^{\text{ét}} \subset M$. □

6.15 The map $\text{Sh}_{K_0} \rightarrow \text{Sh}_{\tilde{K}_0}$ canonically lifts to a map $\text{Sh}_{K_0} \rightarrow Z_{K_0^p}(\Lambda)$. Indeed, it is enough to show that $\text{Sh}_K \rightarrow \text{Sh}_{\tilde{K}}$ lifts canonically to a map $\text{Sh}_K \rightarrow Z_{K^p}(\Lambda)$. This follows from (6.5) and (6.7).

Let ι be the tautological Λ -structure over $Z_{K_0^p}(\Lambda)$ and let $\Lambda_{\text{dR}} = \Lambda \otimes \mathcal{O}_{Z_{K_0^p}(\Lambda)}$. The map ι induces a map of vector bundles:

$$\iota_{\text{dR}} : \Lambda_{\text{dR}} \rightarrow \tilde{\mathbf{L}}_{\text{dR}}|_{Z_{K_0^p}(\Lambda)}.$$

LEMMA 6.16. (i) *There exists an open sub-scheme $Z_{K_0^p}^{\text{pr}}(\Lambda) \subset Z_{K_0^p}(\Lambda)$ such that $T \rightarrow Z_{K_0^p}(\Lambda)$ factors through $Z_{K_0^p}^{\text{pr}}(\Lambda)$ if and only if the restriction of $\text{coker } \iota_{\text{dR}}$ to T is a vector bundle of rank $n + 2$.*

(ii) *$Z_{K_0^p}^{\text{pr}}(\Lambda)$ contains the generic fiber $Z_{K_0^p}(\Lambda)_{\mathbb{Q}}$.*

(iii) *Let $W = W(\overline{\mathbb{F}}_p)$, and suppose that we have $\tilde{s} : \text{Spec } W \rightarrow Z_{K_0^p}(\Lambda)$ such that the restriction to $\text{Spec } W_{\mathbb{Q}}$ factors through Sh_{K_0} . Then \tilde{s} factors through $Z_{K_0^p}^{\text{pr}}(\Lambda)$.*

(iv) *Suppose that T is smooth over $\mathbb{Z}_{(p)}$ and that $f : T \rightarrow Z_{K_0^p}(\Lambda)$ is such that $f|_{T_{\mathbb{Q}}}$ factors through Sh_{K_0} . Then f factors through $Z_{K_0^p}^{\text{pr}}(\Lambda)$.*

(v) *If L has square-free discriminant, then $Z_{K_0^p}^{\text{pr}}(\Lambda) = Z_{K_0^p}(\Lambda)$.*

Proof. We begin with (i): choose a basis element for the top exterior power $\wedge^d \Lambda$, identifying it with $\mathbb{Z}_{(p)}$. Then the map ι_{dR} induces a map

$$\mathcal{O}_{Z_{K_0^p}(\Lambda)} = \wedge^d \Lambda_{\text{dR}} \xrightarrow{\wedge^d \iota_{\text{dR}}} \wedge^d \tilde{\mathbf{L}}_{\text{dR}}$$

and hence a section $e \in H^0(Z_{K_0^p}(\Lambda), \wedge^d \tilde{\mathbf{L}}_{\text{dR}})$. One easily checks now that $Z_{K_0^p}^{\text{pr}}(\Lambda)$ is the open locus where this section does not vanish.

To prove (ii), it suffices to show that $Z_{K_0^p}^{\text{pr}}(\Lambda)$ contains every point $s \in Z_{K_0^p}(\Lambda)(\mathbb{C})$. But the map $\Lambda_{\text{dR},s} \rightarrow \tilde{\mathbf{L}}_{\text{dR},s}$ arises from an isometric embedding of $\mathbb{Z}_{(p)}$ -Hodge structures $\iota_s : \Lambda \hookrightarrow \tilde{\mathbf{L}}_{B,s}$. In particular, it has to be an embedding of \mathbb{C} -vector spaces.

Define $Z_{K_0^p}^{\text{pr}}(\Lambda)$ to be the pre-image in $Z_{K_0^p}(\Lambda)$ of $Z_{K_0^p}^{\text{pr}}(\Lambda)$. It suffices to prove the remaining assertions for $\mathcal{S}_{\tilde{K}}$ -schemes with $Z_{K_0^p}^{\text{pr}}(\Lambda)$ replaced everywhere by $Z_{K_0^p}^{\text{pr}}(\Lambda)$.

We will now consider (iii). Choose an algebraic closure $\overline{E}/W(k)_{\mathbb{Q}}$. By (6.14), it suffices to show that the étale realization of $\iota(\Lambda)$ generates a direct summand of $\tilde{\mathbf{L}}_{p,\tilde{s}_{\overline{E}}}$. But, over Sh_{K_0} , the sub-space generated by this realization is globally a direct summand of $\tilde{\mathbf{L}}_p$. Indeed, its inclusion in the latter is induced by the map of G_0 -representations $\Lambda \hookrightarrow \tilde{\mathbf{L}}$.

Part (iv) now follows: indeed, (i) and (iii) show that $U := f^{-1}(Z_{K_0^p}^{\text{pr}}(\Lambda))$ is an open sub-scheme of T containing $T_{\mathbb{Q}}$, and through which all the $W(\overline{\mathbb{F}}_p)$ -valued points of T factor. Since T is smooth over $\mathbb{Z}_{(p)}$, this implies that all the $\overline{\mathbb{F}}_p$ -points of T factor through U , and so U must be all of T .

Now assume that L has square-free discriminant; then $\text{disc}(\Lambda)$ is either trivial or isomorphic to $\mathbb{Z}/p\mathbb{Z}$. In the first case, $Z_{K_0^p}(\Lambda)$ is smooth, and so we are done by (iv). In the second case, by our assumptions, Λ is of rank 1, and is generated by an element v satisfying $\text{ord}_p(Q(v)) = 1$. In particular, given any $\overline{\mathbb{F}}_p$ -point x_0 of $Z_{K_0^p}(\Lambda)$, the crystalline realization of the associated special endomorphism $f_0 \in L(\tilde{A}_{x_0}^{\text{KS}})$ must necessarily generate a direct summand of $\mathbf{L}_{\text{cris},x_0}$. This shows (v). □

The following result is directly inspired by [Ogu79, Theorem 2.9].

PROPOSITION 6.17. *Let Z^{sm} be the complement in $Z_{K^p}(\Lambda)$ of the non-smooth locus in $Z_{K^p}(\Lambda)_{\mathbb{F}_p}$. Let $\bar{\eta}$ be a generic point of $Z_{\mathbb{F}_p}^{\text{sm}}$ with algebraically closed residue field. Suppose that $n > r = \text{rank}(\Lambda)$. Then the abelian variety $\tilde{A}_{\bar{\eta}}^{\text{KS}}$ is ordinary and the map*

$$\iota : \Lambda \rightarrow L(\tilde{A}_{\bar{\eta}}^{\text{KS}})$$

is an isomorphism.

Proof. It follows from 5.19 that $\Lambda \otimes k(\bar{\eta})$ maps injectively into $\text{gr}_F^0 \tilde{\mathbf{L}}_{\text{dR}, \bar{\eta}}$. Therefore, over a sufficiently small étale neighborhood Z_0 of $\bar{\eta}$ in $Z_{\mathbb{F}_p}^{\text{sm}}$, the de Rham realization $\mathbf{\Lambda}_{\text{dR}}$ of $\iota(\Lambda)$ will map isomorphically onto a direct summand $\bar{\mathbf{\Lambda}}_{\text{dR}}$ of $\text{gr}_F^0 \tilde{\mathbf{L}}_{\text{dR}}|_{Z_0}$.

Now, consider the Frobenius map $\text{Fr}^* \mathbf{H}_{\text{dR}}|_{Z_0} \rightarrow \mathbf{H}_{\text{dR}}|_{Z_0}$: its image is a lagrangian sub-space $\mathbf{C} \subset \mathbf{H}_{\text{dR}}|_{Z_0}$, and the locus where $\tilde{A}_{\bar{\eta}}^{\text{KS}}|_{Z_0}$ is ordinary is precisely the open sub-space where $\mathbf{C} + F^1 \mathbf{H}_{\text{dR}}|_{Z_0} = \mathbf{H}_{\text{dR}}|_{Z_0}$. By the discussion in § 4.19, after localizing on Z_0 if required, the annihilator of \mathbf{C} is a parallel isotropic line $\mathbf{N} \subset \tilde{\mathbf{L}}_{\text{dR}}|_{Z_0}$. So, we can also describe the non-ordinary locus as the closed sub-space where we have

$$F^1 \tilde{\mathbf{L}}_{\text{dR}}|_{Z_0} \subset F^0 \tilde{\mathbf{L}}_{\text{dR}}|_{Z_0} \cap \mathbf{\Lambda}_{\text{dR}}^\perp \cap \mathbf{N}^\perp.$$

Now, by 4.20, we can assume that $\mathbf{N} + \mathbf{\Lambda}_{\text{dR}}$ is a horizontal direct summand of rank $r + 1$ in $\tilde{\mathbf{L}}_{\text{dR}}|_{Z_0}$. Then, arguing as in the proof of [Ogu79, Theorem 2.9], we find that the locus where $F^1 \mathbf{L}_{\text{dR}}|_{Z_0}$ is contained in this direct summand has dimension at most r . Since $n > r$, we can throw this locus out and assume that the summand does not contain $F^1 \mathbf{L}_{\text{dR}}|_{Z_0}$.

Equivalently, we can assume that \mathbf{N} is not contained in $F^1 \mathbf{L}_{\text{dR}}|_{Z_0} + \mathbf{\Lambda}_{\text{dR}}$; or that $F^0 \mathbf{L}_{\text{dR}}|_{Z_0} \cap \mathbf{\Lambda}_{\text{dR}}^\perp$ is not contained in \mathbf{N}^\perp . Again, arguing as in [Ogu79, Theorem 2.9], we now find that the non-ordinary locus is smooth of dimension $n - 1$. This proves that $\tilde{A}_{\bar{\eta}}^{\text{KS}}$ must be ordinary.

Suppose that ι is not an isomorphism. Then we can find a non-zero element $f \in L(\tilde{A}_{\bar{\eta}}^{\text{KS}})$ such that $\iota(\Lambda) + \langle f \rangle \subset L(\tilde{A}_{\bar{\eta}}^{\text{KS}})$ is a direct summand of rank $r + 1$.

Now, by shrinking Z_0 if necessary, we can assume that f extends to an element in $L(\tilde{A}_{Z_0}^{\text{KS}})$. Let \mathbf{f}_{dR} be the de Rham realization of f , and let $\mathbf{\Lambda}_{\text{dR}} \subset \tilde{\mathbf{L}}_{\text{dR}}|_{Z_0}$ be that of $\iota(\Lambda)$.

Since $\tilde{A}_{\bar{\eta}}^{\text{KS}}$ is ordinary, the map

$$L(\tilde{A}_{\bar{\eta}}^{\text{KS}}) \otimes k(\bar{\eta}) \rightarrow \tilde{\mathbf{L}}_{\text{dR}, \bar{\eta}} \tag{6.17.1}$$

is injective. Therefore, after further localizing on Z_0 , we can assume that the sub-sheaf

$$\mathbf{\Lambda}_{\text{dR}} + \langle \mathbf{f}_{\text{dR}} \rangle \subset \tilde{\mathbf{L}}_{\text{dR}}|_{Z_0}$$

is a local direct summand of rank $r + 1$.

Using 5.19 and [Ogu79, Remark 2.8], we find that the locus in Z_0 where $F^1 \tilde{\mathbf{L}}_{\text{dR}}$ is contained in $\mathbf{\Lambda}_{\text{dR}} + \langle \mathbf{f}_{\text{dR}} \rangle$ is a closed sub-space of dimension at most $1(r + 1 - 1) = r$. Since $n > r$ by hypothesis, and since $\dim Z_0 = n$, this implies that we can shrink Z_0 further and assume that $F^1 \tilde{\mathbf{L}}_{\text{dR}}|_{Z_0}$ is not contained in $\mathbf{\Lambda}_{\text{dR}} + \langle \mathbf{f}_{\text{dR}} \rangle$. Since $F^1 \tilde{\mathbf{L}}_{\text{dR}}|_{Z_0}$ and $\langle \mathbf{f}_{\text{dR}} \rangle$ are both local direct summands in $\tilde{\mathbf{L}}_{\text{dR}}|_{Z_0}$ of rank 1, it follows that $\langle \mathbf{f}_{\text{dR}} \rangle$ is not contained in $F^1 \tilde{\mathbf{L}}_{\text{dR}}|_{Z_0} + \mathbf{\Lambda}_{\text{dR}}$.

Therefore, if $\bar{\mathbf{\Lambda}}_{\text{dR}}$ is the image of $\mathbf{\Lambda}_{\text{dR}}$ in $\text{gr}_F^0 \tilde{\mathbf{L}}_{\text{dR}}|_{Z_0}$, we see that the map

$$\langle \mathbf{f}_{\text{dR}} \rangle \rightarrow \frac{\text{gr}_F^0 \tilde{\mathbf{L}}_{\text{dR}}|_{Z_0}}{\bar{\mathbf{\Lambda}}_{\text{dR}}}$$

is non-zero. But, on the other hand, f_{dR} is horizontal for the Gauss–Manin connection, which contradicts the fact (5.19) that the Kodaira–Spencer map

$$\frac{\text{gr}_F^0 \tilde{\mathbf{L}}_{\text{dR}}|_{Z_0}}{\mathbf{\Lambda}_{\text{dR}}} \rightarrow \text{gr}_F^{-1} \tilde{\mathbf{L}}_{\text{dR}} \otimes \Omega_{Z_0/\mathbb{F}_p}^1$$

is an isomorphism. □

COROLLARY 6.18. *Set $\Lambda' = \Lambda \perp \langle m \rangle$ for some $m \in \mathbb{Z}_{(p)}^{>0}$. The map $Z_{K^p}(\Lambda') \rightarrow \mathcal{S}_{\tilde{K}}$ factors through $Z_{K^p}(\Lambda)$ in the obvious way. Suppose that $n > \text{rank}(\Lambda)$; then the scheme*

$$Z_{K^p}(\Lambda') \times_{Z_{K^p}(\Lambda)} Z^{\text{sm}}$$

is flat over $\mathbb{Z}_{(p)}$.

Proof. Let Z' be the fiber product under consideration: it is, étale locally on Z' , an effective Cartier divisor on Z^{sm} . Since Z^{sm} is smooth over $\mathbb{Z}_{(p)}$, Z' fails to be flat precisely when its image in Z^{sm} contains an irreducible component $Z_0 \subset Z_{\mathbb{F}_p}^{\text{sm}}$. Suppose that this is the case, and let $\tilde{\eta}$ be a geometric generic point of Z_0 . Then we find that the map $\Lambda \rightarrow L(\tilde{A}_{\tilde{\eta}}^{\text{KS}})$ extends to an isometry on Λ' . But, by (6.17), this is impossible. □

6.19 Recall from (4.13) that we have a canonical \tilde{G} -torsor $\tilde{\mathcal{P}}_{\text{dR}}$ over $\mathcal{S}_{\tilde{K}}$ consisting of \tilde{G} -structure preserving trivializations of $\tilde{\mathbf{H}}_{\text{dR}}$. Let ι be the tautological Λ -structure over $Z(\Lambda)$, and let $\iota_{\text{dR}} : \mathbf{\Lambda}_{\text{dR}} \hookrightarrow \tilde{\mathbf{L}}_{\text{dR}}$ be the de Rham realization of ι . Define $\mathcal{P}_{\text{dR},\Lambda} \subset \tilde{\mathcal{P}}_{\text{dR},Z_{K^p}(\Lambda)}$ to be the G -equivariant sub-functor such that, for any $Z_{K^p}(\Lambda)$ -scheme T , we have

$$\mathcal{P}_{\text{dR},\Lambda}(T) = \{\xi \in \tilde{\mathcal{P}}_{\text{dR}}(T) : \xi \circ (\iota_0 \otimes 1) = \iota_{\text{dR}}\}.$$

PROPOSITION 6.20.

- (i) *The restriction of $\mathcal{P}_{\text{dR},\Lambda}$ over $Z_{K^p}^{\text{pr}}(\Lambda)$ is a G -torsor.*
- (ii) *The map $p_2 : \mathcal{P}_{\text{dR},\Lambda} \rightarrow M_G^{\text{loc}}$, given, for any $\mathbb{Z}_{(p)}$ -scheme T , by*

$$\begin{aligned} \mathcal{P}_{\text{dR},\Lambda}(T) &\rightarrow M_G^{\text{loc}}(T), \\ (x, \xi) &\mapsto \xi^{-1}(F^1 \tilde{\mathbf{L}}_{\text{dR},x}), \end{aligned}$$

is G -equivariant and smooth of relative dimension $\dim G_{\mathbb{Q}}$. Here, $(x, \xi) \in \mathcal{P}_{\text{dR},\Lambda}(T)$ lies over a point $x \in Z_{K^p}(\Lambda)(T)$.

- (iii) *Around any point $x \in Z_{K^p}^{\text{pr}}(\Lambda)$ there exist an étale neighborhood $U \rightarrow Z_{K^p}^{\text{pr}}(\Lambda)$ and a section $s : U \rightarrow \mathcal{P}_{\text{dR},\Lambda}$ of the G -torsor $\mathcal{P}_{\text{dR},\Lambda}$ such that the induced map $p_2 \circ s : U \rightarrow M_G^{\text{loc}}$ is étale.*

Proof. We first note that the basic ideas for the proof can already be found in [RZ96, §3] and [Pap00, Theorem 2.2].

Part (i) is an easy consequence of the definition of $Z_{K^p}^{\text{pr}}(\Lambda)$ and (2.8). Part (ii) is essentially a consequence of (5.16) and the formal lifting criterion for smoothness of a finitely presented morphism. Here are the details: it is enough to check that p_2 is smooth over the closed points of M_G^{loc} . Suppose therefore that we are given a surjection of $\mathbb{Z}/p^n\mathbb{Z}$ -algebras $\mathcal{O} \rightarrow \bar{\mathcal{O}}$ with square zero kernel. We need to show that the map

$$\varphi : \mathcal{P}_{\text{dR},\Lambda}(\mathcal{O}) \rightarrow \mathcal{P}_{\text{dR},\Lambda}(\bar{\mathcal{O}}) \times_{M_G^{\text{loc}}(\bar{\mathcal{O}})} M_G^{\text{loc}}(\mathcal{O}) \tag{6.20.1}$$

is surjective. So, suppose that we have a pair $(\bar{x}, \bar{\xi}) \in \mathcal{P}_{\text{dR},\Lambda}(\bar{\mathcal{O}})$. We obtain an isotropic line

$$F^1 L_{\bar{\mathcal{O}}} = \bar{\xi}^{-1}(F^1 \tilde{\mathbf{L}}_{\text{dR},\bar{x}}) \subset L_{\bar{\mathcal{O}}}.$$

Let $\tilde{\mathbf{H}}_{\mathcal{O}}$ is the evaluation of the crystal $\tilde{\mathbf{H}}_{\text{cris}}$ on $\text{Spec } \bar{\mathcal{O}} \hookrightarrow \text{Spec } \mathcal{O}$. As usual, if $\tilde{\mathbf{L}}_{\mathcal{O}}$ is the evaluation of the crystal $\tilde{\mathbf{L}}_{\text{cris}}$ along the same thickening, we obtain an embedding $\tilde{\mathbf{L}}_{\mathcal{O}} \subset \text{End}(\tilde{\mathbf{H}}_{\mathcal{O}})$. We also have the crystalline realization $\nu_{\mathcal{O}} : \Lambda \otimes \mathcal{O} \hookrightarrow \tilde{\mathbf{L}}_{\mathcal{O}}$ of ν ; write $\nu_{\bar{\mathcal{O}}}$ for its change of scalars along $\mathcal{O} \rightarrow \bar{\mathcal{O}}$.

There is a canonical isomorphism $\tilde{\mathbf{H}}_{\text{dR},\bar{x}} \xrightarrow{\sim} \tilde{\mathbf{H}}_{\mathcal{O}} \otimes_{\mathcal{O}} \bar{\mathcal{O}}$. Composing this with the trivialization $\bar{\xi}$ gives us a \tilde{G} -structure preserving isomorphism $\xi_{\bar{\mathcal{O}}} : \tilde{H} \otimes \bar{\mathcal{O}} \xrightarrow{\sim} \tilde{\mathbf{H}}_{\bar{\mathcal{O}}}$ carrying $\nu_0 \otimes 1$ to $\nu_{\bar{\mathcal{O}}}$.

Suppose now that we are also given $F^1 L_{\mathcal{O}} \in \text{M}_G^{\text{loc}}(\mathcal{O})$ lifting $F^1 L_{\bar{\mathcal{O}}}$. Let $\mathcal{P}(\mathcal{O} \rightarrow \bar{\mathcal{O}})$ be the set of \tilde{G} -structure preserving isomorphisms $\xi_{\mathcal{O}} : \tilde{H} \otimes \mathcal{O} \xrightarrow{\sim} \tilde{\mathbf{H}}_{\mathcal{O}}$ carrying $\nu_0 \otimes 1$ to $\nu_{\mathcal{O}}$, and lifting $\xi_{\bar{\mathcal{O}}}$.

Observe now that this set is non-empty: we can lift \bar{x} to any \mathcal{O} -valued point $x' \in \mathcal{S}_{\tilde{K}}(\mathcal{O})$. This gives a canonical identification $\tilde{\mathbf{H}}_{\text{dR},x'} \xrightarrow{\sim} \tilde{\mathbf{H}}_{\bar{\mathcal{O}}}$. Since $\tilde{\mathcal{P}}_{\text{dR}}$ is a \tilde{G} -torsor over $\mathcal{S}_{\tilde{K}}$, there now exists a \tilde{G} -structure preserving trivialization $\xi'_{\mathcal{O}} : \tilde{H} \otimes \mathcal{O} \xrightarrow{\sim} \tilde{\mathbf{H}}_{\mathcal{O}}$ lifting $\xi_{\bar{\mathcal{O}}}$. If $\xi'_{\mathcal{O}}$ carries $\nu_0 \otimes 1$ to $\nu_{\mathcal{O}}$, we are done. Otherwise, it follows from (2.8) that we can compose it with an element of $\tilde{G}(\mathcal{O})$ to ensure that it lies in $\mathcal{P}(\mathcal{O} \rightarrow \bar{\mathcal{O}})$. In fact, *loc. cit.* implies that $\mathcal{P}(\mathcal{O} \rightarrow \bar{\mathcal{O}})$ is a non-empty torsor under $\ker(G(\mathcal{O}) \rightarrow G(\bar{\mathcal{O}}))$.

We claim that $\mathcal{P}(\mathcal{O} \rightarrow \bar{\mathcal{O}})$ is in canonical bijection with the fiber $\varphi^{-1}((\bar{x}, \bar{\xi}), F^1 L_{\mathcal{O}})$. This will show that φ is smooth. If we set $\mathcal{O} = k[\epsilon]$ and $\bar{\mathcal{O}} = k$, we find that the fiber of φ over any k -valued point is a torsor under $\ker(G(k[\epsilon]) \rightarrow G(k)) = \text{Lie } G \otimes k$. So, our claim would also show that φ has relative dimension $\dim G_{\mathbb{Q}}$.

Let us prove the claim: given such any $\xi_{\mathcal{O}} \in \mathcal{P}(\mathcal{O} \rightarrow \bar{\mathcal{O}})$, $\xi_{\mathcal{O}}(F^1 L_{\mathcal{O}}) \subset \tilde{\mathbf{L}}_{\mathcal{O}}$ is an isotropic line lifting $F^1 \tilde{\mathbf{L}}_{\text{dR},\bar{x}}$ and isotropic to the image of $\nu_{\mathcal{O}}$. By (5.16), this determines a lift $x \in Z_{K^p}^{\text{pr}}(\Lambda)(\mathcal{O})$ of \bar{x} . Furthermore, there exists a canonical isomorphism $\tilde{\mathbf{H}}_{\mathcal{O}} \xrightarrow{\sim} \tilde{\mathbf{H}}_{\text{dR},x}$ carrying $F^1 \tilde{\mathbf{L}}_{\mathcal{O}}$ onto $F^1 \tilde{\mathbf{L}}_{\text{dR},x}$. The composition ξ of this isomorphism with $\xi_{\mathcal{O}}$ gives us a lift $(x, \xi) \in \varphi^{-1}((\bar{x}, \bar{\xi}), F^1 L_{\mathcal{O}})$. Using the bijectivity of the correspondence in (5.16), one can check that the assignment defined in this fashion is in fact a bijection from $\mathcal{P}(\mathcal{O} \rightarrow \bar{\mathcal{O}})$ to $\varphi^{-1}((\bar{x}, \bar{\xi}), F^1 L_{\mathcal{O}})$.

For (iii), it suffices to prove the result in a neighborhood of a closed point $x \in Z_{K^p}^{\text{pr}}(\Lambda)(\bar{\mathbb{F}}_p)$. Fix any section $s_0 : \text{Spec } \bar{\mathbb{F}}_p \rightarrow \mathcal{P}_{\text{dR},\Lambda}$ over x , and let $\xi_0 : \tilde{H} \otimes \bar{\mathbb{F}}_p \xrightarrow{\sim} \tilde{\mathbf{H}}_{\text{dR},x}$ be the corresponding \tilde{G} -structure preserving isomorphism. The point $y = p_2(s_0(x)) \in \text{M}_G^{\text{loc}}(\bar{\mathbb{F}}_p)$ corresponds to the isotropic line $\xi_0^{-1}(F^1 \tilde{\mathbf{L}}_{\text{dR},x}) \subset L \otimes \bar{\mathbb{F}}_p$.

Let $T = \text{Spec } R$ (respectively $T' = \text{Spec } R'$) be the henselization of $Z_{K^p}^{\text{pr}}(\Lambda)$ at x (respectively M_G^{loc} at y). It is enough to show that there exists a section $s : T \rightarrow \mathcal{P}_{\text{dR},\Lambda}$ lifting s_0 such that the induced map $p_2 \circ s : T \rightarrow T'$ is an isomorphism.

Let $\mathfrak{m}_R \subset R$ (respectively $\mathfrak{m}_{R'} \subset R'$) be the maximal ideal, and set $R_1 = R/(\mathfrak{m}_R^2 + (p))$ (respectively $R'_1 = R'/(\mathfrak{m}_{R'}^2 + (p))$). Also, set $T_1 = \text{Spec } R_1$ and $T'_1 = \text{Spec } R'_1$.

By [RZ96, 3.33], it is enough to find a section s as above such that the induced map $T_1 \rightarrow T'_1$ is an isomorphism.¹⁰

By Hensel’s lemma, it now suffices to find a section $s_1 : T_1 \rightarrow \mathcal{P}_{\text{dR},\Lambda}$ lifting s_0 such that the induced map $p_2 \circ s_1 : T_1 \rightarrow T'_1$ is an isomorphism.

Composing the isomorphism ξ_0 with the obvious identification $\tilde{\mathbf{H}}_{\text{dR},x} \otimes_k R_1 = \tilde{\mathbf{H}}_{\text{dR},R_1}$ gives

¹⁰ In the language of [RZ96], such a section is ‘rigid of the first order’.

us a \tilde{G} -structure preserving isomorphism

$$\xi_1 : \tilde{H} \otimes R_1 \xrightarrow{\cong} \tilde{\mathbf{H}}_{\mathrm{dR}, R_1},$$

which corresponds to a section $s_1 : T_1 \rightarrow \mathcal{P}_{\mathrm{dR}, \Lambda}$ lifting s_0 . It now follows from (5.16) that the induced map $q \circ s_1 : T_1 \rightarrow T'_1$ is an isomorphism. \square

Remark 6.21. The proposition gives us a *local model diagram* in the terminology of [RZ96, DP94, Pap00].

COROLLARY 6.22. $Z_{K^p}^{\mathrm{pr}}(\Lambda)$, and hence $Z_{K_0^p}^{\mathrm{pr}}(\Lambda)$, is lci and flat of relative dimension n over $\mathbb{Z}_{(p)}$. Moreover, $Z_{K^p}^{\mathrm{pr}}(\Lambda)_{\mathbb{F}_p}$ is reduced if $n \geq t$, where t is the dimension of the radical of $L_{\mathbb{F}_p}$. It is normal if $n \geq t + 1$, and is smooth if $t = 0$. In particular, if $n \geq t$, then $Z_{K^p}^{\mathrm{pr}}(\Lambda)$ is normal.

Proof. This follows from (2.11) and (6.20). \square

COROLLARY 6.23. Suppose that L is maximal with $t \leq 1$. Then $Z_{K^p}(\Lambda)$ (and hence $Z_{K_0^p}(\Lambda)$) is regular and locally healthy.

Proof. This is clear from (2.16) and (6.20). \square

6.24 Assume now that L is maximal with $t = 2$. As in §2.17, we will fix a quadratic extension F/\mathbb{Q} in which p is inert, but we will place the additional constraint that F be real.

Fix a self-dual $\mathcal{O}_{F,(p)}$ -lattice $L^\diamond \subset F \otimes L$ containing $\mathcal{O}_{F,(p)} \otimes L$. By *loc. cit.*, this gives us a proper G -equivariant map $M_G^{\mathrm{ref}} \rightarrow M_G^{\mathrm{loc}}$, whose source is regular and locally healthy.

PROPOSITION 6.25. There exist an algebraic space $Z_{K^p}^{\mathrm{ref}}(\Lambda)$ over $\mathbb{Z}_{(p)}$ and a proper morphism $Z_{K^p}^{\mathrm{ref}}(\Lambda) \rightarrow Z_{K^p}^{\mathrm{pr}}(\Lambda)$, determined uniquely up to unique isomorphism, with the following properties.

- (i) There exists a G -equivariant isomorphism

$$\mathcal{P}_{\mathrm{dR}, \Lambda} \times_{Z_{K^p}^{\mathrm{pr}}(\Lambda)} Z_{K^p}^{\mathrm{ref}}(\Lambda) \xrightarrow{\cong} \mathcal{P}_{\mathrm{dR}, \Lambda} \times_{M_G^{\mathrm{loc}}} M_G^{\mathrm{ref}}.$$

Here, the G -action on the left-hand side is via its action on $\mathcal{P}_{\mathrm{dR}, \Lambda}$, and the action on the right is the diagonal action.

- (ii) Every point of $Z_{K^p}^{\mathrm{ref}}(\Lambda)$ has an étale neighborhood $U \rightarrow Z_{K^p}^{\mathrm{ref}}(\Lambda)$ equipped with a section $s : U \rightarrow \mathcal{P}_{\mathrm{dR}, \Lambda} \times_{Z_{K^p}^{\mathrm{pr}}(\Lambda)} Z_{K^p}^{\mathrm{ref}}(\Lambda)$ such that the composition

$$U \xrightarrow{s} \mathcal{P}_{\mathrm{dR}, \Lambda} \times_{Z_{K^p}^{\mathrm{pr}}(\Lambda)} Z_{K^p}^{\mathrm{ref}}(\Lambda) \xrightarrow{\cong} \mathcal{P}_{\mathrm{dR}, \Lambda} \times_{M_G^{\mathrm{loc}}} M_G^{\mathrm{ref}} \rightarrow M_G^{\mathrm{ref}}$$

is étale.

In particular, $Z_{K^p}^{\mathrm{ref}}(\Lambda)$ is regular and locally healthy, and the map $Z_{K^p}^{\mathrm{ref}}(\Lambda) \rightarrow Z_{K^p}^{\mathrm{pr}}(\Lambda)$ is an isomorphism over the regular locus of the target.

Proof. This is a consequence of (6.20) and [Pap00, Proposition 2.4]. The statement of the cited result does not apply directly in our setting, but it is easily seen that its *proof* does. \square

Remark 6.26. The morphism $Z_{K^p}^{\mathrm{ref}}(\Lambda) \rightarrow Z_{K^p}^{\mathrm{pr}}(\Lambda)$ is a *linear modification* in the sense of [Pap00, 2.6]. Although our construction does not fit strictly within the framework of *loc. cit.*, it is inspired by obvious analogy.

6.27 We will continue with the assumption that $t = 2$. As explained in the introduction, in this situation, $Z_{K^p}^{\text{pr}}(\Lambda)$ is strictly contained in $Z_{K^p}(\Lambda)$. In other words, there exist points in $Z_{K^p}(\Lambda)(\overline{\mathbb{F}}_p)$ lying over $s_0 \in \mathcal{S}_{\tilde{K}}(\overline{\mathbb{F}}_p)$ such that the image of the associated map $\iota_0 : \Lambda \rightarrow L(\tilde{A}_{s_0}^{\text{KS}})$ does not generate a saturated sub-space of $\tilde{\mathbf{L}}_{\text{cris},s_0}$.

To understand this phenomenon, it will be helpful to have a better handle on the structure of $\tilde{\mathbf{L}}_{\text{cris},s_0}$. In the language of [Ogu79, §3], this is (up to twist) a $K3$ crystal of rank $\tilde{n} + 2$: that is, it is strongly divisible in the sense of §4.8, and it carries an \mathbf{F}_{s_0} -invariant self-dual quadratic form.

The structure of such objects is well understood. For simplicity, set $\tilde{\mathbf{L}} = \tilde{\mathbf{L}}_{\text{cris},s_0}$ and $\tilde{L}_p(s_0) = \tilde{\mathbf{L}}^{\mathbf{F}_{s_0}=1}$. One associates with $\tilde{\mathbf{L}}$ a Hodge polygon and a Newton polygon; cf. [Kat79, (1.2) and (1.3)]. Essentially, the Hodge polygon encodes information about the Hodge filtration on $\tilde{\mathbf{L}}_{\text{dR},s_0}$ and the Newton polygon encodes information about the slopes of \mathbf{F}_{s_0} . Both polygons are convex by construction and, by a theorem of Mazur [Kat79, Theorem 1.4.1], they have the same end points and the Newton polygon lies above the Hodge polygon.

In our situation, the Hodge polygon is very simple: it begins at $(0, 0)$ and ends at $(\tilde{n} + 2, 0)$, and the slope-0 segment has length \tilde{n} . This reflects the fact that $F^1\tilde{\mathbf{L}}_{\text{dR},s_0}$ has rank 1 and that $\text{gr}_0^F \tilde{\mathbf{L}}_{\text{dR},s_0}$ has rank \tilde{n} .

There are now two possibilities for the Newton polygon, with different flavors.

- The Newton polygon is non-constant: in this case, it admits a break at x -coordinate h , where h is an integer between 1 and $\lfloor(\tilde{n} + 2)/2\rfloor$. We say then that $\tilde{\mathbf{L}}$ has *finite height* h . By a theorem of Katz [Kat79, 1.6.1], $\tilde{\mathbf{L}}$ admits a Newton–Hodge decomposition

$$\tilde{\mathbf{L}} = \tilde{\mathbf{L}}_{-h} \oplus \tilde{\mathbf{L}}_0 \oplus \tilde{\mathbf{L}}_h,$$

where each of the summands is \mathbf{F}_{s_0} -stable after inverting p . The summand $\tilde{\mathbf{L}}_{-h}$ has rank h and the quadratic form on $\tilde{\mathbf{L}}$ induces a perfect pairing between $\tilde{\mathbf{L}}_{-h}$ and $\tilde{\mathbf{L}}_h$. The summand $\tilde{\mathbf{L}}_0$ is a unit root F -crystal (that is, it is generated as a W -module by its \mathbf{F}_{s_0} -invariant elements) orthogonal to $\tilde{\mathbf{L}}_{-h} \oplus \tilde{\mathbf{L}}_h$.

We have $\tilde{L}_p(s_0) = \tilde{\mathbf{L}}_0^{\mathbf{F}_{s_0}=1}$, and $W \otimes_{\mathbb{Z}_p} \tilde{L}_p(s_0) = \tilde{\mathbf{L}}_0$ is a direct summand of $\tilde{\mathbf{L}}$. In particular, any point of $Z_{K^p}(\Lambda)$ lying above s_0 will be in $Z_{K^p}^{\text{pr}}(\Lambda)$.

- The Newton polygon has constant slope 0: in this case, we say that $\tilde{\mathbf{L}}$ is *supersingular*. By a result of Ogus [Ogu79, Theorem 3.4], $\tilde{\mathbf{L}}$ admits an orthogonal decomposition

$$\tilde{\mathbf{L}} = \tilde{\mathbf{L}}_1 \perp \tilde{\mathbf{L}}_0,$$

which is again \mathbf{F}_{s_0} -stable after inverting p . In this decomposition, $\tilde{\mathbf{L}}_0$ is a unit root F -crystal, which is self-dual with respect to the quadratic form. Moreover, $\tilde{L}_p(s_0)$ is a free module of rank $\tilde{n} + 2$ over \mathbb{Z}_p , and it inherits an orthogonal decomposition

$$\tilde{L}_p(s_0) = \tilde{L}_1 \perp \tilde{L}_0,$$

where $\tilde{L}_i = \tilde{\mathbf{L}}_i^{\mathbf{F}_{s_0}=1}$ for $i = 0, 1$. The quadratic form on \tilde{L}_1 is p -times a self-dual form.

We can say more: the sub-space $W \otimes_{\mathbb{Z}_p} \tilde{L}_p(s_0) \subset \tilde{\mathbf{L}}$ has index p^σ , where $2\sigma = \text{rank } \tilde{L}_1$, and the discriminant of $\tilde{L}_p(s_0)$ has p -adic valuation σ . The integer σ , which lies between 1 and $\lfloor(\tilde{n} + 2)/2\rfloor$, is the *Artin invariant* of $\tilde{\mathbf{L}}$.

Now, since Λ^\vee/Λ is isomorphic to $\mathbb{Z}/p\mathbb{Z} \oplus \mathbb{Z}/p\mathbb{Z}$, we can find an orthogonal decomposition $\Lambda = \Lambda_1 \perp \Lambda_0$, where Λ_0 is self-dual, Λ_1 has rank 2 and the restriction of the quadratic form

to Λ_1 is divisible by p . If $2\sigma \geq \tilde{n}$, then one can check that there can never be an isometric embedding $\Lambda_1 \hookrightarrow \tilde{L}_p(s_0)$ whose image generates a direct summand of \tilde{L} . Therefore, any point of $Z_{K^p}(\Lambda)$ lying above s_0 will be outside of $Z_{K^p}^{\text{pr}}(\Lambda)$.

6.28 Consider the special case where $\tilde{n} = 3$, $d = 2$ and \tilde{L} admits a maximal isotropic sub-space of rank 2: in this situation, $\mathcal{S}_{\tilde{K}}$ is a moduli space of polarized abelian surfaces (the Kuga–Satake abelian scheme \tilde{A}^{KS} is isomorphic to a power of the universal abelian surface), and $Z_{K^p}(\Lambda)$ is a union of integral models for compact Shimura curves with non-maximal parahoric level structure. The Artin invariant σ is either 1 or 2: it is 1 when $\tilde{A}_{s_0}^{\text{KS}}$ is isomorphic to a product of supersingular elliptic curves (the *superspecial* case), and it is 2 if $\tilde{A}_{s_0}^{\text{KS}}$ is supersingular, but not superspecial, which should be the ‘generic’ situation on $Z_{K^p}(\Lambda)$. The above discussion shows that the open locus $Z_{K^p}^{\text{pr}}(\Lambda)$ will not contain any of the non-superspecial points, and so misses out on what should be the most interesting part of the geometry of $Z_{K^p}(\Lambda)$. Similar caveats apply in higher dimensions. We intend to return to the general question of the structure of $Z_{K^p}(\Lambda)$ in future work.

7. Integral canonical models II

The notation will be as in the previous section. Recall from § 6.15 that we have canonical maps $\text{Sh}_{K_0} \rightarrow Z_{K_0^p}(\Lambda)_{\mathbb{Q}}$ and $\text{Sh}_K \rightarrow Z_{K^p}(\Lambda)_{\mathbb{Q}}$.

LEMMA 7.1. *Both of these maps are isomorphisms onto open and closed sub-schemes of the targets.*

Proof. By construction, it is enough to prove this for $\text{Sh}_K \rightarrow Z_{K^p}(\Lambda)_{\mathbb{Q}}$. Since both varieties in question are smooth of dimension n and are unramified over $\text{Sh}_{\tilde{K}}$, it is enough to show that the map is injective on \mathbb{C} -valued points.

Pick a point $x \in \text{Sh}_K(\mathbb{C})$ and let $(\mathbf{h}, g) \in X \times G(\mathbb{A}_f^p)$ be a lift of x . We can describe the corresponding point of $Z_{K^p}(\Lambda)(\mathbb{C})$ as follows: first, we can view (\mathbf{h}, g) as a point in $\tilde{X} \times \tilde{G}(\mathbb{A}_f^p)$; write \tilde{x} for its image in $\text{Sh}_{\tilde{K}}(\mathbb{C})$. Then \mathbf{h} induces a Hodge structure $\tilde{H}_{\mathbf{h}}$ on \tilde{H} , which preserves $\tilde{L} \subset \tilde{H}^{\otimes(1,1)}$ and is such that the induced Hodge structure on $\Lambda \subset \tilde{L}$ is trivial. There is a \tilde{G} -structure preserving isomorphism of Hodge structures $j_x : \tilde{H}_{\mathbf{h}} \xrightarrow{\sim} \tilde{H}_{B, \tilde{x}}$ and the composition

$$\iota_x : \Lambda \hookrightarrow \tilde{L}_{\mathbf{h}} \cap (\tilde{L}_{\mathbf{h}} \otimes \mathbb{C})^{0,0} \xrightarrow{j_x} \tilde{L}_{B, \tilde{x}} \cap (\tilde{L}_{B, \tilde{x}} \otimes \mathbb{C})^{0,0} = L(\tilde{A}_{\tilde{x}}^{\text{KS}}) \tag{7.1.1}$$

is the Λ -structure attached to x . Further, the K^p -level structure on (\tilde{x}, ι_x) attached to x is the K^p -orbit $[\tilde{\eta}_x]$ of the isomorphism

$$\tilde{\eta}_x : \tilde{H} \otimes \mathbb{A}_f^p \xrightarrow[g]{\simeq} \tilde{H} \otimes \mathbb{A}_f^p \xrightarrow[\simeq]{j_x \otimes 1} \tilde{H}_{B, \tilde{x}} \otimes \mathbb{A}_f^p \xrightarrow[\simeq]{} \tilde{H}_{\mathbb{A}_f^p, \tilde{x}}. \tag{7.1.2}$$

Note that the triple $(\tilde{x}, \iota_x, [\tilde{\eta}_x])$ determines the image of x in $Z_{K^p}(\Lambda)(\mathbb{C})$.

Suppose that $x' \in \text{Sh}_K(\mathbb{C})$ lifts to $(\mathbf{h}', g') \in X \times G(\mathbb{A}_f^p)$ and is such that $(\tilde{x}', \iota_{x'}, [\tilde{\eta}_{x'}]) = (\tilde{x}', \iota_{x'}, [\tilde{\eta}_{x'}])$. Then, using (7.1.1), we find that $\mathbf{h}' = \gamma \cdot \mathbf{h}$, where $\gamma = j_{x'}^{-1} \circ j_x \in G(\mathbb{Z}_{(p)})$. Also, using (7.1.2), we see that γg and g' are in the same K^p -orbit in $G(\mathbb{A}_f^p)$. This shows that $x' = x$ and finishes the proof of the lemma. \square

DEFINITION 7.2. Let $\mathcal{S}_K^{\text{pr}}$ be the Zariski closure of Sh_K in $Z_{K^p}^{\text{pr}}(\Lambda)$. If L is maximal with $t \leq 1$, set $\mathcal{S}_K := \mathcal{S}_K^{\text{pr}}$. If L is maximal and $t = 2$, fix F and L^\diamond as in § 6.24 and let $\mathcal{S}_K^{\text{ref}}$ be the proper resolution of $\mathcal{S}_K^{\text{pr}}$ obtained by taking the Zariski closure of Sh_K in $Z_{K^p}^{\text{ref}}(\Lambda)$.

7.3 We will now assume that L is maximal with $t \leq 1$.

We set

$$\mathcal{S}_{K_p} = \varprojlim_{K^p \subset \tilde{K}^p} \mathcal{S}_{K_p K^p}.$$

In this inverse system, for $K_1^p \subset K_2^p$, the map $\mathcal{S}_{K_p K_1^p} \rightarrow \mathcal{S}_{K_p K_2^p}$ is induced by the obvious map $Z_{K_1^p}(\Lambda) \rightarrow Z_{K_2^p}(\Lambda)$. Then \mathcal{S}_{K_p} is actually the projective limit of quasi-projective $\mathbb{Z}_{(p)}$ -schemes along finite étale morphisms, and so is itself a scheme over $\mathbb{Z}_{(p)}$.

THEOREM 7.4.

- (i) \mathcal{S}_{K_p} (respectively $\mathcal{S}_{K_{0,p}}$) is an integral canonical model for the pro-Shimura variety Sh_{K_p} (respectively $\text{Sh}_{K_{0,p}}$). The map $\mathcal{S}_{K_p} \rightarrow \mathcal{S}_{K_{0,p}}$ is a pro-étale cover.
- (ii) For any neat compact open $K \subset G(\mathbb{A}_f^p)$ with p -primary part K_p , there exist a projective regular $\mathbb{Z}_{(p)}$ -scheme $\overline{\mathcal{S}}_K$ and a fiberwise dense open immersion $\mathcal{S}_K \hookrightarrow \overline{\mathcal{S}}_K$. Moreover, $\overline{\mathcal{S}}_K$ has reduced special fiber.

Proof. By (7.1), (6.23) and (6.25), \mathcal{S}_{K_p} is regular and locally healthy. We need to show that it has the extension property. We already know that $\mathcal{S}_{\tilde{K}_p}$ has the extension property by [Kis10, 2.3.8]. So, using [FC90, I.2.7], we find that \mathcal{S}_{K_p} also has the extension property. If $t \leq 1$, we are now done.

Now, let $\mathcal{S}_{K_{0,p}}$ be the Zariski closure of $\text{Sh}_{K_{0,p}}$ in

$$\varprojlim_{K_0^p \subset G_0(\mathbb{A}_f^p)} Z_{K_0^p}(\Lambda).$$

Then there is a pro-finite, pro-étale map $\mathcal{S}_{K_p} \rightarrow \mathcal{S}_{K_{0,p}}$.

To show that $\mathcal{S}_{K_{0,p}}$ has the extension property, we can use the argument from the proof of [Moo98, 3.21.4]: given a regular, locally healthy $\mathbb{Z}_{(p)}$ -scheme S_0 and a map $f_0 : S_{0,\mathbb{Q}} \rightarrow \text{Sh}_{K_{0,p}}$, it follows from the Nagata–Zariski purity theorem that there exist a pro-finite pro-étale cover S of S_0 and a map $f : S_{\mathbb{Q}} \rightarrow \text{Sh}_{K_p}$ lifting f_0 . Now, S is also regular and locally healthy, so, by the extension property of \mathcal{S}_{K_p} , f extends to a map $S \rightarrow \mathcal{S}_{K_p}$ that descends to a map $S_0 \rightarrow \mathcal{S}_{K_{0,p}}$.

The existence of the regular projective compactification $\overline{\mathcal{S}}_K$ is a special case of [MaP15, Theorem 1], which says the following: suppose that we have a symplectic embedding $(G_{\mathbb{Q}}, X) \hookrightarrow (\text{GSp}(U), \mathcal{X}(U))$ of Shimura data, and suppose that $U_{(p)} \subset U$ is a symplectic lattice such that $G(\mathbb{Z}_p)$ is the stabilizer of $U_{(p)} \otimes \mathbb{Z}_p$ in $G(\mathbb{Q}_p)$. Fix $\mathcal{K}(U) \subset \text{GSp}(U)(\mathbb{A}_f)$ such that K maps to $\mathcal{K}(U)$. Let $\mathcal{S}_{\mathcal{K}(U)}$ be the associated Mumford integral model over $\mathbb{Z}_{(p)}$ for the Siegel Shimura variety $\text{Sh}_{\mathcal{K}(U)}$.¹¹ Let \mathcal{S} be the unique normal integral model for Sh_K over $\mathbb{Z}_{(p)}$ such that $\text{Sh}_K \rightarrow \text{Sh}_{\mathcal{K}(U)}$ extends to a finite map $\mathcal{S} \rightarrow \mathcal{S}_{\mathcal{K}(U)}$: it does not depend on the choice of $\mathcal{K}(U)$. Then \mathcal{S} admits a fiberwise dense open immersion $\mathcal{S} \hookrightarrow \overline{\mathcal{S}}$, where $\overline{\mathcal{S}}$ is projective over $\mathbb{Z}_{(p)}$ and with singularities no worse than those of \mathcal{S} . In particular, $\overline{\mathcal{S}}$ is regular (respectively has reduced special fiber) whenever \mathcal{S} is regular (respectively has reduced special fiber).

The required hypothesis holds for the model \mathcal{S}_K in our situation: we take $U_{(p)} = H$.

To finish, it is now enough to show that \mathcal{S}_K has reduced special fiber. This follows from (6.22). □

Remark 7.5. In particular, \mathcal{S}_{K_p} and $\mathcal{S}_{K_{0,p}}$ are uniquely determined by their generic fibers: they depend only on L and not on the choice of self-dual quadratic space \tilde{L} containing L .

¹¹ This is defined as a moduli space of polarized abelian schemes as in §3.7.

COROLLARY 7.6. *Suppose that $n \geq t$; then, for any neat compact open sub-group $K \subset G(\mathbb{A}_f^p)$ with p -primary part K_p , there is a canonical bijection between the connected components of $\text{Sh}_{K, \overline{\mathbb{Q}}}$ (respectively $\text{Sh}_{K_0, \overline{\mathbb{Q}}}$) and $\mathcal{S}_{K, \overline{\mathbb{F}}_p}$ (respectively $\mathcal{S}_{K_0, \overline{\mathbb{F}}_p}$).*

Proof. Since \mathcal{S}_{K_0} is a finite étale quotient of \mathcal{S}_K , the result for the latter implies that for the former. For \mathcal{S}_K , it is enough to check that the components of $\overline{\mathcal{S}}_{K, \overline{\mathbb{Q}}}$ and $\overline{\mathcal{S}}_{K, \overline{\mathbb{F}}_p}$ are in bijection.

But, in general, given any flat, projective map $X \rightarrow \text{Spec } \mathbb{Z}_{(p)}$ with reduced special fiber, there is a canonical bijection between the components of $X_{\overline{\mathbb{Q}}}$ and $X_{\overline{\mathbb{F}}_p}$. Indeed, using Stein factorization, we reduce to the case where X is finite flat over $\mathbb{Z}_{(p)}$. Having reduced special fiber implies that X is actually finite étale over $\mathbb{Z}_{(p)}$ and so the result is immediate. \square

For applications, we will need a different kind of canonical model that makes sense in some non-maximal cases.

DEFINITION 7.7. A pro-scheme X over $\mathbb{Z}_{(p)}$ satisfies the *smooth extension property* if, for any regular, formally smooth $\mathbb{Z}_{(p)}$ -scheme S , any map $S \otimes \mathbb{Q} \rightarrow X$ extends to a map $S \rightarrow X$.

Suppose that L is non-maximal with cyclic discriminant. We still have the pro-Shimura variety Sh_{K_p} over \mathbb{Q} , where $K_p = G(\mathbb{Z}_p)$, with G the smooth group scheme attached to L .

DEFINITION 7.8. In analogy with (4.3), we will define a *smooth integral canonical model* of Sh_{K_p} to be a regular, formally smooth model $\mathcal{S}_{K_p}^{\text{sm}}$ (respectively $\mathcal{S}_{K_0, p}^{\text{sm}}$) which has the smooth extension property.

PROPOSITION 7.9. *The pro-Shimura variety Sh_{K_p} (respectively $\text{Sh}_{K_0, p}$) admits a smooth canonical model $\mathcal{S}_{K_p}^{\text{sm}}$ (respectively $\mathcal{S}_{K_0, p}^{\text{sm}}$) such that the map $\mathcal{S}_{K_p}^{\text{sm}} \rightarrow \mathcal{S}_{K_0, p}^{\text{sm}}$ is pro-étale.*

Proof. By (5.21)(ii), we find that, for any level sub-group $K^p \subset G(\mathbb{A}_f^p)$ contained in \tilde{K}^p , the $\mathcal{S}_{\tilde{K}^p, \tilde{K}^p}$ -scheme $Z_{K^p}^{\text{pr}}(\Lambda)$ is smooth at all of its $\overline{\mathbb{F}}_p$ -valued points. In particular, $\mathcal{S}_{K^p, K^p}^{\text{sm}} := \mathcal{S}_{K^p, K^p}^{\text{pr}}$ must be smooth over $\mathbb{Z}_{(p)}$. Now set

$$\mathcal{S}_{K_p}^{\text{sm}} := \varprojlim_{K^p \subset \tilde{K}^p} \mathcal{S}_{K^p, K^p}^{\text{sm}}.$$

It remains to show that $\mathcal{S}_{K_p}^{\text{sm}}$ has the smooth extension property. Since $\mathcal{S}_{\tilde{K}^p}$ has the extension property, by [FC90, I.2.7], the inverse limit

$$\varprojlim_{K^p \subset G(\mathbb{A}_f^p)} Z_{K^p}(\Lambda)$$

also has the extension property.

It is therefore sufficient to make the following observation, which is immediate from (6.16)(iv): suppose that we are given a map $x : S \rightarrow Z_{K^p}(\Lambda)$, with S smooth over $\mathbb{Z}_{(p)}$. Suppose also that the restriction of x to $S_{\mathbb{Q}}$ factors through Sh_{K_p, K^p} . Then x must necessarily factor through $Z_{K^p}^{\text{pr}}(\Lambda)$.

The construction of $\mathcal{S}_{K_0, p}^{\text{sm}}$ from $\mathcal{S}_{K_0, p}^{\text{sm}}$ proceeds as in the maximal case. \square

7.10 Fix $K^p \subset G(\mathbb{A}_f^p)$ sufficiently small with image $K_0^p \subset G_0(\mathbb{A}_f^p)$. Set $K = K_p K^p$ and $K_0 = K_{0, p} K_0^p$ and write $\mathcal{S} = \mathcal{S}_K$ (respectively $\mathcal{S}_0 = \mathcal{S}_{K_0}$) if L is maximal with $t \leq 1$, $\mathcal{S} = \mathcal{S}_K^{\text{ref}}$ (respectively $\mathcal{S}_{K_0}^{\text{ref}}$) if L is maximal with $t = 2$; and $\mathcal{S} = \mathcal{S}_K^{\text{sm}}$ (respectively $\mathcal{S}_0 = \mathcal{S}_{K_0}^{\text{sm}}$) for non-maximal L with cyclic discriminant. All these schemes are regular and locally healthy.

Set $L_{\text{dR}} = \Lambda_{\text{dR}}^\perp \subset \tilde{L}_{\text{dR}}|_{\mathcal{S}}$ and $L_{\text{cris}} = \Lambda_{\text{cris}}^\perp \subset \tilde{L}_{\text{cris}}$. Here, Λ_{cris} is the sub-crystal of $\tilde{L}_{\text{cris}}|_{\mathcal{S}}$ generated by the crystalline realization of $\iota(\Lambda)$.

Suppose that we are given a point $s \in \mathcal{S}(k)$. The evaluation of L_{cris} along $\text{Spec } k \hookrightarrow \text{Spf } W(k)$ is a direct summand $L_{\text{cris},s} = \Lambda_{\text{cris},s}^\perp \subset \tilde{L}_{\text{cris},s}$. Moreover, the F -isocrystal structure on $\tilde{L}_{\text{cris}}[p^{-1}]$ induces one on $L_{\text{cris},s}[p^{-1}]$.

Set $W = W(k)$, let $E/W_{\mathbb{Q}}$ be a finite extension and suppose that $\tilde{s} \in \mathcal{S}(\mathcal{O}_E)$ is a lift of s . Let \bar{E}/E be an algebraic closure and let $\tilde{s}_{\bar{E}}$ be the geometric generic fiber of \tilde{s} . From the definition of the sheaves, (4.7)(ii) and the functoriality of the comparison isomorphism, we obtain the following result.

PROPOSITION 7.11. *There is a canonical isometric comparison isomorphism compatible with additional structures:*

$$L_{p,\tilde{s}_{\bar{E}}} \otimes_{\mathbb{Z}_p} B_{\text{cris}} \xrightarrow{\cong} L_{\text{cris},s} \otimes_W B_{\text{cris}}. \quad \square$$

7.12 Consider the ‘intrinsic’ Kuga–Satake abelian scheme $A_{\text{Sh}_K}^{\text{KS}}$ over Sh_K : we claim that it extends uniquely to an abelian scheme A^{KS} over \mathcal{S} . Indeed, since \mathcal{S} is locally healthy and hence healthy, it suffices to show the following result.

Let $x \rightarrow \mathcal{S}$ be a co-dimension-1 point such that $\mathcal{O}_{\mathcal{S}_K,x}$ is a discrete valuation ring of mixed characteristic $(0,p)$. Set $T = \text{Spec } \mathcal{O}_{\mathcal{S},x}$. Then $A_{T_{\mathbb{Q}}}^{\text{KS}}$ has good reduction over T .

To show this, we first observe that, for $\ell \neq p$, the ℓ -adic cohomology H_ℓ of $A_{\text{Sh}_K}^{\text{KS}}$ is related to \tilde{H}_ℓ via the formula (cf. proof of (6.4))

$$H_\ell \otimes_{C(L)} C(\tilde{L}) \xrightarrow{\cong} \tilde{H}_\ell|_{\text{Sh}_K}.$$

Since $\tilde{A}_{T_{\mathbb{Q}}}^{\text{KS}}$ extends to an abelian scheme over T , the restriction of \tilde{H}_ℓ over $T_{\mathbb{Q}}$ is unramified, which implies that the restriction of H_ℓ to $T_{\mathbb{Q}}$ is also unramified. Our claim now follows from the usual criterion for good reduction of abelian varieties.

7.13 By [FC90, I.2.7], the isomorphism from (6.4) extends to an isomorphism of abelian schemes over \mathcal{S} :

$$i : \tilde{A}_{\mathcal{S}}^{\text{KS}} \xrightarrow{\cong} \underline{\text{Hom}}_C(\tilde{C}, A^{\text{KS}}). \quad (7.13.1)$$

If H_{dR} is the degree-1 de Rham cohomology of A^{KS} over \mathcal{S} , the de Rham realization of i gives us a canonical isomorphism of vector bundles with integrable connection:

$$i_{\text{dR}} : H_{\text{dR}} \otimes_C \tilde{C} \xrightarrow{\cong} \tilde{H}_{\text{dR}}|_{\mathcal{S}}. \quad (7.13.2)$$

Similarly, if H_{cris} is the F -crystal over $(\mathcal{S}_{\mathbb{F}_p}/\mathbb{Z}_p)_{\text{cris}}$ obtained from the degree-1 crystalline cohomology of $A_{\mathbb{F}_p}^{\text{KS}}$, then we have a canonical \tilde{C} -equivariant, $\mathbb{Z}/2\mathbb{Z}$ -graded isomorphism of F -crystals:

$$i_{\text{cris}} : H_{\text{cris}} \otimes_C \tilde{C} \xrightarrow{\cong} \tilde{H}_{\text{cris}}|_{(\mathcal{S}_{\mathbb{F}_p}/\mathbb{Z}_p)_{\text{cris}}}. \quad (7.13.3)$$

It follows from the argument in (2.6) that we have

$$C = \{z^+ + z^- \in \tilde{C}^+ \oplus \tilde{C}^- = \tilde{C} : v \cdot (z^+ + z^-) = (z^+ - z^-) \cdot v, \text{ for all } v \in \Lambda\}. \quad (7.13.4)$$

We can view $\widetilde{\mathbf{H}}_{\text{cris}}^{\otimes(1,1)}$ as the internal endomorphism object $\underline{\text{End}}(\widetilde{\mathbf{H}}_{\text{cris}})$ within the category of crystals over $\mathcal{S}_{\mathbb{F}_p}$. As such, within it we have the sub-object $\underline{\text{End}}_{\widetilde{C}}(\widetilde{\mathbf{H}}_{\text{cris}})$ of \widetilde{C} -equivariant endomorphisms. Similarly, within $\mathbf{H}_{\text{cris}}^{\otimes(1,1)}$ we have the sub-object $\underline{\text{End}}_C(\mathbf{H}_{\text{cris}})$ of C -equivariant endomorphisms. Using (7.13.4) and (7.13.3), we find that we can exhibit $\underline{\text{End}}_C(\mathbf{H}_{\text{cris}})$ as the sub-crystal of $\underline{\text{End}}_{\widetilde{C}}(\widetilde{\mathbf{H}}_{\text{cris}})|_{\mathcal{S}_{\mathbb{F}_p}}$ that consists of endomorphisms that anti-commute with $\mathbf{\Lambda}_{\text{cris}}$. By its definition, \mathbf{L}_{cris} anti-commutes with $\mathbf{\Lambda}_{\text{cris}}$ within $\underline{\text{End}}_{\widetilde{C}}(\widetilde{\mathbf{H}}_{\text{cris}})|_{\mathcal{S}_{\mathbb{F}_p}}$. Therefore, we obtain a commutative diagram of embeddings of crystals as follows.

$$\begin{array}{ccc}
 \mathbf{L}_{\text{cris}} & \hookrightarrow & \mathbf{H}_{\text{cris}}^{\otimes(1,1)} \\
 \downarrow \simeq & & \downarrow \\
 \mathbf{\Lambda}_{\text{cris}}^\perp & \hookrightarrow & \widetilde{\mathbf{L}}_{\text{cris}}|_{\mathcal{S}_{\mathbb{F}_p}} \hookrightarrow \widetilde{\mathbf{H}}_{\text{cris}}^{\otimes(1,1)}|_{\mathcal{S}_{\mathbb{F}_p}}
 \end{array} \tag{7.13.5}$$

7.14 Given (7.13.5), the notions of ℓ -specialness and p -specialness for an endomorphism of A^{KS} carry over verbatim from §5. For any \mathcal{S} -scheme T , denote the space of ℓ -special endomorphisms of A_T^{KS} by $L_\ell(A_T^{\text{KS}})$. Let $L(A_T^{\text{KS}})$ be the space of special endomorphisms, where ‘special’ means ℓ -special for every ℓ . As in (6.5), we obtain the following result.

PROPOSITION 7.15. *For any scheme $T \rightarrow \mathcal{S}$, there exists a canonical isometry*

$$L(A_T^{\text{KS}}) \xrightarrow{\simeq} \iota(\Lambda)^\perp \subset L(\widetilde{A}_T^{\text{KS}})$$

compatible with all cohomological realizations. □

Remark 7.16. Just as in §5.24, the sheaf of endomorphism algebras $\underline{\text{End}}(A^{\text{KS}})_{(p)}$ canonically descends to a sheaf \mathbf{E} over \mathcal{S}_0 . Suppose now that T_0 is a scheme over \mathcal{S}_0 . Then one can canonically attach to T_0 a group of ‘special endomorphisms’ $L(T_0) \subset \mathbf{E}(T_0)$ such that, if T_0 is in fact an \mathcal{S} -scheme, then $L(T_0) = L(A_{T_0}^{\text{KS}})$.

7.17 Given that we have the notion of a special endomorphism of A^{KS} , for any $m \in \mathbb{Z}_{(p)}^{>0}$, just as in §6.12, we can define the finite unramified schemes $Z_{K^p}(m) \rightarrow \mathcal{S}$ and $Z_{K_0^p}(m) \rightarrow \mathcal{S}_0$, which parameterize special endomorphisms of A^{KS} of degree m . From the construction of the schemes \mathcal{S} , (7.15) and (5.17), we find that the deformation theory of a special endomorphism of A^{KS} is locally governed by a single equation. In particular, étale locally on \mathcal{S} , $Z_{K^p}(m)$ is actually an effective Cartier divisor.

PROPOSITION 7.18. *Suppose that $n > 2$. Then the schemes $Z_{K^p}(m)$ and $Z_{K_0^p}(m)$ are flat over $\mathbb{Z}_{(p)}$.*

Proof. It is enough to show that $Z_{K^p}(m)$ is flat over $\mathbb{Z}_{(p)}$. Since, étale locally on \mathcal{S} , $Z_{K^p}(m)$ is an effective Cartier divisor, if it were not $\mathbb{Z}_{(p)}$ -flat, its image in \mathcal{S} would contain an entire component of $\mathcal{S}_{\mathbb{F}_p}$.

On the other hand, it is clear from the description of the local properties of \mathcal{S} via local models that the smooth locus of $\mathcal{S}_{\mathbb{F}_p}$ is a dense open sub-scheme in $\mathcal{S}_{\mathbb{F}_p}$. So, it suffices to show that the restriction of $Z_{K^p}(m)$ to the complement of the non-smooth locus in $\mathcal{S}_{\mathbb{F}_p}$ is flat over $\mathbb{Z}_{(p)}$. This follows directly from (6.18) □

ACKNOWLEDGEMENTS

I thank Fabrizio Andreatta, Ben Howard, Mark Kisin, Ben Moonen, George Pappas and Junecue Suh for valuable comments, insights and corrections. Above all, I owe a deep debt to the referees for their careful reading and their numerous corrections and recommendations that immeasurably improved the paper, in both exposition and content. This work was partially supported by an NSF Postdoctoral Research Fellowship DMS-1204165 and an AMS Simons Travel Grant.

REFERENCES

- BB66 W. L. Baily Jr. and A. Borel, *Compactification of arithmetic quotients of bounded symmetric domains*, Ann. of Math. (2) **84** (1966), 442–528; [MR 0216035](#) (35 #6870).
- Bas74 H. Bass, *Clifford algebras and spinor norms over a commutative ring*, Amer. J. Math. **96** (1974), 156–206; [MR 0360645](#) (50 #13092).
- BM90 P. Berthelot and W. Messing, *Théorie de Dieudonné cristalline. III. Théorèmes d'équivalence et de pleine fidélité*, in *The Grothendieck Festschrift, vol. I*, Progress in Mathematics, vol. 86 (Birkhäuser, Boston, MA, 1990), 173–247; (in French); [MR 1086886](#) (92h:14012).
- Bla94 D. Blasius, *A p -adic property of Hodge classes on abelian varieties*, in *Motives (Seattle, WA, 1991)*, Proceedings of Symposia in Pure Mathematics, vol. 55 (American Mathematical Society, Providence, RI, 1994), 293–308; [MR 1265557](#) (95j:14022).
- Bou07 N. Bourbaki, *Éléments de mathématique. Algèbre* (Springer, Berlin, 2007), ch. 5 (in French); reprint of the 1959 original; [MR 2325344](#) (2008f:15001).
- BT84 F. Bruhat and J. Tits, *Groupes réductifs sur un corps local. II. Schémas en groupes. Existence d'une donnée radicielle valuée*, Publ. Math. Inst. Hautes Études Sci. **60** (1984), 197–376 (in French); [MR 756316](#) (86c:20042).
- CN90 C.-L. Chai and P. Norman, *Bad reduction of the Siegel moduli scheme of genus two with $\Gamma_0(p)$ -level structure*, Amer. J. Math. **112** (1990), 1003–1071, doi:[10.2307/2374734](#); [MR 1081813](#) (91i:14033).
- Del79 P. Deligne, *Variétés de Shimura: interprétation modulaire, et techniques de construction de modèles canoniques*, in *Automorphic Forms, Representations and L-functions (Proc. Symp. Pure Math., Oregon State Univ., Corvallis, OR, 1977), Part 2*, Proceedings of Symposia in Pure Mathematics, vol. XXXIII (American Mathematical Society, Providence, RI, 1979), 247–289 (in French); [MR 546620](#) (81i:10032).
- Del81 P. Deligne, *Relèvement des surfaces K3 en caractéristique nulle*, in *Algebraic surfaces (Orsay, 1976)*, Lecture Notes in Mathematics, vol. 868 (Springer, Berlin, 1981), 58–79 (in French); prepared for publication by Luc Illusie; [MR 638598](#) (83j:14034).
- D MOS82 P. Deligne, J. S. Milne, A. Ogus and K.-Y. Shih, *Hodge Cycles, Motives, and Shimura Varieties*, Lecture Notes in Mathematics, vol. 900 (Springer, Berlin, 1982); [MR 654325](#) (84m:14046).
- DP94 P. Deligne and G. Pappas, *Singularités des espaces de modules de Hilbert, en les caractéristiques divisant le discriminant*, Compositio Math. **90** (1994), 59–79 (in French); [MR 1266495](#) (95a:11041).
- Fal99 G. Faltings, *Integral crystalline cohomology over very ramified valuation rings*, J. Amer. Math. Soc. **12** (1999), 117–144, doi:[10.1090/S0894-0347-99-00273-8](#); [MR 1618483](#) (99e:14022).
- FC90 G. Faltings and C.-L. Chai, *Degeneration of Abelian Varieties*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 22 (Springer, Berlin, 1990), with an appendix by David Mumford; [MR 1083353](#) (92d:14036).
- Gör03 U. Görtz, *On the flatness of local models for the symplectic group*, Adv. Math. **176** (2003), 89–115, doi:[10.1016/S0001-8708\(02\)00062-2](#); [MR 1978342](#) (2004d:14023).

- Har85 M. Harris, *Arithmetic vector bundles and automorphic forms on Shimura varieties. I*, Invent. Math. **82** (1985), 151–189, doi:[10.1007/BF01394784](https://doi.org/10.1007/BF01394784); MR 808114 (88e:11046).
- Har M. Harris, *Arithmetic of the oscillator representation*, www.math.jussieu.fr/~harris/Arithmetictheta.pdf.
- Kat73 N. M. Katz, *p-adic properties of modular schemes and modular forms*, in *Modular functions of one variable, III (Proc. Internat. Summer School, Univ. Antwerp, 1972)*, Lecture Notes in Mathematics, vol. 350 (Springer, Berlin, 1973), 69–190; MR 0447119 (56 #5434).
- Kat79 N. M. Katz, *Slope filtration of F-crystals*, in *Journées de géométrie algébrique de Rennes, (Rennes, 1978)*, Astérisque, vol. 63 (Société Mathématiques de France, Paris, 1979), 113–163; MR 563463 (81i:14014).
- Kat81 N. Katz, *Serre–Tate local moduli*, in *Algebraic surfaces (Orsay, 1976)*, Lecture Notes in Mathematics, vol. 868 (Springer, Berlin, 1981), 138–202; MR 638600 (83k:14039b).
- Kis10 M. Kisin, *Integral models for Shimura varieties of abelian type*, J. Amer. Math. Soc. **23** (2010), 967–1012, doi:[10.1090/S0894-0347-10-00667-3](https://doi.org/10.1090/S0894-0347-10-00667-3); MR 2669706 (2011j:11109).
- Kud97 S. S. Kudla, *Algebraic cycles on Shimura varieties of orthogonal type*, Duke Math. J. **86** (1997), 39–78, doi:[10.1215/S0012-7094-97-08602-6](https://doi.org/10.1215/S0012-7094-97-08602-6); MR 1427845 (98e:11058).
- KR99 S. S. Kudla and M. Rapoport, *Arithmetic Hirzebruch–Zagier cycles*, J. reine angew. Math. **515** (1999), 155–244, doi:[10.1515/crll.1999.076](https://doi.org/10.1515/crll.1999.076); MR 1717613 (2002e:11076a).
- KRY06 S. S. Kudla, M. Rapoport and T. Yang, *Modular Forms and Special Cycles on Shimura Curves*, Annals of Mathematics Studies, vol. 161 (Princeton University Press, Princeton, NJ, 2006); MR 2220359 (2007i:11084).
- Laf80 G. Laffaille, *Groupes p-divisibles et modules filtrés: le cas peu ramifié*, Bull. Soc. Math. France **108** (1980), 187–206 (in French, with English summary); MR 606088 (82i:14028).
- Lan08 K.-W. Lan, *Arithmetic compactifications of PEL-type Shimura varieties*, PhD thesis, Harvard University, ProQuest LLC, Ann Arbor, MI (2008); MR 2711676.
- MaP14 K. Madapusi Pera, *The Tate conjecture for K3 surfaces in odd characteristic*, Invent. Math. **201** (2014), 625–668, doi:[10.1007/s00222-014-0557-5](https://doi.org/10.1007/s00222-014-0557-5).
- MaP15 K. Madapusi Pera, *Toroidal compactifications of integral models of Shimura varieties of Hodge type*, Preprint (2015), pp. 98, available at: <http://www.math.uchicago.edu/~keerthi/papers/toroidal.pdf>.
- MaS11 K. S. Madapusi Sampath, *Toroidal compactifications of integral models of Shimura varieties of Hodge type*, PhD thesis, The University of Chicago, ProQuest LLC, Ann Arbor, MI (2011); MR 2898617.
- Man62 Ju. I. Manin, *On the classification of formal Abelian groups*, Dokl. Akad. Nauk SSSR **144** (1962), 490–492 (in Russian); MR 0162802 (29 #106).
- Mau14 D. Maulik, *Supersingular K3 surfaces for large primes*, Duke Math. J. **163** (2014), 2357–2425, doi:[10.1215/00127094-2804783](https://doi.org/10.1215/00127094-2804783), with an appendix by Andrew Snowden; MR 3265555.
- Mes72 W. Messing, *The Crystals Associated to Barsotti–Tate Groups: with Applications to Abelian Schemes*, Lecture Notes in Mathematics, vol. 264 (Springer, Berlin, 1972); MR 0347836 (50 #337).
- Moo98 B. Moonen, *Models of Shimura varieties in mixed characteristics*, in *Galois Representations in Arithmetic Algebraic Geometry (Durham, 1996)*, London Mathematical Society Lecture Note Series, vol. 254 (Cambridge University Press, Cambridge, 1998), 267–350, doi:[10.1017/CBO9780511662010.008](https://doi.org/10.1017/CBO9780511662010.008); MR 1696489 (2000e:11077).
- Ogu79 A. Ogus, *Supersingular K3 crystals*, in *Journées de géométrie algébrique de Rennes (Rennes, 1978)*, Astérisque, vol. 64 (Société Mathématique de France, Paris, 1979), 3–86; MR 563467 (81e:14024).
- Pap00 G. Pappas, *On the arithmetic moduli schemes of PEL Shimura varieties*, J. Algebraic Geom. **9** (2000), 577–605; MR 1752014 (2001g:14042).

- PR94 V. Platonov and A. Rapinchuk, *Algebraic Groups and Number Theory*, Pure and Applied Mathematics, vol. 139 (Academic Press, Boston, MA, 1994), translated from the 1991 Russian original by Rachel Rowen; [MR 1278263](#) (95b:11039).
- RZ96 M. Rapoport and Th. Zink, *Period Spaces for p -divisible Groups*, Annals of Mathematics Studies, vol. 141 (Princeton University Press, Princeton, NJ, 1996); [MR 1393439](#) (97f:14023).
- Vas99 A. Vasiu, *Integral canonical models of Shimura varieties of preabelian type*, Asian J. Math. **3** (1999), 401–518; [MR 1796512](#) (2002b:11087).
- Vas14 A. Vasiu, *Integral models in unramified mixed characteristic $(0,2)$ of hermitian orthogonal Shimura varieties of PEL type, Part II*, Math. Nachr. **287** (2014), 1756–1773.
- VZ10 A. Vasiu and T. Zink, *Purity results for p -divisible groups and abelian schemes over regular bases of mixed characteristic*, Doc. Math. **15** (2010), 571–599; [MR 2679067](#).

Keerthi Madapusi Pera keerthi@math.harvard.edu
Department of Mathematics, 1 Oxford St, Harvard University,
Cambridge, MA 02118, USA