

was distributed to patients, professionals, and referrers, either using, or associated with the team.

The total number of participants recruited was $n = 45$. These were made up of service users ($n = 17$), referrers to the service ($n = 10$), and other professionals ($n = 18$).

Results. Participants felt they understood the purpose of the SCFT, placed importance on being involved in service evaluation, and were confident their responses would influence service development.

Reasons to refer to the SCFT were the perceived helpfulness of the team, supporting transitions, risk management, teamwork, therapeutic alliance, quality, and clinical knowledge.

Results favoured multi-disciplinary team agreement as being an important factor in the timing of SCFT referral. Upon admission, or granting of unescorted leave, were also cited as appropriate times to refer to the service. Clinically appropriate timing, individual needs, and service user motivation were additional indicators for SCFT referral.

Conclusion. The West Yorkshire SCFT offers previously unavailable pathways from secure services into the community. The clinical model uses a trauma-informed, formulation-driven, collaborative approach to care, treatment, and risk management, which participants found favourable. Improved community pathways and connections offer a sense of improved hope, and a feeling of being helped, which is supportive of personal recovery.

There are recommendations which suggest that a community pathway agenda, embedded into services from admission, will support clinically appropriate timing of SCFT referral, and should be a decision which is made collaboratively, with patients, carers, and the multidisciplinary services around them.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

Service Evaluation of a Boxercise Programme in an Inpatient Rehabilitation Setting

Dr Alexander Graham*, Miss Leah Canning and Dr Meenakshi Lachman

Leeds and York Partnership Foundation Trust, Leeds, United Kingdom

*Presenting author.

doi: 10.1192/bjo.2024.474

Aims. We undertook a service evaluation obtaining feedback from service users in an inpatient rehabilitation setting about a weekly Boxercise class. The aim was to assess the experiences of service users, and the role it has in their recovery.

We hypothesised that the class would be well received by service users in aspects of enjoyment, impact on biopsychosocial wellbeing and recovery based on positive comments made by service users.

There is an increasing trend to utilise physical activity as an adjunct to improve mental health within healthcare settings; to increase motivation, educate on healthier lifestyles and to enhance well-being outcomes. This Boxercise programme has been developed by the Healthy Living Advisor within the rehabilitation inpatient facility at Leeds and York Partnership Trust. The programme has run for one year, and there has been a large uptake of service users who participate in the group. The Boxercise classes aim to encourage discipline, communication, spatial awareness, and cognitive skills in a modality that is interesting to service users.

Methods. Service users who are regular participants in a Boxercise programme at an inpatient rehabilitation centre completed a questionnaire. A five-point Likert scale assessed participant views across seven domains. Participants were then asked to write three words that describe their feelings about the Boxercise programme, complete a drawing showing their thoughts after a Boxercise class and provide suggestions for improvement.

Results. Eleven participants completed the questionnaire. Average scores for the domains were as follows: enjoyability 4.45/5 (89%), physical health 4.55/5 (91%), mental health 4.27/5 (85%), recovery 4.09/5 (82%), socialising 3.91/5 (82%), safety 4.64/5 (93%), continue after discharge 3.36/5 (67%).

The 'three words' were put in a word cloud generator with highest weighted words: 'Fun', 'Good', 'Energetic', 'Confident'.

Common themes from the pictures shown were smiling faces and 'strongman' images.

Six participants gave feedback that more equipment (pads and gloves) would help to improve their experience in the classes.

Conclusion. The Boxercise programme received positive feedback from participants that aligns with the hypothesis; particularly in safety, enjoyability, benefit to physical health and benefit to mental health.

The participants had positive views on the class as an adjunct to the management of their physical and mental wellbeing. The feedback from all the participants is that they felt safe during the classes.

This service evaluation indicates that the participants value the Boxercise classes as an enjoyable activity and as an adjunct to their treatment.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

Mind Over Menopause: Bridging the Gap in Mental Health Care

Dr Alexandra Thatcher¹, Dr Catherine Graham^{2*}, Dr Grace Denton², Dr Katherine Kearly-Shiers³ and Dr Kristyn Manley³

¹Gloucestershire Health and Care NHS Trust, Cheltenham, United Kingdom; ²Avon and Wiltshire Partnership NHS Trust, Bristol, United Kingdom and ³University Hospitals Bristol and Weston NHS Trust, Bristol, United Kingdom

*Presenting author.

doi: 10.1192/bjo.2024.475

Aims. The effect of menopause on mental health is increasingly well recognised. Studies assessing peri- and post-menopausal women report higher incidences of depression and anxiety. Without recognition and treatment, the negative impact on mental health during menopause can lead to long-lasting effects on quality of life. NICE and the British Menopause Society (BMS) guidelines recommend cognitive behavioural therapy (CBT) and an individualised approach, for women experiencing depression and anxiety as a result of menopause. The aim of this project was to collect data relating to the provision of mental health interventions (and how they are accessed) for women seeing menopause specialists across the UK. This data can then be used to inform and promote improvements in the delivery of care for menopause mood symptoms.

Methods. An expert panel of psychiatry, gynaecology and general practice clinicians designed an online survey which considered