

MULTIPLICITIES OF HIGHER LIE CHARACTERS

MANFRED SCHOCKER

(Received 17 June 2001; revised 26 April 2002)

Communicated by Jie Du

Abstract

The higher Lie characters of the symmetric group S_n arise from the Poincaré-Birkhoff-Witt basis of the free associative algebra. They are indexed by the partitions of n and sum up to the regular character of S_n . A combinatorial description of the multiplicities of their irreducible components is given. As a special case the Kraśkiewicz-Weyman result on the multiplicities of the classical Lie character is obtained.

2000 *Mathematics subject classification*: primary 20C30*, 05E10, 17B99.

Keywords and phrases: symmetric group, general linear group, free Lie algebra, tableau, major index.

1. Introduction

At the beginning of the last century Schur studied the structure of the tensor algebra $T(V)$ over a finite dimensional K -vector space V as a $GL(V)$ -module. In his thesis ([13]) and a famous subsequent paper ([14]) he was able to describe the decomposition of the homogeneous components

$$T_n(V) := \underbrace{V \otimes \cdots \otimes V}_n$$

of degree n in $T(V)$ into irreducible $GL(V)$ -modules using the irreducible representations of the symmetric group S_n . The usual Lie bracketing $[x, y] := xy - yx$ turns $T(V)$ into a Lie algebra. The Lie subalgebra $L(V)$ generated by V is free over any basis of V by a classical result of Witt ([17]), and $L_n(V) := T_n(V) \cap L(V)$ is a $GL(V)$ -submodule of $T_n(V)$ for all n . Let $q = q_1 \dots q_k$ be a partition of n , that is, $q_1 \geq \dots \geq q_k$ and $q_1 + \dots + q_k = n$. Then we define

$$L_q(V) := \left\langle \sum_{\pi \in S_k} P_{1\pi} \cdots P_{k\pi} \mid P_i \in L_{q_i}(V) \text{ for } 1 \leq i \leq k \right\rangle_K.$$

By the Poincaré-Birkhoff-Witt theorem, $T_n(V)$ is the direct sum of these subspaces:

$$(1) \quad T_n(V) = \bigoplus_{q \vdash n} L_q(V),$$

and this decomposition is $GL(V)$ -invariant.

Meanwhile, different families of idempotents e_q in the group algebra KS_n indexed by partitions have been introduced such that $L_q(V) \cong e_q T_n(V)$ for all q (see, for example, [2, 3, 11]). For any decomposition $e_q KS_n = \bigoplus_p a_{q,p} M_p$ into irreducible S_n -modules, we now have

$$L_q(V) = e_q T_n(V) \cong e_q KS_n \otimes_{KS_n} T_n(V) = \bigoplus_p a_{q,p} (M_p \otimes_{KS_n} T_n(V)).$$

In this decomposition, by Schur's fundamental result, $M_p \otimes_{KS_n} T_n(V)$ is either 0 or an irreducible $GL(V)$ -module. Hence the $GL(V)$ -module structure of $L_q(V)$ is completely determined by the multiplicities $a_{q,p}$ of the *higher Lie module* $e_q KS_n$ of S_n . In this vein, for the special case of $q = n$, the problem of describing the $GL(V)$ -module structure of $L_n(V)$ formulated by Thrall ([16]) could finally be solved in a satisfying way by works of Klyachko ([8]) and Kraśkiewicz and Weyman ([9]).

The *higher Lie characters* λ_q of S_n corresponding to the modules $e_q KS_n$ sum up to the regular character of S_n , by (1), and it is natural to ask for their multiplicities for arbitrary q . In this paper, a combinatorial description of these multiplicities is given in terms of alternating sums of numbers of standard tableaux with certain major index properties (Section 3). For $q = n$, we obtain the Kraśkiewicz-Weyman result mentioned above. Our approach is based on a generalization of Klyachko's result (Section 2) combined with the calculus of noncommutative character theory introduced in [6] (Section 4).

2. The reduction to partitions of block type

Let q be a partition of n . The higher Lie character λ_q is induced by a certain linear character of the centralizer of an element of cycle type q in S_n . For $q = n$, this result is due to Klyachko ([8]). In full generality, it is implicitly contained in [1] for the first time (for details, see [12, Section 8.5]) and will be briefly recalled in two steps in this section.

Let \mathbb{N} (\mathbb{N}_0 , respectively) be the set of all positive (nonnegative, respectively) integers and $\underline{n} := \{k \in \mathbb{N} \mid k \leq n\}$ for all $n \in \mathbb{N}_0$. Let \mathbb{N}^* be a free monoid over the alphabet \mathbb{N} . We write $q.r$ for the concatenation product of $q, r \in \mathbb{N}^*$ in order to avoid confusion with the ordinary product in \mathbb{N} . Accordingly, we denote by d^k the k -th power of a letter $d \in \mathbb{N}$ in \mathbb{N}^* , for all $k \in \mathbb{N}_0$. If $n \in \mathbb{N}$ and $q = q_1 \dots q_k \in \mathbb{N}^*$ such that

$q_1 + \dots + q_k = n$, we say that q is a *composition* of n of length $|q| := k$, and write $q \models n$. If, additionally, $q_1 \geq \dots \geq q_k$ and hence q is a partition of n , we write $q \vdash n$.

Let K be a field of characteristic 0 containing a primitive n -th root of unity ε_n for all $n \in \mathbb{N}$. For all $n \in \mathbb{N}_0$, we denote by $\text{Cl}_K(S_n)$ the ring of class functions of the symmetric group S_n . Let C_q be the conjugacy class consisting of all permutations π whose cycle partition $z(\pi)$ is a rearrangement of q , for all $q \in \mathbb{N}^*$. Let $\text{ch}_q \in \text{Cl}_K(S_n)$ such that $(\chi, \text{ch}_q)_{S_n} = \chi(C_q)$ is the value of χ on any element $\pi \in C_q$ for all $\chi \in \text{Cl}_K(S_n)$. Then, up to a certain factor, ch_q is the characteristic function of C_q in $\text{Cl}_K(S_n)$, and we have $C_q = C_r$ and $\text{ch}_q = \text{ch}_r$ whenever q is a rearrangement of r , for all $q, r \in \mathbb{N}^*$. The *outer product* \bullet on the direct sum $\text{Cl} := \bigoplus_{n \in \mathbb{N}_0} \text{Cl}_K(S_n)$ may now be defined by

$$(2) \quad \text{ch}_q \bullet \text{ch}_r := \text{ch}_{q,r}$$

for all $q, r \in \mathbb{N}^*$. It corresponds via Frobenius' characteristic mapping to the ordinary multiplication of symmetric functions.

Our starting point is the following part of [12, Theorem 8.23], which already occurs in [16, Section 8].

LEMMA 2.1. *Let $n \in \mathbb{N}$ and $q \vdash n$. Denote by a_i the multiplicity of the letter i in q , for all $i \in \underline{n}$. Then we have $\lambda_q = \lambda_{n,a_n} \bullet \dots \bullet \lambda_{1,a_1}$.*

Hence, with ζ^p denoting the irreducible character of S_n corresponding to p for $p \vdash n$, the problem of describing the multiplicities

$$a_{q,p} := (\lambda_q, \zeta^p)_{S_n}$$

may be reduced to the case that q is of *block type*, that is, $q = d^k$ is the k -th power of a single letter d . Indeed, for partitions $q = q_1 \dots q_k \vdash x$, $r = r_1 \dots r_l \vdash y$ such that $q_k > r_1$ and $x + y = n$, we have

$$(3) \quad (\lambda_{q,r}, \zeta^p)_{S_n} = (\lambda_q \bullet \lambda_r, \zeta^p)_{S_n} = \sum_{s \vdash x} \sum_{t \vdash y} c_{s,t}^p a_{q,s} a_{r,t}$$

by Lemma 2.1, where $c_{s,t}^p = (\zeta^s \bullet \zeta^t, \zeta^p)_{S_n}$ is the well-known Littlewood-Richardson coefficient.

For all $n, m \in \mathbb{N}_0$, $\psi \in S_n$ and $\sigma \in S_m$, we define $\psi \# \sigma \in S_{n+m}$ by

$$i(\psi \# \sigma) := \begin{cases} i\psi & i \leq n; \\ (i - n)\sigma + n & i > n \end{cases}$$

for all $i \in \underline{n+m}$. Furthermore, for $d, k \in \mathbb{N}$, $n := dk$ and $\pi \in S_k$, we define $\pi^{[d^k]} \in S_n$ by

$$(dj - i)\pi^{[d^k]} := d(j\pi) - i$$

for all $j \in \underline{k}$, $i \in \underline{d-1} \cup \{0\}$. That is, $\pi^{[d^k]}$ is permuting the k successive blocks of length d in \underline{n} according to π . Now let $\tau_d := (1, \dots, d) \in S_d$ be the standard cycle of length d in S_d and put

$$\sigma_{d^k} := \underbrace{\tau_d \# \dots \# \tau_d}_k \in C_{d^k} \subseteq S_n.$$

Then the centralizer of σ_{d^k} in S_n is a wreath product of the cyclic group generated by τ_d with S_k and may be described as

$$C^{d^k} := C_{S_n}(\sigma_{d^k}) = \left\{ \pi^{[d^k]}(\tau_d^{i_1} \# \dots \# \tau_d^{i_k}) \mid \pi \in S_k, i_1, \dots, i_k \in \underline{d} \right\}.$$

([5, Section 4.1]). With these notations, the remaining part of Theorem 8.23 in [12], transferred to Cl, reads as follows.


THEOREM 2.2. *Let $d, k \in \mathbb{N}$ and $n := dk$. Then*

$$\psi_{d^k} : C^{d^k} \longrightarrow K, \quad \pi^{[d^k]}(\tau_d^{i_1} \# \dots \# \tau_d^{i_k}) \longmapsto \varepsilon_d^{-(i_1 + \dots + i_k)}$$


is a linear representation of C^{d^k} , and $(\psi_{d^k})^{S_n} = \lambda_{d^k}$.

3. Multiplicities

In order to state our main result (Theorem 3.1), we need the notion of a standard Young tableau and its multi major index corresponding to a composition. Let $n \in \mathbb{N}$ and $p = p_1 \dots p_l \vdash n$. The frame $R(p) := \{(i, j) \in \mathbb{N} \times \mathbb{N} \mid i \in \underline{l}, j \in \underline{p_{ij}}\}$ corresponding to p may be visualized by its Ferrers diagram, an array of boxes with p_1 boxes in the first (top) row, p_2 boxes in the second row and so on. For example, we have



The images $1\pi, \dots, n\pi$ of any permutation $\pi \in S_n$ may be entered into $R(p)$ row by row, starting at bottom left and ending at top right. Let SYT^p be the set of all permutations which are increasing in rows (from left to right) and columns (downwards) when entered into $R(p)$ in this way. The elements of SYT^p are called *standard Young tableaux* of shape p . In the above example, the elements of $\text{SYT}^{3,2}$, entered into $R(3.2)$, are



Accordingly, we obtain

$$\text{SYT}^{3,2} = \left\{ \begin{pmatrix} 12345 \\ 45123 \end{pmatrix}, \begin{pmatrix} 12345 \\ 35124 \end{pmatrix}, \begin{pmatrix} 12345 \\ 25134 \end{pmatrix}, \begin{pmatrix} 12345 \\ 24135 \end{pmatrix}, \begin{pmatrix} 12345 \\ 34125 \end{pmatrix} \right\} \subseteq S_5.$$

For all $\pi \in S_n$, $D(\pi) := \{i \in \underline{n-1} \mid i\pi > (i+1)\pi\}$ is called the *descent set* of π . Let $q = q_1 \dots q_k \vdash n$ and put $s_j := q_1 + \dots + q_j$ for all $j \in \underline{k} \cup \{0\}$. Then the *multi major index* of π corresponding to q is defined as

$$(4) \quad \text{maj}_q \pi := m_1 \dots m_k \in \mathbb{N}^*,$$

where

$$(5) \quad m_j := \sum_{\substack{s_{j-1} < i \leq s_j \\ i \in D(\pi)}} (i - s_{j-1})$$

for all $j \in \underline{k}$. For $q = n$, we obtain the ordinary major index $\text{maj} \pi := \text{maj}_n \pi$ of π . If, additionally, $r = r_1 \dots r_k \in \mathbb{N}^*$, we define

$$(6) \quad \text{syt}_{q,r}^p := \left| \left\{ \pi \in \text{SYT}^p \mid \forall j \in \underline{k} : (\text{maj}_q(\pi^{-1}))_j \equiv r_j \pmod{q_j} \right\} \right|.$$

Here $(\text{maj}_q(\pi^{-1}))_j$ always denotes the j -th letter of $\text{maj}_q(\pi^{-1})$, for all $j \in \underline{k}$. For arbitrary $r = r_1 \dots r_l, q = q_1 \dots q_k \in \mathbb{N}^*$ we write $r \mid q$ if and only if $l = k$ and r_i is a divisor of q_i for all $i \in \underline{k}$. In this case, we define furthermore the following extension of the number theoretic Möbius function μ :

$$(7) \quad \mu(q/r) := \prod_{i=1}^{|q|} \mu(q_i/r_i).$$

Finally, for $k \in \mathbb{N}$ and $r = r_1 \dots r_l \in \mathbb{N}^*$, we put $k \star r := (kr_1) \dots (kr_l)$.

MAIN THEOREM 3.1. *Let $d, k, n \in \mathbb{N}$ such that $dk = n$. Let $p \vdash n$. Then we have*

$$(\lambda_{d^k}, \zeta^p)_{S_n} = \frac{1}{k!} \sum_{q \vdash k} |C_q| \sum_{r \mid q} \mu(q/r) \text{syt}_{d \star q, r}^p.$$

The proof will be given in Section 5. A description of the multiplicity $(\lambda_q, \zeta^p)_{S_n}$ for arbitrary $q \vdash n$ may be obtained from Theorem 3.1 via (3). For $k \leq 3$, we obtain the following specializations of Theorem 3.1, the first of which is due to Kraśkiewicz and Weyman (see the Remark at the end of this section).

COROLLARY 3.2. *Let $d \in \mathbb{N}$.*

- (a) *For all $p \vdash d$, we have $(\lambda_d, \zeta^p)_{S_d} = \text{syt}_{d,1}^p$.*
- (b) *For all $p \vdash 2d$, we have $(\lambda_{d,d}, \zeta^p)_{S_{2d}} = 1/2(\text{syt}_{d,d,1,1}^p + \text{syt}_{2d,2}^p - \text{syt}_{2d,1}^p)$.*

TABLE 1.

π	π^{-1}	$\text{maj}_6 \pi^{-1}$	$\text{maj}_{3,3} \pi^{-1}$	$\text{maj}_{2,2,2} \pi^{-1}$	$\text{maj}_{4,2} \pi^{-1}$
$\begin{array}{ c c } \hline 1 & \underline{2} \\ \hline 3 & \underline{4} \\ \hline 5 & 6 \\ \hline \end{array}$	$\begin{pmatrix} 123456 \\ 563412 \end{pmatrix}$	6	2.1	0.0.0	2.0
$\begin{array}{ c c } \hline 1 & \underline{2} \\ \hline \underline{3} & \underline{5} \\ \hline 4 & 6 \\ \hline \end{array}$	$\begin{pmatrix} 123456 \\ 563142 \end{pmatrix}$	10	2.2	0.1.1	5.1
$\begin{array}{ c c } \hline \underline{1} & \underline{3} \\ \hline 2 & \underline{4} \\ \hline 5 & 6 \\ \hline \end{array}$	$\begin{pmatrix} 123456 \\ 536412 \end{pmatrix}$	8	1.1	1.1.0	4.0
$\begin{array}{ c c } \hline \underline{1} & \underline{3} \\ \hline 2 & \underline{5} \\ \hline 4 & 6 \\ \hline \end{array}$	$\begin{pmatrix} 123456 \\ 536142 \end{pmatrix}$	9	1.2	1.1.1	4.1
$\begin{array}{ c c } \hline \underline{1} & \underline{4} \\ \hline 2 & \underline{5} \\ \hline 3 & 6 \\ \hline \end{array}$	$\begin{pmatrix} 123456 \\ 531642 \end{pmatrix}$	12	3.3	1.0.1	3.1

(c) For all $p \vdash 3d$, we have

$$(\lambda_{d,d,d}, \zeta^p)_{S_{3d}} = \frac{1}{6} (\text{sytt}_{d,d,d,1,1,1}^p + 3(\text{sytt}_{(2d),d,2,1}^p - \text{sytt}_{(2d),d,1,1}^p) + 2(\text{sytt}_{3d,3}^p - \text{sytt}_{3d,1}^p)).$$

We will illustrate Corollary 3.2 in the case of $p = 2.2.2$. The standard Young tableaux π of shape p are listed in Table 1 together with their multi major indices in question. The descents of π^{-1} are underlined in each case.

By Corollary 3.2, we obtain $(\lambda_6, \zeta^{2.2.2})_{S_6} = 0$ and furthermore

$$(\lambda_{3,3}, \zeta^{2.2.2})_{S_6} = \frac{1}{2}(1 + 1 - 0) = 1$$

and

$$(\lambda_{2,2,2}, \zeta^{2.2.2})_{S_6} = \frac{1}{6}(1 + 3(0 - 1) + 2(1 - 0)) = 0.$$

For $p \vdash d \in \mathbb{N}$ and $\pi \in \text{SYT}^p$, note that $i \in \underline{d-1}$ is a descent of π^{-1} if and only if i stands strictly above $i + 1$ in π , entered into $R(p)$. Hence Corollary 3.2 (a) indeed coincides with the original result of Kraśkiewicz and Weyman on the Lie character λ_d ([9]).

4. Noncommutative character theory

Let $n \in \mathbb{N}$. The *descent algebra* \mathcal{D}_n is defined as the linear span of the elements $\delta^D := \sum \{\pi \in S_n \mid D(\pi) = D\} \ (D \subseteq \underline{n-1})$ in KS_n . Due to Solomon ([15]), \mathcal{D}_n is a subalgebra of KS_n , and there exists a certain epimorphism of algebras $c_n : \mathcal{D}_n \rightarrow Cl_K(S_n)$, for all n . The direct sum $KS := \bigoplus_{n \in \mathbb{N}} KS_n$ is a graded algebra with respect to the convolution product \bullet (see [6, 1.3] for a combinatorial description), and $\mathcal{D} := \bigoplus_{n \in \mathbb{N}} \mathcal{D}_n$ is a \bullet -subalgebra of KS (see [12]). In [6], a (noncommutative) \bullet -subalgebra \mathcal{R} of KS and a \bullet -homomorphism $c : \mathcal{R} \rightarrow Cl$ are introduced such that $\mathcal{D} \subseteq \mathcal{R}$ and $c|_{\mathcal{D}_n} = c_n$ for all n . Furthermore, a (bilinear) scalar product (\cdot, \cdot) on KS is defined by

$$(\pi, \sigma) := \begin{cases} 1 & \pi = \sigma^{-1}; \\ 0 & \pi \neq \sigma^{-1} \end{cases}$$

for all permutations π, σ , and it is shown that

$$(8) \quad (\varphi, \psi) = (c(\varphi), c(\psi))_S$$

for all $\varphi, \psi \in \mathcal{R}$, where the scalar product on the right hand side is the canonical orthogonal extension of the ordinary scalar products $(\cdot, \cdot)_{S_n}$ on $Cl_K(S_n)$, $n \in \mathbb{N}$. For any partition $p \in \mathbb{N}^*$, $Z^p := \sum_{\pi \in SYT^p} \pi$ is an element of \mathcal{R} such that

$$(9) \quad c(Z^p) = \zeta^p$$

is the irreducible character of S_n corresponding to p . For example, for $p = 3.2$, we obtain $Z^{3.2} = \binom{12345}{45123} + \binom{12345}{35124} + \binom{12345}{25134} + \binom{12345}{24135} + \binom{12345}{34125}$. These results provide the following general concept for describing multiplicities: Given an arbitrary character $\chi \in Cl_K(S_n)$, any inverse image $\varphi \in \mathcal{R}$ of χ under c may be understood as a *noncommutative character* corresponding to χ . By (8) and (9), for each such φ , it follows that

$$(10) \quad (\chi, \zeta^p)_{S_n} = (c(\varphi), c(Z^p))_{S_n} = (\varphi, Z^p).$$

The right-hand side of (10) gives different combinatorial descriptions of the multiplicity on the left-hand side, according to the choice of φ , simply by the definition of Z^p and the scalar product on \mathcal{R} .

5. Klyachkos’s idempotent and Ramanujan sums

In the sequel, following the concept described in Section 4, an inverse image of $\lambda_{d,k}$ under c in \mathcal{D} is constructed. It leads to a short proof of our main result Theorem 3.1, by means of (10).

Let $n \in \mathbb{N}$. We put $\kappa_n(x) := \sum_{\pi \in \mathcal{S}_n} x^{\text{maj } \pi} \pi$ (x a variable) and

$$M_{n,i} := \sum_{\substack{\pi \in \mathcal{S}_n \\ \text{maj } \pi \equiv i \pmod n}} \pi \in \mathcal{D}_n$$

for all $i \in \mathbb{N}_0$. Then, up to the factor $1/n$, $\kappa_n(\varepsilon_n) = \sum_{i=1}^n \varepsilon_n^i M_{n,i} \in \mathcal{D}_n$ is a Lie idempotent, that is, $\kappa_n^2 = n\kappa_n$ and $L_n(V) = \kappa_n T_n(V)$. This remarkable result is due to Klyachko ([8]).

LEMMA 5.1. *Let $n, i \in \mathbb{N}$ and d be the order of ε_n^i . Then we have*

$$\kappa_n(\varepsilon_n^i) = \underbrace{\kappa_d(\varepsilon_n^i) \bullet \cdots \bullet \kappa_d(\varepsilon_n^i)}_{n/d}.$$

In particular, $c(\kappa_n(\varepsilon_n^i)) = \text{ch}_{d^{n/d}}$.

The main part of the preceding lemma is a special case of [10, Proposition 4.1], while the additional claim on the c -image follows from [7, Proposition 1]. For $n, m \in \mathbb{N}$, we denote by $\text{gcd}(n, m)$ the greatest common divisor of n and m .

COROLLARY 5.2. *Let $n \in \mathbb{N}$ and $i, j \in \mathbb{N}_0$ such that $\text{gcd}(i, n) = \text{gcd}(j, n)$. Then $c(M_{n,i}) = c(M_{n,j})$.*

PROOF. As $\text{gcd}(i, n) = \text{gcd}(j, n)$, we can find an integer $m \in \mathbb{N}$ such that $i \equiv jm$ modulo n and $\text{gcd}(m, n) = 1$. For all $k \in \mathbb{N}$, we have $\text{gcd}(km, n) = \text{gcd}(k, n)$ and hence $c(\kappa_n(\varepsilon_n^k)) = c(\kappa_n(\varepsilon_n^{mk}))$, by Lemma 5.1. It follows that

$$\begin{aligned} nc(M_{n,i}) &= c\left(\sum_{l=1}^n \sum_{k=1}^n (\varepsilon_n^{l-i})^k M_{n,l}\right) = c\left(\sum_{k=1}^n \varepsilon_n^{-ik} \kappa_n(\varepsilon_n^k)\right) \\ &= c\left(\sum_{k=1}^n \varepsilon_n^{-ik} \kappa_n(\varepsilon_n^{mk})\right) = c\left(\sum_{l=1}^n \sum_{k=1}^n (\varepsilon_n^{lm-i})^k M_{n,l}\right) \\ &= c\left(\sum_{l=1}^n \sum_{k=1}^n ((\varepsilon_n^m)^{l-j})^k M_{n,l}\right) = nc(M_{n,j}). \quad \square \end{aligned}$$

Let $n, m \in \mathbb{N}$. The *Ramanujan sum* corresponding to n and m is defined by

$$\varrho(n, m) := \sum \varepsilon^m,$$

where the sum is taken over all primitive n -th roots of unity ε . In the particular case of $m = 1$ ($m = n$, respectively), $\varrho(n, m)$ yields the Möbius function $\mu(n) = \varrho(n, 1)$

(Euler’s function $\varphi(n) = \varrho(n, n)$, respectively). We write $x \mid m$, if $x \in \mathbb{N}$ is a divisor of m , and put

$$(11) \quad R(n, m) := \sum_{x \mid m} \varrho(n, x) \varrho(m/x, 1).$$

Now, for all $d, k \in \mathbb{N}$ and $p = p_1 \cdot \dots \cdot p_l \in \mathbb{N}^*$, let

$$(12) \quad M_d(k) := \sum_{y \mid dk} R(dk/y, d) M_{dk,y}$$

and

$$M_d(p) := M_d(p_1) \bullet \dots \bullet M_d(p_l).$$

Note that $M_d(p) \in \mathcal{D}$, as \mathcal{D} is closed under the convolution product.

LEMMA 5.3. *For all $d, k \in \mathbb{N}$, we have*

$$\lambda_{d^k} = c \left(\frac{1}{k!} \sum_{\pi \in S_k} \frac{1}{d^{|\mathbf{z}(\pi)|}} M_d(\mathbf{z}(\pi)) \right).$$

(Recall that $\mathbf{z}(\pi)$ denotes the cycle partition of π for any permutation π .)

PROOF. We write

$$\mathbf{z}(\pi; i_1, \dots, i_k) := \mathbf{z}(\pi^{[d^k]}(\tau_d^{i_1} \# \dots \# \tau_d^{i_k}))$$

for all $\pi \in S_k$, $i_1, \dots, i_k \in \underline{d-1} \cup \{0\}$. By Theorem 2.2, we then have

$$\begin{aligned} \lambda_{d^k} &= \frac{1}{|C^{d^k}|} \sum_{q \vdash dk} \left(\sum_{\substack{\varphi \in C^{d^k} \\ \mathbf{z}(\varphi)=q}} \psi_{d^k}(\varphi) \right) \text{ch}_q \\ &= \frac{1}{k!} \sum_{\pi \in S_k} \frac{1}{d^k} \sum_{i_1, \dots, i_k=0}^{d-1} \varepsilon_d^{-\sum i_j} \text{ch}_{\mathbf{z}(\pi; i_1, \dots, i_k)}. \end{aligned}$$

By induction on the number $z = |\mathbf{z}(\pi)|$ of cycles in $\pi \in S_k$, we show that

$$(*) \quad \frac{1}{d^k} \sum_{i_1, \dots, i_k=0}^{d-1} \varepsilon_d^{-\sum i_j} \text{ch}_{\mathbf{z}(\pi; i_1, \dots, i_k)} = c \left(\frac{1}{d^z} M_d(\mathbf{z}(\pi)) \right),$$

which implies our claim. We will use some basic facts about cycle partitions of elements of C^{d^k} which can be found in [5, 4.2]. Let $z = 1$. Then $\pi \in S_k$ is a long

cycle. Putting $\eta := \varepsilon_{kd}$ and applying [5, 4.2.17], Lemma 5.1 and Corollary 5.2, we obtain

$$\begin{aligned} & \frac{1}{d^k} \sum_{i_1, \dots, i_k=0}^{d-1} \varepsilon_d^{-\sum i_j} \text{ch}_{z(\pi; i_1, \dots, i_k)} \\ &= \frac{1}{d} \sum_{i=0}^{d-1} \varepsilon_d^{-i} \text{ch}_{k^*z(\tau_d^i)} = \frac{1}{d} \sum_{x|d} \varrho(d/x, 1) \text{ch}_{k^*z(\tau_d^x)} \\ &= c \left(\frac{1}{d} \sum_{x|d} \varrho(d/x, 1) \kappa_{kd}(\eta^x) \right) = c \left(\frac{1}{d} \sum_{x|d} \sum_{j=0}^{dk-1} \varrho(d/x, 1) \eta^{jx} M_{dk}^{(j)} \right) \\ &= c \left(\frac{1}{d} \sum_{y|dk} M_{dk}^{(y)} \sum_{x|d} \varrho(d/x, 1) \varrho(dk/y, x) \right) = c \left(\frac{1}{d} \sum_{y|dk} M_{dk}^{(y)} R(dk/y, d) \right) \\ &= c(M_d(k)/d). \end{aligned}$$

Now let $z > 1$, say, $\pi = \tilde{\pi}\sigma$ for a cycle σ of length l in π . Then we have, by [5, 4.2.19], (2) and our induction hypothesis,

$$\begin{aligned} & \frac{1}{d^k} \sum_{i_1, \dots, i_k=0}^{d-1} \varepsilon_d^{-\sum i_j} \text{ch}_{z(\pi; i_1, \dots, i_k)} \\ &= \left(\frac{1}{d^{k-l}} \sum_{i_1, \dots, i_{k-l}=0}^{d-1} \varepsilon_d^{-\sum i_j} \text{ch}_{z(\tilde{\pi}; i_1, \dots, i_{k-l})} \right) \cdot \left(\frac{1}{d^l} \sum_{i_{k-l+1}, \dots, i_k=0}^{d-1} \varepsilon_d^{-\sum i_j} \text{ch}_{z(\sigma; i_{k-l+1}, \dots, i_k)} \right) \\ &= c \left(\frac{1}{d^{z-1}} M_d(z(\tilde{\pi})) \cdot \frac{1}{d} M_d(z(\sigma)) \right) = c \left(\frac{1}{d^z} M_d(z(\pi)) \right). \end{aligned}$$

This completes the proof of (*). □

The inverse image of λ_{d^k} under c constructed in the preceding lemma may be simplified by means of a short analysis of the numbers $R(n, m)$. This will be done in three steps.

PROPOSITION 5.4. *Let $n_1, n_2, m_1, m_2 \in \mathbb{N}$ such that*

$$\gcd(n_1, n_2) = \gcd(m_1, m_2) = \gcd(n_1, m_2) = \gcd(n_2, m_1) = 1.$$

Then we have $R(n_1 n_2, m_1 m_2) = R(n_1, m_1) R(n_2, m_2)$.

PROOF. By [4, Theorem 67], the Ramanujan sums have the following factorizing property: $\varrho(a_1 a_2, b) = \varrho(a_1, b) \varrho(a_2, b)$ for all $a_1, a_2, b \in \mathbb{N}$ such that $\gcd(a_1, a_2) = 1$. Furthermore, we have $\varrho(a, b_1 b_2) = \varrho(a, b_1)$ for all $a, b_1, b_2 \in \mathbb{N}$ such that $(a, b_2) = 1$,

as in this case taking the b_2 -th power induces an automorphism of the group of a -th roots of unity. These two observations imply that

$$\begin{aligned}
 R(n_1 n_2, m_1 m_2) &= \sum_{x_1 | m_1} \sum_{x_2 | m_2} \varrho(n_1 n_2, x_1 x_2) \varrho\left(\frac{m_1}{x_1} \frac{m_2}{x_2}, 1\right) \\
 &= \sum_{x_1 | m_1} \sum_{x_2 | m_2} \varrho(n_1, x_1 x_2) \varrho(n_2, x_1 x_2) \varrho\left(\frac{m_1}{x_1}, 1\right) \varrho\left(\frac{m_2}{x_2}, 1\right) \\
 &= \sum_{x_1 | m_1} \varrho(n_1, x_1) \varrho\left(\frac{m_1}{x_1}, 1\right) \sum_{x_2 | m_2} \varrho(n_2, x_2) \varrho\left(\frac{m_2}{x_2}, 1\right) \\
 &= R(n_1, m_1) R(n_2, m_2). \quad \square
 \end{aligned}$$

Let \mathbb{P} be the set of all prime numbers.

PROPOSITION 5.5. For all $a, b \in \mathbb{N}_0$ and $p \in \mathbb{P}$, we have

$$R(p^a, p^b) = \begin{cases} \mu(p^{a-b}) p^b & b \leq a; \\ 0 & b > a. \end{cases}$$

PROOF. For all $n, m \in \mathbb{N}$, the Ramanujan sum corresponding to n and m may be expressed in terms of the Möbius and the Euler function as follows:

$$\varrho(n, m) = \mu(n / \gcd(n, m)) \frac{\varphi(n)}{\varphi(n / \gcd(n, m))}$$

([4, Theorem 272]). Let $c := \min\{a, b\}$ and $d := \min\{a, b - 1\}$. Then

$$\begin{aligned}
 R(p^a, p^b) &= \sum_{i=0}^b \varrho(p^a, p^i) \varrho(p^{b-i}, 1) \\
 &= \varrho(p^a, p^b) - \varrho(p^a, p^{b-1}) \\
 &= \mu(p^{a-c}) \frac{\varphi(p^a)}{\varphi(p^{a-c})} - \mu(p^{a-d}) \frac{\varphi(p^a)}{\varphi(p^{a-d})}
 \end{aligned}$$

and hence $R(p^a, p^b) = 0$ for $b > a$, as $c = d = a$ in this case. Let $b \leq a$. Then we have $c = b$ and $d = b - 1$, that is,

$$R(p^a, p^b) = \mu(p^{a-b}) \frac{\varphi(p^a)}{\varphi(p^{a-b})} - \mu(p^{a-b+1}) \frac{\varphi(p^a)}{\varphi(p^{a-b+1})}.$$

For $b < a - 1$, this shows $R(p^a, p^b) = 0$ as asserted. For $b = a - 1$ it follows that $R(p^a, p^b) = -\varphi(p^{b+1})/\varphi(p) = -p^b$, while, for $b = a$, we may conclude that $R(p^a, p^b) = \varphi(p^b) - \varphi(p^b)/\varphi(p) = p^b$. □

LEMMA 5.6. For all $n, m \in \mathbb{N}$, we have

$$R(n, m) = \begin{cases} \mu(n/m)m & m \mid n; \\ 0 & \text{otherwise.} \end{cases}$$

PROOF. Choose $a_p, b_p \in \mathbb{N}_0$ for all $p \in \mathbb{P}$ such that $n = \prod_{p \in \mathbb{P}} p^{a_p}$ and $m = \prod_{p \in \mathbb{P}} p^{b_p}$. Applying Propositions 5.4 and 5.5 we obtain

$$\begin{aligned} R(n, m) &= \prod_{p \in \mathbb{P}} R(p^{a_p}, p^{b_p}) \\ &= \begin{cases} \prod_{p \in \mathbb{P}} \mu(p^{a_p - b_p}) p^{b_p} & \forall p \in \mathbb{P} : b_p \leq a_p; \\ 0 & \text{otherwise} \end{cases} \\ &= \begin{cases} \mu(n/m)m & m \mid n; \\ 0 & \text{otherwise.} \end{cases} \quad \square \end{aligned}$$

COROLLARY 5.7. Let $d, k \in \mathbb{N}$. Then $M_d(k) = d \sum_{y \mid k} \mu(k/y) M_{dk,y}$.

PROOF. Let y be a divisor of dk . Then Lemma 5.6 implies that

$$R(dk/y, d) = \begin{cases} \mu(dk/dy)d & d \mid dk/y; \\ 0 & \text{otherwise} \end{cases} = \begin{cases} \mu(k/y)d & y \mid k; \\ 0 & \text{otherwise.} \end{cases} \quad \square$$

We are now in a position to give the proof of the Main Theorem 3.1.

PROOF OF THE MAIN THEOREM 3.1. By Lemma 5.3 and (10), we have

$$(\lambda_{d^k}, \zeta^p)_{S_n} = \frac{1}{k!} \sum_{\pi \in S_k} \frac{1}{d^{|\zeta(\pi)|}} (M_d(z(\pi)), \mathbb{Z}^p).$$

But, for $\pi \in S_k$ and $q = q_1 \cdot \dots \cdot q_k := z(\pi)$, we may conclude from Corollary 5.7 that

$$\begin{aligned} \frac{1}{d^{|\zeta(\pi)|}} (M_d(z(\pi)), \mathbb{Z}^p) &= \frac{1}{d^k} (M_d(q_1) \bullet \dots \bullet M_d(q_k), \mathbb{Z}^p) \\ &= \sum_{r_1 \mid q_1} \dots \sum_{r_k \mid q_k} \mu(q_1/r_1) \dots \mu(q_k/r_k) (M_{dq_1, r_1} \bullet \dots \bullet M_{dq_k, r_k}, \mathbb{Z}^p) \\ &= \sum_{r \mid q} \mu(q/r) (M_{dq_1, r_1} \bullet \dots \bullet M_{dq_k, r_k}, \mathbb{Z}^p). \end{aligned}$$

This completes the proof, as $(M_{dq_1, r_1} \bullet \dots \bullet M_{dq_k, r_k}, \mathbb{Z}^p) = \text{syt}_{d \star q, r}^p$ for all $r \mid q$, simply by definition of the scalar product (\cdot, \cdot) and the convolution product \bullet in [6, 1.3]. \square

References

- [1] F. Bergeron, N. Bergeron and A. M. Garsia, 'Idempotents for the free Lie algebra and q -enumeration', in: *Invariant theory and tableaux (Minneapolis, MN, 1988)*, IMA Vol. Math. Appl. 19 (Springer New York, 1990) pp. 166–190.
- [2] D. Blessenohl and H. Laue, 'On the descending Loewy series of Solomon's descent algebra', *J. Algebra* **180** (1996), 698–724.
- [3] A. M. Garsia and C. Reutenauer, 'A decomposition of Solomon's descent algebra', *Adv. in Math.* **77** (1989), 189–262.
- [4] G. H. Hardy and E. M. Wright, *An introduction to the theory of numbers*, 4th Edition (Oxford University Press, Oxford, 1960).
- [5] G. James and A. Kerber, *The representation theory of the symmetric group* (Addison-Wesley, Reading, Massachusetts, 1981).
- [6] A. Jöllenbeck, 'Nichtkommutative Charaktertheorie der symmetrischen Gruppen', *Bayr. Math. Schr.* **56** (1999), 1–41.
- [7] A. Jöllenbeck and M. Schocker, 'Cyclic characters of symmetric groups', *J. Algebraic Combin.* **12** (2000), 155–161.
- [8] A. A. Klyachko, 'Lie elements in the tensor algebra', *Siberian Math. J.* **15** (1974), 914–920.
- [9] W. Kraśkiewicz and J. Weyman, 'Algebra of invariants and the action of a Coxeter element', *Bayr. Math. Schr.* **63** (2001), 265–284.
- [10] B. Leclerc, T. Scharf and J.-Y. Thibon, 'Noncommutative cyclic characters of symmetric groups', *J. Combin. Theory Ser. A (1)* **75** (1996), 55–69.
- [11] F. Patras and C. Reutenauer, 'Higher Lie Idempotents', *J. Algebra* **222** (1999), 51–64.
- [12] C. Reutenauer, *Free Lie algebras*, London Math. Soc. Monographs 7 (Oxford University Press, Oxford, 1993).
- [13] I. Schur, *Über eine Klasse von Matrizen, die sich einer gegebenen Matrix zuordnen lassen* (Dissertation, Berlin, 1901).
- [14] ———, 'Über die rationalen Darstellungen der allgemeinen linearen Gruppe', *Sitzungsber. Pr. Akad. Wiss.* (1927), 58–75.
- [15] L. Solomon, 'A Mackey formula in the group ring of a Coxeter group', *J. Algebra* **41** (1976), 255–268.
- [16] R. Thrall, 'On symmetrized Kronecker powers and the structure of the free Lie ring', *Amer. J. Math.* **64** (1942), 371–388.
- [17] E. Witt, 'Treue Darstellung Liescher Ringe', *J. Reine Angew. Math.* **177** (1937), 152–160.

Mathematical Institute

24–29 St Giles

Oxford OX1 3LB

UK

e-mail: schocker@maths.ox.ac.uk