

Review

Cite this article: Chamanga E, Dyson J, Loke J, McKeown E. (2020) Factors influencing the recruitment and retention of registered nurses in adult community nursing services: an integrative literature review. *Primary Health Care Research & Development* 21(e31): 1–9. doi: [10.1017/S1463423620000353](https://doi.org/10.1017/S1463423620000353)

Received: 8 November 2019

Revised: 1 May 2020

Accepted: 20 July 2020

Key words:

community nursing; community nurses; integrated literature review; recruitment; retention

Author for correspondence:

Edwin Chamanga, Department of Health Service Research and Management, City University London, Northampton Square, Clerkenwell, London, EC1V 0BH, UK.
E-mail: dwin.chamanga.1@city.ac.uk

This paper has not been submitted elsewhere for publication and duplicate publication has been avoided.

Factors influencing the recruitment and retention of registered nurses in adult community nursing services: an integrative literature review

Edwin Chamanga¹ , Judith Dyson², Jennifer Loke³ and Eamonn McKeown⁴

¹City University London, London, UK; ²City University London, London, UK; ³University of Hull, Hull, UK and ⁴City University London, London, UK

Abstract

Background: Adult community nursing services are evolving around the world in response to government policies and changing patient demographics. Amidst these changes, recruitment and retention of community nursing staff are proving a challenge. An integrative literature review has identified multiple factors that influence nurse retention in adult community nursing with sparse information on recruitment factors. Although factors impacting retention of community nurses have been identified, their generalisability around the world is a challenge as they are context and co-dependent. Indicating the need for this area of study to be explored at a local level, as the same factors present with different findings globally. **Aim:** To establish factors influencing recruitment and retention of registered nurses in adult community nursing services. **Design:** Integrative literature review. **Data sources:** Four electronic databases were searched in August 2019 from January 2008 to December 2018: CINAHL Complete, Web of Science, MEDLINE and PROQUEST. Both qualitative and quantitative studies focusing on factors influencing community nursing recruitment and retention were included. **Review methods:** An integrative literature review methodology by Whittemore and Knafl (The integrative review: updated methodology. *Journal of Advanced Nursing* 52, 546–553) was followed, supported by Cochrane guidelines on data synthesis and analysis using a narrative synthesis method. The Center for Evidence-Based Management (CEBMA) critical appraisal tools were used for study quality assessment. **Results:** Ten papers met the study inclusion criteria. Data synthesis and analysis revealed individual and organisational factors influencing the retention of community nurses with the following three dominant themes: (1) work pressure, (2) working conditions and (3) lack of appreciation by managers. **Conclusion:** The review identified context-dependent factors that influence adult community nurses' retention with limited generalisability. There is a lack of data on factors influencing recruitment into adult community nursing; further research is needed to explore factors affiliated to community nursing recruitment.

Introduction and background

The demand for primary care, community care and community nursing services is on the increase due to world demographic changes (World Health Organization, 2008; Maybin, Charles and Honeyman, 2016; Kroezen *et al.*, 2015). The needs of community nursing patients are changing, requiring a new skill mix responsive to local patient and population needs (Drennan and Ross, 2019; Drennan *et al.*, 2018; Jackson *et al.*, 2015). The increasing need for community nurses is further confounded by challenges with recruitment and retention of nurses with the World Health Organization reckoning it is nearing a universal challenge (World Health Organization, 2016, 2020).

One in three adults in developed countries have multiple long-term conditions; this is predicted to double by 2035 (Hajat and Kishore, 2018). By 2050, there will be a global increase in nursing demand due to an ageing population, in both developed and developing countries (United Nations, 2015). As nursing home placements are costly, patients are increasingly opting to be nursed at home to mitigate such costs (Maurits *et al.*, 2015a). In most countries, a good death is defined by people dying in a place of their choice whilst receiving optimal care; this has become a social and political priority (Sines *et al.*, 2013; Gomes *et al.*, 2010; Reed, Fitzgerald and Bish, 2018) and most people in both developed and developing nations are opting to die at home in the comfort of family and friends whilst receiving community nursing care.

Governments are urging for the provision of care closer to home (Edwards, 2014; NHS England, 2015; Reed, Fitzgerald and Bish, 2018; Aiken *et al.*, 2014) based on a recognition that quality and evidence-based healthcare in the community can be cheaper than hospital care (Kraszewski and Norris, 2014; Dickson, Gough and Bain, 2011; Maybin, Charles and

© The Author(s) 2020. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.

Honeyman, 2016; Naruse *et al.*, 2012). However, in England, nursing workforce shortage is a greater challenge to community nursing healthcare provision than funding (The Health Foundation, The King's Fund and the Nuffield Trust, 2018) making it difficult to meet such demands.

The English adult community nursing workforce is diverse and made up of different roles (NHS England, 2015). For example, community matrons, specialist nurses, registered general nurses and district nurses to name a few. District nurses have an additional specialist qualification to being a general registered nurse and, in most cases, they hold a team leader role (Queen's Nursing Institute [QNI], 2015). Between 2010 and 2017, district nurses have declined by approximately 45% with an average vacancy rate of 21% (Stephenson, 2015; Nuffieldtrust, 2017; Marangozov *et al.*, 2017). This is further precipitated by most district nurses retiring and an underfunding of the specialist qualification (QNI, 2015). In 2014, 35% of adult community nurses were above the age of 50 years, in comparison to 23.6% of hospital nurses, with both groups planning on retiring within a decade (Royal College of Nursing [RCN], 2012; Ball *et al.*, 2014).

Community nurses' scope of practice is hospital avoidance and an improved quality of life for those with long-term conditions, palliative care for those at the end of life who are housebound (Drennan, 2018). Community nurses are currently employed by different organisations, the government through the National Health Service (NHS), charitable organisations, private sector and community interest companies (QNI, 2014; Maybin *et al.*, 2016). Historically, community nursing services were a single entity provision being provided by district nurses. However, with population changes and disease, progression models aimed at addressing the changes in demand have resulted in new models of care being designed. For instance, in February 1996, community-integrated care teams were introduced, and in September, the same year rapid response teams were introduced (Audit Commission, 1999). These two services aimed at having a group of therapists (nurses, physiotherapists, occupational therapists, social workers and doctors) working as one team from the same office providing a hospital avoidance service. To date, other models of care such as the @home service, a highly specialised service staffed by highly skilled clinicians aimed at hospital avoidance (hospital at home), by administering therapies and requesting tests which are usually accessible in a hospital setting, has been designed (Lee *et al.*, 2017). In addition to the above, Buurtzorg model (originated from the Netherlands) has been evaluated across the country (Drennan *et al.*, 2018). The Buurtzorg model evolves around small teams of nursing staff providing a range of personal care, social and clinical care to people in their own homes in a neighbourhood (a smaller area to the one being currently covered by community nurses). Thus, indicating possible future changes to current community nursing service provision.

Methods

The aim of this integrative literature review was to answer: 'What are the factors that present challenges to community healthcare organisations in recruiting and retaining registered nursing staff into adult community nursing teams?'

Search strategy

A search on PROSPERO database was conducted to establish there were no similar ongoing reviews. The protocol for this review can

Table 1. Search terms

Population
'community health nurs*' OR 'primary care nurs*' OR 'home care' OR 'district nurs*' OR 'public health' OR 'community healthcare' OR 'community health service' OR 'domiciliary healthcare' OR 'domiciliary care' OR 'home visi*' OR 'primary healthcare' OR 'community nurs*'
AND
Exposure/Outcome
'recruit*' OR retain* OR selection OR career* OR employ* OR appoint* OR retention OR turnover OR attrition'

be found on PROSPERO (registration number CRD42018086197). The databases CINAHL Complete, WEB of Science, MEDLINE, and PROQUEST were used as some of the best electronic databases for supporting nursing-related literature searches which are, based on wide coverage of journals, widely used and also covering biomedical literature (Allen *et al.*, 2006). Citation review and snowball approach were applied to identify papers that may have used unusual terms within the inclusion criteria.

Population, Exposure/Outcomes (PEO) underpinned the research question and search terms as it sought to understand a phenomenon, which could have been answered using quantitative or qualitative approach (Bettany-Saltikov, 2012). The search terms chosen to answer the review question are presented in Table 1; both search terms and search process were guided by a subject librarian.

Inclusion and exclusion criteria

The search was restricted to 10 years to maintain a comprehensive yet contemporary review (January 2008–December 2018). Selected studies were peer-reviewed empirical research looking at community nursing for adult patients (over 18 years of age). Studies which looked at community nursing as a subset of other nursing fields were also included due to a dearth solely focusing on community nursing as an independent cohort. All studies were published in English from high- to middle-income countries with a community nursing service comparable to the English community nursing service. As the review process accessed international literature, nurses included were registered nurses of any role licensed to provide community nursing care (in peoples' homes) to adult patients with physical health needs. Opinion papers, editorials, secondary research, commentaries and studies funded by charitable organisations were excluded due to the potential risk of bias.

Study selection, data extraction and quality appraisal

Title and abstract reviews were conducted by EC and JD. JL mediated any study disputes between EC and JD. All the authors reached consensus on all papers which were appropriate for a full-text review and inclusion. Details of the papers selected are shown in the adapted Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) flowchart (Figure 1) (Moher *et al.*, 2009). In total, 1106 subject-related papers were identified as shown in Figure 1, and these were downloaded to Refworks® from each database for record keeping and further analysis. After the removal of duplicates, most of the papers did not meet the inclusion criteria by title (as they were either opinion papers, editorials or not inclusive of adult community nursing or, primary care in general). The remaining papers were reviewed by abstract and 95 did not meet the inclusion criteria as they were non-empirical studies and or irrelevant to this review in terms




Figure 1. Process of paper selection – adapted PRISMA flow diagram.

of focus, which left 27 papers for full-text review against the inclusion criteria, from which only 10 papers were selected. Data extraction included author, aims, location, design, findings and quality appraisal exceptions. The Center for Evidence-Based Management (CEBMA, 2014) critical appraisal of survey and qualitative research tools were used for quality appraisal (Table 2).

Analysis

An integrative literature review approach was chosen as it permits the inclusion of empirical studies conducted with varied methodologies (Whittemore and Knafl, 2005). Specifically, qualitative, and quantitative using different methods, without following a rigorous systematic process on inclusion papers. Data were extracted according to the review question and a Cochrane narrative synthesis was employed as the method of data analysis as it is a robust and systematic method (Ryan, 2013; Snilstveit *et al.*, 2012).

Results

From the selected 10 empirical studies, 3 were from the Netherlands and there was 1 study conducted in each of the following countries: Canada, United States, China, Australia, Japan and England. The tenth paper included was from Europe including eight countries (the Netherlands, Belgium, Finland, France, Germany, Italy, Poland and Slovakia). All but one of the studies were surveys, the exception being focus group interviews.

Synthesis of results and emergent themes

There were no studies found relating to the recruitment of community nurses. Yet, there were 10 reviewed studies focusing on factors which influence retention of community nurses, with emergent and recurring individual and organisational themes that were then selected for synthesis. They were identified by similarities, before being clustered together, either as a main theme or a subtheme as shown in Figure 2. The three key themes were (1) work pressure,

(2) working conditions and (3) lack of appreciation by managers. Each will be discussed along with the included subthemes.

Individual factors

Work pressure

Eight studies identified work pressure (caseload and workload) as a factor that influences the decision for community nurses to remain or leave employment (Maurits *et al.*, 2015b; Estryn-Behar *et al.*, 2010; Ellenbecker *et al.*, 2008; Tummers *et al.*, 2013; Storey *et al.*, 2009; Halcomb and Ashley, 2016; Naruse *et al.*, 2012). In one study, 60–70% of nurses identified work pressure as an important factor that would influence their decision to leave employment (Estryn-Behar *et al.*, 2010). It is worth noting that this study included nurses from other fields of nursing not only community nurses and it was not possible to disaggregate responses. However, this argument was also supported two other studies (Ellenbecker *et al.*, 2008; Tourangeau *et al.*, 2014), which focused only on community nurses. Both the authors argued that a reduction in stress and work pressure supports nurse retention. Storey *et al.* (2009) found caseload pressure linked to early retirement of community nurses who are nearing their retirement age. Participants in one study said they would prefer a lighter workload rather than retire (Storey *et al.*, 2009). The same study found 36% of participants identified staff shortages and increased workload as precipitating factors towards early retirement from community nursing. However, staff retirement is a by-product of staff being overwhelmed nothing to do with recruitment and retention. Unsurprisingly, increased work pressure correlated with reduced occupational commitment and subsequent negative job satisfaction, impacting on patient safety and quality of service being provided (Maurits *et al.*, 2015a). Similarly, study participants in Halcomb and Ashley's (2016) study felt rushed and spending less time as desirable with patients. As rightly pointed out by Naruse *et al.* (2012), 30% of community nurses experience time pressure frequently due to workload, which results in higher levels of emotional exhaustion.

Table 2. Study characteristics

Author	Aim	Country/location	Design and participants	Findings	Quality appraisal exceptions
Ellenbecker <i>et al.</i> (2008)	To examine the contributions of job satisfaction, nurse, agency and market characteristics and intent to stay employed.	United States	Random opportunistic survey ($n = 2459$) (62% response rate) followed by instrument testing ($n = 1900$) (83% response rate) and a further 34% telephone chase up of non-respondents bring a total of the two phases to ($n = 1912$) with complete datasets.	Job retention is influenced by: job tenure, job satisfaction, intent to stay, living arrangements, retirement plan; position of direct patient care, non-profit agency and area wages. Intent to stay within an organisation is influenced by: job satisfaction, direct patient care, educational level and agency size.	Limited response to the mailed survey in phase 1, no pilot of the instrument and no account of ethical consideration. Sample size not justified.
Estryn-Behar <i>et al.</i> (2010)	To examine specific determinant factors that differentiate between 'stayers' and 'leavers' within the nursing profession and identifying factors for premature leaving.	Eight countries (the Netherlands, Belgium, Finland, France, Germany, Italy, Poland and Slovakia).	Cross-national European survey with two time points (baseline) ($n = 34,587$) follow-up ($n = 14,882$).	Main reasons for leaving employment were associated with working conditions, family, health problems and the desire for continuing education.	Self-reported questionnaire bias. Sample size not justified.
Hai-Xia <i>et al.</i> (2015)	To elicit the conditions and factors that cause job burnout in community nurses.	China	Convenience survey ($n = 420$).	Job burnout is intrinsically linked to leaving employment with the following factors impacting significantly on job burnout: emotional exhaustion, income, work satisfaction, educational opportunities, professional title, and years of employment, age, marital status and educational level.	No account of ethical consideration. Sample size not justified.
Halcomb and Ashley (2016)	To identify the most and least satisfying elements of primary healthcare nurse work.	Australia	Online cross-sectional survey using an instrument designed by the research team ($n = 950$).	Most satisfying aspects were helping people, teamwork, autonomy, work-life balance and variety of work. The least satisfying were time constraints, lack of space, lack of respect or recognition and poor remuneration.	Unbalance sample representation as the study had more practice nurses than community nurses. Sample size not justified.
Maurits <i>et al.</i> (2015a)	To examine home care nursing staffs' self-perceived autonomy and how this relates to whether they have considered leaving the healthcare sector.	The Netherlands	Random opportunistic cross-sectional survey ($n = 262$) (67% response rate).	Home care nurses who perceive more autonomy and are more engaged with their work are less likely to consider leaving.	Self-administered questionnaire bias. Sample size not justified.
Maurits <i>et al.</i> (2015b)	To identify job and organisational factors related to self-perceived ability to continue working in the current line of work until the official retirement age.	The Netherlands	Cross-sectional correlational survey with two validated questionnaires ($n = 730$) completed both questionnaires.	Work pressure, appreciation by senior management, autonomy, educational opportunities and communication have a substantial impact on job satisfaction, which is related to self-perceived ability to continue working.	Self-reported assumption bias as reported by the researcher, selection bias. Sample size not justified.
Naruse <i>et al.</i> (2012)	To investigate perceptions between perceived time pressure and burnout among home visiting nurses.	Japan	Self-administered questionnaires with three validated instruments ($n = 177$) (85.1% response rate).	Home nurses experienced a higher emotional exhaustion and depersonalisation as they feel overloaded from paperwork, and anxiety during home visits, due to frequently perceived time pressure, with insufficient rest time.	Self-administered questionnaire bias. Sample size not justified.

Storey <i>et al.</i> (2009)	To examine issues associated with the impact of age on retention of nurses.	England	Cross-sectional survey using semi-structured postal questionnaires (<i>n</i> = 485) (61% response rate).	Nurses are likely to leave in response to high administrative tasks, work and family commitments; and lack of support. They would also like a reduction in working hours near retirement, reduced workload and pension considerations; older nurses felt it was important to be valued and considered when changes are being implemented.	Lack of representativeness in the sample. Sample size not justified.
Tourangeau <i>et al.</i> (2014)	To identify factors influencing home care nurses' intention to remain employed.	Canada	Qualitative focus group interviews (<i>n</i> = 50).	Home care nurses' intention to remain employed is influenced by multiple factors: job characteristics, work structures, relationships and communication, work environment, conditions of employment and nurse response to work.	Participants self-selected to participate in the focus group. No account of interview question pilot study.
Tummers <i>et al.</i> (2013)	To analyse the impact of six job characteristics on the intention to leave employment.	The Netherlands	Survey of (<i>n</i> = 9982) nursing home, care home and home care nurses using comparable validated questionnaires.	Nurses' intention to leave is personal and context-dependent; however, lack of development and career opportunities are the most important factors, followed by a negative working atmosphere.	Sample size not justified.

Poor work-life balance

Community nurses were found to be dissatisfied with their work-life balance with challenges in meeting work and family demands in four studies (Storey *et al.*, 2009; Tourangeau *et al.*, 2014; Estryn-Behar *et al.*, 2010; Naruse *et al.*, 2012). From focus groups conducted by Tourangeau *et al.* (2014), community nurses reported of working from home after hours, with the process being described as stressful, characterised by paperwork to be completed and order of supplies. Eighty-eight per cent of participants in a study by Naruse *et al.* (2012) reported of insufficient rest time. Conversely, Estryn-Behar *et al.* (2010) note starting a family and caring for a family member are challenges associated with achieving a work-life balance, especially for those with full-time contracts.

Emotional exhaustion

A study by Hai-Xia *et al.* (2015) defined community nursing associated with emotional exhaustion as a situation, where community nurses cannot easily address challenges they are facing at work. These challenges were reported to be a result of work environment and workload, which will result in perceived time pressure and job burnout, leading to staff turnover, due to decreased job satisfaction and lack of professional pride (Naruse *et al.*, 2012; Hai-Xia *et al.*, 2015). Emotional exhaustion as a reason for leaving employment in over 50% of participants (Estryn-behar *et al.*, 2010) and Hai-Xia *et al.* (2015) observed a positive relationship between married community nurses and emotional exhaustion. As married community nurses presented with a higher score of emotional exhaustion in comparison to their unmarried counterparts, this highlights family life as a significant factor that can influence community nurses' decision-making in relation to remaining employed.

Depersonalisation

A study by Hai-Xia *et al.* (2015) reported depersonalisation to be (the adoption of negative and indifferent attitudes towards others) positively correlated with job burnout. And Naruse *et al.*, (2012) found that community nurses who perceive job-related time pressure are likely to experience a higher level of depersonalisation. Time pressure is felt by community nurses at patients' homes and during commuting with 95% reporting of anxiety during home visits (Naruse *et al.*, 2012). Emotional exhaustion, poor work-life balance and depersonalisation are positively correlated (Hai-Xia *et al.*, 2015; Naruse *et al.*, 2012).

Organisational factors

Working conditions

In a cross European study with participants from different nursing fields, including community nurses, the main and most frequent reason for leaving an organisation was poor working conditions (Estryn-Behar *et al.*, 2010). A poor work atmosphere was the second highest ranked motive for people to leave employment; conducive working conditions enhance an organisation's ability to retain community nursing workforce (Tummers *et al.*, 2013). The nature of these working conditions was not defined. Conversely, a positive work atmosphere including pleasure at work, good team spirit and collegiality was found necessary for community nurses' retention (Tummers *et al.*, 2013).

Lack of autonomy

Six studies identified community nurses are likely to remain employed if they felt autonomous (empowered/independent)




Figure 2. Themes and subthemes from the reviewed literature.

within their roles (Maurits *et al.*, 2015b; Tummers *et al.*, 2013; Maurits *et al.*, 2015a; Halcomb and Ashley, 2016; Tourangeau *et al.*, 2014; Estryn-Behar *et al.*, 2010). Autonomy encourages job satisfaction which promotes staff retention (Maurits *et al.*, 2015b) and dissatisfaction with a lack of autonomy is a contributing factor towards the decision to leave an organisation (Estryn-Behar *et al.*, 2010; Tummers *et al.*, 2013). Whilst autonomy was valued, some community nurses found it isolating and stressful (Halcomb and Ashley, 2016).

Salaries and wages

There is some evidence to suggest that nurses employed in Eastern European countries are likely to leave employment due to remuneration in comparison to those in other European countries (Estryn-Behar *et al.*, 2010). However, some community nurses felt their low wage rate was mitigated by good working conditions (Halcomb and Ashley, 2016). Some nurses said their salary or wage did not reflect the amount of work and responsibilities of their jobs (Halcomb and Ashley, 2016; Storey *et al.*, 2009). Community nurses earning below a certain threshold are more likely to feel emotionally exhausted and burnout, with a higher intent to leave employment (Hai-Xia *et al.*, 2015). Salaries and wages were used as some of the studies had participants on a monthly salary, whilst others had a wage paid based on the hours worked per week.

Lack of educational opportunities

Educational opportunities (career opportunities/personal development) were identified as important factors in retaining community nurses in five studies (Estryn-Behar *et al.*, 2010; Tummers, Groeneveld and Lankhaar, 2013; Maurits *et al.*, 2015b; Halcomb and Ashley, 2016; Storey *et al.*, 2009). In a study by Halcomb and Ashley (2016), most participants were dissatisfied with access to education and training, leading to staff turnover. Development and career opportunities are important for retaining workforce in community nursing (Estryn-Behar *et al.*, 2010). In contrast, Maurits *et al.* (2015b) argue a lack of educational opportunities alone is not directly significant for staff turnover. However, combined with other factors, it increases one's likelihood of leaving

employment. This is also evidenced by Storey *et al.* (2009) where community nurses nearing retirement were happy to take early retirement as they felt, there were no further educational opportunities; they had reached the 'ceiling' of their career.

Increased administration

Four studies identified community nurses' administration (paperwork) as a risk factor that has a negative impact on workforce retention (Halcomb and Ashley, 2016; Storey *et al.*, 2009; Tourangeau *et al.*, 2014; Naruse *et al.*, 2012). Over 90% of participants who took part in the community nurses' job burnout study (Naruse *et al.*, 2012) felt overloaded by paperwork. Similarly, Tourangeau *et al.* (2014) and Storey *et al.* (2009) identified issues not only with the quantity of community nurses' paperwork but also how this had to be done in their own time with no remuneration and duplication of effort in having to paper then electronic records.

Lack of appreciation by managers

Four studies identified a lack of appreciation (recognition) by managers as a motive for community nurses to consider leaving employment (Halcomb and Ashley, 2016; Tummers, Groeneveld and Lankhaar, 2013; Storey *et al.*, 2009; Maurits *et al.*, 2015b). Occupational commitment, job satisfaction and staff retention are positively linked to appreciation by managers Maurits *et al.* (2015b). Although, Tummers *et al.* (2013) found appreciation by managers had a modest but nonetheless important impact on community nursing retention. Nurses feeling particularly undervalued, especially when being managed by non-clinicians as they believed they had no understanding of their work (Halcomb and Ashley, 2016). Similarly, Storey *et al.* (2009) found job satisfaction diminishes when there is a lack of appreciation by managers.

Application of knowledge and skill

Two studies identified community nurses were dissatisfied when they were unable to use of their competencies and this influenced their decision to consider leaving employment (Estryn-Behar *et al.*, 2010, Tourangeau *et al.*, 2014). The application of knowledge and

skill with a variety of patients is essential to job satisfaction and retaining community nurses (Tourangeau *et al.*, 2014).

Provision of pension and benefits

A good exit pension was a factor influencing the retention of community nursing staff who are nearing retirement, though few nurses understood their pension options (Storey *et al.*, 2009). The pension was not enough to the remaining in the workplace, other factors such as an opportunity for reduced working hours and workload were also important.

Discussion

Our exploration of literature identified a multitude of factors which influence the retentions of nurses as highlighted above at both individual and organisational level. Interestingly, the World Health Organization (2020) suggests that in some situations, nurse retention can be solved by improving salaries, specifically pay equity, working conditions with developmental opportunities and lastly enabling nurses to work to the full extent of their scope of practice. Some of these factors have already been identified as influencing factors towards the retention of community nurses as shown above. However, drawing from the wider literature, a number of studies have considered why hospital nurses leave employment and many of these studies cite job satisfaction (Lu *et al.*, 2012), effectiveness of nurse managers (Halter *et al.*, 2017), staffing levels (Hairr *et al.*, 2014), work autonomy and collegial relationships (Duffield *et al.*, 2014) as influencing factors. No doubt, some of these factors mirror those highlighted above. However, contextual differences should be considered when analysing these factors. As the setting or the context may have an influence on the level of resilience for the nurse.

Community nurses as autonomous practitioners, in most cases, they are lone workers, working in isolation and this may influence their decision-making. For example, one significant difference for community nursing compared with hospital nursing is the nurse–patient ratio related to safe staffing levels and its impact on capacity modelling (Keller *et al.*, 2013; Hairr *et al.*, 2014). A hospital ward has a manageable flow of patients usually with the same levels of acuity, and the ward can control the number of admissions which means the nurse–patient ratio can be managed; this is not the case in community nursing and there is no provision of establishing a nurse–patient ratio (Jackson *et al.*, 2016). Thus, putting more demand on community nurses not only based on skill mix but an expectation to be able to manage all patients with varying levels of acuity. As a result, community nurses become task-focused to complete a task, or in some cases, patient visits are reassigned to another day, impacting on continuity of care (Maybin *et al.*, 2016) and such practices do influence one's decision to remain employed (RCN, 2018). It is recommended that patient acuity and skill mix are considered when addressing nursing workforce issues (Tevington, 2011; Hairr *et al.*, 2014). This results in increased work pressure on community nurses, which will negatively impact on their work–life balance, resulting in poor emotional well-being and may result in depersonalisation.

Despite some of the nurse retention factors being similar between hospital and community nursing, the supply and demand of nurses is a challenge for both sectors (Lee *et al.*, 2017) with a huge impact on community nursing. As community nursing is a subset of the nursing workforce after hospital nursing, competing for the same supply of nurses (Drennan *et al.*, 2015). Furthermore, in England, there has been a steady decline of community nurses

between 2008 and 2018, equivalent to one in three and this decline is currently estimated at 37% (Rolewicz and Palmer, 2019). Therefore, the presentation of similar retention factors to an already depleted workforce will have different implications between hospital and community nursing workforce. It is worth noting economies of scale which are enjoyed by most hospital trusts have an influence on how much extra administrative workload is done by nurses related to caring. This is based on differences in their funding and contractual obligations. As hospitals can either specifically create nursing roles to address such workloads, which may be a luxury for community nursing organisations, given the reduction in the pool of staff they can recruit from and financial commitments (Everhart *et al.*, 2013; Beech *et al.*, 2019).

The English nursing register indicates that nurses leave employment before the age of retirement, due to alternative career opportunities, which may attract flexibility and better working conditions (Nursing and Midwifery Council, 2017) and these factors have been identified here too. In addition to this, nurses have been reported of using the profession as 'stepping stone' (McCrae *et al.*, 2014). This perhaps is related to poor provision of pension and benefits, and remuneration. However, it is unknown how many community nurses and how many hospital nurses have taken this approach or have identified opportunities elsewhere. Overall, literature shows the absence of effective and efficient models considering organisational, professional and personal factors, impinging the development of interventions capable of improving the retention of nursing workforce (Halter *et al.*, 2017).

Current workforce strategies

Evidence suggests that current nursing workforce retention strategies have been mainly focusing on hospital nursing staff, with nothing specific for community nursing, subsequently community nurses are regarded as 'the invisible workforce' (QNI, 2009, 2014; RCN, 2013;). This is not the only flaw, there is also evidence suggesting that these retention models are not co-created with the frontline staff, they are a top-down approach (The King's Fund, 2014), and in some cases, they are unevaluated and their impact is unknown (Ellenbecker *et al.*, 2007). For example, in England, a series of measures were put in place in the early 2000s, to date, and it is still unclear if these measures have been a success at retaining staff or if they were even implemented (Halter *et al.*, 2017). Schemes such as workplace nurseries and 'golden handshakes' have been implemented to entice nurse to join an organisation and remain employed within the organisation (Drennan *et al.*, 2011; Kleebauer, 2016; Jeffrey, 2017). In 2017, a national nurse retention initiative was implemented in England with seven steps for organisations to help improve their staff retention (NHS Improvement, 2017).

To address nurse retention, some organisations have developed mentorship/preceptorship schemes, mainly aimed at newly qualified nurses (Forlines, 2018; Camarena, 2018; Sherrod *et al.*, 2020). As a result, nurses have been noted to have improved job satisfaction, thus leading to retention. However, longitudinal studies are needed to evaluate the effectiveness of these schemes with staff retention (Camarena, 2018).

In some cases, there are regional disparities in terms of what is being implemented as a way of staff retention. For example, in London, there is the Capital Nurse project aimed at returning student nurses who were trained in London to work in London (Longhurst, 2016). Therefore, there is a need for the co-creation of community nursing-specific retention strategies which are

informed by lived experiences of community nurses to inform policy.

Limitations

There were several limitations associated with this review. First, research on community nursing recruitment and retention, is lacking for the former, scant and diverse for the latter. Second, the heterogeneity of methods, instruments and surveys (and included constructs or items).

Third, we cannot be confident we have captured all studies relating to community nurse retention due to the range of job descriptions and the potential for a range in or job expectations. However, we thoroughly reviewed citations and used a snowball approach to identify other terms or potential papers.

Fourth, the reviewed papers' quality of included papers was variable. Included papers were mostly surveys offering no more than a snapshot of information and providing no depth of understanding of the issues relating to retention of community nurses or the potential solutions to the problems. As a result, these methodological issues hampered the extent to which conclusions could be drawn.

Fifth, ethical considerations were considered in all but a few studies. However, it is unknown whether the absence of reference to ethics approval represents researchers' failure to observe the process or failure to report the process.

Future research

Further research exploring both recruitment and retention into community nursing will be able to inform on factors impacting on community nursing recruitment and retention challenges. And this needs to take a qualitative approach, in order to fully explore recruitment and retention factors in detail in comparison to using a predetermined questionnaire, which may not be able to provide detailed information on either individual or organisational influences.

Conclusion

There are multitude of factors which influence community nurses' retention with a lack of information on recruitment factors, and it is difficult to draw a conclusive argument as some of the factors which influence retention also influence recruitment. From the explored studies, factors influencing retention are context-dependent, co-dependent and vary from one clinician to another, and one country to another.

Acknowledgements. The authors thank Fiona Ware, subject librarian and skills advisor at the University of Hull, for her assistance with retrieving some of the reviewed research papers included in this review.

Author contributions. All the authors have agreed on the final version.

Conflict of interest. No conflict of interest.

References

Aiken LH, Sloane DM, Bruyneel L, Van den Heede K, Griffiths P, Busse R, Diomidous M, Kinnunen J, Kozka M, Lesaffre E, McHugh MD, Moreno-Casbas MT, Rafferty AM, Schwendimann R, Scott A, Tishelman C, Achterberg T and Sermeus W (2014) Nurse staffing and education and hospital mortality in nine European countries: a retrospective observational study. *The Lancet* **383**, 1824–1830.

Allen M, Jacobs SK and Levy JR (2006) Mapping the literature of nursing: 1996–2000. *Journal of the Medical Library Association* **94** (2 Suppl), 206–220.

Audit Commission (1999) *First assessment: a review of district nursing services in England and Wales*. London: Audit Commission.

Ball J, Philippou J, Pike G and Sethi J (2014) *Survey of district and community nurses in 2013 Report to the Royal College of Nursing*. London.

Beech J, Bottery S, Charlesworth A, Evans H, Gershlick B, Hemmings N, Imison C, Kahtan P, McKenna H, Murray R and Palmer B (2019) Closing the Gap: Key areas for action on health and care workforce. [Online]. Retrieved 7 May 2019 from <https://www.kingsfund.org.uk/sites/default/files/2019-03/closing-the-gap-health-care-workforce-full-report.pdf>

Bettany-Saltikov J (2012) *How to do a systematic literature review in nursing: a step-by-step guide*. UK: McGraw-Hill Education.

Camarena L (2018) Supporting nurses to increase retention: A Literature Review. DNP Qualifying Manuscripts. 7. [Online]. Retrieved 8 April 2020 from <https://core.ac.uk/display/216990642?source=4>

Dickson CA, Gough H and Bain H. (2011) Meeting the policy agenda, part 2: is a 'Cinderella service' sufficient? *British Journal of Community Nursing* **16**, 540–545.

Drennan V, Goodman C, Manthorpe J, Davies S, Scott C, Gage H and Illiffe S (2011) Establishing new nursing roles: a case study of the English community matron initiative. *Journal of Clinical Nursing* **20**, 2948–2957.

Drennan V and Ross F (2019) Global nurse shortages—the facts, the impact and action for change. *British Medical Bulletin* **130**, 25–37.

Drennan VM (2018) More care out of hospital? A qualitative exploration of the factors influencing the development of the district nursing workforce in England. *Journal of Health Services Research & Policy* **13**, 1355819618769082.

Drennan VM, Caletani M, Ross F, Saunders M and West P (2018) Tackling the workforce crisis in district nursing: can the Dutch Buurtzorg model offer a solution and a better patient experience? A mixed methods case study. *BMJ Open* **8**, e021931–2018-021931.

Drennan VM, Halter M, Grant RL, Gale J, Harris R and Gourlay S (2015) Adult nurse turnover and retention: South London project report, [Online]. Retrieved 16 November 2016 from <https://eprints.kingston.ac.uk/33053/7/Drennan-V-33053.pdf>

Duffield CM, Roche MA, Homer C, Buchan J and Dimitrelis S (2014) A comparative review of nurse turnover rates and costs across countries. *Journal of Advanced Nursing* **70**, 2703–2712.

Edwards N (2014) *Community services: how they can transform care*. London: King's Fund.

Ellenbecker CH, Samia L, Cushman MJ and Porell FW (2007) Employer retention strategies and their effect on nurses' job satisfaction and intent to stay. *Home Health Care Services Quarterly* **26**, 43–58.

Ellenbecker CH, Porell FW, Samia L, Byleckie JJ and Milburn M (2008) Predictors of home healthcare nurse retention/ *Journal of Nursing Scholarship* **40**, 151–160.

Estry-Behar M, van der Heijden BIJM, Fry C and Hasselhorn H (2010) Longitudinal analysis of personal and work-related factors associated with turnover among nurses. *Recherche En Soins Infirmiers* **103**, 29–45.

Everhart D, Neff O, Al-Amin M, Nogle J and Welch-Maldonado R (2013) The effects of Nurse staffing on hospital financial performance: competitive versus less competitive markets. *Health Care Management Review*.

Forlines KL (2018) Increasing registered nurse retention using a mentorship program. Doctoral Dissertations and projects. 1943. [Online]. Retrieved 25 April 2020 from <https://digitalcommons.liberty.edu/doctoral/1943>

Gomes B, McCrone P, Hall S, Koffman J and Higginson IJ (2010) Variations in the quality and costs of end-of-life care, preferences and palliative outcomes for cancer patients by place of death: the QUALYCARE study. *BMC Cancer* **10**, 400.

Hairr DC, Salisbury H, Johannsson M and Redfern-Vance N (2014) Nurse staffing and the relationship to job satisfaction and retention. *Nursing Economic* **32**, 142–147.

Hai-Xia H, Li-Ting L, Feng-Jun Z, Yao-Yao Y, Yu-Xia G and Gui-Ru W (2015) Factors related to job burnout among community nurses in Changchun, China. *Journal of Nursing Research (Lippincott Williams & Wilkins)* **23**, 172–180.

Hajat C and Kishore SP (2018) The case for a global focus on multiple chronic conditions. *BMJ Global Health* **3**, e000874.

- Halcomb E and Ashley C** (2016) Australian primary health care nurses most and least satisfying aspects of work. *Journal of Clinical Nursing* **26**, 535–545.
- Halter M, Pelone F, Boiko O, Beighton C, Harris R, Gale J, Gourlay S and Drennan V** (2017) Interventions to reduce adult nursing turnover: a systematic review of systematic reviews. *The Open Nursing Journal* **11**, 108–123.
- Jackson C, Leadbetter T, Martin A, Wright T and Manley K** (2015) Making the complexity of community nursing visible: the Cassandra project. *British Journal of Community Nursing* **20**, 126–133.
- Jackson C, Wright T and Martin A** (2016) Safe caseloads for adult community nursing services—an updated review of the evidence [Online]. Retrieved 16 November 2017 from https://improvement.nhs.uk/documents/817/Final_Version_Managing_Safe_Caseloads.pdf
- Jeffrey C** (2017) RN recruitment Strategy. Retrieved 15 April 2020 from <http://www.sompar.nhs.uk/media/5466/3-nurse-recruitment-strategy-november-2017.pdf>: Somerset Partnership NHS Foundation Trust.
- Keller AE, Dulle M, Kwiecinski S, Altmier L and Owen C** (2013) Professional collaboration: who should determine safe staffing for nurses?, *Newborn and Infant Nursing Reviews* **13**, 113–116.
- Kleebauer A** (2016) Trust offers band 5 nurses £1,000 ‘golden hello’. [Online]. Retrieved 12 December 2016 from <https://rcni.com/nursing-management/newsroom/news/trust-offers-band-5-nurses-%C2%A31000-golden-hello-bonus-65556>: Nursing Management.
- Kraszewski S and Norris K** (2014) The modern renaissance of the district nurse. *Primary Health Care* (2014) **24**, 26.
- Kroezen M, Dussault G, Craveiro I, Dieleman M, Jansen C, Bucham J, Barriball L, Rafferty A, Bremner J and Sermeus W** (2015) Recruitment and retention of health professionals across Europe: a literature review and multiple case study research. *Health Policy* **119**, 1517–1528.
- Lee G, Pickstone N, Facultad J and Titchener K** (2017) The future of community nursing: Hospital in the Home. *British Journal of Community Nursing* **22**, 174–180.
- Longhurst C** (2016) London student nurse job scheme should address cost of living. Retrieved 27 December 2016 from <https://rcni.com/newsroom/news/london-student-nurse-job-scheme-should-address-cost-living-40401>: RCNi.
- Lu H, Barriball KL, Zhang X and While AE** (2012) Job satisfaction among hospital nurses revisited: a systematic review. *International Journal of Nursing Studies* **49**, 1017–1038.
- Marangozov R, Huxley C, Manzoni C and Pike G** (2017) *Royal College of Nursing Employment Survey 2017*. London: RCN.
- Maurits EEM, de Veer AJE, van der Hoek LS and Francke AL** (2015a) Autonomous home-care nursing staff are more engaged in their work and less likely to consider leaving the healthcare sector: a questionnaire survey. *International Journal of Nursing Studies* **52**, 1816–1823.
- Maurits EEM, de Veer AJE, van der Hoek LS and Francke AL** (2015b) Factors associated with the self-perceived ability of nursing staff to remain working until retirement: a questionnaire survey. *BMC Health Services Research* **15**, 356.
- Maybin J, Charles A and Honeyman M** (2016) *Understanding quality in district nursing services*. London: The King’s Fund. Retrieved 22 September 2016 from <http://www.kingsfund.org.uk/publications/quality-district-nursing> (accessed 22 September 2016).
- McCrae N, Askey-Jones S and Laker C** (2014) Merely a stepping stone? Professional identity and career prospects following postgraduate mental health nurse training. *Journal of Psychiatric and Mental Health Nursing* **21**, 767–773.
- Moher D, Liberati A, Tetzlaff J and Altman DG** (2009) Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Annals of Internal Medicine* **151**, 264–269.
- Naruse T, Taguchi A, Kuwahara Y, Nagata S, Watai I and Murashima S** (2012) Relationship between perceived time pressure during visits and burn-out among home visiting nurses in Japan. *Japan Journal of Nursing Science* **9**, 185–194.
- NHS England** (2015) Framework for commissioning community nursing [Online]. Retrieved 11 June 2016 from <https://www.england.nhs.uk/wp-content/uploads/2015/10/Framework-for-commissioning-community-nursing.pdf>
- NHS Improvement** (2017) Retaining your clinical staff: a practical improvement resource. [Online]. Retrieved 31 December 2017 from https://improvement.nhs.uk/documents/2070/170807_Retention_Improvement_Pack_V13.pdf
- NMC** (2017) *The NMC register 2012/13-2016/17*. London: Nursing Midwifery Council.
- Nuffieldtrust** (2017) Evidence to the House of Lords Committee – The long-term sustainability of the NHS: Creating a sustainable workforce.
- QNI** (2009) *2020 Vision: Focusing on the future of district nursing*. London: QNI.
- QNI** (2014) *2020 Vision 5 Years On: Reassessing the Future of District Nursing*. London: Queen’s Nursing Institute.
- QNI** (2015) The value of the District Nurse Specialist Practitioner Qualification: A report by the Queen’s Nursing Institute. [Online]. Retrieved 2 April 2020 from https://www.qni.org.uk/wp-content/uploads/2016/09/SPQDN_Report_WEB2.pdf
- RCN** (2012) *Mandatory nursing staffing levels*. London: RCN.
- RCN** (2013) *District Nursing: Harnessing the Potential. The RCN’s UK Position on District Nursing*. RCN.
- RCN** (2018) *Staffing for safe and effective care: Nursing on the brink*. London: RCN.
- Reed FM, Fitzgerald L and Bish MR** (2018) A practice model for rural district nursing success in end-of-life advocacy care. *Scandinavian Journal of Caring Sciences* **32**, 746–755.
- Rolewicz L and Palmer B** (2019) Nuffieldtrust: The NHS workforce in numbers. [Online]. Retrieved 8 May 2019 from <https://www.nuffieldtrust.org.uk/resource/the-nhs-workforce-in-numbers>
- Ryan R** (2013) Cochrane Consumers and Communication Review Group. ‘Cochrane Consumers and Communication Review Group: data synthesis and analysis [Online]. Retrieved 14 July 2017 from <https://cccr.cochrane.org/sites/cccr.cochrane.org/files/public/uploads/Analysis.pdf>
- Sherrod D, Holland C and Battle LH** (2020) Nurse preceptors: a valuable resource for adapting staff to change. *Nursing Management* **51**, 50–53.
- Sines D, Aldridge-Bent S, Fanning A, Farrelly P, Potter K and Wright J** (2013) *Community and public health nursing*. London: John Wiley & Sons.
- Sniltveit B, Oliver S and Vojtkova M** (2012) ‘Narrative approaches to systematic review and synthesis of evidence for international development policy and practice. *Journal of Development Effectiveness* **4**, 409–429.
- Stephenson J** (2015) Specialist district nurse role ‘vital’ to safe care in community. *Nursing Times* **16**, 5.
- Storey C, Cheater F, Ford J and Leese B** (2009) Retaining older nurses in primary care and the community. *Journal of Advanced Nursing* **65**, 1400–1411.
- Tevington P** (2011) ‘Mandatory nurse-patient ratios’. *Medsurg Nursing* **20**, 265.
- The Health Foundation, The King’s Fund and the Nuffield Trust** (2018) The health care workforce in England: Make or break? [Online]. Retrieved 2 December 2018 from <https://www.kingsfund.org.uk/sites/default/files/2018-11/The%20health%20care%20workforce%20in%20England.pdf>
- The King’s Fund** (2014) *Culture and leadership in the NHS: The King’s Fund 2014 Survey*. London: The King’s Fund.
- Tourangeau A, Patterson E, Rowe A, Saari M, Thompson H, Macdonald G, Cranley L and Squires M** (2014) Factors influencing home care nurse intention to remain employed. *Journal of Nursing Management* **22**, 1015–1026.
- Tummers LG, Groeneveld SM and Lankhaar M** (2013) Why do nurses intend to leave their organization? A large-scale analysis in long-term care. *Journal of Advanced Nursing* **69**, 2826–2838.
- United Nations** (2015) World population Ageing 2015. [Online]. Retrieved 15 August 2017 from https://www.un.org/en/development/desa/population/publications/pdf/ageing/WPA2015_Report.pdf
- Whittemore R and Knaf K** (2005) The integrative review: updated methodology. *Journal of Advanced Nursing* **52**, 546–553.
- World Health Organization** (2008) The World Health Report 2008: Primary Health Care Now More than Ever: Introduction and Overview [Online]. Retrieved 3 June 2015 from https://www.who.int/whr/2008/whr08_en.pdf
- World Health Organization** (2016) Global strategy on human resources for health: workforce 2030 [Online]. Retrieved 12 January 2017 from <https://www.who.int/hrh/resources/globstrathrh-2030/en/>
- World Health Organisation** (2020) State of the World’s Nursing 2020: Investing in education, jobs and leadership. [Online]. Retrieved 13 April 2020 from <https://www.who.int/publications-detail/nursing-report-2020>