

Of?p stars: a class of slowly rotating magnetic massive stars

Gregg A. Wade¹, Jason H. Grunhut¹, Wagner L. F. Marcolino²,
Fabrice Martins³, Ian D. Howarth⁴, Yael Nazé⁵, Nolan R. Walborn⁶
and the MiMeS Collaboration

¹RMC, Canada; ²ONB, Brazil; ³GRAAL, France; ⁴UCL, UK; ⁵Liège, Belgium; ⁶STScI, USA

Abstract. Only 5 Of?p stars have been identified in the Galaxy. Of these, 3 have been studied in detail, and within the past 5 years magnetic fields have been detected in each of them. The observed magnetic and spectral characteristics are indicative of organised magnetic fields, likely of fossil origin, confining their supersonic stellar winds into dense, structured magnetospheres. The systematic detection of magnetic fields in these stars strongly suggests that the Of?p stars represent a general class of magnetic O-type stars.

Keywords. techniques: spectroscopic, stars: magnetic fields, stars: individual (HD 191612, HD 108, HD 148937)

1. Introduction

The enigmatic Of?p stars are identified by a number of peculiar and outstanding observational properties. The classification was first introduced by Walborn (1972) according to the presence of C III λ 4650 emission with a strength comparable to the neighbouring N III lines. Well-studied Of?p stars are now known to exhibit recurrent, and apparently periodic, spectral variations (in Balmer, He I, C III and Si III lines) with periods ranging from days to decades, strong C III λ 4650 in emission, narrow P Cygni or emission components in the Balmer lines and He I lines, and UV wind lines weaker than those of typical Of supergiants (see Nazé *et al.* 2010 and references therein).

Only 5 Galactic Of?p stars are known (Walborn *et al.* 2010): HD 108, HD 148937, HD 191612, NGC 1624-2 and CPD-28°2561. Three of these stars - HD 108, HD 148937 and HD 191612 - have been studied in detail. In recent years, HD 191612 was carefully examined for the presence of magnetic fields (Donati *et al.* 2006), and was clearly detected. Recent observations, obtained chiefly within the context of the Magnetism in Massive Stars (MiMeS) Project (Martins *et al.* 2010; Wade *et al.*, in prep) have furthermore detected magnetic fields in HD 108 and HD 148937, thereby confirming the view of Of?p stars as a class of slowly rotating, magnetic massive stars.

2. HD 191612

HD 191612 was the first Of?p star in which a magnetic field was detected (Donati *et al.* 2006). Subsequent MiMeS observations with ESPaDOnS@CFHT (Wade *et al.*, in prep) confirm the existence of the field, and demonstrate the sinusoidal variability of the longitudinal field with the H α and photometric period of 537.6 d. As shown in Fig. 1, the longitudinal field, H α and photometric extrema occur simultaneously when folded according to the 537.6 d period. This implies a clear relationship between the magnetic field and the circumstellar envelope. We interpret these observations in the context of the oblique rotator model, in which the stellar wind couples to the kilogauss dipolar magnetic




Figure 1. Longitudinal field (top), H α EW (middle) and Hipparcos mag (bottom) of the Of?p star HD 191612, all phased according to the 537.6 d period. From Wade *et al.*, in preparation.

field, generating a dense, structured magnetosphere, resulting in all observables varying according to the stellar rotation period.

3. HD 108

HD 108 was the second Of?p star in which a magnetic field was detected (Martins *et al.* 2010). Based on long-term photometric and spectroscopic monitoring, HD 108 is suspected to vary on a timescale of 50-60 y (Nazé *et al.* 2001). The magnetic observations acquired by Martins *et al.* from 2007-2009 show at most a marginal increase of the longitudinal field during more than 2 years of observation. This supports the proposal that the variation timescale is in fact the stellar rotational period, and that HD 108 is a magnetic oblique rotator that has undergone extreme magnetic braking.

4. HD 148937

HD 148937 was recently observed intensely by the MiMeS Collaboration, resulting in the detection of circular polarisation within line profiles indicative of the presence of an organised magnetic field of kilogauss strength (Wade *et al.*, in prep). Although the field is consistently detected in the observations, no variability is observed, in particular according to the 7.03 d spectral period. This result supports the proposal by Nazé *et al.* (2010) that HD 148937 is observed with our line-of-sight near the stellar rotational pole.

References

- Donati, J.-F., Howarth, I. D., Bouret, J.-C., Petit, P. *et al.* 2006, *MNRAS*, 365, L6
 Martins, F., Donati, J.-F., Marcolino, W. L. F., Bouret, J.-C. *et al.* 2010, *MNRAS* 407, 1423
 Nazé, Y., Ud-Doula, A., Spano, M., Rauw, G. *et al.* 2010, *A&A* 520A, 59
 Nazé, Y., Vreux, J.-M., & Rauw, G. 2001, *A&A*, 372, 195
 Walborn, N. R. 1972, *JRASC* 66, 71
 Walborn, N. R., Sota, A., Maíz Apellániz, J., Alfaro, E. J. *et al.* 2010, *ApJ* (Letters), 711, L143