



COMPOSITIO MATHEMATICA

A motivic version of the theorem of Fontaine and Wintenberger

Alberto Vezzani

Compositio Math. **155** (2019), 38–88.

[doi:10.1112/S0010437X18007595](https://doi.org/10.1112/S0010437X18007595)



FOUNDATION
COMPOSITIO
MATHEMATICA



LONDON
MATHEMATICAL
SOCIETY
EST. 1865



A motivic version of the theorem of Fontaine and Wintenberger

Alberto Vezzani

ABSTRACT

We establish a tilting equivalence for rational, homotopy-invariant cohomology theories defined over non-archimedean analytic varieties. More precisely, we prove an equivalence between the categories of motives of rigid analytic varieties over a perfectoid field K of mixed characteristic and over the associated (tilted) perfectoid field K^b of equal characteristic. This can be considered as a motivic generalization of a theorem of Fontaine and Wintenberger, claiming that the Galois groups of K and K^b are isomorphic.

Contents

1	Generalities on adic spaces	41
2	Semi-perfectoid spaces	45
3	Categories of adic motives	49
4	Motivic interpretation of approximation results	58
5	The de-perfectoidification functor in characteristic 0	61
6	The de-perfectoidification functor in characteristic p	65
7	The main theorem	68
	Acknowledgements	78
	Appendix A. An implicit function theorem and approximation results	78
	References	85

Introduction

A theorem of Fontaine and Wintenberger [FW79], later expanded by Scholze [Sch12], states that there is an isomorphism between the Galois groups of a perfectoid field K and the associated (tilted) perfect field K^b of positive characteristic. The standard example of such a pair is formed by the completions of the fields $\mathbb{Q}_p(p^{1/p^\infty})$ and $\mathbb{F}_p((t))(t^{1/p^\infty})$. This theorem is a cornerstone of p -adic Hodge theory, providing a striking equivalence between objects in mixed characteristic (finite étale algebras over K) and objects in equal characteristic p (finite étale algebras over K^b).

In the same spirit, the ‘tilting equivalence’ of Scholze [Sch12] promotes the equivalence above to a certain kind of space of higher dimension, namely the *perfectoid spaces* over the two fields. These objects are non-noetherian analytic spaces, which can typically be thought as certain infinite covers of classical rigid analytic varieties obtained by adding all p th roots of local coordinates, and their introduction has had many different applications in arithmetic geometry (see [Sch14]).

Received 8 November 2017, accepted in final form 3 October 2018, published online 23 November 2018.

2010 Mathematics Subject Classification 14F42, 14G22 (primary).

Keywords: rigid analytic geometry, motives, perfectoid spaces.

This journal is © Foundation Compositio Mathematica 2018.

The aim of this paper is to ‘descend’ Scholze’s tilting equivalence to the level of classical rigid analytic varieties defined over K and K^b , at least from a (co-)homological point of view. What we actually prove, is an equivalence between the two associated categories of (mixed, effective, with rational coefficients) rigid analytic motives **RigDM** defined over the two fields, introduced by Ayoub [Ayo15] adapting the classic construction of **DM** by Voevodsky [Voe96].

THEOREM 7.10. *Let K be a perfectoid field with tilt K^b and let Λ be a \mathbb{Q} -algebra. There is a monoidal triangulated equivalence of categories*

$$\mathfrak{F} : \mathbf{RigDM}_{\acute{e}t}^{\text{eff}}(K^b, \Lambda) \xrightarrow{\sim} \mathbf{RigDM}_{\acute{e}t}^{\text{eff}}(K, \Lambda).$$

We remark that, from a motivic point of view, the theorem of Fontaine and Wintenberger can be rephrased by saying that the categories of *Artin motives* over the two fields are equivalent. Conversely, our theorem shows that not only the absolute Galois groups of a perfectoid field and its tilt are isomorphic, but also their *absolute local motivic Galois groups* (in the sense of the introduction of [Ayo15]). By [Vez17], it is also possible to state the result above with respect to motives *without transfers* **RigDA** defined over the two fields (the rigid analytic analog of the category **DA** see [Ayo15]).

The statement above involves only rigid analytic varieties and its proof uses Scholze’s theory of perfectoid spaces only in an auxiliary way. Nonetheless, we can restate our main result highlighting the role of perfectoid spaces as follows.

THEOREM 7.11. *Let K be a perfectoid field and let Λ be a \mathbb{Q} -algebra. There is a monoidal triangulated equivalence of categories*

$$\mathbf{RigDM}_{\acute{e}t}^{\text{eff}}(K, \Lambda) \xrightarrow{\sim} \mathbf{PerfDA}_{\acute{e}t}^{\text{eff}}(K, \Lambda).$$

The category **PerfDA**_{ét}^{eff}(K, Λ) is built in the same way as **RigDA**_{ét}^{eff}(K, Λ) using as a starting point the big étale site of smooth, small perfectoid spaces, i.e., those that are locally étale over some perfectoid ball $\widehat{\mathbb{B}}^n$.

This last theorem provides a way to ‘perfectoidify’ (and ‘de-perfectoidify’) canonically a rigid analytic variety into a perfectoid motive (and vice versa). The first theorem gives a way to ‘tilt’ (or ‘untilt’) canonically also a rigid analytic varieties over K into a rigid analytic motive defined on the field K^b (and vice versa).

We now make a rough sketch on how the motivic ‘untilting’ and ‘de-perfectoidification’ procedures work. Start from a smooth rigid variety X over K^b and associate to it a perfectoid space \widehat{X} obtained by taking the perfection of X . This operation can be performed canonically since K^b has positive characteristic. We then use Scholze’s theorem to tilt \widehat{X} obtaining a perfectoid space \widehat{Y} in mixed characteristic. Suppose now that \widehat{Y} is the limit of a tower of rigid analytic varieties

$$\cdots \rightarrow Y_{h+1} \rightarrow Y_h \rightarrow \cdots \rightarrow Y_1 \rightarrow Y_0$$


such that Y_0 is étale over the Tate ball $\mathbb{B}^n = \text{Spa } K\langle v_1, \dots, v_n \rangle$ and each Y_h is obtained as the pullback of Y_0 by the map $\mathbb{B}^n \rightarrow \mathbb{B}^n$ defined by taking the p^h -powers of the coordinates $v_i \mapsto v_i^{p^h}$. Under such hypotheses (we will actually need slightly stronger conditions on the tower above) we then ‘de-perfectoidify’ \widehat{Y} by associating to it $Y_{\bar{h}}$ for a sufficiently big index \bar{h} .

The main technical problem of this construction is to show that it is independent of the choice of the tower, and on the index \bar{h} . It is also by definition a local procedure, which is not canonically extendable to arbitrary varieties by gluing. In order to overcome these obstacles, we

use in a crucial way some techniques of approximating maps between spaces up to homotopy, which are obtained by a generalization of the implicit function theorem in the non-archimedean setting.

We point out that many deep motivic theorems are widely used throughout the article, and they are crucial to prove that the recipe sketched above gives rise to an equivalence (see the proof of 7.11) and not only for the construction of explicit homotopies. Among them, the Cancellation Theorem [Voe10] and the dualizability of compact motives [Rio05]. These results admit a rigid analytic version, proved by Ayoub in [Ayo15]. We also use the parallel between **DA** and **DM** and in order to do so, we introduce a finer topology in characteristic p (the Frobét topology, also called quiet in [FJ13]) giving rise to transfers by means of a localization procedure (see [Vez17]).

The following diagram of categories of motives summarizes the situation. The equivalence in the bottom line follows easily from the ‘tilting equivalence’ of Scholze. All notations introduced in the theorems and in the diagram will be described in later sections.



The main result also extends to the categories of non-effective motives, as follows.

THEOREM 7.26. *Let K be a perfectoid field with tilt K^{\flat} and let Λ be a \mathbb{Q} -algebra. There is a monoidal triangulated equivalence of categories*

$$\mathfrak{F}^{\text{st}} : \mathbf{RigDM}_{\text{ét}}(K^{\flat}, \Lambda) \xrightarrow{\sim} \mathbf{RigDM}_{\text{ét}}(K, \Lambda).$$

Due to the intricate construction of the functor \mathfrak{F} , the theorem above is not immediately obtained from the effective case and requires a little extra inspection of the categories of spectra.

As for concrete applications, our theorem allows the ‘(un-)tilting’ and ‘(de-)perfectoidifying’ of any motivic cohomology theory on rigid or perfectoid spaces satisfying étale descent and homotopy invariance, having coefficients over \mathbb{Q} (it is expected to extend this over $\mathbb{Z}[1/p]$ as well, see Remark 7.28). Such an example is the overconvergent de Rham cohomology for rigid analytic varieties over K . It gives rise to new cohomology theories ‘à la de Rham’ for (small) perfectoid spaces as well as for rigid varieties over local fields of equi-characteristic p , satisfying descent, homotopy invariance, finite-dimensionality and further formal properties (see [Vez18]), which are compatible with rigid cohomology [Vez19]. The relation with p -adic periods is an object of future research. Also, we underline that such realization functors might finally give rise (thanks to the derived tannakian formalism developed in [Ayo14b]) to the absolute local motivic Galois group of K for any valued field K , the case of equi-characteristic zero being described in the introduction of [Ayo15].

It is possible to use our main theorem to answer positively to a conjecture of Ayoub [Ayo15, Conjecture 2.5.72] giving an explicit description of rigid analytic motives of good reduction over a local field K as well as morphisms between compact rigid analytic motives over K in terms of motives and morphisms defined over the residue field, extending these results from the equi-characteristic case to the mixed characteristic case (see [Vez19]).

We also point out that the motivic tilting equivalence can be used to ‘replace’ the base field K with another one K' having the same tilt. A classic example is formed by the completions K and K' of the fields $\mathbb{Q}_p(p^{1/p^\infty})$ and $\mathbb{Q}_p(\mu_{p^\infty})$, respectively. Similar techniques are exploited by Kedlaya and Liu in [KL16].

The article is organized as follows. In §1 we recall the basic definitions and the language of adic spaces while in §2 we define the environment in which we will perform our construction, namely the category of semi-perfectoid spaces $\widehat{\text{RigSm}}$ and we define the étale topology on it. In §3 we define the categories of motives for RigSm , $\widehat{\text{RigSm}}$ and PerfSm adapting the constructions of Voevodsky and Ayoub. Thanks to the general model-categorical tools introduced in this section, we give in §4 a motivic interpretation of some approximation results of maps valid for non-archimedean Banach algebras. In §§5 and 6 we prove the existence of the de-perfectoidification functor $\mathbb{L}\mu$ from perfectoid motives to rigid motives in zero and positive characteristics, respectively. Finally, we give in §7 the proof of our main result, both in its effective and stable form.

In the appendix, we collect some technical theorems that are used in our proof. Specifically, we first present a generalization of the implicit function theorem in the rigid setting, and conclude a result about the approximation of maps modulo homotopy as well as its geometric counterpart. We also prove the existence of compatible approximations of a collection of maps $\{f_1, \dots, f_N\}$ from a variety in $\widehat{\text{RigSm}}$ of the form $X \times \mathbb{B}^n$ to a rigid variety Y such that the compatibility conditions among the maps f_i on the faces $X \times \mathbb{B}^{n-1}$ are preserved. This fact has important consequences for computing maps to \mathbb{B}^1 -local complexes of presheaves in the motivic setting.

1. Generalities on adic spaces

We start by recalling the language of adic spaces, as introduced by Huber [Hub93, Hub94] and generalized by Scholze and Weinstein [SW13]. We will always work with adic spaces over a non-archimedean valued field K in the following sense.

DEFINITION 1.1. A *non-archimedean field* is a topological field K whose topology is induced by a non-trivial valuation of rank one. The associated norm is a multiplicative map that we denote by $|\cdot|: K \rightarrow \mathbb{R}_{\geq 0}$ and its valuation ring is denoted by K° .

From now on, we fix a non-archimedean field K and we pick a non-zero element $\pi \in K$ with $|\pi| < 1$.

DEFINITION 1.2. A *Tate K -algebra* is a topological K -algebra R for which there exists a subring R_0 such that the set $\{\pi^k R_0\}$ forms a basis of neighborhoods of 0. A subring R_0 with the above property is called a *ring of definition*.

DEFINITION 1.3. Let R be a Tate K -algebra.

- A subset S of R is *bounded* if it is contained in the set $\pi^{-N} R_0$ for some integer N . An element x of R is *power-bounded* if the set $\{x^n\}_{n \in \mathbb{N}}$ is bounded. The set of power-bounded elements is a subring of R that we denote by R° .

- An element x of R is *topologically nilpotent* if $\lim_{n \rightarrow +\infty} x^n = 0$. The set of topologically nilpotent elements is an ideal of R° that we denote by $R^{\circ\circ}$.

DEFINITION 1.4. An *affinoid K -algebra* is a pair (R, R^+) where R is a Tate K -algebra and R^+ is an open and integrally closed K° -subalgebra of R° . A morphism $(R, R^+) \rightarrow (S, S^+)$ of affinoid K -algebras is a pair of compatible K° -linear continuous maps of rings (f, f^+) . An affinoid K -algebra (R, R^+) is called *complete* if R (and hence also R^+) is complete.

Remark 1.5. By [Wed12, Proposition 5.30] if R is a Tate K -algebra, then (R, R°) is an affinoid K -algebra.

DEFINITION 1.6. For any complete Tate K -algebra R we denote by $R\langle v_1, \dots, v_n \rangle$ the Banach algebra of strictly convergent power series in $R[[v_1, \dots, v_n]]$ endowed with the sup-norm (see [BGR84, §1.4.1]). A *topologically of finite type Tate algebra* (or simply *tft Tate algebra*) is a Banach K -algebra R isomorphic to a quotient of the normed K -algebra $K\langle v_1, \dots, v_n \rangle$ for some n .

If R is a tft Tate algebra, the pair (R, R°) is an affinoid K -algebra, and R° is a ring of definition whenever R is reduced (see [BGR84, Theorem 6.2.4/1]).

We now recall the definition of perfectoid pairs, introduced in [Sch12].

DEFINITION 1.7. A *perfectoid field* K is a complete non-archimedean field whose rank one valuation is non-discrete, whose residue characteristic is p and such that the Frobenius is surjective on K°/p . In case $\text{char } K = p$ this last condition amounts to saying that K is perfect.

DEFINITION 1.8. Let K be a perfectoid field.

- A *perfectoid algebra* is a Banach K -algebra R such that R° is bounded and the Frobenius map is surjective on R°/p .
- A *perfectoid affinoid K -algebra* is an affinoid K -algebra (R, R^+) over a perfectoid field K such that R is perfectoid.

Remark 1.9. If R is a perfectoid algebra, then (R, R°) is a perfectoid affinoid K -algebra.

Example 1.10. Suppose that K is a perfectoid field. A basic example of a perfectoid algebra is the following: let $\underline{v} = (v_1, \dots, v_N)$ be a N -tuple of coordinates and $K^\circ[\underline{v}^{1/p^\infty}]$ be the ring $\varinjlim_h K^\circ[\underline{v}^{1/p^h}]$ endowed with the sup-norm induced by the norm on K . We also denote by $K^\circ\langle \underline{v}^{1/p^\infty} \rangle$ its π -adic completion. By [Sch12, Proposition 5.20], the ring $K^\circ\langle \underline{v}^{1/p^\infty} \rangle[\pi^{-1}]$ is a perfectoid K -algebra, which we will denote by $K\langle \underline{v}^{1/p^\infty} \rangle$. The pair $(K\langle \underline{v}^{1/p^\infty} \rangle, K^\circ\langle \underline{v}^{1/p^\infty} \rangle)$ is a perfectoid affinoid K -algebra. We also define in the same way the perfectoid affinoid K -algebra $(K\langle \underline{v}^{\pm 1/p^\infty} \rangle, K^\circ\langle \underline{v}^{\pm 1/p^\infty} \rangle)$ (see [Sch13, Example 4.4]).

Remark 1.11. $K\langle \underline{v}^{1/p^\infty} \rangle$ is isomorphic as a $K\langle \underline{v} \rangle$ -topological module to the completion $\widehat{\bigoplus} K\langle \underline{v} \rangle$ of the free module $\bigoplus K\langle \underline{v} \rangle$ with a basis indexed by the set $I = (\mathbb{Z}[1/p] \cap [0, 1))^N$. By [BGR84, Proposition 2.1.5/7] there is an explicit description of this ring as the subring of $\prod_{i \in I} K\langle \underline{v} \rangle$ whose elements are those $(x_i)_{i \in I}$ such that for any $\varepsilon > 0$ the inequality $\|x_i\| < \varepsilon$ holds for almost all i (that is, for all i except for a finite number of them).

The following theorem summarizes some results of Scholze, including the *tilting equivalence* of perfectoid algebras, which will play a crucial role in our construction.

THEOREM 1.12 [Sch12]. *Let K be a perfectoid field.*

- (i) [Sch12, Lemma 3.4] *The multiplicative monoid $\varprojlim_{x \rightarrow x^p} K$ can be given a structure K^{\flat} of perfectoid field with the norm induced by the multiplicative map $\sharp : K^{\flat} \rightarrow K$. The field K^{\flat} has characteristic p and coincides with K in case $\text{char } K = p$.*
- (ii) [Sch12, Theorem 3.7] *The functor $L \mapsto L^{\flat}$ for L finite étale over K induces an isomorphism $\text{Gal}(K) \cong \text{Gal}(K^{\flat})$.*
- (iii) [Sch12, Lemma 6.2] *There is an equivalence of categories, the tilting equivalence, from perfectoid affinoid K -algebras to perfectoid affinoid K^{\flat} -algebras denoted by $(R, R^+) \mapsto (R^{\flat}, R^{\flat+})$ such that R^{\flat} is multiplicatively isomorphic to $\varprojlim_{x \rightarrow x^p} R$ and $R^{\flat+}$ is multiplicatively isomorphic to $\varprojlim_{x \rightarrow x^p} R^+$.*
- (iv) [Sch12, Proposition 5.20 and Corollary 6.8] *The tilting equivalence associates the perfectoid K -algebra $(K\langle \underline{v}^{1/p^\infty} \rangle, K^{\circ}\langle \underline{v}^{1/p^\infty} \rangle)$ to $(K^{\flat}\langle \underline{v}^{1/p^\infty} \rangle, K^{\flat\circ}\langle \underline{v}^{1/p^\infty} \rangle)$ and the perfectoid K -algebra $(K\langle \underline{v}^{\pm 1/p^\infty} \rangle, K^{\circ}\langle \underline{v}^{\pm 1/p^\infty} \rangle)$ to $(K^{\flat}\langle \underline{v}^{\pm 1/p^\infty} \rangle, K^{\flat\circ}\langle \underline{v}^{\pm 1/p^\infty} \rangle)$.*

We now recall Huber’s construction of the spectrum of a valuation ring (see [Hub94]).

CONSTRUCTION 1.13. Let (R, R^+) be an affinoid K -algebra. The set $\text{Spa}(R, R^+)$ is the set of equivalence classes of continuous multiplicative valuations $|\cdot|$ (see [Hub93, §3] and [Sch12, Definitions 2.2, 2.5]). It is endowed with the topology generated by the basis of *rational subsets* $\{U(f_1, \dots, f_n | g)\}$ by letting f_1, \dots, f_n, g vary among elements in R such that f_1, \dots, f_n generate R as an ideal and where the set $U(f_1, \dots, f_n | g)$ is the set of those valuations $|\cdot|$ satisfying $|f_i| \leq |g|$ for all i .

If we define maps of valuation fields over K (see [Sch12, Definition 2.26]) as maps of affinoid K -algebras $(L, L^+) \rightarrow (L', L'^+)$ such that $L'^+ \cap L = L^+$ then $\text{Spa}(R, R^+)$ has the following alternative description: it is the set $\varinjlim \text{Hom}((R, R^+), (L, L^+))$ by letting (L, L^+) vary in the category of valuation fields over K (see [Sch12, Proposition 2.27]). Its topology can be defined by declaring the sets $\{\phi : 0 \neq |\phi(f)| \leq |\phi(g)|\}$ to be open, for all pairs of elements f, g in R .

We can associate to a rational subset $U(f_1, \dots, f_n | g)$ the affinoid K -algebra the affinoid K -algebra

$$(\mathcal{O}(U), \mathcal{O}^+(U)) = (R\langle f_1/g, \dots, f_n/g \rangle, R\langle f_1/g, \dots, f_n/g \rangle^+)$$

defined in [Hub94, §1] and [Sch12, Definition 2.13]. This way, we define a presheaf of affinoid K -algebras $(\mathcal{O}_X, \mathcal{O}_X^+)$ on a basis of $X = \text{Spa}(R, R^+)$.

By [Hub94, Lemma 1.5, Proposition 1.6] for any $x \in X = \text{Spa}(R, R^+)$ the valuation at x extends to a valuation on $\mathcal{O}_{X,x}$ and the stalk $\mathcal{O}_{X,x}^+$ is local and corresponds to $\{f \in \mathcal{O}_{X,x} : |f(x)| \leq 1\}$.

By [Hub94, Proposition 1.6] there holds $\mathcal{O}^+(U) = \{f \in \mathcal{O}(U) : |f(x)| \leq 1 \text{ for all } x \in U\}$ for any rational subset U of $\text{Spa}(R, R^+)$ so that \mathcal{O}^+ is a sheaf if and only if \mathcal{O} is a sheaf.

Sadly enough, $\mathcal{O}, \mathcal{O}^+$ are not sheaves in general as shown at the end of [Hub94, §1]. By Tate’s acyclicity theorem [BGR84, Theorem 8.2.1/1] and Scholze’s acyclicity theorem [Sch12, Theorem 6.3], if (R, R^+) is a tft Tate algebra or a perfectoid affinoid K -algebra, then $\mathcal{O}, \mathcal{O}^+$ are sheaves (see [Hub94, §2]). Also, if $\mathcal{O}^+(U)$ is bounded for all rational subspaces U of X (in which case we say that X is *stably uniform*) then $\mathcal{O}, \mathcal{O}^+$ are sheaves (see [BV18]).

DEFINITION 1.14. Objects of \mathcal{V} are triples $(X, \mathcal{O}_X, \{|\cdot|_x\}_{x \in X})$, where X is a topological space, \mathcal{O}_X is a sheaf of complete topological K -algebras, and $|\cdot|_x$ is a continuous valuation on $\mathcal{O}_{X,x}$. Maps are morphisms of ringed spaces, which induce continuous K -algebra morphisms of sheaves and are compatible with the valuations on stalks.

Remark 1.15. By abuse of notation, whenever R is a tft Tate algebra we sometimes denote by $\text{Spa } R$ the object $\text{Spa}(R, R^\circ)$ of \mathcal{V} .

DEFINITION 1.16. Let X be an object of \mathcal{V} .

- We say that X is an *affinoid adic space* if it is isomorphic to $\text{Spa}(R, R^+)$ for some affinoid K -algebra (R, R^+) .
- We say that X is an *affinoid rigid variety* if it is isomorphic to $\text{Spa}(R, R^\circ)$ for some tft Tate algebra R .
- We say that X is a *perfectoid affinoid space* if it is isomorphic to $\text{Spa}(R, R^+)$ for some perfectoid affinoid K -algebra (R, R^+) .
- We say that X is an *adic space* if it is locally isomorphic to an affinoid adic space.
- We say that X is a *rigid variety* if it is locally isomorphic to an affinoid rigid variety.
- We say that X is a *perfectoid space* if it is locally isomorphic to a perfectoid affinoid space.

There is an apparent clash of definitions between rigid varieties as presented above, and as defined by Tate. In fact, the two categories are canonically isomorphic. We refer to [Hub94, § 4] and [Sch12, § 2] for a more detailed collection of results on the comparison between these theories.

Remark 1.17. By [Hub94, Proposition 2.1(ii)] if X is an adic space and $Y = \text{Spa}(R, R^+)$ is an affinoid adic space then $\text{Hom}(X, Y) \cong \text{Hom}((R, R^+), (\mathcal{O}_X(X), \mathcal{O}_X^+(X)))$. Moreover, as shown in [Hub93, § 3] and [Hub94, § 2], if $(\widehat{R}, \widehat{R}^+)$ is the completion of (R, R^+) then it is an affinoid K -algebra and $\text{Spa}(\widehat{R}, \widehat{R}^+) \cong \text{Spa}(R, R^+)$.

ASSUMPTION 1.18. From now on, we will always assume that K is a perfectoid field. We also make the extra assumption that the invertible element π of K satisfies $|p| \leq |\pi| < 1$ and coincides with $(\pi^b)^\sharp$ for a chosen π^b in K^b . In particular, π is equipped with a compatible system of p -power roots π^{1/p^h} (see [Sch12, Remark 3.5]).

We now consider some basic examples and fix some notation.

Example 1.19. Let $\underline{v} = (v_1, \dots, v_N)$ be a N -tuple of coordinates. The Tate N -ball $\text{Spa}(K\langle \underline{v} \rangle, K^\circ\langle \underline{v} \rangle)$ will be denoted by \mathbb{B}^N and the N -torus $\text{Spa}(K\langle \underline{v}^{\pm 1} \rangle, K^\circ\langle \underline{v}^{\pm 1} \rangle)$ by \mathbb{T}^N . It is the rational subspace $U(1 | v_1 \dots v_N)$ of \mathbb{B}^N . The map of spaces induced by the inclusion $(K\langle \underline{v} \rangle, K^\circ\langle \underline{v} \rangle) \rightarrow (K\langle \underline{v}^{1/p^h} \rangle, K^\circ\langle \underline{v}^{1/p^h} \rangle)$ will be denoted by $\mathbb{B}^N\langle \underline{v}^{1/p^h} \rangle \rightarrow \mathbb{B}^N$. We use the analogous notation $\mathbb{T}^N\langle \underline{v}^{1/p^h} \rangle \rightarrow \mathbb{T}^N$ for the torus. These maps are clearly isomorphic to the endomorphism of \mathbb{B}^N respectively \mathbb{T}^N induced by $v_i \mapsto v_i^{p^h}$.

The space defined by the perfectoid affinoid K -algebra $(K\langle \underline{v}^{1/p^\infty} \rangle, K^\circ\langle \underline{v}^{1/p^\infty} \rangle)$ will be denoted by $\widehat{\mathbb{B}}^N$ and referred to as the *perfectoid N -ball*. The space defined by the perfectoid affinoid K -algebra $(K\langle \underline{v}^{\pm 1/p^\infty} \rangle, K^\circ\langle \underline{v}^{\pm 1/p^\infty} \rangle)$ coincides with the rational subspace $U(1 | v_1 \dots v_N)$ of $\widehat{\mathbb{B}}^N$ and will be denoted by $\widehat{\mathbb{T}}^N$ and will be referred to as the *perfectoid N -torus*.

We now recall the definition of étale maps on the category of adic spaces, taken from [Sch12, § 7].

DEFINITION 1.20. A map of affinoid adic spaces $f : \mathrm{Spa}(S, S^+) \rightarrow \mathrm{Spa}(R, R^+)$ is *finite étale* if the associated map $R \rightarrow S$ is a finite étale map of rings, and if S^+ is the integral closure of R^+ in S . A map of adic spaces $f : X \rightarrow Y$ is *étale* if for any point $x \in X$ there exists an open neighborhood U of x and an affinoid open subset V of Y containing $f(U)$ such that $f|_U : U \rightarrow V$ factors as an open embedding $U \rightarrow W$ and a finite étale map $W \rightarrow V$ for some affinoid adic space W .

The previous definitions, when restricted to the case of tft Tate varieties, coincide with the usual ones, as proved in [FvdP04, Proposition 8.1.2].

Remark 1.21. Suppose we are given a diagram of affinoid K -algebras.

$$\begin{array}{ccc} (R, R^+) & \longrightarrow & (S, S^+) \\ & & \downarrow \\ & & (T, T^+) \end{array}$$

In general, it is not possible to define a push-out in the category of affinoid K -algebras. Nonetheless, this can be performed under some hypothesis. For example, if the affinoid K -algebras are tft Tate algebras then the push-out exists and it is the tft Tate algebra associated to the completion $S \widehat{\otimes}_R T$ of $S \otimes_R T$ endowed with the norm of the tensor product (see [BGR84, § 3.1.1]). In case the affinoid K -algebras are perfectoid affinoid, then the push-out exists and is also perfectoid affinoid. It coincides with the completion of (L, L^+) , where L is the ring $S \otimes_R T$ endowed with the norm of the tensor product and L^+ is the integral closure of $S^+ \otimes_{R^+} T^+$ in L (see [Sch12, Proposition 6.18]). The same construction holds in case the map $(R, R^+) \rightarrow (S, S^+)$ is finite étale and (T, T^+) is a perfectoid affinoid (see [Sch12, Lemma 7.3]). By Remark 1.17, the constructions above give rise to fiber products in the category of adic spaces.

2. Semi-perfectoid spaces

We can now introduce a convenient generalization of both smooth rigid varieties and smooth perfectoid spaces. We recall that our base field K is a perfectoid field (see Assumption 1.18).

PROPOSITION 2.1. Let $\underline{v} = v_1, \dots, v_N$ and $\underline{\nu} = \nu_1, \dots, \nu_M$ be two systems of coordinates. Let (R_0, R_0°) be a tft Tate algebra and let $f : \mathrm{Spa}(R_0, R_0^\circ) \rightarrow \mathbb{T}^N \times \mathbb{T}^M = \mathrm{Spa} K \langle \underline{v}^{\pm 1}, \underline{\nu}^{\pm 1} \rangle$ be a map, which is a composition of finite étale maps and rational embeddings. Also let $\mathrm{Spa}(R_h, R_h^\circ)$ be the affinoid rigid variety $\mathrm{Spa}(R_0, R_0^\circ) \times_{\mathbb{T}^N} \mathbb{T}^N \langle \underline{v}^{1/p^h} \rangle$. The π -adic completion (T, T^+) of $(\varinjlim_h R_h, \varinjlim_h R_h^\circ)$ represents the fiber product $\mathrm{Spa}(R_0, R_0^\circ) \times_{\mathbb{T}^N} \widehat{\mathbb{T}}^N$ and defines a bounded affinoid adic space. Moreover, (T, T^+) is isomorphic to the completion of (L, L^+) , where L is the ring $R_0 \otimes_{K \langle \underline{v} \rangle} K \langle \underline{v}^{1/p^\infty} \rangle$ endowed with the norm of the tensor product and L^+ is the integral closure of R_0° in L .

Proof. Let (T, T^+) be as in the last claim. We let W' be the fiber product of $\mathrm{Spa}(T, T^+)$ and $\widehat{\mathbb{T}}^N \times \widehat{\mathbb{T}}^M$ over $\widehat{\mathbb{T}}^N \times \mathbb{T}^M$. If $\mathrm{char} K = 0$ by [Sch13, Lemma 4.5(i)] it exists and is affinoid perfectoid, represented by a perfectoid pair (T', T'^+) . The same is true if $\mathrm{char} K = p$ as in this

case it coincides with the completed perfection of X_0 (indeed, the pullback of an étale map over Frobenius is isomorphic to Frobenius, see for example [Sta18, Lemma 0EBS]).

From the strict inclusion $(T, T^+) \rightarrow (T', T'^\circ)$ we deduce that T^+ is bounded since T'^+ is, being T' perfectoid. By considering rational subspaces of $\text{Spa}(T, T^+)$ we deduce that this space is stably uniform, and hence sheafy (by [BV18]).

The proof of the alternative description of (T, T^+) follows in the same way as [Sch13, Lemma 4.5(i)]. \square

Remark 2.2. The statement of the previous proposition is an instance when the second term T^+ of some affinoid K -algebra (T, T^+) may *not* be equal to the ring T° .

COROLLARY 2.3. *Let X be a reduced rigid variety with an étale map*

$$f : X \rightarrow \mathbb{T}^N \times \mathbb{T}^M = \text{Spa } K\langle \underline{v}^{\pm 1}, \underline{v}'^{\pm 1} \rangle.$$

Then the fiber product $X \times_{\mathbb{T}^N} \widehat{\mathbb{T}}^N$ exists in the category of adic spaces.

Proof. This follows from Proposition 2.1 and the fact that every étale map is locally (on the source) a composition of rational embeddings and finite étale maps. \square

DEFINITION 2.4. We denote by $\widehat{\text{RigSm}}^{\text{gc}}/K$ the full subcategory of adic spaces whose objects are isomorphic to spaces $X = X_0 \times_{\mathbb{T}^N} \widehat{\mathbb{T}}^N$ with respect to a map of affinoid rigid varieties $f : X_0 \rightarrow \mathbb{T}^N \times \mathbb{T}^M$ that is a composition of rational embeddings and finite étale maps. Such spaces will be called *smooth semi-perfectoid spaces with good coordinates*. Because of Proposition 2.1, such fiber products $X = X_0 \times_{\mathbb{T}^N} \widehat{\mathbb{T}}^N$ exist and are affinoid. Whenever $N = 0$ these varieties are rigid analytic varieties and the full subcategory they form will be denoted by $\text{RigSm}^{\text{gc}}/K$ and referred to as *smooth affinoid rigid varieties with good coordinates*. Whenever $M = 0$ these varieties are perfectoid affinoid spaces and the full subcategory they form will be denoted by $\text{PerfSm}^{\text{gc}}/K$ and referred to as *smooth affinoid perfectoids with good coordinates*. A perfectoid space X in $\widehat{\text{RigSm}}^{\text{gc}}/K$ will be sometimes denoted with \widehat{X} .

When $X = X_0 \times_{\mathbb{T}^N} \widehat{\mathbb{T}}^N$ is in $\widehat{\text{RigSm}}^{\text{gc}}/K$ we denote by X_h the fiber product $X_0 \times_{\mathbb{T}^N} \mathbb{T}^N\langle \underline{v}^{1/p^h} \rangle$ and we will write $X = \varprojlim_h X_h$. We say that a presentation $X = \varprojlim_h X_h$ of an object X in $\widehat{\text{RigSm}}^{\text{gc}}/K$ has *good reduction* if the map $X_0 \rightarrow \mathbb{T}^n \times \mathbb{T}^m$ has an étale formal model $\mathfrak{X} \rightarrow \text{Spf}(K^\circ\langle \underline{v}^{\pm 1}, \underline{v}'^{\pm 1} \rangle)$. We say that a presentation $X = \varprojlim_h X_h$ of an object X in $\widehat{\text{RigSm}}^{\text{gc}}/K$ has *potentially good reduction* if there exists a finite separable field extension L/K such that $X_L = \varprojlim_h (X_h)_L$ has good reduction in $\widehat{\text{RigSm}}^{\text{gc}}/L$. We warn the reader that the association $X \mapsto X_0$ is not functorial and the varieties X_h are not uniquely determined by X in general.

We denote by $\widehat{\text{RigSm}}/K$ the full subcategory of adic spaces which are locally isomorphic to objects in $\widehat{\text{RigSm}}^{\text{gc}}/K$ and its objects will be called *smooth semi-perfectoid spaces*. We denote by RigSm/K the full subcategory of adic spaces, which are locally isomorphic to objects in $\text{RigSm}^{\text{gc}}/K$ and by PerfSm/K the one of adic spaces, which are locally isomorphic to objects in $\text{PerfSm}^{\text{gc}}/K$. Its objects will be called *smooth perfectoid spaces*. Whenever the context allows it, we omit K from the notation.

Remark 2.5. Any smooth rigid variety (see for example [Ayo15, Definition 1.1.41]) has locally good coordinates over \mathbb{T}^N by [Ayo15, Corollary 1.1.51]. Hence, RigSm coincides with the category of smooth rigid varieties.

We remark that the presentations defined above having good reduction are a special case of the objects considered in [And06].

The notation $X = \varprojlim_h X_h$ is justified by the following corollary, which is inspired by [SW13, Proposition 2.4.5].

COROLLARY 2.6. *Let $Y = \text{Spa}(S, S^+)$ be an affinoid such that S^+ is a ring of definition and let $X = \varprojlim_h X_h$ be in $\widehat{\text{RigSm}}^{\text{gc}}$. Then $\text{Hom}(Y, X) \cong \varprojlim_h \text{Hom}(Y, X_h)$.*

Proof. Suppose that $X_h = \text{Spa}(R_h, R_h^\circ)$ and $X = \text{Spa}(R, R^+)$. By Proposition 2.1 R^+ is a ring of definition and is the completion of $\varinjlim R_h^+$. A map from (R, R^+) to (S, S^+) is uniquely determined by a K° -linear map from $\varinjlim R_h^\circ$ to S^+ . Similarly, a map from (R_h, R_h°) to (S, S^+) is uniquely determined by a K° -linear map from R_h° to S^+ . From the isomorphism $\text{Hom}_{K^+}(\varinjlim R_h^\circ, S^+) \cong \varprojlim_h \text{Hom}_{K^+}(R_h^\circ, S^+)$ we then deduce the claim. \square

Let $\{X_h, f_h\}_{h \in I}$ be a cofiltered diagram of rigid varieties and let $\{X \rightarrow X_h\}_{h \in I}$ be a collection of compatible maps of adic spaces. We recall that, according to [Hub96, Remark 2.4.5], one writes $X \sim \varprojlim_h X_h$ when the following two conditions are satisfied.

- (i) The induced map on topological spaces $|X| \rightarrow \varprojlim_h |X_h|$ is a homeomorphism.
- (ii) For any $x \in X$ with images $x_h \in X_h$ the map of residue fields $\varinjlim_h k(x_h) \rightarrow k(x)$ has a dense image.

The apparent clash of notations is solved by the following fact.

PROPOSITION 2.7. *Let $X = \varprojlim_h X_h$ be in $\widehat{\text{RigSm}}^{\text{gc}}$. Then $X \sim \varprojlim_h X_h$.*

Proof. This follows from $\widehat{\mathbb{T}}^N \sim \varprojlim_h \text{Spa } K\langle v^{\pm 1/p^h} \rangle$ and from [Sch12, Proposition 7.16]. \square

Étale maps define a topology on $\widehat{\text{RigSm}}$ in the following way.

DEFINITION 2.8. A collection of étale maps of adic spaces $\{U_i \rightarrow X\}_{i \in I}$ is an *étale cover* if the induced map $\bigsqcup_{i \in I} U_i \rightarrow X$ is surjective. These covers define a Grothendieck topology on $\widehat{\text{RigSm}}$ called the *étale topology*.

We pin down the following facts on the topology of the objects $X = \varprojlim_h X_h$.

PROPOSITION 2.9. *Let $X = \varprojlim_h X_h$ be an object of $\widehat{\text{RigSm}}^{\text{gc}}$.*

- (i) Any finite étale map $U \rightarrow X$ is isomorphic to $U_{\bar{h}} \times_{X_{\bar{h}}} X$ for some integer \bar{h} and some finite étale map $U_{\bar{h}} \rightarrow X_{\bar{h}}$.
- (ii) Any rational subspace $U \subset X$ is isomorphic to $U_{\bar{h}} \times_{X_{\bar{h}}} X$ for some integer H and some rational subspace $U_{\bar{h}} \subset X_{\bar{h}}$.

Proof. The first statement follows from [Sch12, Lemma 7.5]. For the second, we remark that according to [Hub93, Lemma 3.10] any rational subspace $U = (f_1, \dots, f_n | g)$ of X can be defined by means of elements f_i, g lying in $\varinjlim_h \mathcal{O}(X_h)$ since it is dense in $\mathcal{O}(X)$, and hence the claim. \square

The proposition above can also be used to prove that a finite étale extension of an object $\text{Spa}(T, T^+)$ in $\widehat{\text{RigSm}}^{\text{gc}}/K$ also lies in $\widehat{\text{RigSm}}^{\text{gc}}/K$, and hence arbitrary fiber products of étale maps in $\widehat{\text{RigSm}}/K$ exist, and are étale.

COROLLARY 2.10. Let $X = \varprojlim_h X_h$ be an object of $\widehat{\text{RigSm}}^{\text{gc}}$ and let $\mathcal{U} := \{f_i : U_i \rightarrow X\}$ be an étale covering of adic spaces. There exists an integer \bar{h} and a finite affinoid refinement $\{V_j \rightarrow X\}$ of \mathcal{U} , which is obtained by pullback of an étale covering $\{V_{\bar{h}j} \rightarrow X_{\bar{h}}\}$ of $X_{\bar{h}}$ and such that $V = \varprojlim_h V_{hj}$ lies in $\widehat{\text{RigSm}}^{\text{gc}}$ by letting V_{hj} be $V_{\bar{h}j} \times_{X_{\bar{h}}} X_h$ for all $h \geq \bar{h}$.

Proof. Any étale map of adic spaces is locally a composition of rational embeddings and finite étale maps and they descend because of Proposition 2.9. We, therefore, obtain an affinoid refinement $\{V_j \rightarrow X\}$ of \mathcal{U} such that each V_i descends to some X_h . Since X is quasi-compact, we can also refine this covering by a finite one, and choose a common index \bar{h} where each V_j descends to. □

COROLLARY 2.11. A perfectoid space X lies in PerfSm if and only if it is locally étale over $\widehat{\mathbb{T}}^N$.

Proof. Let X be locally étale over $\widehat{\mathbb{T}}^N$. Then it is locally open in a finite étale space over a rational subspace of $\widehat{\mathbb{T}}^N = \varprojlim_h \mathbb{T}^N \langle \underline{v}^{\pm 1/p^h} \rangle$. By Proposition 2.9, we conclude it is locally of the form $X_0 \times_{\mathbb{T}^N} \widehat{\mathbb{T}}^N$ for some étale map $X_0 \rightarrow \mathbb{T}^N = \text{Spa}(K \langle \underline{v}^{\pm 1} \rangle, K^\circ \langle \underline{v}^{\pm 1} \rangle)$, which is the composition of rational embeddings and finite étale maps. □

Remark 2.12. If X is a smooth affinoid perfectoid space, then it has a finite number of connected components. Indeed, it is quasi-compact and locally isomorphic to a rational subspace of a perfectoid space, which is finite étale over a rational subspace of $\widehat{\mathbb{T}}^N$.

For later use, we record the following simple example of a space $X = \varprojlim_h X_h$ for which the varieties X_h are easy to understand.

PROPOSITION 2.13. Consider the smooth affinoid variety with good coordinates

$$X_0 = U(v - 1 \mid \pi) \hookrightarrow \mathbb{T}^1 = \text{Spa}(K \langle v^{\pm 1} \rangle).$$

One has $X_h \cong \mathbb{B}^1$ for all h and $\widehat{X} = \varprojlim_h X_h \cong \widehat{\mathbb{B}}^1$.

Proof. By direct computation, the variety X_h is isomorphic to $\text{Spa}(K \langle v, \omega \rangle / (\omega^{p^h} - (\pi v + 1)))$. Since $|p| \leq |\pi|$ we deduce that $|\binom{p^h}{i}| \leq |\pi|$ for all $0 < i < p^h$. In particular, in the ring $K \langle v, \omega \rangle / (\omega^{p^h} - (\pi v + 1))$ one has

$$|(\omega - 1)^{p^h}| = \left| \pi v + \sum_{i=1}^{p^h-1} \binom{p^h}{i} \omega^i \right| = |\pi|.$$

Analogously, in the ring $K \langle \chi \rangle$ one has

$$|(\chi + \pi^{-1/p^h})^{p^h} - \pi^{-1}| = \left| \chi^{p^h} + \sum_{i=1}^{p^h-1} \binom{p^h}{i} \chi^{p^h-1} \pi^{-i/p^h} \right| = 1.$$

The following maps are, therefore, well defined and clearly mutually inverse:

$$\begin{aligned} X_h = \text{Spa}(K \langle v, \omega \rangle / (\omega^{p^h} - (\pi v + 1))) &\xleftrightarrow{\sim} \text{Spa}(K \langle \chi \rangle) = \mathbb{B}^1 \\ (v, \omega) &\mapsto ((\chi + \pi^{-1/p^h})^{p^h} - \pi^{-1}, \pi^{1/p^h} \chi + 1) \\ \pi^{-1/p^h} (\omega - 1) &\leftrightarrow \chi. \end{aligned}$$

Consider the multiplicative map $\sharp: K^b\langle v^{1/p^\infty} \rangle = (K\langle v^{1/p^\infty} \rangle)^b \rightarrow K\langle v^{1/p^\infty} \rangle$ defined in [Sch12, Proposition 5.17]. By our assumptions on π the element $(v - 1)^\sharp - (v - 1)$ is divisible by π in $K^\circ\langle v^{1/p^\infty} \rangle$ and, therefore, the rational subspace $\widehat{X} \cong U(v - 1 | \pi)$ of $\widehat{\mathbb{T}}^1$ coincides with $U((v - 1)^\sharp | \pi^\sharp)$. From [Sch12, Theorem 6.3] we conclude $\widehat{X}^b \cong U(v - 1 | \pi^b) \hookrightarrow \widehat{\mathbb{T}}^{b1}$, which is isomorphic to \mathbb{B}^{b1} , and hence the claim. \square

From the previous proposition we conclude, in particular, that the perfectoid space $\widehat{\mathbb{B}}^1$ lies in $\text{PerfSm}^{\text{gc}}$.

3. Categories of adic motives

From now on, we fix a commutative ring Λ and work with Λ -enriched categories. In particular, the term ‘presheaf’ should be understood as ‘presheaf of Λ -modules’ and similarly for the term ‘sheaf’. The presheaf $\Lambda(X)$ represented by an object X of a category \mathbf{C} sends an object Y of \mathbf{C} to the free Λ -module $\Lambda \text{Hom}(Y, X)$.

ASSUMPTION 3.1. Unless otherwise stated, we assume from now on that Λ is a \mathbb{Q} -algebra and we omit it from the notations.

We make extensive use of the theory of model categories and localization, following the approach of Ayoub in [Ayo15, Ayo07]. Fix a site (\mathbf{C}, τ) . In our situation, this will be the étale site of RigSm or $\widehat{\text{RigSm}}$. The category of complexes of presheaves $\mathbf{Ch}(\mathbf{Psh}(\mathbf{C}))$ can be endowed with the *projective model structure* for which weak equivalences are quasi-isomorphisms and fibrations are maps $\mathcal{F} \rightarrow \mathcal{F}'$ such that $\mathcal{F}(X) \rightarrow \mathcal{F}'(X)$ is a surjection for all X in \mathbf{C} (cfr [Hov99, § 2.3] and [Ayo07, Proposition 4.4.16]).

Also the category of complexes of sheaves $\mathbf{Ch}(\mathbf{Sh}_\tau(\mathbf{C}))$ can be endowed with the *projective model structure* defined in [Ayo07, Definition 4.4.40]. In this structure, weak equivalences are quasi-isomorphisms of complexes of sheaves.

Remark 3.2. Let \mathbf{C} be a category. As shown in [Fau06] any projectively cofibrant complex \mathcal{F} in $\mathbf{ChPsh}(\mathbf{C})$ is a retract of a complex that is the filtered colimit of bounded above complexes, each constituted by presheaves that are direct sums of representable ones.

Just like in [Jar87, MVW06, MV99] or [Rio07], we consider the left Bousfield localization of the model category $\mathbf{Ch}(\mathbf{Psh}(\mathbf{C}))$ with respect to the topology we select, and a chosen ‘contractible object’. We recall that left Bousfield localizations with respect to a class of maps S (see [Hir03, ch. 3]) is the universal model categories in which the maps in S become weak equivalences. The existence of such structures is granted only under some technical hypotheses, as shown in [Hir03, Theorem 4.1.1] and [Ayo07, Theorem 4.2.71].

PROPOSITION 3.3. *Let (\mathbf{C}, τ) be a site with finite direct products and let \mathbf{C}' be a full subcategory of \mathbf{C} such that every object of \mathbf{C} has a covering by objects of \mathbf{C}' . Also let I be an object of \mathbf{C}' .*

(i) *The projective model category $\mathbf{ChPsh}(\mathbf{C})$ admits a left Bousfield localization $\mathbf{Ch}_I\mathbf{Psh}(\mathbf{C})$ with respect to the set S_I of all maps $\Lambda(I \times X)[i] \rightarrow \Lambda(X)[i]$ as X varies in \mathbf{C} and i varies in \mathbb{Z} .*

(ii) *The projective model categories $\mathbf{ChPsh}(\mathbf{C})$ and $\mathbf{ChPsh}(\mathbf{C}')$ admit left Bousfield localizations $\mathbf{Ch}_\tau\mathbf{Psh}(\mathbf{C})$ and $\mathbf{Ch}_\tau\mathbf{Psh}(\mathbf{C}')$ with respect to the class S_τ of maps $\mathcal{F} \rightarrow \mathcal{F}'$ inducing*

isomorphisms on the ét-sheaves associated to $H_i(\mathcal{F})$ and $H_i(\mathcal{F}')$ for all $i \in \mathbb{Z}$. Moreover, the two localized model categories are Quillen equivalent and the sheafification functor induces a Quillen equivalence to the projective model category $\mathbf{ChSh}_\tau(\mathbf{C})$.

(iii) The model categories $\mathbf{Ch}_\tau\mathbf{Psh}(\mathbf{C})$ and $\mathbf{Ch}_\tau\mathbf{Psh}(\mathbf{C}')$ admit left Bousfield localizations $\mathbf{Ch}_{\tau,I}\mathbf{Psh}(\mathbf{C})$ and $\mathbf{Ch}_{\tau,I}\mathbf{Psh}(\mathbf{C}')$ with respect to the set S_I defined above. Moreover, the two localized model categories are Quillen equivalent.

Proof. According to [Hir03, Theorem 4.1.1], any model category that is left proper and cellular (some technical properties that are defined in [Hir03, Definitions 12.1.1 and 13.1.1]) admits a left Bousfield localization with respect to a set of maps. The model structure on complexes is left proper and cellular (see [SS00, p. 7]). It follows that the projective model structures in the statement are also left proper and cellular (see [Hir03, Propositions 12.1.5 and 13.1.14]), and hence the first claim.

For the first part of the second claim, it suffices to apply [Ayo07, Proposition 4.4.31, Lemma 4.4.34] showing that the localization over S_τ is equivalent to a localization over a set of maps. The second part is a restatement of [Ayo07, Corollary 4.4.42, Proposition 4.4.55].

Since by [Ayo07, Proposition 4.4.31] the τ -localization coincides with the Bousfield localization with respect to a set, we conclude by [Ayo07, Theorem 4.2.71] that the model category $\mathbf{Ch}_\tau\mathbf{Psh}(\mathbf{C})$ is still left proper and cellular. The last statement then follows from [Hir03, Theorem 4.1.1] and the second claim. \square

In the situation above, we will denote by $S_{(\tau,I)}$ the union of the class S_τ and the set S_I .

Remark 3.4. A geometrically relevant situation is induced when I is endowed with a multiplication map $\mu: I \times I \rightarrow I$ and maps i_0 and i_1 from the terminal object to I satisfying the relations of a monoidal object with 0 as in the definition of an interval object (see [MV99, §2.3]). Under these hypotheses, we say that the triple (\mathbf{C}, τ, I) is a *site with an interval*.

Example 3.5. The affinoid rigid variety with good coordinates $\mathbb{B}^1 = \text{Spa } K\langle\chi\rangle$ is an interval object with respect to the natural multiplication μ and maps i_0 and i_1 induced by the substitution $\chi \mapsto 0$ and $\chi \mapsto 1$, respectively.

We now apply the constructions above to the sites introduced in the previous sections. We recall that we consider adic spaces defined over a perfectoid field K .

COROLLARY 3.6. *The following pairs of model categories are Quillen equivalent.*

- $\mathbf{Ch}_{\text{ét}}\mathbf{Psh}(\text{RigSm})$ and $\mathbf{Ch}_{\text{ét}}\mathbf{Psh}(\text{RigSm}^{\text{gc}})$.
- $\mathbf{Ch}_{\text{ét},\mathbb{B}^1}\mathbf{Psh}(\text{RigSm})$ and $\mathbf{Ch}_{\text{ét},\mathbb{B}^1}\mathbf{Psh}(\text{RigSm}^{\text{gc}})$.
- $\mathbf{Ch}_{\text{ét}}\mathbf{Psh}(\widehat{\text{RigSm}})$ and $\mathbf{Ch}_{\text{ét}}\mathbf{Psh}(\widehat{\text{RigSm}}^{\text{gc}})$.
- $\mathbf{Ch}_{\text{ét},\mathbb{B}^1}\mathbf{Psh}(\widehat{\text{RigSm}})$ and $\mathbf{Ch}_{\text{ét},\mathbb{B}^1}\mathbf{Psh}(\widehat{\text{RigSm}}^{\text{gc}})$.

Proof. It suffices to apply Proposition 3.3 to the sites with interval $(\text{RigSm}, \text{ét}, \mathbb{B}^1)$ and $(\widehat{\text{RigSm}}, \text{ét}, \mathbb{B}^1)$, where \mathbf{C}' is in both cases the subcategory of varieties with good coordinates. \square

DEFINITION 3.7. For $\eta \in \{\text{ét}, \mathbb{B}^1, (\text{ét}, \mathbb{B}^1)\}$ we say that a map in $\mathbf{ChPsh}(\text{RigSm})$ [respectively $\mathbf{ChPsh}(\widehat{\text{RigSm}})$] is a η -weak equivalence if it is a weak equivalence in the model structure

$\mathbf{Ch}_\eta\mathbf{Psh}(\mathbf{RigSm})$ [respectively $\mathbf{Ch}_\eta\mathbf{Psh}(\widehat{\mathbf{RigSm}})$]. The triangulated homotopy category associated to the localization $\mathbf{Ch}_{\acute{e}t, \mathbb{B}^1}\mathbf{Psh}(\mathbf{RigSm})$ [respectively to the localization $\mathbf{Ch}_{\acute{e}t, \mathbb{B}^1}\mathbf{Psh}(\widehat{\mathbf{RigSm}})$] is denoted by $\mathbf{RigDA}_{\acute{e}t}^{\text{eff}}(K, \Lambda)$ [respectively $\widehat{\mathbf{RigDA}}_{\acute{e}t, \mathbb{B}^1}^{\text{eff}}(K, \Lambda)$]. We omit Λ from the notation whenever the context allows it. The image of a variety X in one of these categories is denoted by $\Lambda(X)$. We say that an object \mathcal{F} of the derived category $\mathbf{D} = \mathbf{D}(\mathbf{Psh}(\mathbf{RigSm}))$ [respectively $\mathbf{D} = \mathbf{D}(\mathbf{Psh}(\widehat{\mathbf{RigSm}}))$] is η -local if the functor $\text{Hom}_{\mathbf{D}}(\cdot, \mathcal{F})$ sends maps in S_η (see Proposition 3.3) to isomorphisms. This amounts to saying that \mathcal{F} is quasi-isomorphic to a η -fibrant object.

We need to keep track of \mathbb{B}^1 in the notation of $\widehat{\mathbf{RigDA}}_{\acute{e}t, \mathbb{B}^1}^{\text{eff}}(K, \Lambda)$ since later we will perform a localization on $\mathbf{Ch}\mathbf{Psh}(\widehat{\mathbf{RigSm}})$ with respect to a different interval object.

Remark 3.8. Using the language of [BV08], the localizations defined above induce endofunctors C^η of the derived categories $\mathbf{D}(\mathbf{Psh}(\mathbf{RigSm}))$, $\mathbf{D}(\mathbf{Psh}(\mathbf{RigSm}^{\text{gc}}))$, $\mathbf{D}(\mathbf{Psh}(\widehat{\mathbf{RigSm}}))$ and $\mathbf{D}(\mathbf{Psh}(\widehat{\mathbf{RigSm}}^{\text{gc}}))$ such that $C^\eta\mathcal{F}$ is η -local for all \mathcal{F} and there is a natural transformation $C^\eta \rightarrow \text{id}$, which is a pointwise η -weak equivalence. The functor C^η restricts to a triangulated equivalence on the objects \mathcal{F} that are η -local and one can compute the Hom set $\text{Hom}(\mathcal{F}, \mathcal{F}')$ in the homotopy category of the η -localization as $\mathbf{D}(\mathcal{F}, C^\eta\mathcal{F}')$.

Remark 3.9. According to Proposition 3.3, for any X in $\widehat{\mathbf{RigSm}}$ and any integer i , one has $\text{Hom}_{\mathbf{D}}(\Lambda(X)[-i], C^{\acute{e}t}\mathcal{F}) \cong \text{Hom}_{\mathbf{D}(\mathbf{Sh}_{\acute{e}t}(\widehat{\mathbf{RigSm}}))}(\Lambda(X)[-i], \mathcal{F})$. The latter group is the étale hypercohomology group $\mathbb{H}_{\acute{e}t}^i(X, \mathcal{F})$, which can be computed with respect to the *small* étale site over X . The property $\text{Hom}_{\mathbf{D}}(\Lambda(X)[-i], C^{\acute{e}t}\mathcal{F}) \cong \mathbb{H}^i(X_{\acute{e}t}, \mathcal{F})$ characterizes $C^{\acute{e}t}\mathcal{F}$ up to quasi-isomorphisms (and holds true for more general topologies, see [Ayo07, Proposition 4.4.58]).

We now show that the étale localization can alternatively be described in terms of étale hypercoverings $\mathcal{U}_\bullet \rightarrow X$ (see for example [DHI04]). Any such datum defines a simplicial presheaf $n \mapsto \bigoplus_i \Lambda(U_{ni})$ whenever $U_n = \bigsqcup_i h_{U_{ni}}$ is the sum of the presheaves of sets $h_{U_{ni}}$ represented by U_{ni} . This simplicial presheaf can be associated to a normalized chain complex that we denote by $\Lambda(\mathcal{U}_\bullet)$. It is endowed with a map to $\Lambda(X)$.

PROPOSITION 3.10. *The localization over $S_{\acute{e}t}$ on $\mathbf{Ch}\mathbf{Psh}(\mathbf{RigSm}^{\text{gc}})$ [respectively $\mathbf{Ch}\mathbf{Psh}(\widehat{\mathbf{RigSm}}^{\text{gc}})$] coincides with the localization over the set $\Lambda(\mathcal{U}_\bullet)[i] \rightarrow \Lambda(X)[i]$ as $\mathcal{U}_\bullet \rightarrow X$ varies among bounded étale hypercoverings of the objects X of $\mathbf{RigSm}^{\text{gc}}$ [respectively $\widehat{\mathbf{RigSm}}^{\text{gc}}$] and i varies in \mathbb{Z} .*

Proof. Any ét-local object \mathcal{F} is also local with respect to the maps of the statement. We are left to prove that a complex \mathcal{F} , which is local with respect to the maps of the statement, is also ét-local.

Since Λ contains \mathbb{Q} the étale cohomology of an étale sheaf \mathcal{F} coincides with the Nisnevich cohomology (the same proof of [MVW06, Proposition 14.23] holds also here). By means of [Ayo15, Corollary 1.2.21] we conclude that any rigid variety X has a finite cohomological dimension. By [SGA4, Theorem V.7.4.1] and [SV00, Theorem 0.3], we obtain for any rigid variety X and any complex of presheaves \mathcal{F} an isomorphism

$$\mathbb{H}_{\acute{e}t}^n(X, \mathcal{F}) \cong \varinjlim_{\mathcal{U}_\bullet \in \text{HR}_\infty(X)} H_{-n}\text{Hom}_\bullet(\Lambda(\mathcal{U}_\bullet), \mathcal{F}),$$

where $HR_\infty(X)$ is the category of bounded étale hypercoverings of X (see [SGA4, V.7.3]) and Hom_\bullet is the Hom-complex computed in the unbounded derived category of presheaves. Suppose now \mathcal{F} is local with respect to the maps of the statement. Then $\text{Hom}_\bullet(\Lambda(\mathcal{U}_\bullet), \mathcal{F})$ is quasi-isomorphic to $\text{Hom}_\bullet(X, \mathcal{F})$ for every bounded hypercovering \mathcal{U}_\bullet , and hence $H_{-n}\mathcal{F}(X) \cong \mathbb{H}_{\text{ét}}^n(X, \mathcal{F})$ by the formula above. We then conclude that the map $\mathcal{F} \rightarrow C^{\text{ét}}\mathcal{F}$ is a quasi-isomorphism, proving the proposition. \square

As the following proposition shows, there are also alternative presentations of the homotopy categories introduced so far, which we will later use.

PROPOSITION 3.11. *Let Λ be a \mathbb{Q} -algebra. The natural inclusion induces a Quillen equivalence*

$$L_S\mathbf{Ch}(\mathbf{Psh}(\widehat{\text{RigSm}}^{\text{gc}})) \rightleftarrows \mathbf{Ch}_{\text{ét}}\mathbf{Psh}(\widehat{\text{RigSm}}^{\text{gc}}),$$

where L_S denotes the Bousfield localization with respect to the set S of shifts of the maps of complexes induced by étale Čech hypercoverings $\mathcal{U}_\bullet \rightarrow X$ of objects X in $\widehat{\text{RigSm}}^{\text{gc}}$ such that for some presentation $X = \varprojlim_h X_h$ the covering $\mathcal{U}_0 \rightarrow X$ descends to a covering of X_0 .

Proof. Using Proposition 3.10, it suffices to prove that the map $\Lambda(\mathcal{U}_\bullet) \rightarrow \Lambda(X)$ is an isomorphism in the homotopy category $L_S\mathbf{Ch}(\mathbf{Psh}(\widehat{\text{RigSm}}^{\text{gc}}))$ for a fixed bounded étale hypercovering \mathcal{U}_\bullet of an object X in $\widehat{\text{RigSm}}^{\text{gc}}$.

Since the inclusion functor $\mathbf{Ch}_{\geq 0} \rightarrow \mathbf{Ch}$ is a Quillen functor, it suffices to prove that $\Lambda(\mathcal{U}_\bullet) \rightarrow \Lambda(X)$ is a weak equivalence in $L_T\mathbf{Ch}_{\geq 0}(\mathbf{Psh}(\widehat{\text{RigSm}}^{\text{gc}}))$, where T is the set of shifts of the maps of complexes induced by étale Čech hypercoverings descending at finite level. Let $L_{\tilde{T}}\mathbf{sPsh}(\widehat{\text{RigSm}}^{\text{gc}})$ be the Bousfield localization of the projective model structure on simplicial presheaves of sets with respect to the set \tilde{T} formed by maps induced by étale Čech hypercoverings $\mathcal{U}_\bullet \rightarrow X$ descending at finite level. We remark that the Dold–Kan correspondence (see [SS03, §4.1]) and the Λ -enrichment also define a left Quillen functor from $L_{\tilde{T}}\mathbf{sPsh}(\widehat{\text{RigSm}}^{\text{gc}})$ to the category $L_T\mathbf{Ch}_{\geq 0}(\mathbf{Psh}(\widehat{\text{RigSm}}^{\text{gc}}))$. It, therefore, suffices to prove that $\mathcal{U}_\bullet \rightarrow X$ is a weak equivalence in $L_{\tilde{T}}\mathbf{sPsh}(\widehat{\text{RigSm}}^{\text{gc}})$ and this follows from the fact that bounded hypercoverings define the same localization as Čech hypercoverings (see [DHI04, Theorem A.6]) together with the fact that coverings descending to finite level define the same topology (Corollary 2.10), and hence the same localization [DHI04, Corollary A.8]. We remark that [DHI04, Corollary A.8] applies in our case even if the coverings $\mathcal{U} \rightarrow X$ descending to the finite level do not form a basis of the topology, as their pullback via an arbitrary map $Y \rightarrow X$ may not have the same property. However, the proof of the statement relies on [DHI04, Proposition A.2], where it is only used that the chosen family of coverings $\mathcal{U} \rightarrow X$ generates the topology and that the fiber product $\mathcal{U} \times_X \mathcal{U}$ is defined. \square

Remark 3.12. It is shown in the proof that the statements of Propositions 3.10 and 3.11 hold true without any assumptions on Λ under the condition that all varieties X have finite cohomological dimension with respect to the étale topology.

As we pointed out in Remark 3.9, there is a characterization of $C^{\text{ét}}\mathcal{F}$ for any complex \mathcal{F} . This is also true for the \mathbb{B}^1 -localization, described in the following part.

DEFINITION 3.13. We denote by \square the Σ -enriched cocubical object (see [Ayo14b, Appendix A]) defined by putting $\square^n = \mathbb{B}^n = \text{Spa } K \langle \tau_1, \dots, \tau_n \rangle$ and considering the morphisms $d_{r,\epsilon}$ induced

by the maps $\mathbb{B}^n \rightarrow \mathbb{B}^{n+1}$ corresponding to the substitution $\tau_r = \epsilon$ for $\epsilon \in \{0, 1\}$ and the morphisms p_r induced by the projections $\mathbb{B}^n \rightarrow \mathbb{B}^{n-1}$. For any variety X and any presheaf \mathcal{F} with values in an abelian category, we can therefore consider the Σ -enriched cubical object $\mathcal{F}(X \times \square)$ (see [Ayo14b, Appendix A]). Associated to any Σ -enriched cubical object \mathcal{F} there are the following complexes: the complex $C_{\bullet}^{\sharp}\mathcal{F}$ defined as $C_n^{\sharp}\mathcal{F} = \mathcal{F}_n$ and with differential $\sum(-1)^r(d_{r,1}^* - d_{r,0}^*)$; the *simple complex* $C_{\bullet}\mathcal{F}$ defined as $C_n\mathcal{F} = \bigcap_{r=1}^n \ker d_{r,0}^*$ and with differential $\sum(-1)^r d_{r,1}^*$; the *normalized complex* $N_{\bullet}\mathcal{F}$ defined as $N_n\mathcal{F} = C_n \cap \mathcal{F} \bigcap_{r=2}^n \ker d_{r,1}^*$ and with differential $-d_{1,1}^*$. By [Ayo14c, Lemma A.3, Propositions A.8, A.11], the inclusion $N_{\bullet}\mathcal{F} \hookrightarrow C_{\bullet}\mathcal{F}$ is a quasi-isomorphism and both inclusions $C_{\bullet}\mathcal{F} \hookrightarrow C_{\bullet}^{\sharp}\mathcal{F}$ and $N_{\bullet}\mathcal{F} \hookrightarrow C_{\bullet}\mathcal{F}$ split.

For any complex of presheaves \mathcal{F} we define the *singular complex of \mathcal{F}* , denoted by $\text{Sing}^{\mathbb{B}^1}\mathcal{F}$, to be the total complex of the simple complex associated to the $\underline{\text{Hom}}(\Lambda(\square), \mathcal{F})$. It sends the object X to the total complex of the simple complex associated to $\mathcal{F}(X \times \square)$.

The following lemma is the cocubical version of [MVW06, Lemma 2.18].

LEMMA 3.14. *For any presheaf \mathcal{F} the two maps of cubical sets $i_0^*, i_1^*: \mathcal{F}(\square \times \mathbb{B}^1) \rightarrow \mathcal{F}(\square)$ induce chain homotopic maps on the associated simple and normalized complexes.*

Proof. Consider the isomorphism $s_n: \mathbb{B}^{n+1} \rightarrow \mathbb{B}^n \times \mathbb{B}^1$ defined on points by separating the last coordinate and let s_n^* be the induced map $\mathcal{F}(\square^n \times \mathbb{B}^1) \rightarrow \mathcal{F}(\square^{n+1})$. We have $s_{n-1}^* \circ d_{r,\epsilon}^* = d_{r,\epsilon}^* \circ s_n^*$ for all $1 \leq r \leq n$ and $\epsilon \in \{0, 1\}$. We conclude that

$$\begin{aligned} & s_{n-1}^* \circ \sum_{r=1}^n (-1)^r (d_{r,1}^* - d_{r,0}^*) + \sum_{r=1}^{n+1} (-1)^r (d_{r,1}^* - d_{r,0}^*) \circ (-s_n^*) \\ &= (-1)^n (d_{n+1,1}^* \circ s_n^* - d_{n+1,0}^* \circ s_n^*) = (-1)^n (i_1^* - i_0^*). \end{aligned}$$

Therefore, the maps $\{(-1)^n s_n^*\}$ define a chain homotopy from i_0^* to i_1^* as maps of complexes $C_{\bullet}^{\sharp}\mathcal{F}(\square \times \mathbb{B}^1) \rightarrow C_{\bullet}^{\sharp}\mathcal{F}(\square)$.

We automatically deduce that if an inclusion $C'_{\bullet}\mathcal{F} \rightarrow C_{\bullet}^{\sharp}\mathcal{F}$ has a functorial retraction, then the maps $i_0^*, i_1^*: C'_{\bullet}\mathcal{F}(\square \times \mathbb{B}^1) \rightarrow C'_{\bullet}\mathcal{F}(\square)$ are also chain homotopic. \square

The following proposition is the rigid analytic analog of [Ayo14b, Theorem 2.23], or the cocubical analog of [Ayo15, Lemma 2.5.31].

PROPOSITION 3.15. *Let \mathcal{F} be a complex in $\mathbf{Ch Psh}(\widehat{\text{RigSm}})$. Then $\text{Sing}^{\mathbb{B}^1}\mathcal{F}$ is \mathbb{B}^1 -local and \mathbb{B}^1 -weak equivalent to \mathcal{F} in $\mathbf{Ch Psh}(\widehat{\text{RigSm}})$.*

Proof. In order to prove that $\text{Sing}^{\mathbb{B}^1}\mathcal{F}$ is \mathbb{B}^1 -local in $\mathbf{Ch Psh}(\widehat{\text{RigSm}})$ we need to check that each homology presheaf $H_n(\text{Sing}^{\mathbb{B}^1}\mathcal{F})$ is homotopy-invariant. By means of [Ayo15, Proposition 2.2.46] it suffices to show that the maps $i_0^*, i_1^*: N_{\bullet}\mathcal{F}(\square \times \mathbb{B}^1) \rightarrow N_{\bullet}\mathcal{F}(\square)$ are chain homotopic, and this follows from Lemma 3.14.

We now prove that $\text{Sing}^{\mathbb{B}^1}\mathcal{F}$ is \mathbb{B}^1 -weak equivalent to \mathcal{F} . We first prove that the canonical map $a: \mathcal{F} \rightarrow \underline{\text{Hom}}(\Lambda(\square^n), \mathcal{F})$ has an inverse up to homotopy for a fixed n . Consider the map $b: \underline{\text{Hom}}(\Lambda(\square^n), \mathcal{F}) \rightarrow \mathcal{F}$ induced by the zero section of \square^n . It holds that $b \circ a = \text{id}$ and $a \circ b$ is homotopic to id via the map

$$H: \Lambda(\mathbb{B}^1) \otimes \underline{\text{Hom}}(\Lambda(\square^n), \mathcal{F}) \rightarrow \underline{\text{Hom}}(\Lambda(\square^n), \mathcal{F}),$$

which is deduced from the adjunction $(\Lambda(\mathbb{B}^1) \otimes \cdot, \underline{\text{Hom}}(\Lambda(\mathbb{B}^1), \cdot))$ and the map

$$\underline{\text{Hom}}(\Lambda(\square^n), \mathcal{F}) \rightarrow \underline{\text{Hom}}(\Lambda(\mathbb{B}^1 \times \square^n), \mathcal{F})$$

defined via the homothety of \mathbb{B}^1 on \square^n . As \mathbb{B}^1 -weak equivalences are stable under filtered colimits and cones, we also conclude that the total complex associated to the simple complex of $\underline{\text{Hom}}(\Lambda(\square), \mathcal{F})$ is \mathbb{B}^1 -equivalent to the one associated to the constant cubical object \mathcal{F} (see, for example, the argument of [Ayo15, Corollary 2.5.36]), which is in turn quasi-isomorphic to \mathcal{F} . \square

COROLLARY 3.16. *Let Λ be a \mathbb{Q} -algebra. For any \mathcal{F} in $\mathbf{Ch Psh}(\widehat{\text{RigSm}})$ the localization $C^{\mathbb{B}^1} \mathcal{F}$ is quasi-isomorphic to $\text{Sing}^{\mathbb{B}^1} \mathcal{F}$ and the localization $C^{\text{ét}, \mathbb{B}^1} \mathcal{F}$ is quasi-isomorphic to $\text{Sing}^{\mathbb{B}^1}(C^{\text{ét}} \mathcal{F}^\bullet)$.*

Proof. The first claim follows from Proposition 3.15. We are left to prove that the complex $\text{Sing}^{\mathbb{B}^1}(C^{\text{ét}} \mathcal{F}^\bullet)$ is ét-local. To this aim, we use the description given in Proposition 3.10 and we show that $\text{Sing}^{\mathbb{B}^1}(C^{\text{ét}} \mathcal{F}^\bullet)$ is local with respect to shifts of maps $\Lambda(\mathcal{U}_\bullet) \rightarrow \Lambda(X)$ induced by bounded hypercoverings $\mathcal{U}_\bullet \rightarrow X$.

Fix such a hypercovering $\mathcal{U}_\bullet \rightarrow X$. From the isomorphisms

$$H_p \text{Hom}_\bullet(\Lambda(\mathcal{U}_\bullet \times \square^q), C^{\text{ét}} \mathcal{F}) \cong H_p \text{Hom}_\bullet(\Lambda(X \times \square^q), C^{\text{ét}} \mathcal{F})$$

valid for all p, q and a spectral sequence argument (see [SV00, Theorem 0.3]) we deduce that

$$\mathbf{D}(\Lambda(X)[n], \text{Sing}^{\mathbb{B}^1} C^{\text{ét}} \mathcal{F}) \cong \mathbf{D}(\Lambda(\mathcal{U}_\bullet)[n], \text{Sing}^{\mathbb{B}^1} C^{\text{ét}} \mathcal{F})$$

for all n as wanted. \square

We now investigate some of the natural Quillen functors that arise between the model categories introduced so far. We start by considering the natural inclusion of categories $\text{RigSm} \rightarrow \widehat{\text{RigSm}}$

PROPOSITION 3.17. *The inclusion $\text{RigSm} \hookrightarrow \widehat{\text{RigSm}}$ induces a Quillen adjunction*

$$\iota^* : \mathbf{Ch}_{\text{ét}, \mathbb{B}^1} \mathbf{Psh}(\text{RigSm}) \rightleftarrows \mathbf{Ch}_{\text{ét}, \mathbb{B}^1} \mathbf{Psh}(\widehat{\text{RigSm}}) : \iota_*$$

Moreover, the functor $\mathbb{L}\iota^* : \mathbf{RigDA}_{\text{ét}}^{\text{eff}}(K) \rightarrow \widehat{\mathbf{RigDA}}_{\text{ét}, \mathbb{B}^1}^{\text{eff}}(K)$ is fully faithful.

Proof. The first claim is a special instance of [Ayo07, Proposition 4.4.45].

We prove the second claim by showing that $\mathbb{R}\iota_* \mathbb{L}\iota^*$ is isomorphic to the identity. Let \mathcal{F} be a cofibrant object in $\mathbf{Ch}_{\text{ét}, \mathbb{B}^1} \mathbf{Psh}(\text{RigSm})$. We need to prove that the map $\mathcal{F} \rightarrow \iota_*(\text{Sing}^{\mathbb{B}^1} C^{\text{ét}}(\iota^* \mathcal{F}))$ is an $(\text{ét}, \mathbb{B}^1)$ -weak equivalence. Since ι_* commutes with $\text{Sing}^{\mathbb{B}^1}$ we are left to prove that the map $\iota_* \iota^* \mathcal{F} = \mathcal{F} \rightarrow \iota_* C^{\text{ét}}(\iota^* \mathcal{F})$ is an ét-weak equivalence. This follows since ι_* preserves ét-weak equivalences, as it commutes with ét-sheafification. \square

We are now interested in finding a convenient set of compact objects that generate the categories above, as triangulated categories with small sums. This will simplify many definitions and proofs in what follows.

PROPOSITION 3.18. *The category $\mathbf{RigDA}_{\text{ét}}^{\text{eff}}(K)$ [respectively $\widehat{\mathbf{RigDA}}_{\text{ét}, \mathbb{B}^1}^{\text{eff}}(K)$] is compactly generated (as a triangulated category with small sums) by motives $\Lambda(X)$ associated to rigid varieties X , which are in RigSm^{gc} [respectively $\widehat{\text{RigSm}}^{\text{gc}}$].*

Proof. The statements are analogous, and we only consider the case of $\widehat{\mathbf{RigDA}}_{\acute{e}t, \mathbb{B}^1}^{\text{eff}}(K)$. It is clear that the set of functors $H_i \text{Hom}_{\bullet}(\Lambda(X), \cdot)$ detect quasi-isomorphisms between étale local objects, by letting X vary in $\widehat{\mathbf{RigSm}}^{\text{gc}}$ and i vary in \mathbb{Z} . We are left to prove that the motives $\Lambda(X)$ with X in $\widehat{\mathbf{RigSm}}^{\text{gc}}$ are compact. Since $\Lambda(X)$ is compact in $\mathbf{D}(\mathbf{Psh}(\widehat{\mathbf{RigSm}}^{\text{gc}}))$ and $\text{Sing}^{\mathbb{B}^1}$ commutes with direct sums, it suffices to prove that if $\{\mathcal{F}_i\}_{i \in I}$ is a family of ét-local complexes, then also $\bigoplus_i \mathcal{F}_i$ is ét-local. If I is finite, the claim follows from the isomorphisms $H_{-n} \text{Hom}_{\bullet}(X, \bigoplus_i \mathcal{F}_i) \cong \bigoplus_i H_{-n} \text{Hom}_{\bullet}(X, \mathcal{F}_i) \cong \mathbb{H}^n(X, \bigoplus_i \mathcal{F}_i)$. A coproduct over an arbitrary family is a filtered colimit of finite coproducts, and hence the claim follows from [Ayo07, Proposition 4.5.62]. \square

Remark 3.19. The above proof shows that the statement of Proposition 3.18 holds true without any assumptions on Λ under the condition that all varieties X have finite cohomological dimension with respect to the étale topology.

We now introduce the category of motives associated to smooth perfectoid spaces, using the same formalism as before. In this category, the canonical choice of the ‘interval object’ for defining homotopies is the perfectoid ball $\widehat{\mathbb{B}}^1$.

Example 3.20. The perfectoid ball $\widehat{\mathbb{B}}^1 = \text{Spa}(K\langle\chi^{1/p^\infty}\rangle, K^\circ\langle\chi^{1/p^\infty}\rangle)$ is an interval object with respect to the natural multiplication μ and maps i_0 and i_1 induced by the substitution $\chi^{1/p^h} \mapsto 0$ and $\chi^{1/p^h} \mapsto 1$, respectively.

The perfectoid variety $\widehat{\mathbb{B}}^1$ naturally lives in $\widehat{\mathbf{RigSm}}$ and has good coordinates by Proposition 2.13. It can therefore be used to define another homotopy category out of $\mathbf{ChPsh}(\widehat{\mathbf{RigSm}})$ and $\mathbf{ChPsh}(\widehat{\mathbf{RigSm}}^{\text{gc}})$.

COROLLARY 3.21. *The following pairs of model categories are Quillen equivalent.*

- $\mathbf{Ch}_{\acute{e}t} \mathbf{Psh}(\text{PerfSm})$ and $\mathbf{Ch}_{\acute{e}t} \mathbf{Psh}(\text{PerfSm}^{\text{gc}})$.
- $\mathbf{Ch}_{\acute{e}t, \widehat{\mathbb{B}}^1} \mathbf{Psh}(\text{PerfSm})$ and $\mathbf{Ch}_{\acute{e}t, \widehat{\mathbb{B}}^1} \mathbf{Psh}(\text{PerfSm}^{\text{gc}})$.
- $\mathbf{Ch}_{\acute{e}t} \mathbf{Psh}(\widehat{\mathbf{RigSm}})$ and $\mathbf{Ch}_{\acute{e}t} \mathbf{Psh}(\widehat{\mathbf{RigSm}}^{\text{gc}})$.
- $\mathbf{Ch}_{\acute{e}t, \widehat{\mathbb{B}}^1} \mathbf{Psh}(\widehat{\mathbf{RigSm}})$ and $\mathbf{Ch}_{\acute{e}t, \widehat{\mathbb{B}}^1} \mathbf{Psh}(\widehat{\mathbf{RigSm}}^{\text{gc}})$.

Proof. It suffices to apply Proposition 3.3 to the sites with interval $(\text{PerfSm}, \acute{e}t, \widehat{\mathbb{B}}^1)$ and $(\widehat{\mathbf{RigSm}}, \acute{e}t, \widehat{\mathbb{B}}^1)$, where \mathbf{C}' is in both cases the subcategory of affinoid rigid varieties with good coordinates. \square

DEFINITION 3.22. For $\eta \in \{\acute{e}t, \widehat{\mathbb{B}}^1, (\acute{e}t, \widehat{\mathbb{B}}^1)\}$ we say that a map in $\mathbf{ChPsh}(\text{PerfSm})$ [respectively $\mathbf{ChPsh}(\widehat{\mathbf{RigSm}})$] is a η -weak equivalence if it is a weak equivalence in the model structure $\mathbf{Ch}_\eta \mathbf{Psh}(\text{PerfSm})$ [respectively $\mathbf{Ch}_\eta \mathbf{Psh}(\widehat{\mathbf{RigSm}})$]. We say that an object \mathcal{F} of the derived category $\mathbf{D} = \mathbf{D}(\mathbf{Psh}(\text{PerfSm}))$ [respectively $\mathbf{D} = \mathbf{D}(\mathbf{Psh}(\widehat{\mathbf{RigSm}}))$] is η -local if the functor $\text{Hom}_{\mathbf{D}}(\cdot, \mathcal{F})$ sends maps in S_η (see Proposition 3.3) to isomorphisms. This amounts to say that \mathcal{F} is quasi-isomorphic to a η -fibrant object. The triangulated homotopy category associated to the localization $\mathbf{Ch}_{\acute{e}t, \widehat{\mathbb{B}}^1} \mathbf{Psh}(\text{PerfSm})$ [respectively $\mathbf{Ch}_{\acute{e}t, \widehat{\mathbb{B}}^1} \mathbf{Psh}(\widehat{\mathbf{RigSm}})$] will be denoted by $\mathbf{PerfDA}_{\acute{e}t}^{\text{eff}}(K, \Lambda)$ [respectively $\widehat{\mathbf{RigDA}}_{\acute{e}t, \widehat{\mathbb{B}}^1}^{\text{eff}}(K, \Lambda)$]. We will omit Λ whenever the context allows it. The image of a variety X in one of these categories will be denoted by $\Lambda(X)$.

We recall one of the main results of Scholze [Sch12], reshaped in our derived homotopical setting. It will constitute the bridge to pass from characteristic p to characteristic 0. As summarized in Theorem 1.12 there is an equivalence of categories between perfectoid affinoid K -algebras and perfectoid affinoid K^b -algebras, extending to an equivalence between the categories of perfectoid spaces over K and over K^b (see [Sch12, Proposition 6.17]). We refer to this equivalence as the *tilting equivalence*.

PROPOSITION 3.23. *There exists an equivalence of triangulated categories*

$$(-)^\sharp : \mathbf{PerfDA}_{\text{ét}}^{\text{eff}}(K^b) \rightleftarrows \mathbf{PerfDA}_{\text{ét}}^{\text{eff}}(K) : (-)^b$$

induced by the tilting equivalence.

Proof. The tilting equivalence (see Theorem 1.12) induces an equivalence of the étale sites on perfectoid spaces over K and over K^b (see [Sch12, Theorem 7.12]). Moreover $(\widehat{\mathbb{T}^n})^b = \widehat{\mathbb{T}^n}$ and $(\widehat{\mathbb{B}^n})^b = \widehat{\mathbb{B}^n}$. It therefore induces an equivalence of sites with interval $(\mathbf{PerfSm}/K, \text{ét}, \widehat{\mathbb{B}^1}) \cong (\mathbf{PerfSm}/K^b, \text{ét}, \widehat{\mathbb{B}^1})$, and hence the claim. \square

We now investigate the triangulated functor between the categories of motives induced by the natural embedding $\mathbf{PerfSm} \rightarrow \widehat{\mathbf{RigSm}}$ in the same spirit of what we did previously in Proposition 3.17.

PROPOSITION 3.24. *The inclusion $\mathbf{PerfSm} \hookrightarrow \widehat{\mathbf{RigSm}}$ induces a Quillen adjunction*

$$j^* : \mathbf{Ch}_{\text{ét}, \widehat{\mathbb{B}^1}} \mathbf{Psh}(\mathbf{PerfSm}) \rightleftarrows \mathbf{Ch}_{\text{ét}, \widehat{\mathbb{B}^1}} \mathbf{Psh}(\widehat{\mathbf{RigSm}}) : j_*$$

Moreover, the functor $\mathbb{L}j^* : \mathbf{PerfDA}_{\text{ét}}^{\text{eff}}(K) \rightarrow \widehat{\mathbf{RigDA}}_{\text{ét}, \widehat{\mathbb{B}^1}}^{\text{eff}}(K)$ is fully faithful.

Proof. The result follows in the same way as Proposition 3.17. \square

Also in this framework, the $\widehat{\mathbb{B}^1}$ -localization has a very explicit construction. Most proofs are straightforward analogs of those relative to the \mathbb{B}^1 -localizations, and will therefore be omitted.

DEFINITION 3.25. We denote by $\widehat{\square}$ the Σ -enriched cocubical object (see [Ayo14c, Appendix A]) defined by putting $\widehat{\square}^n = \widehat{\mathbb{B}^n} = \text{Spa } K \langle \tau_1^{1/p^\infty}, \dots, \tau_n^{1/p^{1/\infty}} \rangle$ and considering the morphisms $d_{r,\epsilon}$ induced by the maps $\widehat{\mathbb{B}^n} \rightarrow \widehat{\mathbb{B}^{n+1}}$ corresponding to the substitution $\tau_r^{1/p^h} = \epsilon$ for $\epsilon \in \{0, 1\}$ and the morphisms p_r induced by the projections $\widehat{\mathbb{B}^n} \rightarrow \widehat{\mathbb{B}^{n-1}}$. For any complex of presheaves \mathcal{F} we let $\text{Sing}^{\widehat{\mathbb{B}^1}} \mathcal{F}$ be the total complex of the simple complex associated to $\underline{\text{Hom}}(\widehat{\square}, \mathcal{F})$. It sends the object X to the total complex of the simple complex associated to $\mathcal{F}(X \times \widehat{\square})$.

PROPOSITION 3.26. *Let \mathcal{F} be a complex in $\mathbf{Ch Psh}(\mathbf{PerfSm})$ [respectively in $\mathbf{Ch Psh}(\widehat{\mathbf{RigSm}})$]. Then $\text{Sing}^{\widehat{\mathbb{B}^1}} \mathcal{F}$ is $\widehat{\mathbb{B}^1}$ -local and $\widehat{\mathbb{B}^1}$ -weak equivalent to \mathcal{F} .*

Proof. The fact that $\text{Sing}^{\widehat{\mathbb{B}^1}} \mathcal{F}$ is $\widehat{\mathbb{B}^1}$ -local in $\mathbf{Ch Psh}(\widehat{\mathbf{RigSm}})$ can be deduced by Lemmas 3.27 and 3.28. We are left to prove that $\text{Sing}^{\widehat{\mathbb{B}^1}} \mathcal{F}$ is $\widehat{\mathbb{B}^1}$ -weak equivalent to \mathcal{F} and this follows in the same way as in the proof of Proposition 3.15. \square

The following lemmas are used in the previous proof.

LEMMA 3.27. A presheaf \mathcal{F} in $\mathbf{Psh}(\mathrm{Sm}\mathrm{Perf})$ [respectively in $\mathbf{Psh}(\widehat{\mathrm{RigSm}})]$ is \mathbb{B}^1 -invariant if and only if $i_0^* = i_1^* : \mathcal{F}(X \times \mathbb{B}^1) \rightarrow \mathcal{F}(X)$ for all X in $\mathrm{Sm}\mathrm{Perf}$ [respectively in $\widehat{\mathrm{RigSm}}$].

Proof. This follows in the same way as [MVW06, Lemma 2.16]. □

LEMMA 3.28. For any presheaf \mathcal{F} the two maps of cubical sets $i_0^*, i_1^* : \mathcal{F}(\widehat{\square} \times \widehat{\mathbb{B}}^1) \rightarrow \mathcal{F}(\widehat{\square})$ induce chain homotopic maps on the associated simple and normalized complexes.

Proof. This follows in the same way as Lemma 3.14. □

COROLLARY 3.29. Let \mathcal{F} be in $\mathbf{Ch}\mathbf{Psh}(\mathrm{PerfSm})$ [respectively in $\mathbf{Ch}\mathbf{Psh}(\widehat{\mathrm{RigSm}})]$ the $(\acute{e}t, \mathbb{B}^1)$ -localization $C^{\acute{e}t, \mathbb{B}^1} \mathcal{F}$ is quasi-isomorphic to $\mathrm{Sing}^{\mathbb{B}^1}(C^{\acute{e}t} \mathcal{F})$.

Proof. This follows in the same way as Corollary 3.16. □

PROPOSITION 3.30. The category $\mathbf{PerfDA}_{\acute{e}t}^{\mathrm{eff}}(K)$ [respectively $\widehat{\mathbf{RigDA}}_{\acute{e}t, \mathbb{B}^1}^{\mathrm{eff}}(K)$] is compactly generated (as a triangulated category with small sums) by motives $\Lambda(X)$ associated to rigid varieties X , which are in $\mathrm{PerfSm}^{\mathrm{gc}}$ [respectively $\widehat{\mathrm{RigSm}}^{\mathrm{gc}}$].

Proof. This follows in the same way as Proposition 3.18. □

Remark 3.31. The above proof shows that the statement of Proposition 3.30 holds true without any assumptions on Λ under the condition that all varieties X have finite cohomological dimension with respect to the étale topology.

So far, we have defined two different Bousfield localizations on complexes of presheaves on $\widehat{\mathrm{RigSm}}$ according to two different choices of intervals: \mathbb{B}^1 and $\widehat{\mathbb{B}}^1$. We remark that the second constitutes a further localization of the first, in the following sense.

PROPOSITION 3.32. \mathbb{B}^1 -weak equivalences in $\mathbf{Ch}\mathbf{Psh}(\widehat{\mathrm{RigSm}})$ are $\widehat{\mathbb{B}}^1$ -weak equivalences.

Proof. It suffices to prove that $X \times \mathbb{B}^1 \rightarrow X$ induces a $\widehat{\mathbb{B}}^1$ -weak equivalence, for any variety X in $\widehat{\mathrm{RigSm}}$. This follows as the multiplicative homothety $\widehat{\mathbb{B}}^1 \times \mathbb{B}^1 \rightarrow \mathbb{B}^1$ induces a homotopy between the zero map and the identity on \mathbb{B}^1 . □

COROLLARY 3.33. The triangulated category $\widehat{\mathbf{RigDA}}_{\acute{e}t, \widehat{\mathbb{B}}^1}^{\mathrm{eff}}(K)$ is equivalent to the full triangulated subcategory of $\widehat{\mathbf{RigDA}}_{\acute{e}t, \mathbb{B}^1}^{\mathrm{eff}}(K)$ formed by $\widehat{\mathbb{B}}^1$ -local objects.

Proof. Because of Proposition 3.32, the triangulated category $\widehat{\mathbf{RigDA}}_{\acute{e}t, \widehat{\mathbb{B}}^1}^{\mathrm{eff}}(K)$ coincides with the localization of $\widehat{\mathbf{RigDA}}_{\acute{e}t, \mathbb{B}^1}^{\mathrm{eff}}(K)$ with respect to the set generated by the maps $\Lambda(\widehat{\mathbb{B}}^1_X)[n] \rightarrow \Lambda(X)[n]$ as X varies in $\widehat{\mathrm{RigSm}}$ and n in \mathbb{Z} . □

We end this section by recalling the definition of rigid motives with transfers. The notion of finite correspondence plays an important role in Voevodsky’s theory of motives. In the case of rigid varieties over a field K correspondences give rise to the category $\mathrm{RigCor}(K)$ as defined in [Ayo15, Definition 2.2.29].

DEFINITION 3.34. Additive presheaves over $\text{RigCor}(K)$ are called *presheaves with transfers*, and the category they form is denoted by $\mathbf{PST}(\text{RigSm}/K, \Lambda)$ or simply by $\mathbf{PST}(\text{RigSm})$ when the context allows it.

By [Ayo15, Definition 2.5.15], the projective model category $\mathbf{ChPST}(\text{RigSm})$ admits a Bousfield localization $\mathbf{Ch}_{\text{ét}, \mathbb{B}^1} \mathbf{PST}(\text{RigSm})$ with respect to the union of the class of maps $\mathcal{F} \rightarrow \mathcal{F}'$ inducing isomorphisms on the ét-sheaves associated to $H_i(\mathcal{F})$ and $H_i(\mathcal{F}')$ for all $i \in \mathbb{Z}$ and the set of all maps $\Lambda(\mathbb{B}_X^1)[i] \rightarrow \Lambda(X)[i]$ as X varies in RigSm and i varies in \mathbb{Z} .

DEFINITION 3.35. The triangulated homotopy category associated to $\mathbf{Ch}_{\text{ét}, \mathbb{B}^1} \mathbf{PST}(\text{RigSm})$ will be denoted by $\mathbf{RigDM}_{\text{ét}}^{\text{eff}}(K, \Lambda)$. We will omit Λ from the notation whenever the context allows it. The image of a variety X in will be denoted by $\Lambda_{\text{tr}}(X)$.

Remark 3.36. Since Λ is a \mathbb{Q} -algebra, one can equivalently consider the Nisnevich topology in the definition above and obtain a homotopy category $\mathbf{RigDM}_{\text{Nis}}^{\text{eff}}(K, \Lambda)$, which is equivalent to $\mathbf{RigDM}_{\text{ét}}^{\text{eff}}(K, \Lambda)$.

Remark 3.37. The faithful embedding of categories $\text{RigSm} \rightarrow \text{RigCor}$ induces a Quillen adjunction (see [Ayo15, Proposition 2.5.19]):

$$a_{\text{tr}} : \mathbf{Ch}_{\text{ét}, \mathbb{B}^1} \mathbf{Psh}(\text{RigSm}) \rightleftarrows \mathbf{Ch}_{\text{ét}, \mathbb{B}^1} \mathbf{PST}(\text{RigSm}) : o_{\text{tr}},$$

such that $a_{\text{tr}}\Lambda(X) = \Lambda_{\text{tr}}(X)$ for any $X \in \text{RigSm}$ and o_{tr} is the functor of forgetting transfers. These functors induce an adjoint pair:

$$\mathbb{L}a_{\text{tr}} : \mathbf{RigDA}_{\text{ét}}^{\text{eff}}(K) \rightleftarrows \mathbf{RigDM}_{\text{ét}}^{\text{eff}}(K) : \mathbb{R}o_{\text{tr}},$$

which is investigated in [Vez17].

4. Motivic interpretation of approximation results

Throughout this section, K is a perfectoid field of arbitrary characteristic. We begin by presenting an approximation result whose proof is postponed to Appendix A.

PROPOSITION 4.1. Let $X = \varprojlim_h X_h$ be in $\widehat{\text{RigSm}}^{\text{gc}}$. Also let Y be an affinoid rigid variety endowed with an étale map $Y \rightarrow \mathbb{B}^m$. For a given finite set of maps $\{f_1, \dots, f_N\}$ in $\text{Hom}(X \times \mathbb{B}^n, Y)$ we can find corresponding maps $\{H_1, \dots, H_N\}$ in $\text{Hom}(X \times \mathbb{B}^n \times \mathbb{B}^1, Y)$ and an integer \bar{h} such that the following are true.

- (i) For all $1 \leq k \leq N$ it holds that $i_0^* H_k = f_k$ and $i_1^* H_k$ factors over the canonical map $X \rightarrow X_{\bar{h}}$.
- (ii) If $f_k \circ d_{r,\epsilon} = f_{k'} \circ d_{r,\epsilon}$ for some $1 \leq k, k' \leq N$ and some $(r, \epsilon) \in \{1, \dots, n\} \times \{0, 1\}$ then $H_k \circ d_{r,\epsilon} = H_{k'} \circ d_{r,\epsilon}$.
- (iii) If for some $1 \leq k \leq N$ and some $h \in \mathbb{N}$ the map $f_k \circ d_{1,1} \in \text{Hom}(X \times \mathbb{B}^{n-1}, Y)$ lies in $\text{Hom}(X_h \times \mathbb{B}^{n-1}, Y)$ then the element $H_k \circ d_{1,1}$ of $\text{Hom}(X \times \mathbb{B}^{n-1} \times \mathbb{B}^1, Y)$ is constant on \mathbb{B}^1 equal to $f_k \circ d_{1,1}$.

The statement above has the following interpretation in terms of complexes.

PROPOSITION 4.2. Let $X = \varprojlim_h X_h$ be in $\widehat{\text{RigSm}}^{\text{gc}}$ and let Y be in RigSm^{gc} . The natural map

$$\phi : \varinjlim_h (\text{Sing}^{\mathbb{B}^1} \Lambda(Y))(X_h) \rightarrow (\text{Sing}^{\mathbb{B}^1} \Lambda(Y))(X)$$

is a quasi-isomorphism.

Proof. We need to prove that the natural map

$$\phi : \varinjlim_h C_\bullet \Lambda \text{Hom}(X_h \times \square, Y) \rightarrow C_\bullet \Lambda \text{Hom}(X \times \square, Y)$$

defines bijections on homology groups.

We start by proving surjectivity. As \square is a Σ -enriched cocubical object, the complexes above are quasi-isomorphic to the associated normalized complexes N_\bullet , which we consider instead. Suppose that $\beta \in \Lambda \text{Hom}(X \times \square^n, Y)$ defines a cycle in N_n , i.e., $\beta \circ d_{r,\epsilon} = 0$ for $1 \leq r \leq n$ and $\epsilon \in \{0, 1\}$. This means that $\beta = \sum \lambda_k f_k$ with $\lambda_k \in \Lambda$, $f_k \in \text{Hom}(X \times \square^n, Y)$ and $\sum \lambda_k f_k \circ d_{r,\epsilon} = 0$. This amounts to saying that for every k, r, ϵ the sum $\sum \lambda_{k'}$ over the indices k' such that $f_{k'} \circ d_{r,\epsilon} = f_k \circ d_{r,\epsilon}$ is zero. By Proposition 4.1, we can find an integer h and maps $H_k \in \text{Hom}(X \times \square^n \times \mathbb{B}^1, Y)$ such that $i_0^* H = f_k$, $i_1^* H = \phi(f_k)$ with $\tilde{f}_k \in \text{Hom}(X_h \times \square^n, Y)$ and $H_k \circ d_{r,\epsilon} = H_{k'} \circ d_{r,\epsilon}$ whenever $f_k \circ d_{r,\epsilon} = f_{k'} \circ d_{r,\epsilon}$. If we denote by H the cycle $\sum \lambda_k H_k \in \Lambda \text{Hom}(X \times \square^n \times \mathbb{B}^1, Y)$ we therefore have $d_{r,\epsilon}^* H = 0$ for all r, ϵ .

By Lemma 3.14, we conclude that $i_1^* H$ and $i_0^* H$ define the same homology class, and therefore β defines the same class as $i_1^* H$, which is the image of a class in $\Lambda \text{Hom}(X_h \times \square^n, Y)$, as desired.

We now turn to injectivity. Consider an element $\alpha \in \Lambda \text{Hom}(X_0 \times \square^n, Y)$ such that $\alpha \circ d_{r,\epsilon} = 0$ for all r, ϵ and suppose there exists an element $\beta = \sum \lambda_i f_i \in \Lambda \text{Hom}(X \times \square^{n+1}, Y)$ such that $\beta \circ d_{r,0} = 0$ for $1 \leq r \leq n+1$, $\beta \circ d_{r,1} = 0$ for $2 \leq r \leq n+1$ and $\beta \circ d_{1,1} = \phi(\alpha)$. Again, by Proposition 4.1, we can find an integer \bar{h} and maps $H_k \in \text{Hom}(X \times \square^{n+1} \times \mathbb{B}^1, Y)$ such that $H := \sum \lambda_k H_k$ satisfies $i_1^* H = \phi(\gamma)$ for some $\gamma \in \Lambda \text{Hom}(X_{\bar{h}} \times \square^{n+1}, Y)$, $H \circ d_{r,0} = 0$ for $1 \leq r \leq n+1$, $H \circ d_{r,1} = 0$ for $2 \leq r \leq n+1$ and $H \circ d_{1,1}$ is constant on \mathbb{B}^1 and coincides with $\phi(\alpha)$. We conclude that $\gamma \in N_n$ and $d\gamma = \alpha$. In particular, $\alpha = 0$ in the homology group, as desired. \square

COROLLARY 4.3. Let \mathcal{F} be a projectively cofibrant complex in $\mathbf{Ch Psh}(\text{RigSm}^{\text{gc}})$. For any $X = \varprojlim_h X_h$ in $\widehat{\text{RigSm}}^{\text{gc}}$ the natural map

$$\phi : \varinjlim_h (\text{Sing}^{\mathbb{B}^1} \mathcal{F})(X_h) \rightarrow (\text{Sing}^{\mathbb{B}^1} \iota^* \mathcal{F})(X)$$

is a quasi-isomorphism.

Proof. As homology commutes with filtered colimits, by means of Remark 3.2 we can assume that \mathcal{F} is a bounded above complex formed by sums of representable presheaves. For any X in $\widehat{\text{RigSm}}$ the homology of $\text{Sing}^{\mathbb{B}^1} \mathcal{F}(X)$ coincides with the homology of the total complex associated to $C_\bullet(\mathcal{F}(X \times \square))$. The result then follows from Proposition 4.2 and the convergence of the spectral sequence associated to the double complex above, which is concentrated in one quadrant. \square

The following technical proposition is actually a crucial point of our proof, as it allows some explicit computations of morphisms in the category $\widehat{\mathbf{RigDA}}_{\text{ét}}^{\text{eff}}(K)$.

PROPOSITION 4.4. *Let \mathcal{F} be a cofibrant and $(\mathbb{B}^1, \text{ét})$ -fibrant complex in $\mathbf{Ch Psh}(\widehat{\text{RigSm}}^{\text{gc}})$. Then $\text{Sing}^{\mathbb{B}^1}(\iota^*\mathcal{F})$ is $(\mathbb{B}^1, \text{ét})$ -local in $\mathbf{Ch Psh}(\widehat{\text{RigSm}}^{\text{gc}})$.*

Proof. The difficulty lies in showing that the object $\text{Sing}^{\mathbb{B}^1}(\iota^*\mathcal{F})$ is ét-local. By Propositions 3.11 and 3.15, it suffices to prove that $\text{Sing}^{\mathbb{B}^1}(\iota^*\mathcal{F})$ is local with respect to the étale-Cech hypercoverings $\mathcal{U}_\bullet \rightarrow X$ in $\widehat{\text{RigSm}}^{\text{gc}}$ of $X = \varprojlim_h X_h$ descending at finite level. Let $\mathcal{U}_\bullet \rightarrow X$ be one of them. Without loss of generality, we assume that it descends to an étale covering of X_0 . In particular, we conclude that $\mathcal{U}_n = \varprojlim_h \mathcal{U}_{nh}$ is a disjoint union of objects in $\widehat{\text{RigSm}}^{\text{gc}}$.

We need to show that $\text{Hom}_\bullet(\Lambda(\mathcal{U}_\bullet), \text{Sing}^{\mathbb{B}^1}(\iota^*\mathcal{F}))$ is quasi-isomorphic to $\text{Sing}^{\mathbb{B}^1}(\iota^*\mathcal{F})(X)$. Using Corollary 4.3, we conclude that for each $n \in \mathbb{N}$ the complex $(\text{Sing}^{\mathbb{B}^1} \iota^*\mathcal{F})(\mathcal{U}_n)$ is quasi-isomorphic to $\varinjlim_h (\text{Sing}^{\mathbb{B}^1} \iota^*\mathcal{F})(\mathcal{U}_{nh})$. Passing to the homotopy limit on n on both sides, we deduce that $\text{Hom}_\bullet(\Lambda(\mathcal{U}_\bullet), \text{Sing}^{\mathbb{B}^1} \iota^*\mathcal{F})$ is quasi-isomorphic to $\varinjlim_h \text{Hom}_\bullet(\Lambda(\mathcal{U}_{\bullet h}), \text{Sing}^{\mathbb{B}^1} \iota^*\mathcal{F})$. Using again Corollary 4.3, we also obtain that $(\text{Sing}^{\mathbb{B}^1} \iota^*\mathcal{F})(X)$ is quasi-isomorphic to $\varinjlim_h (\text{Sing}^{\mathbb{B}^1} \iota^*\mathcal{F})(X_h)$.

From the exactness of \varinjlim it suffices then to prove that the maps

$$\text{Hom}_\bullet(\Lambda(\mathcal{U}_{\bullet h}), \text{Sing}^{\mathbb{B}^1} \mathcal{F}) \rightarrow \text{Hom}_\bullet(\Lambda(X_h), \text{Sing}^{\mathbb{B}^1} \mathcal{F})$$

are quasi-isomorphisms. This follows once we show that the complex $\text{Sing}^{\mathbb{B}^1} \mathcal{F}$ is ét-local.

We point out that since \mathcal{F} is \mathbb{B}^1 -local, then the canonical map $\mathcal{F} \rightarrow \text{Sing}^{\mathbb{B}^1} \mathcal{F}$ is a quasi-isomorphism. As \mathcal{F} is ét-local we conclude that $\text{Sing}^{\mathbb{B}^1} \mathcal{F}$ also is, and hence the claim. \square

We are finally ready to state the main result of this section.

PROPOSITION 4.5. *Let $X = \varprojlim_h X_h$ be in $\widehat{\text{RigSm}}^{\text{gc}}$. For any complex of presheaves \mathcal{F} on RigSm^{gc} the natural map*

$$\varinjlim_h \mathbf{RigDA}_{\text{ét}}^{\text{eff}}(K)(\Lambda(X_h), \mathcal{F}) \rightarrow \widehat{\mathbf{RigDA}}_{\text{ét}, \mathbb{B}^1}^{\text{eff}}(K)(\Lambda(X), \mathbb{L}\iota^*\mathcal{F})$$

is an isomorphism.

Proof. Since any complex \mathcal{F} has a fibrant-cofibrant replacement in $\mathbf{Ch}_{\text{ét}, \mathbb{B}^1} \mathbf{Psh}(\text{RigSm}^{\text{gc}})$ we can assume that \mathcal{F} is cofibrant and $(\text{ét}, \mathbb{B}^1)$ -fibrant. Since it is \mathbb{B}^1 -local, it is quasi-isomorphic to $\text{Sing}^{\mathbb{B}^1} \mathcal{F}$. By Corollary 4.3, for any integer i one has

$$\varinjlim_h \text{Hom}(\Lambda(X_h)[i], \text{Sing}^{\mathbb{B}^1} \mathcal{F}) \cong \text{Hom}(\Lambda(X)[i], \text{Sing}^{\mathbb{B}^1} \iota^*\mathcal{F}).$$

As $\Lambda(X)$ is a cofibrant object in $\mathbf{Ch Psh}(\widehat{\text{RigSm}}^{\text{gc}})$ and $\text{Sing}^{\mathbb{B}^1} \iota^*\mathcal{F}$ is a $(\mathbb{B}^1, \text{ét})$ -local replacement of \mathcal{F} in $\mathbf{Ch}_{\text{ét}, \mathbb{B}^1} \mathbf{Psh}(\widehat{\text{RigSm}}^{\text{gc}})$ by Proposition 4.4, we conclude that the previous isomorphism can be rephrased in the following way:

$$\varinjlim_h \mathbf{RigDA}_{\text{ét}}^{\text{eff}}(K)(\Lambda(X_h)[i], \mathcal{F}) \cong \widehat{\mathbf{RigDA}}_{\text{ét}, \mathbb{B}^1}^{\text{eff}}(K)(\Lambda(X)[i], \mathbb{L}\iota^*\mathcal{F}),$$

proving the claim. \square

5. The de-perfectoidification functor in characteristic 0

The results proved in § 4 are valid both for char $K = 0$ and char $K = p$. In contrast, the results of this section require that char $K = 0$. We will present later their variant for the case char $K = p$.

We start by considering the adjunction between motives with and without transfers (see Remark 3.37). Thanks to the following theorem, we are allowed to add or ignore transfers according to the situation.

THEOREM 5.1 [Vez17]. *Suppose that char $K = 0$. The functors $(a_{\text{tr}}, o_{\text{tr}})$ induce an equivalence:*

$$\mathbb{L}a_{\text{tr}} : \mathbf{RigDA}_{\text{ét}}^{\text{eff}}(K) \rightleftarrows \mathbf{RigDM}_{\text{ét}}^{\text{eff}}(K) : \mathbb{R}o_{\text{tr}}.$$

Remark 5.2. The proof of the statement above uses in a crucial way the fact that the ring of coefficients Λ is a \mathbb{Q} -algebra.

PROPOSITION 5.3. *Suppose char $K = 0$. Let $X = \varprojlim_h X_h$ be in $\widehat{\text{RigSm}}^{\text{gc}}$. If h is big enough, then the map $\Lambda(X_{h+1}) \rightarrow \Lambda(X_h)$ is an isomorphism in $\mathbf{RigDA}_{\text{ét}}^{\text{eff}}(K)$.*

Proof. By means of Theorem 5.1, we can equally prove the statement in the category $\mathbf{RigDM}_{\text{ét}}^{\text{eff}}(K)$. We claim that we can also make an arbitrary finite field extension L/K . Indeed the transpose of the natural map $Y_L \rightarrow Y$ is a correspondence from Y to Y_L . Since Λ is a \mathbb{Q} -algebra, we conclude that $\Lambda_{\text{tr}}(Y)$ is a direct factor of $\Lambda_{\text{tr}}(Y_L) = \mathbb{L}e_{\sharp}\Lambda_{\text{tr}}(Y_L)$ for any variety Y , where $\mathbb{L}e_{\sharp}$ is the functor $\mathbf{RigDM}_{\text{ét}}^{\text{eff}}(L) \rightarrow \mathbf{RigDM}_{\text{ét}}^{\text{eff}}(K)$ induced by restriction of scalars. In particular, if $\Lambda_{\text{tr}}((X_{h+1})_L) \rightarrow \Lambda_{\text{tr}}((X_h)_L)$ is an isomorphism in $\mathbf{RigDM}_{\text{ét}}^{\text{eff}}(L)$ then $\Lambda_{\text{tr}}((X_{h+1})_L) \rightarrow \Lambda_{\text{tr}}((X_h)_L)$ is an isomorphism in $\mathbf{RigDM}_{\text{ét}}^{\text{eff}}(K)$ and therefore also $\Lambda_{\text{tr}}(X_{h+1}) \rightarrow \Lambda_{\text{tr}}(X_h)$ is.

By lemma [Ayo15, 1.1.52], we can suppose that $X_0 = \text{Spa}(R_0, R_0^\circ)$ with $R_0 = S\langle\sigma, \tau\rangle/(P(\sigma, \tau))$, where $S = \mathcal{O}(\mathbb{T}^M)$, $\sigma = (\sigma_1, \dots, \sigma_N)$ is a N -tuple of coordinates, $\tau = (\tau_1, \dots, \tau_m)$ is a m -tuple of coordinates and P is a set of m polynomials in $S[\sigma, \tau]$ with $\det(\partial P/\partial \tau) \in R_0^\times$. In particular, $X_1 = \text{Spa}(R_1, R_1^\circ)$ with $R_1 = S\langle\sigma, \tau\rangle/(P(\sigma^p, \tau))$ and the map $f : X_1 \rightarrow X_0$ is induced by $\sigma \mapsto \sigma^p$, $\tau \mapsto \tau$. Since the map f is finite and surjective, we can also consider the transpose correspondence $f^T \in \text{RigCor}(X_0, X_1)$. The composition $f \circ f^T$ is associated to the correspondence $X_0 \xleftarrow{f} X_1 \xrightarrow{f} X_0$, which is the cycle $\text{deg}(f)X_0 = p^N \cdot \text{id}_{X_0}$. The composition $f^T \circ f$ is associated to the correspondence $X_1 \xleftarrow{p_1} X_1 \times_{X_0} X_1 \xrightarrow{p_2} X_1$. Since $\mathbb{T}^N \langle\sigma^{1/p}\rangle \times_{\mathbb{T}^N} \mathbb{T}^N \langle\sigma^{1/p}\rangle \cong \mathbb{T}^N \langle\sigma^{1/p}\rangle \times \mu_p^N$ we conclude that the above correspondence is $X_1 \xleftarrow{p_1} X_1 \times (\mu_p)^N \xrightarrow{\eta} X_1$, where η is induced by the multiplication map $\mathbb{T}^N \times \mu_p^N \rightarrow \mathbb{T}^N$. Up to a finite field extension, we can assume that K has p th roots of unity. The above correspondence is then equal to $\sum f_\zeta$, where each f_ζ is a map $X_1 \rightarrow X_1$ defined by $\sigma_i \mapsto \zeta_i \sigma_i$, $\tau \mapsto \tau$ for each N -tuple $\zeta = (\zeta_i)$ of p th roots of unity. If we prove that each f_ζ is homotopically equivalent to id_{X_1} then we get $(1/p^N)f^T \circ f = \text{id}$, $f \circ (1/p^N)f^T = \text{id}$ in $\mathbf{RigDM}_{\text{ét}}^{\text{eff}}$, as desired.

We are left to find a homotopy between id and f_ζ for a fixed $\zeta = (\zeta_1, \dots, \zeta_n)$ up to considering higher indices h . For the sake of clarity, we consider them as maps $\text{Spa } \bar{R}_1 \rightarrow \text{Spa } R_1$, where we put $\bar{R}_h = S\langle\bar{\sigma}, \bar{\tau}\rangle/(P(\bar{\sigma}^{p^h}, \bar{\tau}))$ for any integer h . The first map is induced by $\sigma \mapsto \bar{\sigma}$, $\tau \mapsto \bar{\tau}$ and the second induced by $\sigma \mapsto \zeta\bar{\sigma}$, $\tau \mapsto \bar{\tau}$. Let $F_h = \sum_n a_n(\sigma - \bar{\sigma})^n$ be the unique array of formal power series in $\bar{R}_h[[\sigma - \bar{\sigma}]]$ centered in $\bar{\sigma}$ associated to the polynomials $P(\sigma^{p^h}, \tau)$ in $\bar{R}_h[\sigma, \tau]$ via Corollary A.2. Also let ϕ_h be the map $\bar{R}_h \rightarrow \bar{R}_{h+1}$. From the formal equalities $P(\sigma^{p^{(h+1)}}, F_{h+1}(\sigma)) = 0$, $P(\sigma^{p^h}, \phi(F_h(\sigma))) = \phi_h(P(\sigma^{p^h}, F_h(\sigma))) = 0$ and the uniqueness of F_{h+1} we deduce $F_{h+1}(\sigma) = \phi_h(F_h(\sigma^p))$.

We therefore have

$$\begin{aligned}
 F_{h+1}(\sigma) &= \sum_n \phi_h(a_n)(\sigma^p - \bar{\sigma}^p)^n \\
 &= \sum_n \phi_h(a_n) \left((\sigma - \bar{\sigma})^{p-1} + \sum_{j=1}^{p-1} \binom{p}{j} (\sigma - \bar{\sigma})^{j-1} \bar{\sigma}^{p-j} \right)^n (\sigma - \bar{\sigma})^n.
 \end{aligned}$$

The expression

$$Q(x) = x^p + \sum_{j=1}^{p-1} \binom{p}{j} x^j \bar{\sigma}^{p-j}$$

is a polynomial in x and it is easy to show that the mapping $x \mapsto Q(x)$ extends to a map $\bar{R}_{h+1}\langle x \rangle \rightarrow \bar{R}_{h+1}\langle x \rangle$. We deduce that we can read off the convergence in the circle of radius 1 around $\bar{\sigma}$ and the values of F_{h+1} on its expression given above.

We remark that the norm of $Q(\sigma - \bar{\sigma})$ in the circle of radius $\rho \leq 1$ around $\bar{\sigma}$ is bounded by $\max\{\rho^p, |p|\} \leq \max\{\rho, |p|\}$. Suppose that F_h converges in a circle of radius ρ with $0 < \rho \leq 1$ around $\bar{\sigma}$ and in there it takes values in power-bounded elements. By the expression above, the same holds true for F_{h+1} in the circle of radius $\min\{\rho|p|^{-1}, 1\}$ around $\bar{\sigma}$. By induction we conclude that for a sufficiently big h the power series F_h converges in a circle of radius $\delta > |p|^{1/(p-1)}$ around $\bar{\sigma}$ and its values in it are power bounded. Up to rescaling indices, we suppose that this holds for $h = 1$.

From the relation $F_{h+1}(\sigma) = \phi_h(F_h(\sigma^p))$ we also conclude $F_1(\zeta\bar{\sigma}) = F_1(\bar{\sigma}) = \bar{\tau}$. Therefore, since $|\zeta_i - 1| = |p|^{1/(p-1)}$ for all i , the map

$$\begin{aligned}
 X_1 = \text{Spa}(S\langle \sigma, \tau \rangle / P(\sigma^p, \tau)) &\leftarrow X_1 \times \mathbb{B}^1 = \text{Spa}(S\langle \bar{\sigma}, \bar{\tau}, \chi \rangle / (P(\bar{\sigma}^p, \bar{\tau}))) \\
 (\sigma_i, \tau_j) &\mapsto (\bar{\sigma}_i + (\zeta_i - 1)\bar{\sigma}_i\chi, F_1(\bar{\sigma} + (\zeta - 1)\bar{\sigma}\chi))
 \end{aligned}$$

is a well-defined map, inducing a homotopy between id_{X_1} and f_{ζ} , as claimed. □

It cannot be expected that all maps $X_{h+1} \rightarrow X_h$ are isomorphisms in $\mathbf{RigDA}_{\text{ét}}^{\text{eff}}(K)$: consider, for example, $X_0 = \mathbb{T}^1\langle \nu^{1/p} \rangle \rightarrow \mathbb{T}^1$. Then X_0 is a connected variety, while X_1 is not. That said, there is a particular class of objects $X = \varprojlim_h X_h$ in $\widehat{\mathbf{RigSm}}^{\text{gc}}$ for which this happens: this is the content of the following proposition, which nevertheless will not be used in the following.

We recall that a presentation $X = \varprojlim_h X_h$ of an object in $\widehat{\mathbf{RigSm}}^{\text{gc}}$ is of good reduction if the map $X_0 \rightarrow \mathbb{T}^N \times \mathbb{T}^M$ has a formal model, which is an étale map over $\text{Spf } K^\circ\langle \underline{\nu}^{\pm 1}, \underline{\nu}^{\pm 1} \rangle$ and is of potentially good reduction if this happens after base change by a separable finite field extension L/K .

PROPOSITION 5.4. *Let $\text{char } K = 0$ and let $X = \varprojlim_h X_h$ be a presentation of a variety in $\widehat{\mathbf{RigSm}}^{\text{gc}}$ of potentially good reduction. The maps $\Lambda(X_{h+1}) \rightarrow \Lambda(X_h)$ are isomorphisms in $\mathbf{RigDA}_{\text{ét}}^{\text{eff}}(K)$ for all h .*

Proof. If the map $X_0 \rightarrow \mathbb{T}^N \times \mathbb{T}^M$ has an étale formal model, then also the map $X_h \rightarrow \mathbb{T}^N\langle \underline{\nu}^{1/p^h} \rangle \times \mathbb{T}^M$ does. It is then sufficient to consider only the case $h = 0$. Since L/K is finite and Λ is a \mathbb{Q} -algebra, by the same argument of the proof of Proposition 5.3 we can assume that $\varprojlim_h X_h$ has good reduction. Also, by means of Theorem 5.1 and the Cancellation Theorem [Ayo15, Corollary 2.5.49], we can equally prove the statement in the stable category $\mathbf{RigDA}_{\text{ét}}(K)$ defined in [Ayo15, Definition 1.3.19].

Let $\mathfrak{X}_0 \rightarrow \mathrm{Spf} K^\circ \langle \underline{\nu}^{\pm 1}, \underline{\nu}^{\pm 1} \rangle$ be a formal model of the map $X_0 \rightarrow \mathbb{T}^n \times \mathbb{T}^m$. We let \bar{X}_0 be the special fiber over the residue field k of K . The variety X_1 has also a smooth formal model \mathfrak{X}_1 whose special fiber is \bar{X}_1 . By definition, the natural map $\bar{X}_1 \rightarrow \bar{X}_0$ is the push-out of the (relative) Frobenius map $\mathbb{A}_k^{\dim X} \rightarrow \mathbb{A}_k^{\dim X}$, which is isomorphic to the relative Frobenius map, and hence an isomorphism of correspondences as p is invertible in Λ . We conclude that $\Lambda_{\mathrm{tr}}(\bar{X}_1) \rightarrow \Lambda_{\mathrm{tr}}(\bar{X}_0)$ is an isomorphism in $\mathbf{DM}_{\mathrm{ét}}(k)$.

Let $\mathbf{FormDA}_{\mathrm{ét}}(K^\circ)$ be the stable category of motives of formal varieties $\mathbf{FSH}_{\mathfrak{M}}(K^\circ)$ defined in [Ayo15, Definition 1.4.15] associated to the model category $\mathfrak{M} = \mathbf{Ch}(\Lambda - \mathbf{Mod})$. Using [Ayo14a, Theorem B.1] we deduce that the map $\Lambda(\bar{X}_1) \rightarrow \Lambda(\bar{X}_0)$ is an isomorphism in $\mathbf{DA}_{\mathrm{ét}}(k)$ as is its image via the following functor (see [Ayo15, Remark 1.4.30]) induced by the special fiber functor and the generic fiber functor:

$$\mathbf{DA}_{\mathrm{ét}}(k) \xleftarrow[\sim]{(-)_\sigma} \mathbf{FormDA}_{\mathrm{ét}}(K^\circ) \xrightarrow{(-)_\eta} \mathbf{RigDA}_{\mathrm{ét}}(K).$$

This morphism is precisely the map $\Lambda(X_1) \rightarrow \Lambda(X_0)$ proving the claim. □

We are now ready to present the main result of this section.

THEOREM 5.5. *Let $\mathrm{char} K = 0$. The functor $\mathbb{L}\iota^* : \mathbf{RigDA}_{\mathrm{ét}}^{\mathrm{eff}}(K) \rightarrow \widehat{\mathbf{RigDA}}_{\mathrm{ét}}^{\mathrm{eff}}(K)$ has a left adjoint $\mathbb{L}\iota_!$ and the counit map $\mathrm{id} \rightarrow \mathbb{L}\iota_!\mathbb{L}\iota^*$ is invertible. Whenever $X = \varprojlim_h X_h$ is an object of $\widehat{\mathbf{RigSm}}^{\mathrm{gc}}$ then $\mathbb{L}\iota_!\Lambda(X) \cong \Lambda(X_h)$ for a sufficiently large index h . If moreover $X = \varprojlim_h X_h$ is of potentially good reduction, then $\mathbb{L}\iota_!\Lambda(X) \cong \Lambda(X_0)$.*

Proof. We start by proving that the canonical map

$$\mathbf{RigDA}_{\mathrm{ét}}^{\mathrm{eff}}(K)(\Lambda(X_{\bar{h}}), \mathcal{F}) \rightarrow \widehat{\mathbf{RigDA}}_{\mathrm{ét}}^{\mathrm{eff}}(K)(\Lambda(X), \mathbb{L}\iota^*\mathcal{F})$$

is an isomorphism, for every $X = \varprojlim_h X_h$ and for \bar{h} big enough. By Proposition 4.5, it suffices to prove that the natural map

$$\mathbf{RigDA}^{\mathrm{eff}}(K)(\Lambda(X_{\bar{h}}), \mathbb{L}a_{\mathrm{tr}}\mathcal{F}) \rightarrow \varinjlim_h \mathbf{RigDA}^{\mathrm{eff}}(K)(\Lambda(X_h), \mathbb{L}a_{\mathrm{tr}}\mathcal{F})$$

is an isomorphism for some \bar{h} . This follows from Proposition 5.3 since the maps $\Lambda(X_{h+1}) \rightarrow \Lambda(X_h)$ are isomorphisms if $h \geq \bar{h}$ for some big enough \bar{h} . In case $\varprojlim_h X_h$ is of potentially good reduction, then Proposition 5.4 ensures that we can choose $\bar{h} = 0$.

We conclude that the subcategory \mathbf{T} of $\widehat{\mathbf{RigDA}}_{\mathrm{ét}, \mathbb{B}^1}^{\mathrm{eff}}(K)$ formed by the objects M such that the functor $N \mapsto \widehat{\mathbf{RigDA}}_{\mathrm{ét}, \mathbb{B}^1}^{\mathrm{eff}}(K)(M, \mathbb{L}\iota^*N)$ is corepresentable contains all motives $\Lambda(X)$ with X any object of $\widehat{\mathbf{RigSm}}^{\mathrm{gc}}$. Since these objects form a set of compact generators of $\widehat{\mathbf{RigDA}}_{\mathrm{ét}, \mathbb{B}^1}^{\mathrm{eff}}(K)$ by Proposition 3.18, we deduce the existence of the functor $\mathbb{L}\iota_!$ by Lemma 5.6.

The formula $\mathbb{L}\iota_!\mathbb{L}\iota^* \cong \mathrm{id}$ is a formal consequence of the fact that $\mathbb{L}\iota_!$ is the left adjoint of a fully faithful functor $\mathbb{L}\iota^*$ (see Proposition 3.17). □

LEMMA 5.6. *Let $\mathfrak{G} : \mathbf{T} \rightarrow \mathbf{T}'$ be a triangulated functor of triangulated categories. The full subcategory \mathbf{C} of \mathbf{T}' of objects M such that the functor $a_M : N \mapsto \mathrm{Hom}(M, \mathfrak{G}N)$ is corepresentable is closed under cones and small direct sums.*

Proof. For any object M in \mathbf{C} we denote by $\mathfrak{F}M$ the object corepresenting the functor a_M . Now let $\{M_i\}_{i \in I}$ be a set of objects in \mathbf{C} . It is immediate to check that $\bigoplus_i \mathfrak{F}M_i$ corepresents the functor $a_{\bigoplus_i M_i}$.

Now let M_1, M_2 be two objects of \mathbf{C} and $f: M_1 \rightarrow M_2$ be a map between them. There are canonical maps $\eta_i: M_i \rightarrow \mathfrak{G}\mathfrak{F}M_i$ induced by the identity $\mathfrak{F}M_i \rightarrow \mathfrak{F}M_i$ and the universal property of $\mathfrak{F}M_i$. By composing with η_2 we obtain a morphism $\text{Hom}(M_1, M_2) \rightarrow \text{Hom}(M_1, \mathfrak{G}\mathfrak{F}M_2) \cong \text{Hom}(\mathfrak{F}M_1, \mathfrak{F}M_2)$ sending f to a map $\mathfrak{G}\mathfrak{F}f$. Let C be the cone of f and D be the cone of $\mathfrak{F}f$. We claim that D represents a_C . From the triangulated structure we obtain a map of distinguished triangles

$$\begin{array}{ccccccc} M_1 & \xrightarrow{f} & M_2 & \longrightarrow & C & \longrightarrow & \\ \downarrow \eta_1 & & \downarrow \eta_2 & & \downarrow & & \\ \mathfrak{G}\mathfrak{F}M_1 & \xrightarrow{\mathfrak{G}\mathfrak{F}f} & \mathfrak{G}\mathfrak{F}M_2 & \longrightarrow & \mathfrak{G}D & \longrightarrow & \end{array}$$

inducing for any object N of \mathbf{T} the following maps of long exact sequences.

$$\begin{array}{ccccccc} \longleftarrow & \text{Hom}(M_1, \mathfrak{G}N) & \longleftarrow & \text{Hom}(M_2, \mathfrak{G}N) & \longleftarrow & \text{Hom}(C, \mathfrak{G}N) & \longleftarrow \\ & \uparrow & & \uparrow & & \uparrow & \\ \longleftarrow & \text{Hom}(\mathfrak{G}\mathfrak{F}M_1, \mathfrak{G}N) & \longleftarrow & \text{Hom}(\mathfrak{G}\mathfrak{F}M_2, \mathfrak{G}N) & \longleftarrow & \text{Hom}(\mathfrak{G}D, \mathfrak{G}N) & \longleftarrow \\ & \uparrow & & \uparrow & & \uparrow & \\ \longleftarrow & \text{Hom}(\mathfrak{F}M_1, N) & \longleftarrow & \text{Hom}(\mathfrak{F}M_2, N) & \longleftarrow & \text{Hom}(D, N) & \longleftarrow \end{array}$$

Since the vertical compositions are isomorphisms for M_1 and M_2 we deduce that they all are, proving that D corepresents a_C , as desired. \square

We remark that we used the fact that Λ is a \mathbb{Q} -algebra at least twice in the proof of Theorem 5.5: to allow for field extensions and correspondences using Theorem 5.1 as well as to invert the map defined by multiplication by p .

The following fact is a straightforward corollary of Theorem 5.5.

PROPOSITION 5.7. *Let $\text{char } K = 0$. The motive $\mathbb{L}\iota_1\Lambda(\widehat{\mathbb{B}^1})$ is isomorphic to Λ .*

Proof. In order to prove the claim, it suffices to prove that $\mathbb{L}\iota_1\Lambda(\widehat{\mathbb{B}^1}) \cong \Lambda(\mathbb{B}^1)$. This follows from Proposition 2.13 and the description of $\mathbb{L}\iota_1$ given in Theorem 5.5. \square

We recall that all the homotopy categories we consider are monoidal (see [Ayo07, Propositions 4.2.76 and 4.4.62]), and the tensor product $\Lambda(X) \otimes \Lambda(X')$ of two motives associated to varieties X and Y coincides with $\Lambda(X \times X')$. The unit object is obviously the motive Λ . Due to the explicit description of the functor $\mathbb{L}\iota_1$ we constructed above, it is easy to prove that it respects the monoidal structures.

PROPOSITION 5.8. *Let $\text{char } K = 0$. The functor $\mathbb{L}\iota_1: \widehat{\mathbf{RigDA}}_{\text{ét}, \mathbb{B}^1}^{\text{eff}}(K) \rightarrow \mathbf{RigDA}_{\text{ét}}^{\text{eff}}(K)$ is a monoidal functor.*

Proof. Since $\mathbb{L}\iota_1$ is the left adjoint of a monoidal functor $\mathbb{L}\iota^*$ there is a canonical natural transformation of bifunctors $\mathbb{L}\iota_1(M \otimes M') \rightarrow \mathbb{L}\iota_1 M \otimes \mathbb{L}\iota_1 M'$. In order to prove it is an isomorphism, it suffices to check it on a set of generators of $\widehat{\mathbf{RigDA}}_{\text{ét}, \mathbb{B}^1}^{\text{eff}}$ such as motives of

semi-perfectoid varieties $X = \varprojlim_h X_h$, $X' = \varprojlim_h X'_h$. Up to rescaling, we can suppose that $\mathbb{L}\iota_! \Lambda(X) = \Lambda(X_0)$ and $\mathbb{L}\iota_! \Lambda(X') = \Lambda(X'_0)$ by Theorem 5.5. In this case, by definition of the tensor product, we obtain the following isomorphisms:

$$\mathbb{L}\iota_!(\Lambda(X) \otimes \Lambda(X')) \cong \mathbb{L}\iota_!\Lambda(X \times X') \cong \Lambda(X_0 \times X'_0) \cong \Lambda(X_0) \otimes \Lambda(X'_0) \cong \mathbb{L}\iota_!\Lambda(X) \otimes \mathbb{L}\iota_!\Lambda(X'),$$

proving our claim. □

The following proposition can be considered to be a refinement of Theorem 5.5.

PROPOSITION 5.9. *Let $\text{char } K = 0$. The functor $\mathbb{L}\iota_!$ factors through $\widehat{\mathbf{RigDA}}_{\text{ét}, \mathbb{B}^1}^{\text{eff}} \rightarrow \widehat{\mathbf{RigDA}}_{\text{ét}, \widehat{\mathbb{B}^1}}^{\text{eff}}$ and the image of the functor $\mathbb{L}\iota^* : \mathbf{RigDA}_{\text{ét}}^{\text{eff}}(K) \rightarrow \widehat{\mathbf{RigDA}}_{\text{ét}, \mathbb{B}^1}^{\text{eff}}(K)$ lies in the subcategory of $\widehat{\mathbb{B}^1}$ -local objects. In particular, the triangulated adjunction*

$$\mathbb{L}\iota_! : \widehat{\mathbf{RigDA}}_{\text{ét}, \mathbb{B}^1}^{\text{eff}}(K) \rightleftarrows \mathbf{RigDA}_{\text{ét}}^{\text{eff}}(K) : \mathbb{L}\iota^*$$

restricts to a triangulated adjunction

$$\mathbb{L}\iota_! : \widehat{\mathbf{RigDA}}_{\text{ét}, \widehat{\mathbb{B}^1}}^{\text{eff}}(K) \rightleftarrows \mathbf{RigDA}_{\text{ét}}^{\text{eff}}(K) : \mathbb{L}\iota^*.$$

Proof. By Propositions 5.7 and 5.8, $\mathbb{L}\iota_!$ is a monoidal functor sending $\Lambda(\widehat{\mathbb{B}^1})$ to Λ . This proves the first claim.

From the adjunction $(\mathbb{L}\iota_!, \mathbb{L}\iota^*)$ we then obtain the following isomorphisms, for any X in $\widehat{\mathbf{RigSm}}^{\text{gc}}$ and any M in $\mathbf{RigDA}_{\text{ét}}^{\text{eff}}(K)$:

$$\begin{aligned} \widehat{\mathbf{RigDA}}_{\text{ét}, \mathbb{B}^1}^{\text{eff}}(K)(\Lambda(X \times \widehat{\mathbb{B}^1}), \mathbb{L}\iota^* M) &\cong \mathbf{RigDA}_{\text{ét}}^{\text{eff}}(K)(\mathbb{L}\iota_!\Lambda(X) \otimes \Lambda, M) \\ &\cong \mathbf{RigDA}_{\text{ét}}^{\text{eff}}(K)(\mathbb{L}\iota_!\Lambda(X), M) \\ &\cong \widehat{\mathbf{RigDA}}_{\text{ét}, \mathbb{B}^1}^{\text{eff}}(K)(\Lambda(X), \mathbb{L}\iota^* M), \end{aligned}$$

proving the second claim. □

Remark 5.10. In the statement of the proposition above, we make a slight abuse of notation when denoting with $(\mathbb{L}\iota_!, \mathbb{L}\iota^*)$ both adjoint pairs. It will be clear from the context which one we consider at each instance.

6. The de-perfectoidification functor in characteristic p

We now consider the case of a perfectoid field K^\flat of characteristic p and try to generalize the results of § 5. We will need to perform an extra localization on the model structure, and in return we will prove a stronger result. In this section, we always assume that the base perfectoid field has characteristic p . In order to emphasize this hypothesis, we denote it with K^\flat .

In positive characteristic, we are not able to prove Theorem 5.1 as it is stated, and it is therefore not clear that the maps $X_{h+1} \rightarrow X_h$ associated to an object $X = \varprojlim_h X_h$ of $\widehat{\mathbf{RigSm}}$ are isomorphisms in $\mathbf{RigDA}_{\text{ét}}^{\text{eff}}(K^\flat)$ for a sufficiently big h . In order to overcome this obstacle, we localize our model category further.

For any variety X over K^\flat we denote by $X^{(1)}$ the pullback of X over the Frobenius map $\Phi : K^\flat \rightarrow K^\flat, x \mapsto x^p$. The absolute Frobenius morphism induces a K^\flat -linear map $X \rightarrow X^{(1)}$. Since K^\flat is perfect, we can also denote by $X^{(-1)}$ the pullback of X over the inverse of the Frobenius map $\Phi^{-1} : K^\flat \rightarrow K^\flat$ and $X \cong (X^{(-1)})^{(1)}$. There is, in particular, a canonical map $X^{(-1)} \rightarrow X$, which is isomorphic to the map $X' \rightarrow X$ induced by the absolute Frobenius, where we denote by X' the same variety X endowed with the structure map $X \rightarrow \text{Spa } K \xrightarrow{\Phi} \text{Spa } K$.

PROPOSITION 6.1. *The model category $\mathbf{Ch}_{\text{ét}, \mathbb{B}^1} \mathbf{Psh}(\text{RigSm}/K^b)$ admits a left Bousfield localization denoted by $\mathbf{Ch}_{\text{Frobét}, \mathbb{B}^1} \mathbf{Psh}(\text{RigSm}/K^b)$ with respect to the set S_{Frob} of relative Frobenius maps $\Phi : \Lambda(X^{(-1)})[i] \rightarrow \Lambda(X)[i]$ as X varies in RigSm and i varies in \mathbb{Z} .*

Proof. Since by [Ayo07, Proposition 4.4.31] the τ -localization coincides with the Bousfield localization with respect to a set, we conclude by [Ayo07, Theorem 4.2.71] that the model category $\mathbf{Ch}_{\text{ét}, \mathbb{B}^1} \mathbf{Psh}(\text{RigSm}/K^b)$ is still left proper and cellular. We can then apply [Hir03, Theorem 4.1.1]. \square

DEFINITION 6.2. We denote by $\mathbf{RigDA}_{\text{Frobét}}^{\text{eff}}(K^b, \Lambda)$ the homotopy category associated to $\mathbf{Ch}_{\text{Frobét}, \mathbb{B}^1} \mathbf{Psh}(\text{RigSm}/K^b)$. We omit Λ whenever the context allows it. The image of a rigid variety X in this category is denoted by $\Lambda(X)$.

The triangulated category $\mathbf{RigDA}_{\text{Frobét}}^{\text{eff}}(K)$ is canonically isomorphic to the full triangulated subcategory of $\mathbf{RigDA}_{\text{ét}}^{\text{eff}}(K)$ formed by Frob-local objects, i.e., objects that are local with respect to the maps in S_{Frob} . Modulo this identification, there is an obvious functor $\mathbf{RigDA}_{\text{ét}}^{\text{eff}}(K^b) \rightarrow \mathbf{RigDA}_{\text{Frobét}}^{\text{eff}}(K^b)$ associating to \mathcal{F} a Frob-local object $C^{\text{Frob}} \mathcal{F}$.

Inverting Frobenius morphisms is enough to obtain an analog of Theorem 5.1 in characteristic p .

THEOREM 6.3 [Vez17]. *Let $\text{char } K^b = p$. The functors $(a_{\text{tr}}, o_{\text{tr}})$ induce an equivalence of triangulated categories:*

$$\mathbb{L}a_{\text{tr}} : \mathbf{RigDA}_{\text{Frobét}}^{\text{eff}}(K^b) \cong \mathbf{RigDM}_{\text{ét}}^{\text{eff}}(K^b).$$

Remark 6.4. The proof of the statement above uses in a crucial way the fact that the ring of coefficients Λ is a \mathbb{Q} -algebra.

We now investigate the relations between the category $\mathbf{RigDA}_{\text{Frobét}}^{\text{eff}}(K^b)$ we have just defined, and the other categories of motives introduced so far.

PROPOSITION 6.5. *Let X_0 be in RigSm/K^b endowed with an étale map $X_0 \rightarrow \mathbb{T}^N \times \mathbb{T}^M = \text{Spa}(K^b \langle \underline{v}^{\pm 1}, \underline{v}^{\pm 1} \rangle)$. The map $X_1 = X_0 \times_{\mathbb{T}^N} \mathbb{T}^N \langle \underline{v}^{\pm 1/p} \rangle \rightarrow X_0$ is invertible in $\mathbf{RigDA}_{\text{Frobét}}^{\text{eff}}(K^b)$.*

Proof. The map of the claim is a factor of $X_0 \times_{(\mathbb{B}^N \times \mathbb{B}^M)} (\mathbb{B}^N \langle \underline{v}^{1/p} \rangle \times \mathbb{B}^M \langle \underline{v}^{1/p} \rangle) \rightarrow X_0$, which is isomorphic to the relative Frobenius map $X_0^{(-1)} \rightarrow X_0$ (see for example [GR03, Theorem 3.5.13]). If we consider the diagram

$$X_1^{(-1)} \xrightarrow{a} X_0^{(-1)} \xrightarrow{b} X_1 \xrightarrow{c} X_0$$

we conclude that the two compositions ba and cb are isomorphisms, and hence c is also an isomorphism, as claimed. \square

PROPOSITION 6.6. *The image via $\mathbb{L}i^*$ of a Frob-local object of $\mathbf{RigDA}_{\text{ét}}^{\text{eff}}(K^b)$ is $\widehat{\mathbb{B}}^1$ -local. In particular, the functor $\mathbb{L}i^*$ restricts to a functor $\mathbb{L}i^* : \mathbf{RigDA}_{\text{Frobét}}^{\text{eff}}(K^b) \rightarrow \widehat{\mathbf{RigDA}}_{\text{ét}, \widehat{\mathbb{B}}^1}^{\text{eff}}(K^b)$.*

Proof. Let $X' = \varprojlim_h X'_h$ be in $\widehat{\text{RigSm}}^{\text{gc}}$. We consider the object $X' \times \widehat{\mathbb{B}}^1 = \varprojlim_h (X'_h \times X_h)$, where we use the description $\widehat{\mathbb{B}}^1 = \varprojlim_h X_h$ of Proposition 2.13. Let M be a Frob-local object of $\mathbf{RigDA}_{\text{ét}}^{\text{eff}}(K^b)$. From Propositions 4.5 and 6.5, we then deduce the following isomorphisms:

$$\begin{aligned} \widehat{\mathbf{RigDA}}_{\acute{e}t, \mathbb{B}^1}^{\text{eff}}(K^b)(X' \times \widehat{\mathbb{B}}^1, \mathbb{L}^* M) &\cong \varinjlim_h \mathbf{RigDA}_{\acute{e}t}^{\text{eff}}(K^b)(X'_h \times X_h, M) \\ &\cong \mathbf{RigDA}_{\acute{e}t}^{\text{eff}}(K^b)(X'_0 \times \mathbb{B}^1, M) \\ &\cong \mathbf{RigDA}_{\acute{e}t}^{\text{eff}}(K^b)(X'_0, M) \\ &\cong \varinjlim_h \mathbf{RigDA}_{\acute{e}t}^{\text{eff}}(K^b)(X'_h, M) \\ &\cong \widehat{\mathbf{RigDA}}_{\acute{e}t, \mathbb{B}^1}^{\text{eff}}(K^b)(X', \mathbb{L}^* M), \end{aligned}$$

proving the claim. □

We remark that in positive characteristic the perfection $\text{Perf} : X \mapsto \varprojlim X^{(-i)}$ is functorial. This makes the description of various functors a lot easier. We recall that we denote by

$$\mathbb{L}j_* : \mathbf{PerfDA}_{\acute{e}t}^{\text{eff}}(K^b) \rightleftarrows \widehat{\mathbf{RigDA}}_{\acute{e}t, \widehat{\mathbb{B}}^1}^{\text{eff}}(K) : \mathbb{R}j_*$$

the adjoint pair induced by the inclusion of categories $j : \text{PerfSm} \rightarrow \widehat{\mathbf{RigSm}}$.

PROPOSITION 6.7. *The perfection functor $\text{Perf} : \widehat{\mathbf{RigSm}} \rightarrow \text{PerfSm}$ induces an adjunction*

$$\mathbb{L}\text{Perf}^* : \widehat{\mathbf{RigDA}}_{\acute{e}t, \mathbb{B}^1}^{\text{eff}}(K^b) \rightleftarrows \mathbf{PerfDA}_{\acute{e}t}^{\text{eff}}(K^b) : \mathbb{R}\text{Perf}_*$$

and $\mathbb{L}\text{Perf}^*$ factors through $\widehat{\mathbf{RigDA}}_{\acute{e}t, \mathbb{B}^1}^{\text{eff}}(K^b) \rightarrow \widehat{\mathbf{RigDA}}_{\acute{e}t, \widehat{\mathbb{B}}^1}^{\text{eff}}(K^b)$. Moreover, the functor $\mathbb{L}\text{Perf}^*$ coincides with $\mathbb{R}j_*$ on $\widehat{\mathbf{RigDA}}_{\acute{e}t, \mathbb{B}^1}^{\text{eff}}(K^b)$.

Proof. The perfection functor is continuous with respect to the étale topology and maps \mathbb{B}^1 and $\widehat{\mathbb{B}}^1$ to $\widehat{\mathbb{B}}^1$, and hence the first claim.

We now consider the functors $j : \text{PerfSm} \rightarrow \widehat{\mathbf{RigSm}}$ and $\text{Perf} : \widehat{\mathbf{RigSm}} \rightarrow \text{PerfSm}$. They induce two Quillen pairs (j^*, j_*) and $(\text{Perf}^*, \text{Perf}_*)$ on the associated $(\acute{e}t, \widehat{\mathbb{B}}^1)$ -localized model categories of complexes. Since Perf is a right adjoint of j we deduce that Perf^* is a right adjoint of j^* , and hence we obtain an isomorphism $j_* \cong \text{Perf}^*$, which shows the second claim. □

PROPOSITION 6.8. *Let Λ be a \mathbb{Q} -algebra. The functor*

$$\mathbb{L}\text{Perf}^* \mathbb{L}^* : \mathbf{RigDA}_{\acute{e}t}^{\text{eff}}(K^b) \rightarrow \mathbf{PerfDA}_{\acute{e}t}^{\text{eff}}(K^b)$$

factors over $\mathbf{RigDA}_{\text{Frob}\acute{e}t}^{\text{eff}}(K^b)$ and is isomorphic to $\mathbb{R}j_ \mathbb{L}^* C^{\text{Frob}}$.*

Proof. The first claim follows as the perfection of $X^{(-1)}$ is canonically isomorphic to the perfection of X for any object X in \mathbf{RigSm} .

The second part of the statement follows from the first claim and the commutativity of the following diagram, which is ensured by Propositions 6.6 and 6.7.

$$\begin{array}{ccc} \mathbf{RigDA}_{\text{Frob}\acute{e}t}^{\text{eff}}(K^b) & \xrightarrow{\mathbb{L}^*} & \widehat{\mathbf{RigDA}}_{\acute{e}t, \widehat{\mathbb{B}}^1}^{\text{eff}}(K^b) \\ \downarrow & & \downarrow \mathbb{R}j_* \\ \mathbf{RigDA}_{\acute{e}t}^{\text{eff}}(K^b) & \xrightarrow{\mathbb{L}^*} & \widehat{\mathbf{RigDA}}_{\acute{e}t, \mathbb{B}^1}^{\text{eff}}(K^b) \\ & & \uparrow \mathbb{L}\text{Perf}^* \\ & & \mathbf{PerfDA}_{\acute{e}t}^{\text{eff}}(K^b) \end{array}$$

□

THEOREM 6.9. *Let Λ be a \mathbb{Q} -algebra. The functor*

$$\mathbb{L} \text{Perf}^* : \mathbf{RigDA}_{\text{Frob}\acute{e}t}^{\text{eff}}(K^b) \rightarrow \mathbf{PerfDA}_{\acute{e}t}^{\text{eff}}(K^b)$$

defines a monoidal, triangulated equivalence of categories.

Proof. Let X_0 and Y be objects of RigSm^{gc} . Suppose X_0 is endowed with an étale map over \mathbb{T}^N , which is a composition of finite étale maps and inclusions, and let \widehat{X} be $\varprojlim_h X_h$. We can identify \widehat{X} with $\text{Perf } X_0$. Since $C^{\text{Frob}}\Lambda(Y)$ is Frob-local, by Proposition 6.5 the maps

$$\mathbf{RigDA}_{\acute{e}t}^{\text{eff}}(K^b)(\Lambda(X_h), C^{\text{Frob}}\Lambda(Y)) \rightarrow \mathbf{RigDA}_{\acute{e}t}^{\text{eff}}(K^b)(\Lambda(X_{h+1}), C^{\text{Frob}}\Lambda(Y))$$

are isomorphisms for all h . Using Propositions 4.5 and 6.6, we obtain the following sequence of isomorphisms for any $n \in \mathbb{Z}$:

$$\begin{aligned} \mathbf{RigDA}_{\text{Frob}\acute{e}t}^{\text{eff}}(K^b)(\Lambda(X_0), \Lambda(Y)[n]) &\cong \mathbf{RigDA}_{\acute{e}t}^{\text{eff}}(K^b)(\Lambda(X_0), C^{\text{Frob}}\Lambda(Y)[n]) \\ &\cong \varinjlim_h \mathbf{RigDA}_{\acute{e}t}^{\text{eff}}(K^b)(\Lambda(X_h), C^{\text{Frob}}\Lambda(Y)[n]) \\ &\cong \widehat{\mathbf{RigDA}}_{\acute{e}t, \mathbb{B}^1}^{\text{eff}}(K^b)(\Lambda(\widehat{X}), \mathbb{L}\iota^* C^{\text{Frob}}\Lambda(Y)[n]) \\ &\cong \widehat{\mathbf{RigDA}}_{\acute{e}t, \widehat{\mathbb{B}^1}}^{\text{eff}}(K^b)(\Lambda(\widehat{X}), \mathbb{L}\iota^* C^{\text{Frob}}\Lambda(Y)[n]) \\ &\cong \mathbf{PerfDA}_{\acute{e}t}^{\text{eff}}(K^b)(\Lambda(\widehat{X}), \mathbb{R}j_* \mathbb{L}\iota^* C^{\text{Frob}}\Lambda(Y)[n]) \\ &\cong \mathbf{PerfDA}_{\acute{e}t}^{\text{eff}}(K^b)(\mathbb{L} \text{Perf}^*(X_0), \mathbb{L} \text{Perf}^*(Y)[n]), \end{aligned}$$

where the last isomorphism follows from the identification $\widehat{X} \cong \text{Perf } X_0$ and Proposition 6.8. In particular, we deduce that the triangulated functor $\mathbb{L} \text{Perf}^*$ maps a set of compact generators to a set of compact generators (see Propositions 3.18 and 3.30) and on these objects it is fully faithful. By means of [Ayo15, Lemma 1.3.32], we then conclude it is a triangulated equivalence of categories, as claimed. \square

Remark 6.10. From the proof of the previous claim, we also deduce that the inverse $\mathbb{R} \text{Perf}_*$ of $\mathbb{L} \text{Perf}^*$ sends the motive associated to an object $X = \varprojlim_h X_h$ to the motive of X_0 . This functor is then analogous to the de-perfectoidification functor $\mathbb{L}j^* \circ \mathbb{L}\iota_!$ of Theorem 5.5.

7. The main theorem

Thanks to the results of the previous sections, we can reformulate Theorem 5.5 in terms of motives of rigid varieties. We will always assume that $\text{char } K = 0$ since the results of this section are tautological when $\text{char } K = p$.

COROLLARY 7.1. *There exists a triangulated adjunction of categories*

$$\mathfrak{F} : \mathbf{RigDM}_{\acute{e}t}^{\text{eff}}(K^b) \rightleftarrows \mathbf{RigDM}_{\acute{e}t}^{\text{eff}}(K) : \mathfrak{G}$$

such that \mathfrak{F} is a monoidal functor.

Proof. From Theorem 5.5 and Proposition 5.8, we can define an adjunction

$$\mathfrak{F}' : \mathbf{RigDA}_{\text{Frob}\acute{e}t}^{\text{eff}}(K^b) \rightleftarrows \mathbf{RigDA}_{\acute{e}t}^{\text{eff}}(K) : \mathfrak{G}'$$

by putting $\mathfrak{F}' := \mathbb{L}\iota_! \circ \mathbb{L}j^* \circ (-)^\sharp \circ \mathbb{L} \text{Perf}^*$. We remark that by Proposition 5.8, \mathfrak{F}' is also monoidal. The claim then follows from the equivalence of motives with and without transfers (Theorems 5.1 and 6.3). \square

Our goal is to prove that the adjunction of Corollary 7.1 is an equivalence of categories. To this aim, we recall the construction of the stable versions of the rigid motivic categories given in [Ayo15, Definition 2.5.27].

DEFINITION 7.2. Let T be the cokernel in $\mathbf{PST}(\mathbf{RigSm}/K)$ of the unit map $\Lambda_{\mathrm{tr}}(K) \rightarrow \Lambda_{\mathrm{tr}}(\mathbb{T}^1)$. We denote by $\mathbf{RigDM}_{\acute{\mathrm{e}}\mathrm{t}}(K, \Lambda)$ or simply by $\mathbf{RigDM}_{\acute{\mathrm{e}}\mathrm{t}}(K)$ the homotopy category of the stable $(\acute{\mathrm{e}}\mathrm{t}, \mathbb{B}^1)$ -local model structure on symmetric spectra $\mathbf{Spt}_T^{\Sigma}(\mathbf{Ch}_{\acute{\mathrm{e}}\mathrm{t}, \mathbb{B}^1} \mathbf{PST}(\mathbf{RigSm}/K))$.

As explained in [Ayo15, § 2.5], T is cofibrant and the cyclic permutation induces the identity on $T^{\otimes 3}$ in $\mathbf{RigDM}_{\acute{\mathrm{e}}\mathrm{t}}^{\mathrm{eff}}$. Moreover, by [Hov01, Theorem 9.3], $T \otimes -$ is a Quillen equivalence in this category, which is actually the universal model category where this holds (in some weak sense made precise by [Hov01, Theorem 5.1, Proposition 5.3 and Corollary 9.4]). We recall that the canonical functor $\mathbf{RigDM}_{\acute{\mathrm{e}}\mathrm{t}}^{\mathrm{eff}}(K) \rightarrow \mathbf{RigDM}_{\acute{\mathrm{e}}\mathrm{t}}(K)$ is fully faithful, as proved in [Ayo15, Corollary 2.5.49] as a corollary of the cancellation theorem [Ayo15, Theorem 2.5.38].

DEFINITION 7.3. We denote by $\Lambda(1)$ the motive $T[-1]$ in $\mathbf{RigDM}_{\acute{\mathrm{e}}\mathrm{t}}^{\mathrm{eff}}(K)$. For any positive integer d we let $\Lambda(d)$ be $\Lambda(1)^{\otimes d}$. The functor $(\cdot)(d) := (\cdot) \otimes \Lambda(d)$ is an auto-equivalence of $\mathbf{RigDM}_{\acute{\mathrm{e}}\mathrm{t}}(K)$ and its inverse will be denoted with $(\cdot)(-d)$.

DEFINITION 7.4. We denote by $\mathbf{RigDM}_{\acute{\mathrm{e}}\mathrm{t}}^{\mathrm{ct}}(K, \Lambda)$ or simply by $\mathbf{RigDM}_{\acute{\mathrm{e}}\mathrm{t}}^{\mathrm{ct}}(K)$ the full triangulated subcategory of $\mathbf{RigDM}_{\acute{\mathrm{e}}\mathrm{t}}(K, \Lambda)$ whose objects are the compact ones. They are of the form $M(d)$ for some compact object M in $\mathbf{RigDM}_{\acute{\mathrm{e}}\mathrm{t}}^{\mathrm{eff}}(K)$ and some d in \mathbb{Z} . This category is called the category of *constructible motives*.

We now present an important result that is a crucial step toward the proof of our main theorem. The motivic property it induces will be given right afterwards.

PROPOSITION 7.5. *Let \widehat{X} be a smooth affinoid perfectoid. The natural map of complexes*

$$\mathrm{Sing}^{\widehat{\mathbb{B}}^1}(\Lambda(\widehat{\mathbb{T}}^d))(\widehat{X}) \rightarrow \mathrm{Sing}^{\widehat{\mathbb{B}}^1}(\Lambda(\mathbb{T}^d))(\widehat{X})$$

is a quasi-isomorphism.

Proof. We let \widehat{X} be $\mathrm{Spa}(R, R^+)$. A map f in $\mathrm{Hom}(\widehat{X} \times \widehat{\mathbb{B}}^n, \mathbb{T}^d)$ [respectively in $\mathrm{Hom}(\widehat{X} \times \widehat{\mathbb{B}}^n, \widehat{\mathbb{T}}^d)$] corresponds to d elements f_1, \dots, f_d in the group $(R^+ \langle \tau_1^{1/p^\infty}, \dots, \tau_n^{1/p^\infty} \rangle)^\times$ [respectively in the group $(R^{b+} \langle \tau_1^{1/p^\infty}, \dots, \tau_n^{1/p^\infty} \rangle)^\times$] and the map between the two objects is induced by the multiplicative tilt map $R^{b+} \langle \tau_1^{1/p^\infty}, \dots, \tau_n^{1/p^\infty} \rangle \rightarrow R^+ \langle \tau_1^{1/p^\infty}, \dots, \tau_n^{1/p^\infty} \rangle$.

We now present some facts about homotopy theory for cubical objects, which mirror classical results for simplicial objects (see for example [May67, ch. IV]). We remark that the map of the statement is induced by a map of enriched cubical Λ -vector spaces (see [Ayo14b, Definition A.6]), which is obtained by adding Λ -coefficients to a map of enriched cubical sets

$$\mathrm{Hom}(\widehat{X} \times \widehat{\square}, \widehat{\mathbb{T}}^d) \rightarrow \mathrm{Hom}(\widehat{X} \times \widehat{\square}, \mathbb{T}^d).$$

Any enriched cubical object has connections in the sense of [BH81, § 1.2], induced by the maps m_i in [Ayo14b, Definition A.6]. We recall that the category of cubical sets with connections can be endowed with a model structure by which all objects are cofibrant and weak equivalences are defined through the geometric realization (see [Jar02]). Moreover, its homotopy category is

canonically equivalent to the one of simplicial sets, as cubical sets with connections form a strict test category by [Mal09].

The two cubical sets appearing above are abelian groups on each level and the maps defining their cubical structure are group homomorphisms. They therefore are cubical groups. By [Ton92], they are fibrant objects and their homotopy groups π_i coincide with the homology $H_i N$ of the associated normalized complexes of abelian groups (see Definition 3.13). The Λ -enrichment functor is tensorial with respect to the monoidal structure of cubical sets introduced in [BHS11, § 11.2]. Also, the cubical Dold–Kan functor that associates to a cubical Λ -module with connections its normalized complex (see [BHS11, § 14.8]) is a left Quillen functor. We deduce that in order to prove the statement of the proposition it suffices to show that the two normalized complexes of abelian groups are quasi-isomorphic. We also remark that it suffices to consider the case $d = 1$.

We prove the following claim: the n th homology of the complex $N((R \widehat{\otimes} \mathcal{O}(\widehat{\square}))^{+\times})$ is 0 for $n > 0$. Let f be invertible in $R^+ \langle \tau_1^{1/p^\infty}, \dots, \tau_n^{1/p^\infty} \rangle$ with $d_{r,\epsilon} f = 1$ for all (r, ϵ) . We claim that $f - 1$ is topologically nilpotent. Up to adding a topological nilpotent element, we can assume that $f \in R^+[\tau]$. Since f is invertible, its image in $(R^+/R^{\circ\circ})[\tau^{1/p^\infty}]$ is invertible as well. Invertible elements in this ring are just the invertible constants. We deduce that all coefficients of $f - f(0) = f - 1$ are topologically nilpotent, and hence $f - 1$ is topologically nilpotent. In particular, the element $H = f + \tau_{n+1}(1 - f)$ in $R^+ \langle \tau^{1/p^\infty}, \tau_{n+1}^{1/p^\infty} \rangle$ is invertible, satisfies $d_{r,\epsilon} H = 1$ for all ϵ and all $1 \leq r \leq n$ and determines a homotopy between f and 1. This proves the claim.

We can also prove that the 0th homology of the complex $N((R \widehat{\otimes} \mathcal{O}(\widehat{\square}))^{+\times})$ coincides with $R^{+\times}/(1 + R^{\circ\circ})$. This amounts to showing that the image of the ring map

$$\begin{aligned} \{f \in R^+ \langle \tau^{1/p^\infty} \rangle^\times : f(0) = 1\} &\rightarrow R^{+\times} \\ f &\mapsto f(1) \end{aligned}$$

coincides with $1 + R^{\circ\circ}$. Let f be invertible in $R^+ \langle \tau^{1/p^\infty} \rangle$ with $f(0) = 1$. As proved above, $f - 1$ is topologically nilpotent so that also $f(1) - 1$ is. Vice versa if $a \in R$ is topologically nilpotent then the element $1 + a\tau \in R^+ \langle \tau^{1/p^\infty} \rangle$ is invertible, satisfies $f(0) = 1$ and $f(1) = 1 + a$, proving the claim.

We are left to prove that the multiplicative map \sharp induces an isomorphism $(R^{b+})^\times/(1 + R^{b\circ\circ}) \rightarrow (R^+)^\times/(1 + R^{\circ\circ})$. We start by proving it is injective. Let $a \in R^{b+}$ such that $(a^\sharp - 1)$ is topologically nilpotent. Since $(a^\sharp - 1) = (a - 1)^\sharp$ in R^+/π we deduce that the element $(a - 1)^\sharp - (a^\sharp - 1)$ is also topologically nilpotent. We conclude that $(a - 1)^\sharp$, as well as $(a - 1)$, are topologically nilpotent, as desired.

We now prove surjectivity. Let a be invertible in R^+ . In particular, both a and a^{-1} are power-bounded. From the isomorphism $R^{b+}/\pi^b \cong R^+/\pi$ we deduce that there exists an element $b \in R^{b+}$ such that $b^\sharp = a + \pi\alpha = a(1 + \pi\alpha a^{-1})$ for some (power-bounded) element $\alpha \in R^+$. We deduce that $(1 + \pi\alpha a^{-1})$ lies in $1 + R^{\circ\circ}$ and that b^\sharp is invertible. Since the multiplicative structure of R^b is isomorphic to $\varprojlim_{x \mapsto x^p} R$ and \sharp is given by the projection to the last component, we deduce that as b^\sharp is invertible, then also b is. In particular, the image of $b \in (R^{b+})^\times$ in $(R^+)^\times/(1 + R^{\circ\circ})$ is equal to a , as desired. \square

We recall that by Corollary 7.1 there is an adjunction

$$\mathfrak{F} : \mathbf{RigDM}_{\text{ét}}^{\text{eff}}(K^b) \rightleftarrows \mathbf{RigDM}_{\text{ét}}^{\text{eff}}(K) : \mathfrak{G}$$

and our goal is to prove it is an equivalence.

PROPOSITION 7.6. *The motive $\mathfrak{G}\Lambda(d)$ is isomorphic to $\Lambda(d)$ for any positive integer d .*

Proof. The natural map $\Lambda(d) \rightarrow \mathfrak{G}\Lambda(d)$ is induced by the isomorphism $\mathfrak{F}\Lambda(d) \cong \Lambda(d)$. We need to prove it is an isomorphism. The motive $\Lambda(d)$ is a direct factor of the motive $\Lambda(\mathbb{T}^d)[-d]$ and the map above is induced by $\Lambda(\mathbb{T}^d) \rightarrow \mathfrak{G}\Lambda(\mathbb{T}^d)$. It suffices then to prove that the map $\Lambda(\mathbb{T}^d) \rightarrow \mathfrak{G}\Lambda(\mathbb{T}^d)$ is an isomorphism.

By the definition of the adjoint pair $(\mathfrak{F}, \mathfrak{G})$ given in Corollary 7.1, we can equivalently consider the adjunction

$$\mathbb{L}l_! \mathbb{L}j^* : \mathbf{PerfDA}_{\acute{e}t}^{\text{eff}}(K) \rightleftarrows \mathbf{RigDA}_{\acute{e}t}^{\text{eff}}(K) : \mathbb{R}j_* \mathbb{L}l^*$$

and prove that $\Lambda(\widehat{\mathbb{T}}^d) \rightarrow (\mathbb{R}j_* \circ \mathbb{L}l^*)\Lambda(\mathbb{T}^d)$ is an isomorphism in $\mathbf{PerfDA}_{\acute{e}t}^{\text{eff}}(K)$.

From Proposition 7.5 we deduce that the complexes $\text{Sing}^{\widehat{\mathbb{B}}^1} \Lambda(\widehat{\mathbb{T}}^d)$ and $j_* \text{Sing}^{\widehat{\mathbb{B}}^1} \Lambda(\mathbb{T}^d)$ are quasi-isomorphic in $\mathbf{ChPsh}(\text{PerfSm})$. Since j_* commutes with $\text{Sing}^{\widehat{\mathbb{B}}^1}$ and with ét-sheafification, the quasi-isomorphism above can be restated as

$$\text{Sing}^{\widehat{\mathbb{B}}^1} \Lambda(\widehat{\mathbb{T}}^d) \cong \mathbb{R}j_* \text{Sing}^{\widehat{\mathbb{B}}^1} \Lambda(\mathbb{T}^d).$$

Due to Proposition 3.26 and the isomorphism $\mathbb{L}l^* \Lambda(\mathbb{T}^d) \cong \Lambda(\widehat{\mathbb{T}}^d) \cong \mathbb{R}j_* \mathbb{L}l^* \Lambda(\mathbb{T}^d)$, as desired. \square

Remark 7.7. Along the proof of the previous proposition, we showed, in particular, that $\mathbb{R}j_* \mathbb{L}l^* \Lambda(\mathbb{T}^d) \cong \Lambda(\widehat{\mathbb{T}}^d)$.

Remark 7.8. Since j_* commutes with ét-sheafification, it preserves ét-weak equivalences. It also commutes with $\text{Sing}^{\widehat{\mathbb{B}}^1}$ and therefore preserves \mathbb{B}^1 -weak equivalences. We conclude that $\mathbb{R}j_* = j_*$ and, in particular, $\mathbb{R}j_*$ commutes with small direct sums.

DEFINITION 7.9. A rigid analytic variety with good coordinates $X_0 \rightarrow \mathbb{T}^N$, such that the induced maps $\Lambda(X_{h+1}) \rightarrow \Lambda(X_h)$ are invertible in $\mathbf{RigDM}_{\acute{e}t}^{\text{eff}}(K, \mathbb{Q})$, is called a *variety with very good coordinates*.

We are finally ready to present the proof of our main result.

THEOREM 7.10. *Let K be a perfectoid field and Λ be a \mathbb{Q} -algebra. The adjunction*

$$\mathfrak{F} : \mathbf{RigDM}_{\acute{e}t}^{\text{eff}}(K^{\flat}) \rightleftarrows \mathbf{RigDM}_{\acute{e}t}^{\text{eff}}(K) : \mathfrak{G}$$

is a monoidal, triangulated equivalence of categories.

Before the proof, we remark that by putting Theorems 3.23 and 6.9 together, the theorem above has the following restatement.

THEOREM 7.11. *Let K be a perfectoid field and Λ be a \mathbb{Q} -algebra. The adjunction*

$$\mathbb{L}l_! \mathbb{L}j^* : \mathbf{PerfDA}_{\acute{e}t}^{\text{eff}}(K) \rightleftarrows \mathbf{RigDM}_{\acute{e}t}^{\text{eff}}(K) : \mathbb{R}j_* \mathbb{L}l^*$$

is a monoidal, triangulated equivalence of categories.

Proof. By Theorem 5.5 the functor $\mathbb{L}l_! \mathbb{L}j^* : \mathbf{PerfDA}_{\text{ét}}^{\text{eff}}(K) \rightarrow \mathbf{RigDA}_{\text{ét}}^{\text{eff}}(K)$ sends the motive $\Lambda(\widehat{X})$ associated to a perfectoid $\widehat{X} = \varprojlim_h X_h$ to the motive $\Lambda(X_0)$ associated to X_0 up to rescaling indices. It is triangulated, commutes with sums, and its essential image contains motives $\Lambda(X_0)$ of varieties X_0 having very good coordinates $X_0 \rightarrow \mathbb{T}^N$ (see Definition 7.9). By Proposition 5.3, for every rigid variety with good coordinates $X_0 \rightarrow \mathbb{T}^N$ there exists an index h such that $X_h = X_0 \times_{\mathbb{T}^N} \mathbb{T}^N \langle \underline{v}^{\pm 1/p^h} \rangle$ has very good coordinates. Since $\text{char } K = 0$ the map $\mathbb{T}^N \langle \underline{v}^{\pm 1/p^h} \rangle \rightarrow \mathbb{T}^N$ is finite étale, and therefore also the map $X_h \rightarrow X_0$ is. We conclude that any rigid variety with good coordinates has a finite étale covering with very good coordinates, and hence the motives associated to varieties with very good coordinates generate the étale topos. In particular, the motives associated to them generate $\mathbf{RigDA}_{\text{ét}}^{\text{eff}}(K)$, and hence the functor $\mathbb{L}l_! \circ \mathbb{L}j^*$ maps a set of compact generators to a set of compact generators.

Since \mathfrak{F} is monoidal and $\mathfrak{F}(\Lambda(1)) = \Lambda(1)$ it extends formally to a monoidal functor from the category $\mathbf{RigDA}_{\text{ét}}^{\text{ct}}(K^b)$ to $\mathbf{RigDA}_{\text{ét}}(K)$ by putting $\mathfrak{F}(M(-d)) = \mathfrak{F}(M)(-d)$. Now let M, N in $\mathbf{RigDM}_{\text{ét}}(K^b)$ be twists of the motives associated to the analytification of smooth projective varieties X , respectively, X' . They are strongly dualizable objects of $\mathbf{RigDM}_{\text{ét}}(K^b)$ since $\Lambda_{\text{tr}}(X)$ and $\Lambda_{\text{tr}}(X')$ are strongly dualizable in $\mathbf{DM}_{\text{ét}}(K^b)$. Fix an integer d such that $N^\vee(d)$ lies in $\mathbf{RigDM}_{\text{ét}}^{\text{eff}}(K^b)$. The objects M, N, M^\vee and N^\vee lie in $\mathbf{RigDM}_{\text{ét}}^{\text{ct}}(K^b)$ and moreover $\mathfrak{F}(N^\vee) = \mathfrak{F}(N)^\vee$. From Lemma 7.12 we also deduce that the functor \mathfrak{F} induces a bijection

$$\mathbf{RigDM}_{\text{ét}}^{\text{ct}}(K^b)(M \otimes N^\vee, \Lambda) \cong \mathbf{RigDM}_{\text{ét}}(K)(\mathfrak{F}(M) \otimes \mathfrak{F}(N)^\vee, \Lambda).$$

By means of the Cancellation Theorem [Ayo15, Corollary 2.5.49], the first set is isomorphic to $\mathbf{RigDM}_{\text{ét}}^{\text{eff}}(K^b)(M, N)$ and the second is isomorphic to $\mathbf{RigDM}_{\text{ét}}^{\text{eff}}(K)(\mathfrak{F}(M), \mathfrak{F}(N))$. We then deduce that all motives M associated to the analytification of smooth projective varieties lie in the left orthogonal of the cone of the map $N \rightarrow \mathfrak{G}\mathfrak{F}N$, which is closed under direct sums and cones. Since Λ is a \mathbb{Q} -algebra, such motives generate $\mathbf{RigDM}_{\text{ét}}^{\text{eff}}(K^b)$ by means of [Ayo15, Theorem 2.5.35]. We conclude that $N \cong \mathfrak{G}\mathfrak{F}N$. Therefore, the category \mathbf{T} of objects N such that $N \cong \mathfrak{G}\mathfrak{F}N$ contains all motives associated to the analytification of smooth projective varieties. It is clear that \mathbf{T} is closed under cones. The functors \mathfrak{F} and $\mathbb{L}l^*$ commute with direct sums as they are left adjoint functors. As pointed out in Remark 7.8 also the functor $\mathbb{R}j_*$ does. Since \mathfrak{G} is a composite of $\mathbb{R}j_* \mathbb{L}l^*$ with equivalences of categories, it commutes with small sums as well. We conclude that \mathbf{T} is closed under direct sums. Using again [Ayo15, Theorem 2.5.35] we deduce $\mathbf{T} = \mathbf{RigDM}_{\text{ét}}^{\text{eff}}(K^b)$, proving that \mathfrak{F} is fully faithful. This is enough to prove that it is an equivalence of categories, by applying [Ayo15, Lemma 1.3.32]. \square

LEMMA 7.12. *Let M be an object of $\mathbf{RigDA}_{\text{ét}}^{\text{ct}}(K^b)$. The functor \mathfrak{F} induces an isomorphism*

$$\mathbf{RigDM}_{\text{ét}}^{\text{ct}}(K^b)(M, \Lambda) \cong \mathbf{RigDM}_{\text{ét}}(K)(\mathfrak{F}(M), \Lambda).$$

Proof. Suppose that d is an integer such that $M(d)$ lies in $\mathbf{RigDA}_{\text{ét}}^{\text{eff}}(K^b)$. One has $\mathfrak{F}\Lambda(d) \cong \Lambda(d)$ and by Proposition 7.6 the unit map $\eta : \Lambda(d) \rightarrow \mathfrak{G}\mathfrak{F}\Lambda(d)$ is an isomorphism. In particular, from the adjunction $(\mathfrak{F}, \mathfrak{G})$ we obtain a commutative square

$$\begin{array}{ccc} \mathbf{RigDM}_{\text{ét}}^{\text{eff}}(K^b)(M(d), \Lambda(d)) & \xrightarrow{\mathfrak{F}} & \mathbf{RigDM}_{\text{ét}}^{\text{eff}}(K)(\mathfrak{F}M(d), \mathfrak{F}\Lambda(d)) \\ \downarrow = & & \downarrow \sim \\ \mathbf{RigDM}_{\text{ét}}^{\text{eff}}(K^b)(M(d), \Lambda(d)) & \xrightarrow[\sim]{\eta} & \mathbf{RigDM}_{\text{ét}}^{\text{eff}}(K)(M(d), (\mathfrak{G}\mathfrak{F})\Lambda(d)) \end{array}$$

in which the top arrow is then an isomorphism. By the Cancellation Theorem [Ayo15, Corollary 2.5.49] we also obtain the following commutative square

$$\begin{CD} \mathbf{RigDM}_{\acute{e}t}^{\text{ct}}(K^b)(M(d), \Lambda(d)) @>\mathfrak{F}>> \mathbf{RigDM}_{\acute{e}t}(K)(\mathfrak{F}M(d), \Lambda(d)) \\ @V\sim VV @VV\sim V \\ \mathbf{RigDM}_{\acute{e}t}^{\text{eff}}(K^b)(M(d), \Lambda(d)) @>\mathfrak{F}>> \mathbf{RigDM}_{\acute{e}t}^{\text{eff}}(K)(\mathfrak{F}M(d), \Lambda(d)), \end{CD}$$

and hence also the top arrow is an isomorphism. We conclude the claim from the following commutative square, whose vertical arrows are isomorphisms since the functor $(\cdot)(d)$ is invertible in $\mathbf{RigDM}_{\acute{e}t}(K)$.

$$\begin{CD} \mathbf{RigDM}_{\acute{e}t}^{\text{ct}}(K^b)(M, \Lambda) @>\mathfrak{F}>> \mathbf{RigDM}_{\acute{e}t}(K)(\mathfrak{F}M, \Lambda) \\ @V(\cdot)(d)\sim VV @VV(\cdot)(d)\sim V \\ \mathbf{RigDM}_{\acute{e}t}^{\text{ct}}(K^b)(M(d), \Lambda(d)) @>\mathfrak{F}>> \mathbf{RigDM}_{\acute{e}t}(K)(\mathfrak{F}M(d), \Lambda(d)) \quad \square \end{CD}$$

Remark 7.13. In the proof of Theorem 7.10 we again used the hypothesis that Λ is a \mathbb{Q} -algebra in order to apply [Ayo15, Theorem 2.5.35], which states that the motives associated to the analytification of smooth projective varieties generate $\mathbf{RigDM}_{\acute{e}t}^{\text{eff}}(K^b)$.

Remark 7.14. In the proof, we also showed that the motives $\Lambda(X)[i]$ where X has very good coordinates and $i \in \mathbb{Z}$ generate $\mathbf{RigDM}_{\acute{e}t}^{\text{eff}}(K)$ as a triangulated category with small sums.

COROLLARY 7.15. *Let $X_0 \rightarrow \mathbb{T}^N$ be a variety with very good coordinates. Then $\mathbb{R}j_*\mathbb{L}l^*\Lambda(X_0) \cong \Lambda(\varprojlim_h X_h)$.*

Proof. By the description given in Theorem 5.5, we conclude that $\mathbb{R}l_!\mathbb{L}j^*(\Lambda(\varprojlim_h X_h)) \cong \Lambda(X_0)$. The claim then follows from Theorem 7.10, which shows that $\mathbb{R}l_! \circ \mathbb{L}j^*$ is a quasi-inverse of $\mathbb{R}j_* \circ \mathbb{L}l^*$. □

We remark that the proof of Theorem 7.10 also induces the following result.

COROLLARY 7.16. *The functor*

$$\begin{aligned} \mathfrak{F} : \mathbf{RigDM}_{\acute{e}t}^{\text{ct}}(K^b) &\rightarrow \mathbf{RigDM}_{\acute{e}t}^{\text{ct}}(K) \\ M(d) &\mapsto (\mathfrak{F}M)(d) \end{aligned}$$

is a monoidal equivalence of categories.

We can refine the previous corollary by stating the stable version of our main result (Theorem 7.26), which is based on the following intermediate results. For the definitions of spectra of model categories, we refer to [Ayo07, § 4.3] and [Hov01].

DEFINITION 7.17. Let T be the cokernel of the unit map $\Lambda \rightarrow \Lambda(\mathbb{T}^1)$ in $\mathbf{Psh}(\mathbf{RigSm}/K)$ and let \widehat{T} be the cokernel of the unit map $\Lambda \rightarrow \Lambda(\widehat{\mathbb{T}}^1)$ in $\mathbf{Psh}(\mathbf{PerfSm}/K)$. They are direct factors of cofibrant objects, and hence cofibrant.

(i) We denote by $\mathbf{RigDA}_{\acute{e}t}(K, \Lambda)$ the homotopy category of the stable $(\acute{e}t, \mathbb{B}^1)$ -local model structure on spectra $\mathbf{Spt}_T(\mathbf{Ch}_{\acute{e}t, \mathbb{B}^1} \mathbf{Psh}(\mathbf{RigSm}/K))$. We omit Λ whenever the context allows it.

(ii) We denote by $\widehat{\mathbf{RigDA}}_{\acute{e}t, \mathbb{B}^1}(K, \Lambda)$ the homotopy category of the stable $(\acute{e}t, \widehat{\mathbb{B}^1})$ -local model structure on spectra $\mathbf{Spt}_{\iota^*T}(\mathbf{Ch}_{\acute{e}t, \mathbb{B}^1} \mathbf{Psh}(\widehat{\mathbf{RigSm}}/K))$. We omit Λ whenever the context allows it.

(iii) We denote by $\mathbf{PerfDA}_{\acute{e}t}(K, \Lambda)$ the homotopy category of the stable $(\acute{e}t, \widehat{\mathbb{B}^1})$ -local model structure on spectra $\mathbf{Spt}_{\widehat{T}}(\mathbf{Ch}_{\acute{e}t, \mathbb{B}^1} \mathbf{Psh}(\mathbf{PerfSm}/K))$. We omit Λ whenever the context allows it.

(iv) For \mathbf{C} equal to \mathbf{RigSm} , $\widehat{\mathbf{RigSm}}$ or \mathbf{PerfSm} , we denote by Sus_k the k th *suspension functor* from the model category of complexes of presheaves on \mathbf{C} to the associated spectra, which is the left adjoint of the functor

$$\text{Ev}_k : (M_n)_{n \in \mathbb{N}} \mapsto M_k.$$

Remark 7.18. The functor $- \otimes T$ [respectively $- \otimes \iota^*T$ respectively $- \otimes \widehat{T}$] has a prolongation to a left Quillen endofunctor on the associated spectra, which is furthermore a Quillen equivalence. The category of spectra is the universal model category with this property, in a weak sense made precise by [Hov01, Theorem 5.1 and Proposition 5.3].

Remark 7.19. In contrast with Definition 7.2, we use above the categories of *non-symmetric* spectra. On the other hand, we remark that by [Ayo07, Proposition 4.3.47] the model categories of symmetric and non-symmetric spectra are Quillen equivalent. We then conclude by Proposition 3.23, Theorem 6.3, Proposition 6.9 and [Hov01, Theorem 5.5] that the canonical adjunctions

$$\begin{aligned} \mathbf{RigDA}_{\acute{e}t}(K, \Lambda) &\rightleftarrows \mathbf{RigDM}_{\acute{e}t}(K, \Lambda), \\ \mathbf{RigDM}_{\acute{e}t}(K^b, \Lambda) &\rightleftarrows \mathbf{PerfDA}_{\acute{e}t}(K, \Lambda) \end{aligned}$$

are equivalences of categories, and the prolongation of $- \otimes T$ [respectively $- \otimes \widehat{T}$] corresponds to the functor $M \mapsto M(1)$.

Remark 7.20. We point out that the functor Sus_0 has an explicit description. For example, in the case of $\mathbf{Ch}_{\acute{e}t, \mathbb{B}^1} \mathbf{Psh}(\mathbf{RigSm}/K)$ it sends a complex \mathcal{F} to the spectrum $(\mathcal{F} \otimes T^{\otimes n})_{n \in \mathbb{N}}$ with the obvious transition maps. The functor Sus_k is the composition $t^k \text{Sus}_0$, where t is the *shift functor* such that $\text{Ev}_0 tM = 0$ and $\text{Ev}_{k+1} tM = \text{Ev}_k M$ for any spectrum M . The pair $(\text{Sus}_k, \text{Ev}_k)$ is a Quillen adjunction (see [Ayo07, Lemma 4.3.24]) as well as the pair (t, s) , where s is such that $\text{Ev}_k sM = \text{Ev}_{k+1} M$ for any spectrum M (see [Hov01, Definition 3.7]).

DEFINITION 7.21. We denote by

$$\mathbf{Spt}_{\iota^*} : \mathbf{Spt}_T(\mathbf{Ch}_{\acute{e}t, \mathbb{B}^1} \mathbf{Psh}(\mathbf{RigSm})) \rightleftarrows \mathbf{Spt}_{\iota^*T}(\mathbf{Ch}_{\acute{e}t, \widehat{\mathbb{B}^1}} \mathbf{Psh}(\widehat{\mathbf{RigSm}})) : \mathbf{Spt}_{\iota_*}$$

the Quillen adjunction induced by the pair (ι^*, ι_*) by means of [Hov01, Proposition 5.3].

The natural map $\widehat{\mathbb{T}^1} \rightarrow \iota^* \mathbb{T}^1$ induces a map $\widehat{T} \rightarrow j_* \iota^* T$, which, in turn, defines a natural transformation $\tau : j^*(- \otimes \widehat{T}) \rightarrow (j^* -) \otimes \iota^* T$. Nevertheless, it is not clear that for a smooth affinoid perfectoid \widehat{X} the map $\tau : j^*(\Lambda(\widehat{X}) \otimes \widehat{T}) \rightarrow (j^* \Lambda(\widehat{X})) \otimes \iota^* T$ is a weak equivalence, so that we can not apply the criterion of [Hov01, Proposition 5.3] to define a Quillen adjunction between $\mathbf{Spt}_{\widehat{T}}(\mathbf{Ch}_{\acute{e}t, \widehat{\mathbb{B}^1}} \mathbf{Psh}(\mathbf{PerfSm}/K))$ and $\mathbf{Spt}_{\iota^*T}(\mathbf{Ch}_{\acute{e}t, \widehat{\mathbb{B}^1}} \mathbf{Psh}(\widehat{\mathbf{RigSm}}/K))$. Therefore, the following proposition, albeit easy to prove, has a non-trivial content.

PROPOSITION 7.22. *The functor*

$$\begin{aligned} \mathrm{Spt} j_* : \mathbf{Spt}_{\iota^*T}(\mathbf{Ch}_{\acute{e}t, \mathbb{B}^1} \mathbf{Psh}(\widehat{\mathbf{RigSm}}/K)) &\rightarrow \mathbf{Spt}_{\widehat{T}}(\mathbf{Ch}_{\acute{e}t, \mathbb{B}^1} \mathbf{Psh}(\mathrm{PerfSm}/K)) \\ (M_n)_{n \in \mathbb{N}} &\mapsto (j_* M_n)_{n \in \mathbb{N}} \end{aligned}$$

is well defined, preserves stable weak equivalences and induces a triangulated functor

$$\mathbb{R} \mathrm{Spt} j_* : \widehat{\mathbf{RigDA}}_{\acute{e}t, \mathbb{B}^1}(K, \Lambda) \rightarrow \mathbf{PerfDA}_{\acute{e}t}(K, \Lambda).$$

Proof. If (M_n) is a spectrum we can define the transition maps $j_* M_n \otimes \widehat{T} \rightarrow j_* M_{n+1}$ with the following composition

$$j_* M_n \otimes \widehat{T} \xrightarrow{\tau} j_* M_n \otimes j_* \iota^* T \cong j_*(M_n \otimes \iota^* T) \rightarrow j_* M_{n+1}$$

deduced by the transition maps $M_n \otimes \iota^* T \rightarrow M_{n+1}$. Moreover, as shown in the first part of the proof of [Hov01, Proposition 5.3] $\mathrm{Spt} j_*$ is a right Quillen functor with respect to the projective model structures on spectra, whose weak equivalences are level-wise weak equivalences. It also preserves such weak equivalences, as proved in Remark 7.8.

The stable model structure on spectra is obtained as a left Bousfield localization with respect to the maps $\mathrm{Sus}_{n+1}(\mathcal{F} \otimes \iota^* T) \rightarrow \mathrm{Sus}_n \mathcal{F}$ [respectively $\mathrm{Sus}_{n+1}(\mathcal{F} \otimes \widehat{T}) \rightarrow \mathrm{Sus}_n \mathcal{F}$] as \mathcal{F} runs among cofibrant objects (see for example [Ayo07, Definition 4.3.29]). We now prove that $\mathrm{Spt} j_*$ also preserves stable weak equivalences.

Consider the natural map

$$\mathrm{Ev}_k(\mathrm{Spt} j_* \mathrm{Sus}_{n+1}(\mathcal{F} \otimes \iota^* T)) \rightarrow \mathrm{Ev}_k(\mathrm{Spt} j_* \mathrm{Sus}_n \mathcal{F}).$$

For $k > n$ it is an equality, and hence the map

$$\mathrm{Spt} j_* \mathrm{Sus}_{n+1}(\mathcal{F} \otimes \iota^* T) \rightarrow \mathrm{Spt} j_* \mathrm{Sus}_n \mathcal{F}$$

is a stable weak equivalence by [Ayo07, Lemma 4.3.59]. We then deduce that $\mathrm{Spt} j_*$ preserves stable weak equivalences as claimed.

The fact that $\mathbb{R} \mathrm{Spt} j_*$ is triangulated follows from the fact that it coincides with the restriction to stably local spectra of the functor induced by $\mathrm{Spt} j_*$ on the homotopy categories of the projective model structures on spectra, which is triangulated being a right derived Quillen functor. \square

Remark 7.23. The functor $\mathbb{R} \mathrm{Spt} j_*$ commutes with small direct sums by its explicit description and Remark 7.8.

PROPOSITION 7.24. *There is an invertible natural transformation of functors from the category $\mathbf{RigDA}_{\acute{e}t}^{\mathrm{eff}}(K, \Lambda)$ to $\mathbf{PerfDA}_{\acute{e}t}(K, \Lambda)$:*

$$\eta : \mathbb{L} \mathrm{Sus}_0 \circ \mathbb{R} j_* \circ \mathbb{L} \iota^* \xrightarrow{\sim} \mathbb{R} \mathrm{Spt} j_* \circ \mathbb{L} \mathrm{Spt} \iota^* \circ \mathbb{L} \mathrm{Sus}_0.$$

Proof. For any cofibrant object \mathcal{F} in $\mathbf{Ch} \mathbf{Psh}(\mathbf{RigSm})$ one has

$$\begin{aligned} \mathrm{Ev}_k(\mathbb{L} \mathrm{Sus}_0 \circ \mathbb{R} j_* \circ \mathbb{L} \iota^*)(\mathcal{F}) &\cong Q(j_* \iota^* \mathcal{F}) \otimes \widehat{T}^{\otimes k}, \\ \mathrm{Ev}_k(\mathbb{R} \mathrm{Spt} j_* \circ \mathbb{L} \mathrm{Spt} \iota^* \circ \mathbb{L} \mathrm{Sus}_0)(\mathcal{F}) &\cong j_* \iota^* \mathcal{F} \otimes j_*(\iota^* T)^{\otimes k}, \end{aligned}$$

where Q is a cofibrant-replacement functor on $\mathbf{ChPsh}(\widehat{\mathbf{RigSm}})$. The natural transformation η of the statement is then induced by the canonical maps $Q(j_*\iota^*\mathcal{F}) \rightarrow j_*\iota^*\mathcal{F}$ and $\widehat{T} \rightarrow j_*\iota^*T$.

The two functors of the statement are triangulated and commute with small direct sums by Remarks 7.8, 7.23 and Proposition 7.22. We deduce that the subcategory of $\mathbf{RigDA}_{\acute{e}t}^{\text{eff}}(K, \Lambda)$ on which η is invertible is triangulated and closed under direct sums.

By Proposition 3.18 and Remark 7.14 we conclude that it suffices to prove that η is invertible on a fixed motive $\Lambda(X)[i]$ with X with very good coordinates and $i \in \mathbb{Z}$. For such an object, using Corollary 7.15, we have an explicit description of a cofibrant replacement of $j_*\iota^*\Lambda(X)[i]$ namely $\Lambda(\widehat{X})[i]$, where $\widehat{X} = \varprojlim_h X_h$ is built with respect to some very good coordinates $X \rightarrow \mathbb{T}^N$ of X . We are left to prove that the maps $\Lambda(\widehat{X}) \otimes \widehat{T}^{\otimes k}[i] \rightarrow j_*\iota^*\Lambda(X) \otimes j_*(\iota^*T)^{\otimes k}[i]$ induce a weak equivalence of spectra. We claim that each one of them is a weak equivalence.

The object T [respectively \widehat{T}] is a direct summand of $\Lambda(\mathbb{T}^1)$ [respectively $\Lambda(\widehat{\mathbb{T}}^1)$] and the decomposition is compatible with the map $\widehat{T} \rightarrow j_*\iota^*T$. Therefore, in order to prove the claim we can show that the maps

$$\Lambda(\widehat{X} \times \widehat{\mathbb{T}}^k) \cong \Lambda(\widehat{X}) \otimes \Lambda(\widehat{\mathbb{T}}^k) \rightarrow j_*\iota^*\Lambda(X) \otimes j_*\Lambda(\mathbb{T}^k) \cong j_*\iota^*\Lambda(X \times \mathbb{T}^k)$$

are weak equivalences, and this follows from Corollary 7.15 and Remark 7.7. □

We define the following composite functor (see Remark 7.19):

$$\mathfrak{G}^{\text{st}} : \mathbf{RigDM}_{\acute{e}t}(K) \cong \mathbf{RigDA}_{\acute{e}t}(K) \xrightarrow{\mathbb{R}\text{Spt } j_* \circ \mathbb{L}\text{Spt } \iota^*} \mathbf{PerfDA}_{\acute{e}t}(K) \cong \mathbf{RigDM}_{\acute{e}t}(K^b),$$

which can be inserted in the diagram

$$\begin{array}{ccc} \mathbf{RigDM}_{\acute{e}t}^{\text{eff}}(K, \Lambda) & \xrightarrow[\sim]{\mathfrak{G}} & \mathbf{RigDM}_{\acute{e}t}^{\text{eff}}(K^b, \Lambda) \\ \downarrow & & \downarrow \\ \mathbf{RigDM}_{\acute{e}t}(K, \Lambda) & \xrightarrow{\mathfrak{G}^{\text{st}}} & \mathbf{RigDM}_{\acute{e}t}(K^b, \Lambda) \end{array}$$

whose vertical functors are fully faithful by the Cancellation Theorem [Ayo15, Corollary 2.5.49]. It is commutative, up to a natural transformation, by means of Proposition 7.24.

PROPOSITION 7.25. *The category $\mathbf{RigDM}_{\acute{e}t}(K, \Lambda)$ [respectively $\mathbf{RigDM}_{\acute{e}t}^{\text{ct}}(K, \Lambda)$] is generated, as a triangulated category with small sums, by the objects of $\mathbf{RigDM}_{\acute{e}t}^{\text{ct}}(K, \Lambda)$ [respectively $\mathbf{RigDM}_{\acute{e}t}^{\text{ct}}(K^b, \Lambda)$].*

Proof. We prove the statement only for K . From the Cancellation Theorem [Ayo15, Corollary 2.5.49] we deduce that any object M of $\mathbf{RigDM}_{\acute{e}t}(K, \Lambda)$ is isomorphic to $\text{hocolim}_n M_n(-n)$ with M_n in $\mathbf{RigDM}_{\acute{e}t}^{\text{eff}}(K, \Lambda)$ (see the proof of [Kah13, Proposition 7.4.1]). In particular, it sits in a distinguished triangle

$$\bigoplus M_n(-n) \longrightarrow \bigoplus M_n(-n) \longrightarrow M \longrightarrow .$$

Since $\mathbf{RigDM}_{\acute{e}t}^{\text{eff}}(K, \Lambda)$ is generated by compact objects by Proposition 3.18, we conclude that each $M_n(-n)$ lies in the triangulated category with small sums generated by the objects of $\mathbf{RigDM}_{\acute{e}t}^{\text{ct}}(K, \Lambda)$ so that M also does. □

We can finally prove the stable version of the main result of this article.

THEOREM 7.26. *The functor*

$$\mathfrak{G}^{\text{st}} : \mathbf{RigDM}_{\text{ét}}(K, \Lambda) \rightarrow \mathbf{RigDM}_{\text{ét}}(K^{\flat}, \Lambda)$$

is a monoidal triangulated equivalence of categories and, up to a natural transformation, it restricts to the equivalence

$$\mathfrak{G} : \mathbf{RigDM}_{\text{ét}}^{\text{eff}}(K, \Lambda) \rightarrow \mathbf{RigDM}_{\text{ét}}^{\text{eff}}(K^{\flat}, \Lambda)$$

of Theorem 7.10 on the subcategories of effective motives.

Proof. By its definition, the functor $\mathbb{R}\text{Spt } j_*$ commutes with the derived shift functor $\mathbb{L}t$ (see Remark 7.20). We claim that $\mathbb{L}\text{Spt } \iota^*$ also does. We can equivalently show that $\mathbb{R}\text{Spt } \iota_*$ commutes with $\mathbb{R}s$ and this again follows from their definitions (see Remark 7.20 and [Hov01, Theorem 5.3]).

By [Hov01, Theorem 3.8] the functor $\mathbb{L}t$ on $\mathbf{RigDA}_{\text{ét}}(K, \Lambda)$ [respectively on $\mathbf{PerfDA}_{\text{ét}}(K, \Lambda)$] is a quasi-inverse of the prolongation of $\mathbb{L}(- \otimes T)$ [respectively of $\mathbb{L}(- \otimes \widehat{T})$]. Therefore, we conclude that for any M in $\mathbf{RigDA}_{\text{ét}}(K, \Lambda)$:

$$(\mathbb{R}\text{Spt } j_* \circ \mathbb{L}\text{Spt } \iota^*)(\mathbb{L}(- \otimes T))^{-1}M \cong (\mathbb{L}(- \otimes \widehat{T}))^{-1}(\mathbb{R}\text{Spt } j_* \circ \mathbb{L}\text{Spt } \iota^*)M,$$

which implies for any M in $\mathbf{RigDM}_{\text{ét}}(K, \Lambda)$ the following canonical isomorphism:

$$\mathfrak{G}^{\text{st}}(M(-1)) \cong (\mathfrak{G}^{\text{st}}M)(-1).$$

Since we already showed in Proposition 7.24 that \mathfrak{G}^{st} restricts to \mathfrak{G} on effective motives, we conclude that it restricts on $\mathbf{RigDM}_{\text{ét}}^{\text{ct}}(K, \Lambda)$ to a quasi-inverse of the functor \mathfrak{F} of Corollary 7.16. In particular, this restriction is fully faithful and its essential image contains $\mathbf{RigDM}_{\text{ét}}^{\text{ct}}(K^{\flat}, \Lambda)$. By Remark 7.23 we also deduce that \mathfrak{G}^{st} commutes with small direct sums. The claim of the theorem then follows from Proposition 7.25 and [Ayo15, Lemma 1.3.32]. \square

Remark 7.27. It is worth noticing that along the proof of our main theorem, we have not used any result on almost algebra (which nonetheless has a critical role in the theory of perfectoid spaces).

Remark 7.28. The reader may wonder if the equivalence of categories $\mathbf{RigDM}_{\text{ét}}(K, \Lambda) \cong \mathbf{RigDM}_{\text{ét}}(K^{\flat}, \Lambda)$ still holds true for more general rings of coefficients Λ . The hypothesis $\mathbb{Q} \subset \Lambda$ has been used several times along the proof of the main statement, and most crucially in the following two instances. First, it was used to invoke the results of [Vez17] on the equivalence of motives with and without transfers (see Remarks 5.2 and 6.4). Secondly, the hypothesis $\mathbb{Q} \subset \Lambda$ was used in order to apply the result of Ayoub [Ayo15, Theorem 2.5.35] about a generating set for the triangulated category $\mathbf{RigDM}(K^{\flat}, \Lambda)$ (see Remark 7.13).

On the other hand, we conjecture that the tilting equivalence $\mathbf{RigDM}_{\text{ét}}(K, \Lambda) \cong \mathbf{RigDM}_{\text{ét}}(K^{\flat}, \Lambda)$ can be generalized to a ring of coefficients Λ over $\mathbb{Z}[1/p]$. As a matter of fact, for any prime $\ell \neq p$ we expect the category $\mathbf{RigDM}_{\text{ét}}(K, \mathbb{Z}/\ell)$ to be equivalent to the derived category of Galois representations over \mathbb{Z}/ℓ , similarly to the case of $\mathbf{DM}_{\text{ét}}^{\text{eff}}(K, \mathbb{Z}/\ell)$ (see [MVW06, Theorem 9.35]). The tilting equivalence in the case $\Lambda = \mathbb{Z}/\ell$ would then follow from the classic theorem of Fontaine and Wintenberger.

ACKNOWLEDGEMENTS

This paper is part of my PhD thesis, carried out in a co-tutelle program between the University of Milan and the University of Zurich. I am incredibly indebted to my advisor Joseph Ayoub for his suggestion of pursuing the present project, his constant and generous guidance, the outstanding amount of insight that he kindly shared with me, and his endless patience. I wish to express my gratitude to my co-advisor Luca Barbieri Viale, for his invaluable encouragement and the numerous mathematical discussions throughout the development of my thesis. I am grateful to Peter Scholze for answering many questions and also to Yves André, Fabrizio Andreatta and an anonymous referee for their remarks and suggestions. I had the opportunity to give a detailed series of talks on this work at the University of Zurich: I kindly thank Andrew Kresch for this chance and the various discussions on this topic. For the plentiful remarks on preliminary versions of this work and his precious and friendly support, I also wish to thank Simon Pepin Lehalleur.

Appendix A. An implicit function theorem and approximation results

The aim of this appendix is to prove Proposition 4.1, which will be obtained as a corollary of several intermediate approximation results for maps defined from objects of $\widehat{\text{RigSm}}^{\text{gc}}$ to rigid analytic varieties.

Along this section, we assume that K is a complete non-archimedean field. We begin our analysis with the analog of the inverse mapping theorem, which is a variant of [Igu00, Theorem 2.1.1].

PROPOSITION A.1. *Let R be a K -algebra, let $\sigma = (\sigma_1, \dots, \sigma_n)$ and $\tau = (\tau_1, \dots, \tau_m)$ be two systems of coordinates and let $P = (P_1, \dots, P_m)$ be a collection of polynomials in $R[\sigma, \tau]$ such that $P(\sigma = 0, \tau = 0) = 0$ and $\det(\partial P_i / \partial \tau_j)(\sigma = 0, \tau = 0) \in R^\times$. There exists a unique collection $F = (F_1, \dots, F_m)$ of m formal power series in $R[[\sigma]]$ such that $F(\sigma = 0) = 0$ and $P(\sigma, F(\sigma)) = 0$ in $R[[\sigma]]$.*

Moreover, if R is a Banach K -algebra, then F_1, \dots, F_m have a positive radius of convergence.

Proof. Let f be the polynomial $\det(\partial P_i / \partial \tau_j)$ in $R[\sigma, \tau]$ and let S be the ring $R[\sigma, \tau]_f / (P)$. The induced map $R[\sigma] \rightarrow S$ is étale, and from the hypothesis $f(0, 0) \in R^\times$ we conclude that the map $R[\sigma, \tau] / (P) \rightarrow R, (\sigma, \tau) \mapsto 0$ factors through S .

Suppose given a factorization as $R[\sigma]$ -algebras $S \rightarrow R[\sigma] / (\sigma)^n \rightarrow R$ of the map $S \rightarrow R$. By the étale lifting property (see [EGAIV, Definition IV.17.1.1 and Corollary IV.17.6.2]) applied to the square

$$\begin{array}{ccc}
 R[\sigma] & \longrightarrow & R[\sigma] / (\sigma)^{n+1} \\
 \downarrow & \nearrow \exists! & \downarrow \\
 S & \longrightarrow & R[\sigma] / (\sigma)^n
 \end{array}$$

we obtain a uniquely defined $R[\sigma]$ -linear map $S \rightarrow R[\sigma] / (\sigma)^{n+1}$ factoring $S \rightarrow R$, and hence by induction a uniquely defined $R[\sigma]$ -linear map $R[\sigma, \tau] / (P) \rightarrow R[[\sigma]]$ factoring $R[\sigma, \tau] / (P) \rightarrow R$, as desired. The power series F_i is the image of τ_i via this map.

Assume now that R is a Banach K -algebra. We want to prove that the array $F = (F_1, \dots, F_m)$ of formal power series in $R[[\sigma]]$ constructed above is convergent around 0. As R is complete, this amounts to proving estimates on the valuation of the coefficients of F . To this aim, we now try to give an explicit description of them, depending on the coefficients of P . Whenever I is

a n -multi-index $I = (i_1, \dots, i_n)$ we denote by σ^I the product $\sigma_1^{i_1} \cdot \dots \cdot \sigma_n^{i_n}$ and we adopt the analogous notation for τ .

We remark that the claim is not affected by any invertible R -linear transformation of the polynomials P_i . Therefore, by multiplying the column vector P by the matrix $(\partial P_i / \partial \tau_j)(0, 0)^{-1}$ we reduce to the case in which $(\partial P_i / \partial \tau_j)(0, 0) = \delta_{ij}$. We can then write the polynomials P_i in the following form:

$$P_i(\sigma, \tau) = \tau_i - \sum_{|J|+|H|>0} c_{iJH} \sigma^J \tau^H,$$

where J is an n -multi-index, H is an m -multi-index and the coefficients c_{iJH} equal 0 whenever $|J| = 0$ and $|H| = 1$.

We will determine the functions $F_i(\sigma)$ explicitly. We start by writing them as

$$F_i(\sigma) = \sum_{|I|>0} d_{iI} \sigma^I$$

with unknown coefficients d_{iI} for any n -multi-index I . We denote their q -homogeneous parts by

$$F_{iq}(\sigma) := \sum_{|I|=q} d_{iI} \sigma^I.$$

We need to solve the equation $P(\sigma, F(\sigma)) = 0$, which can be rewritten as

$$F_i(\sigma) = \sum_{J,H} c_{iJH} \sigma^J \left(\prod_{r=1}^m F_r(\sigma)^{h_r} \right),$$

where we denote by h_r the components of the m -multi-index H .

By comparing the q -homogeneous parts we get

$$F_{iq}(\sigma) = \sum_{(J,H,\Phi) \in \Sigma_{iq}} c_{iJH} \sigma^J \prod_{r=1}^m \prod_{s=1}^{h_r} F_{r,\Phi(r,s)}(\sigma),$$

where the set Σ_{iq} consists of triples (J, H, Φ) in which J is a n -multi-index, H is a m -multi-index and Φ is a function that associates to any element (r, s) of the set

$$\{(r, s) : r = 1, \dots, m; s = 1, \dots, h_r\}$$

a positive (non-zero!) integer $\Phi(r, s)$ such that $\sum \Phi(r, s) = q - |J|$.

If $\Phi(r, s) \geq q$ for some r we see by definition that $|J| = 0$, $|H| = 1$ and we know that in this case $c_{i0H} = 0$. In particular, we conclude that the right-hand side of the formula above involves only $F_{rq'}$ with $q' < q$. Hence, we can determine the coefficients d_{iI} by induction on $|I|$. Moreover, by construction, each coefficient d_{iI} can be expressed as

$$d_{iI} = Q_{iI}(c_{iJH}), \tag{A.1}$$

where each Q_{iI} is a polynomial in c_{iJH} for $|J| + |H| \leq |I|$ with coefficients in \mathbb{N} .

We can fix a non-zero topological nilpotent element π such that $\|c_{iJK}\| \leq |\pi|^{-1}$ for all i, J, H . From the argument above, we deduce inductively that each coefficient d_{iI} is a finite sum of products of the form $\prod c_{k,JH}$ with $\sum |J| \leq |I|$. In particular, each product has at most $|I|$ factors, and hence $\|d_{iI}\| \leq |\pi|^{-|I|}$. We conclude $\|d_{iI} \pi^{2|I} \| \leq |\pi|^{|I|}$, which tends to 0 as $|I| \rightarrow \infty$. \square

The previous statement has an immediate generalization.

COROLLARY A.2. *Let R be a non-archimedean Banach K -algebra, let $\sigma = (\sigma_1, \dots, \sigma_n)$ and $\tau = (\tau_1, \dots, \tau_m)$ be two systems of coordinates, let $\bar{\sigma} = (\bar{\sigma}_1, \dots, \bar{\sigma}_n)$ and $\bar{\tau} = (\bar{\tau}_1, \dots, \bar{\tau}_m)$ two sequences of elements of R and let $P = (P_1, \dots, P_m)$ be a collection of polynomials in $R[\sigma, \tau]$ such that $P(\sigma = \bar{\sigma}, \tau = \bar{\tau}) = 0$ and $\det(\partial P_i / \partial \tau_j)(\sigma = \bar{\sigma}, \tau = \bar{\tau}) \in R^\times$. There exists a unique collection $F = (F_1, \dots, F_m)$ of m formal power series in $R[[\sigma - \bar{\sigma}]]$ such that $F(\sigma = \bar{\sigma}) = \bar{\tau}$ and $P(\sigma, F(\sigma)) = 0$ in $R[[\sigma - \bar{\sigma}]]$ and they have a positive radius of convergence around $\bar{\sigma}$.*

Proof. If we apply Proposition A.1 to the polynomials $P'_i := P(\bar{\sigma} + \eta, \bar{\tau} + \theta)$ we obtain an array of formal power series $F' = (F'_1, \dots, F'_m)$ in $R[[\eta]]$ with positive radius of convergence such that $P'(\eta, F'(\eta)) = 0$. If we now put $\sigma := \bar{\sigma} + \eta$ and $F := \bar{\tau} + F'$ we get $P(\sigma, F(\sigma - \bar{\sigma})) = 0$ in $R[[\sigma - \bar{\sigma}]]$, as desired. \square

We now assume that K is perfectoid and we come back to the category $\widehat{\text{RigSm}}^{\text{gc}}$ that we introduced above (see Definition 2.4). We recall that an object $X = \varprojlim_h X_h$ of this category is the pullback over $\widehat{\mathbb{T}}^N \rightarrow \mathbb{T}^N$ of a map $X_0 \rightarrow \mathbb{T}^N \times \mathbb{T}^M$ that is a composition of rational embeddings and finite étale maps from an affinoid tft adic space X_0 to a torus $\mathbb{T}^N \times \mathbb{T}^M = \text{Spa } K\langle \underline{v}^{\pm 1}, \underline{\nu}^{\pm 1} \rangle$ and X_h denotes the pullback of X_0 by $\mathbb{T}^N \langle \underline{v}^{1/p^h} \rangle \rightarrow \mathbb{T}^N$.

PROPOSITION A.3. *Let $X = \varprojlim_h X_h$ be an object of $\widehat{\text{RigSm}}^{\text{gc}}$. If an element ξ of $\mathcal{O}^+(X)$ is algebraic and separable over each generic point of $\text{Spec } \mathcal{O}(X_0)$ then it lies in $\mathcal{O}^+(X_{\bar{h}})$ for some \bar{h} .*

Proof. Let X_0 be $\text{Spa}(R_0, R_0^\circ)$, let X_h be $\text{Spa}(R_h, R_h^\circ)$ and X be $\text{Spa}(R, R^+)$. For any $h \in \mathbb{N}$ one has $R_h = R_0 \widehat{\otimes}_{K\langle \underline{v}^{\pm 1} \rangle} K\langle \underline{v}^{\pm 1/p^h} \rangle$ and R^+ coincides with the π -adic completion of $\varinjlim_h R_h^\circ$ by Proposition 2.1. The proof is divided into several steps.

Step 1. We can suppose that R is perfectoid. Indeed, we can consider the refined tower $X'_h = X_0 \times_{\mathbb{T}^N \times \mathbb{T}^M} (\mathbb{T}^N \langle \underline{v}^{1/p^h} \rangle \times \mathbb{T}^M \langle \underline{\nu}^{1/p^h} \rangle)$ whose limit \widehat{X} is perfectoid. If the claim is true for this tower, we conclude that ξ lies in the intersection of $\mathcal{O}(X'_h)$ and $\mathcal{O}(X)$ inside $\mathcal{O}(\widehat{X})$ for some h . By Remark 1.11 this is the intersection

$$\left(\widehat{\bigoplus}_{I \in (\mathbb{Z}[1/p] \cap [0,1])^N} R_0 \underline{v}^I \right) \cap \left(\widehat{\bigoplus}_{\substack{I \in \{a/p^h : 0 \leq a < p^h\}^N \\ J \in \{a/p^h : 0 \leq a < p^h\}^M}} R_0 \underline{v}^I \underline{\nu}^J \right),$$

which coincides with

$$\widehat{\bigoplus}_{I \in \{a/p^h : 0 \leq a < p^h\}^N} R_0 \underline{v}^I = R_h.$$

Step 2. We can always assume that each R_h is an integral domain. Indeed, the number of connected components of $\text{Spa } R_h$ may rise, but it is bounded by the number of connected components of the affinoid perfectoid X , which is finite by Remark 2.12.

We deduce that the number of connected components of $\text{Spa } R_h$ stabilizes for h large enough. Up to shifting indices, we can then suppose that $\text{Spa } R_0$ is the finite disjoint union of irreducible rigid varieties $\text{Spa } R_{i0}$ for $i = 1, \dots, k$ such that $R_{ih} = R_{i0} \widehat{\otimes}_{K\langle \underline{v}^{\pm 1} \rangle} K\langle \underline{v}^{\pm 1/p^h} \rangle$ is a domain for all h . We denote by R_i the ring $R_{i0} \widehat{\otimes}_{K\langle \underline{v}^{\pm 1} \rangle} K\langle \underline{v}^{\pm 1/p^\infty} \rangle$. Now let $\xi = (\xi_i)$ be an element in $R^+ = \prod R_i^+$ that is separable over $\prod \text{Frac } R_i$, i.e., each ξ_i is separable over $\text{Frac } R_i$. If the proposition holds for R_i we then conclude that ξ_i lies in R_{ih}° for some large enough h so that $\xi \in R_h^\circ$ as claimed.

Step 3. We prove that we can consider a non-empty rational subspace $U_0 = \text{Spa } R_0 \langle f_i/g \rangle$ of X_0 instead. Indeed, using Remark 1.11 if the result holds for U_0 assuming $\bar{h} = 0$ we deduce that ξ lies in the intersection of $R \cong \bigoplus R_0$ and of $R_0 \langle f_i/g \rangle$ inside $R \langle f_i/g \rangle \cong \bigoplus R_0 \langle f_i/g \rangle$, which coincides with R_0 .

Step 4. We prove that we can assume ξ to be integral over R_0 . Indeed, let P_ξ be its minimal polynomial over $\text{Frac}(R_0)$. We can suppose there is a common denominator d such that P_ξ has coefficients in $R_0[1/d][x]$. By [BGR84, Proposition 6.2.1/4(ii)] we can also assume that $|d| = 1$. In particular, by [BGR84, Proposition 7.2.6/3], the rational subspace associated to $R_0 \langle 1/d \rangle$ is not empty. By step 3, we can then restrict to it and assume ξ to be integral over R_0 and $R_0[\xi] \cong R_0[x]/P_\xi(x)$.

Step 5. We can suppose that $P_\xi(x)$ is the minimal polynomial of ξ with respect to all non-empty rational subspaces of X_h for all h . If it is not the case, from the previous steps we can rescale indices and restrict to a rational subspace with respect to which the degree of $P_\xi(x)$ is lower. Since the degree is bounded from below, we conclude the claim.

Step 6. We now conclude the claim. By steps 3 and 4 we may suppose that the map $R_0 \rightarrow R_0[\xi]$ is finite étale. Since this map splits after tensoring with R , we can deduce by [Sch12, Lemma 7.5] that it splits already at some level h . By step 5, this implies that $P_\xi(x)$ is of degree 1, and hence $\xi \in R_0$. □

We introduce now the geometric application of Propositions A.1 and A.3. It states that a map from $\varprojlim_h X_h \in \widehat{\text{RigSm}}$ to a rigid variety factors, up to \mathbb{B}^1 -homotopy, over one of the intermediate varieties X_h . Analogous statements are widely used in [Ayo15] (see for example [Ayo15, Theorem 2.2.58]): there, these are obtained as corollaries of Popescu’s theorem [Pop85, Pop86], which is not available in our non-noetherian setting.

PROPOSITION A.4. *Let $X = \varprojlim_h X_h$ be in $\widehat{\text{RigSm}}^{\text{gc}}$. Let Y be an affinoid rigid variety endowed with an étale map $Y \rightarrow \mathbb{B}^n$ and let $f : X \rightarrow Y$ be a map of adic spaces.*

- (i) *There exist m polynomials Q_1, \dots, Q_m in $K[\sigma_1, \dots, \sigma_n, \tau_1, \dots, \tau_m]$ such that $Y \cong \text{Spa } A$ with $A \cong K \langle \sigma, \tau \rangle / (Q)$ and $\det(\partial Q_i / \partial \tau_j) \in A^\times$.*
- (ii) *There exists a map $H : X \times \mathbb{B}^1 \rightarrow Y$ such that $H \circ i_0 = f$ and $H \circ i_1$ factors over the canonical map $X \rightarrow X_h$ for some integer h .*

Moreover, if f is induced by the map $K \langle \sigma, \tau \rangle \rightarrow \mathcal{O}(X)$, $\sigma \mapsto s, \tau \mapsto t$ the map H can be defined via

$$(\sigma, \tau) \mapsto (s + (\tilde{s} - s)\chi, F(s + (\tilde{s} - s)\chi)),$$

where F is the unique array of formal power series in $\mathcal{O}(X)[[\sigma - s]]$ associated to the polynomials $P(\sigma, \tau)$ by Corollary A.2, and \tilde{s} is any element in $\varinjlim_h \mathcal{O}^+(X_h)$ such that the radius of convergence of F is larger than $\|\tilde{s} - s\|$ and $F(\tilde{s})$ lies in $\mathcal{O}^+(X)$.

Proof. The first claim follows from [Ayo15, Lemma 1.1.52]. We turn to the second claim. Let X_0 be $\text{Spa}(R_0, R_0^\circ)$ and X be $\text{Spa}(R, R^+)$. For any $h \in \mathbb{N}$ we denote $R_0 \widehat{\otimes}_{K(\underline{v})} K \langle \underline{v}^{\pm 1/p^h} \rangle$ with R_h so that R^+ coincides with the π -adic completion of $\varinjlim_h R_h^\circ$ by Proposition 2.1.

The map f is determined by the choice of n elements $s = (s_1, \dots, s_n)$ and m elements $t = (t_1, \dots, t_m)$ of R^+ such that $P(s, t) = 0$. We prove that the formula for H provided in the statement defines a map H with the required properties.

By Corollary A.2 there exists a collection $F = (F_1, \dots, F_m)$ of m formal power series in $R[[\sigma - s]]$ with a positive radius of convergence such that $F(s) = t$ and $P(\sigma, F(\sigma)) = 0$. As $\varinjlim_h R_h^\circ$ is dense in R^+ we can find an integer \bar{h} and elements $\tilde{s}_i \in R_{\bar{h}}^\circ$ such that $\|\tilde{s} - s\|$ is smaller than the convergence radius of F . By renaming the indices, we can assume that $\bar{h} = 0$. As F is continuous and R^+ is open, we can also assume that the elements $F_j(\tilde{s})$ lie in R^+ . We are left to prove that they actually lie in $\varinjlim_h R_h^\circ$. Since the determinant of $(\partial P_i / \partial \tau_j)(\tilde{s}, F(\tilde{s}))$ is invertible, the field $L := \text{Frac}(R_0)(F_1(\tilde{s}), \dots, F_m(\tilde{s}))$ is algebraic and separable over $\text{Frac}(R_0)$. We can then apply Proposition A.3 to conclude that each element $F_j(\tilde{s})$ lies in R_h° for a sufficiently big integer h . \square

The goal of the rest of this section is to prove Proposition 4.1. To this aim, we present a generalization of the results above for collections of maps. As before, we start with an algebraic statement and then translate it into a geometrical fact for our specific purposes.

PROPOSITION A.5. *Let R be a Banach K -algebra and let $\{R_h\}_{h \in \mathbb{N}}$ be a collection of nested complete subrings of R such that $\varinjlim R_h$ is dense in R . Let s_1, \dots, s_N be elements of $R\langle\theta_1, \dots, \theta_n\rangle$. For any $\varepsilon > 0$ there exists an integer h and elements $\tilde{s}_1, \dots, \tilde{s}_N$ of $R_h\langle\theta_1, \dots, \theta_n\rangle$ satisfying the following conditions.*

- (i) $|s_\alpha - \tilde{s}_\alpha| < \varepsilon$ for each α .
- (ii) For any $\alpha, \beta \in \{1, \dots, N\}$ and any $k \in \{1, \dots, n\}$ such that $s_\alpha|_{\theta_k=0} = s_\beta|_{\theta_k=0}$ we also have $\tilde{s}_\alpha|_{\theta_k=0} = \tilde{s}_\beta|_{\theta_k=0}$.
- (iii) For any $\alpha, \beta \in \{1, \dots, N\}$ and any $k \in \{1, \dots, n\}$ such that $s_\alpha|_{\theta_k=1} = s_\beta|_{\theta_k=1}$ we also have $\tilde{s}_\alpha|_{\theta_k=1} = \tilde{s}_\beta|_{\theta_k=1}$.
- (iv) For any $\alpha \in \{1, \dots, N\}$ if $s_\alpha|_{\theta_1=1} \in R_{h'}\langle\theta\rangle$ for some h' then $\tilde{s}_\alpha|_{\theta_1=1} = s_\alpha|_{\theta_1=1}$.

Proof. We will actually prove a stronger statement, namely that we can reinforce the previous conditions with the following.

- (v) For any $\alpha, \beta \in \{1, \dots, N\}$ any subset T of $\{1, \dots, n\}$ and any map $\sigma : T \rightarrow \{0, 1\}$ such that $s_\alpha|_\sigma = s_\beta|_\sigma$ then $\tilde{s}_\alpha|_\sigma = \tilde{s}_\beta|_\sigma$.
- (vi) For any $\alpha \in \{1, \dots, N\}$ any subset T of $\{1, \dots, n\}$ containing 1 and any map $\sigma : T \rightarrow \{0, 1\}$ such that $s_\alpha|_\sigma \in R_h\langle\theta\rangle$ for some h then $\tilde{s}_\alpha|_\sigma = s_\alpha|_\sigma$.

Above we denote by $s|_\sigma$ the image of s via the substitution $(\theta_t = \sigma(t))_{t \in T}$. We proceed by induction on N , the case $N = 0$ being trivial.

Consider the conditions we want to preserve that involve the index N . They are of the form

$$s_i|_\sigma = s_N|_\sigma$$

and are indexed by some pairs (σ, i) , where i is an index and σ varies in a set of maps Σ . Our procedure consists in determining by induction the elements $\tilde{s}_1, \dots, \tilde{s}_{N-1}$ first, and then deducing the existence of \tilde{s}_N by means of Lemma A.8 by lifting the elements $\{\tilde{s}_i|_\sigma\}_{(\sigma, i)}$. Therefore, we first define $\varepsilon' := (1/C)\varepsilon$ where $C = C(\Sigma)$ is the constant introduced in Lemma A.8 and then apply the induction hypothesis to the first $N - 1$ elements with respect to ε' .

By the induction hypothesis, the elements $\tilde{s}_i|_\sigma$ satisfy the compatibility condition of Lemma A.8 and lie in $R_h\langle\theta\rangle$ for some integer h . Without loss of generality, we assume $h = 0$. By Lemma A.8 we can find an element \tilde{s}_N of $R_h\langle\theta\rangle$ lifting them such that $|\tilde{s}_N - s_N| < C\varepsilon' = \varepsilon$, as desired. \square

The following lemmas are used in the proof of the previous proposition.

LEMMA A.6. For any normed ring R and any map $\sigma : T_\sigma \rightarrow \{0, 1\}$ defined on a subset T_σ of $\{1, \dots, n\}$ we denote by I_σ the ideal of $R\langle \underline{\theta} \rangle$ generated by $\theta_i - \sigma(i)$ as i varies in T_σ . For any finite set Σ of such maps and any such map η one has $(\bigcap_{\sigma \in \Sigma} I_\sigma) + I_\eta = \bigcap_{\sigma \in \Sigma} (I_\sigma + I_\eta)$.

Proof. We only need to prove the inclusion $\bigcap (I_\sigma + I_\eta) \subseteq (\bigcap I_\sigma) + I_\eta$. We can make induction on the cardinality of T_η and restrict to the case in which T_η is a singleton. By changing variables, we can suppose $T_\eta = \{1\}$ and $\eta(1) = 0$ so that $I_\eta = (\theta_1)$.

We first suppose that $1 \notin T_\sigma$ for all $\sigma \in \Sigma$. Let s be an element of $\bigcap (I_\sigma + (\theta_1))$. This means we can find elements $s_\sigma \in I_\sigma$ and polynomials $p_\sigma \in R\langle \underline{\theta} \rangle$ such that $s = s_\sigma + p_\sigma \theta_1$. Since I_σ is generated by polynomials of the form $\theta_i - \epsilon$ with $i \neq 1$ we can suppose that s_σ contains no θ_1 by eventually changing p_σ . Now let σ, σ' be in Σ . From the equality

$$s_\sigma = (s_\sigma + p_\sigma \theta_1)|_{\theta_1=0} = (s_{\sigma'} + p_{\sigma'} \theta_1)|_{\theta_1=0} = s_{\sigma'}$$

we conclude that $s_\sigma \in \bigcap I_\sigma$. Therefore, $s \in \bigcap I_\sigma + (\theta_1)$, as claimed.

We now move to the general case. Suppose $\bar{\sigma}(1) = 1$ for some $\bar{\sigma} \in \Sigma$. Then $I_{\bar{\sigma}} + I_\eta = R\langle \underline{\theta} \rangle$ and if $f \in \bigcap_{\sigma \neq \bar{\sigma}} I_\sigma$ then $f = -f(\theta_1 - 1) + f\theta_1 \in \bigcap_{\sigma} I_\sigma + (\theta_1)$. Therefore, the contribution of $I_{\bar{\sigma}}$ is trivial on both sides and we can erase it from Σ . We can, therefore, suppose that $\sigma(1) = 0$ whenever $1 \in T_\sigma$.

For any $\sigma \in \Sigma$ let σ' be its restriction to $T_\sigma \setminus \{1\}$. We have $I_{\sigma'} \subseteq I_\sigma$ and $I_{\sigma'} + (\theta_1) = I_\sigma + (\theta_1)$ for all $\sigma \in \Sigma$. By what we already proved, the statement holds for the set $\Sigma' := \{\sigma' : \sigma \in \Sigma\}$. Therefore,

$$\bigcap_{\sigma \in \Sigma} (I_\sigma + (\theta_1)) = \bigcap_{\sigma' \in \Sigma'} (I_{\sigma'} + (\theta_1)) = \bigcap_{\sigma' \in \Sigma'} I_{\sigma'} + (\theta_1) \subseteq \bigcap_{\sigma \in \Sigma} I_\sigma + (\theta_1),$$

proving the claim. □

We recall (see [BGR84, Definition 1.1.9/1]) that a morphism of normed groups $\phi : G \rightarrow H$ is *strict* if the homomorphism $G/\ker \phi \rightarrow \phi(G)$ is a homeomorphism, where the former group is endowed with the quotient topology and the latter with the topology inherited from H . In particular, we say that a sequence of normed K -vector spaces

$$R \xrightarrow{f} S \xrightarrow{g} T$$

is *strict and exact* at S if it is exact at S and if f is strict, i.e., the quotient norm and the norm induced by S on $R/\ker(f) \cong \ker(g)$ are equivalent.

LEMMA A.7. For any map $\sigma : T_\sigma \rightarrow \{0, 1\}$ defined on a subset T_σ of $\{1, \dots, n\}$ we denote by I_σ the ideal of $R\langle \underline{\theta} \rangle = R\langle \theta_1, \dots, \theta_n \rangle$ generated by $\theta_i - \sigma(i)$ as i varies in T_σ . For any finite set Σ of such maps and any complete normed K -algebra R the following sequence of Banach K -algebras is strict and exact

$$0 \rightarrow R\langle \underline{\theta} \rangle \Big/ \bigcap_{\sigma \in \Sigma} I_\sigma \rightarrow \prod_{\sigma \in \Sigma} R\langle \underline{\theta} \rangle \Big/ I_\sigma \rightarrow \prod_{\sigma, \sigma' \in \Sigma} R\langle \underline{\theta} \rangle \Big/ (I_\sigma + I_{\sigma'})$$

and the ideal $\bigcap_{\sigma \in \Sigma} I_\sigma$ is generated by a finite set of polynomials in the θ_i with coefficients in \mathbb{Z} .

Proof. We follow the notation and the proof of [Kle89]. For a collection of ideals $\mathcal{I} = \{I_\sigma\}$ we let $A(\mathcal{I})$ be the kernel of the map $\prod_\sigma R(\underline{\theta})/I_\sigma \rightarrow \prod_{\sigma,\sigma'} R(\underline{\theta})/(I_\sigma + I_{\sigma'})$ and $O(\mathcal{I})$ be the cokernel of $R(\underline{\theta})/\bigcap_\sigma I_\sigma \rightarrow A(\mathcal{I})$. We make induction on the cardinality m of \mathcal{I} . The case $m = 1$ is obvious.

Let \mathcal{I}' be $\mathcal{I} \cup \{I_\eta\}$. From the diagram

$$\begin{array}{ccccccc} & & 0 & \longrightarrow & R(\underline{\theta}) & \xrightarrow{\text{id}} & R(\underline{\theta}) & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & W & \longrightarrow & A(\mathcal{I}') & \longrightarrow & A(\mathcal{I}) & & \end{array}$$

we obtain by the snake lemma the exact sequence

$$0 \rightarrow I_\eta \cap \bigcap I_\sigma \rightarrow \bigcap I_\sigma \rightarrow W \rightarrow O(\mathcal{I}') \rightarrow O(\mathcal{I}).$$

By direct computation, it holds that $W = \bigcap(I_\sigma + I_\eta)/I_\eta$. By the induction hypothesis, we obtain $O(\mathcal{I}) = 0$. Moreover, since $\bigcap I_\sigma + I_\eta = \bigcap(I_\sigma + I_\eta)$ by Lemma A.6, we conclude that the map $\bigcap I_\sigma \rightarrow W$ is surjective, and hence $O(\mathcal{I}') = 0$ proving the main claim.

We remark that the sequence of the statement is obtained from the sequence with $R = K$, by tensoring with R over K and completing. The latter is a (strict) exact sequence of affinoid algebras in the sense of Tate [BGR84, Definition 6.1.1/1] so we deduce that the sequence is strict for any R by means of [BGR84, Proposition 2.1.8/6]. We also remark that the sequence can be deduced from the analogous sequence for $\mathbb{Z}[\underline{\theta}]$ in place of $R(\underline{\theta})$, by tensoring with R and completing. In particular, the ideal $\bigcap_{\sigma \in \Sigma} I_\sigma$ is already defined over $\mathbb{Z}[\underline{\theta}]$, as claimed. \square

Let σ and σ' be maps defined from two subsets T_σ , respectively, $T_{\sigma'}$ of $\{1, \dots, n\}$ to $\{0, 1\}$. We say that they are *compatible* if $\sigma(i) = \sigma'(i)$ for all $i \in T_\sigma \cap T_{\sigma'}$ and in this case we denote by (σ, σ') the map from $T_\sigma \cup T_{\sigma'}$ extending them.

LEMMA A.8. *Let $X = \varprojlim_h X_h$ be an object in $\widehat{\text{RigSm}}$ and Σ a set as in Lemma A.7. We denote $\mathcal{O}(X)$ by R and $\mathcal{O}(X_h)$ by R_h . For any $\sigma \in \Sigma$ let \bar{f}_σ be an element of $R(\underline{\theta})/I_\sigma$ such that $\bar{f}_\sigma|_{(\sigma,\sigma')} = \bar{f}_{\sigma'}|_{(\sigma,\sigma')}$ for any couple $\sigma, \sigma' \in \Sigma$ of compatible maps.*

(i) *There exists an element $f \in R(\underline{\theta})$ such that $f|_\sigma = \bar{f}_\sigma$.*

(ii) *There exists a constant $C = C(\Sigma)$ such that if for some $g \in R(\underline{\theta})$ one has $|\bar{f}_\sigma - g|_\sigma| < \varepsilon$ for all σ then the element f can be chosen so that $|f - g| < C\varepsilon$. Moreover, if $\bar{f}_\sigma \in R_0(\underline{\theta})/I_\sigma$ for all σ then the element f can be chosen inside $R_h(\underline{\theta})$ for some integer h .*

Proof. The first claim and the first part of the second are simply a restatement of Lemma A.7, where $C = C(\Sigma)$ is the constant defining the compatibility $\|\cdot\|_1 \leq C\|\cdot\|_2$ between the norm $\|\cdot\|_1$ on $R(\underline{\theta})/\bigcap I_\sigma$ induced by the quotient and the norm $\|\cdot\|_2$ induced by the embedding in $\prod R(\underline{\theta})/I_\sigma$. We now turn to the last sentence of the second claim.

We apply Lemma A.7 to each R_h and to R . We then obtain exact sequences of Banach spaces:

$$\begin{aligned} 0 \rightarrow R_h(\underline{\theta}) \Big/ \bigcap_{\sigma \in \Sigma} I_\sigma &\rightarrow \prod_{\sigma \in \Sigma} R_h(\underline{\theta}) \Big/ I_\sigma \rightarrow \prod_{\sigma, \sigma' \in \Sigma} R_h(\underline{\theta}) \Big/ (I_\sigma + I_{\sigma'}) \\ 0 \rightarrow R(\underline{\theta}) \Big/ \bigcap_{\sigma \in \Sigma} I_\sigma &\rightarrow \prod_{\sigma \in \Sigma} R(\underline{\theta}) \Big/ I_\sigma \rightarrow \prod_{\sigma, \sigma' \in \Sigma} R(\underline{\theta}) \Big/ (I_\sigma + I_{\sigma'}), \end{aligned}$$

where all ideals that appear are finitely generated by polynomials with \mathbb{Z} -coefficients, depending only on Σ .

In particular, there exist two lifts of $\{\bar{f}_\sigma\}$: an element f_1 of $R_0\langle\theta\rangle$ and an element f_2 of $R\langle\theta\rangle$ such that $|f_2 - g| < C\varepsilon$ and their difference lies in $\bigcap I_\sigma$. Hence, we can find coefficients $\gamma_i \in R\langle\theta\rangle$ such that $f_1 = f_2 + \sum_i \gamma_i p_i$, where $\{p_1, \dots, p_M\}$ are generators of $\bigcap I_\sigma$ which have coefficients in K . Now let $\tilde{\gamma}_i$ be elements of $R_h\langle\theta\rangle$ with $|\tilde{\gamma}_i - \gamma_i| < C\varepsilon/M|p_i|$. The element $f_3 := f_1 - \sum_i \tilde{\gamma}_i p_i$ lies in $\varinjlim_h (R_h\langle\theta\rangle)$ is another lift of $\{\bar{f}_\sigma\}$ and satisfies $|f_3 - g| \leq \max\{|f_2 - g|, |f_2 - f_3|\} < C\varepsilon$, proving the claim. \square

We can now finally prove the approximation result that played a crucial role in § 4.

Proof of Proposition 4.1. For any $h \in \mathbb{Z}$ we will denote $\mathcal{O}(X_h)\langle\theta_1, \dots, \theta_n\rangle$ by R_h . We also denote the π -adic completion of $\varinjlim_h R_h^\circ$ by R^+ and $R^+[\pi^{-1}]$ by R .

By Proposition A.4 we conclude that there exist integers m and n and a m -tuple of polynomials $P = (P_1, \dots, P_m)$ in $K\langle\sigma, \tau\rangle$, where $\sigma = (\sigma_1, \dots, \sigma_n)$ and $\tau = (\tau_1, \dots, \tau_m)$ are systems of variables such that $K\langle\sigma, \tau\rangle/(P) \cong \mathcal{O}(Y)$ and each f_k is induced by maps $(\sigma, \tau) \mapsto (s_k, t_k)$ from $K\langle\sigma, \tau\rangle/(P)$ to R for some m -tuples s_k and n -tuples t_k in R . Moreover, there exists a sequence of power series $F_k = (F_{k1}, \dots, F_{km})$ associated to each f_k such that

$$(\sigma, \tau) \mapsto (s_k + (\tilde{s}_k - s_k)\chi, F_k(s_k + (\tilde{s}_k - s_k)\chi)) \in R\langle\chi\rangle \cong \mathcal{O}(X \times \mathbb{B}^n \times \mathbb{B}^1)$$

defines a map H_k satisfying the first claim, for any choice of $\tilde{s}_k \in \varinjlim_h R_h^\circ$ such that \tilde{s}_k is in the convergence radius of F_k and $F_k(\tilde{s}_k)$ is in R^+ .

Now let ε be a positive real number, smaller than all radii of convergence of the series F_{kj} and such that $F(a) \in R^+$ for all $|a - s| < \varepsilon$. Denote by \tilde{s}_{ki} the elements associated to s_{ki} by applying Proposition A.5 with respect to the chosen ε . In particular, they induce a well-defined map H_k and the elements \tilde{s}_{ki} lie in $R_{\bar{h}}^\circ\langle\theta_1, \dots, \theta_n\rangle$ for some integer \bar{h} . We show that the maps H_k induced by this choice also satisfy the second and third claims of the proposition.

Suppose that $f_k \circ d_{r,\varepsilon} = f_{k'} \circ d_{r,\varepsilon}$ for some $r \in \{1, \dots, n\}$ and $\varepsilon \in \{0, 1\}$. This means that $\bar{s} := s_k|_{\theta_r=\varepsilon} = s_{k'}|_{\theta_r=\varepsilon}$ and $\bar{t} := t_k|_{\theta_r=\varepsilon} = t_{k'}|_{\theta_r=\varepsilon}$. This implies that both $F_k|_{\theta_r=\varepsilon}$ and $F_{k'}|_{\theta_r=\varepsilon}$ are two m -tuples of formal power series \bar{F} with coefficients in $\mathcal{O}(X \times \mathbb{B}^{n-1})$ converging around \bar{s} and such that $P(\sigma, \bar{F}(\sigma)) = 0$, $\bar{F}(\bar{s}) = \bar{t}$. By the uniqueness of such power series stated in Corollary A.2, we conclude that they coincide.

Moreover, by our choice of the elements \tilde{s}_k it follows that $\tilde{\bar{s}} := \tilde{s}_k|_{\theta_r=\varepsilon} = \tilde{s}_{k'}|_{\theta_r=\varepsilon}$. In particular, one has

$$F_k((\tilde{s}_k - s_k)\chi)|_{\theta_r=\varepsilon} = \bar{F}((\tilde{\bar{s}} - \bar{s})\chi) = F_{k'}((\tilde{s}_{k'} - s_{k'})\chi)|_{\theta_r=\varepsilon}$$

and therefore $H_k \circ d_{r,\varepsilon} = H_{k'} \circ d_{r,\varepsilon}$ proving the second claim.

The third claim follows immediately since the elements \tilde{s}_{ki} satisfy the condition (iv) of Proposition A.5. \square

REFERENCES

And06 F. Andreatta, *Generalized ring of norms and generalized (ϕ, Γ) -modules*, Ann. Sci. Éc. Norm. Supér. (4) **39** (2006), 599–647.
 Ayo07 J. Ayoub, *Les six opérations de Grothendieck et le formalisme des cycles évanescents dans le monde motivique, II*, Astérisque **315** (2007).

- Ayo14a J. Ayoub, *La réalisation étale et les opérations de Grothendieck*, Ann. Sci. Éc. Norm. Supér. (4) **47** (2014), 1–145.
- Ayo14b J. Ayoub, *L’algèbre de Hopf et le groupe de Galois motiviques d’un corps de caractéristique nulle, I*, J. Reine Angew. Math. **693** (2014), 1–149.
- Ayo14c J. Ayoub, *L’algèbre de Hopf et le groupe de Galois motiviques d’un corps de caractéristique nulle, II*, J. Reine Angew. Math. **693** (2014), 151–226.
- Ayo15 J. Ayoub, *Motifs des variétés analytiques rigides*, Mém. Soc. Math. Fr. (N.S.) **140–141** (2015).
- BV08 A. Beilinson and V. Vologodsky, *A DG guide to Voevodsky’s motives*, Geom. Funct. Anal. **17** (2008), 1709–1787.
- BGR84 S. Bosch, U. Güntzer and R. Remmert, *Non-Archimedean analysis, A systematic approach to rigid analytic geometry*, Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 261 (Springer, Berlin, 1984).
- BH81 R. Brown and P. J. Higgins, *On the algebra of cubes*, J. Pure Appl. Algebra **21** (1981), 233–260.
- BHS11 R. Brown, P. J. Higgins and R. Sivera, *Nonabelian algebraic topology*, EMS Tracts in Mathematics, vol. 15 (European Mathematical Society, Zürich, 2011); filtered spaces, crossed complexes, cubical homotopy groupoids, with contributions by Christopher D. Wensley and Sergei V. Soloviev.
- BV18 K. Buzzard and A. Verberkmoes, *Stably uniform affinoids are sheafy*, J. Reine Angew. Math. **740** (2018), 25–39.
- DHI04 D. Dugger, S. Hollander and D. C. Isaksen, *Hypercovers and simplicial presheaves*, Math. Proc. Cambridge Philos. Soc. **136** (2004), 9–51.
- EGAIV A. Grothendieck, *Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas IV*, Publ. Math. Inst. Hautes Études Sci. **32** (1967).
- Fau06 H. Fausk, *T-model structures on chain complexes of presheaves*, Preprint (2006), [arXiv:math/0612414](https://arxiv.org/abs/math/0612414) [math.AG].
- FJ13 J.-M. Fontaine and U. Jannsen, *Frobenius gauges and a new theory of p-torsion sheaves in characteristic p. I*, Preprint (2013), [arXiv:1304.3740](https://arxiv.org/abs/1304.3740) [math.AG].
- FW79 J.-M. Fontaine and J.-P. Wintenberger, *Extensions algébriques et corps des normes des extensions APF des corps locaux*, C. R. Acad. Sci. Paris Sér. A-B **288** (1979), A441–A444.
- FvdP04 J. Fresnel and M. van der Put, *Rigid analytic geometry and its applications*, Progress in Mathematics, vol. 218 (Birkhäuser, Boston, MA, 2004).
- GR03 O. Gabber and L. Ramero, *Almost ring theory*, Lecture Notes in Mathematics, vol. 1800 (Springer, Berlin, 2003).
- Hir03 P. S. Hirschhorn, *Model categories and their localizations*, Mathematical Surveys and Monographs, vol. 99 (American Mathematical Society, Providence, RI, 2003).
- Hov99 M. Hovey, *Model categories*, Mathematical Surveys and Monographs, vol. 63 (American Mathematical Society, Providence, RI, 1999).
- Hov01 M. Hovey, *Spectra and symmetric spectra in general model categories*, J. Pure Appl. Algebra **165** (2001), 63–127.
- Hub93 R. Huber, *Continuous valuations*, Math. Z. **212** (1993), 455–477.
- Hub94 R. Huber, *A generalization of formal schemes and rigid analytic varieties*, Math. Z. **217** (1994), 513–551.
- Hub96 R. Huber, *Étale cohomology of rigid analytic varieties and adic spaces*, Aspects of Mathematics, vol. E30 (Friedr. Vieweg & Sohn, Braunschweig, 1996).
- Igu00 J.-I. Igusa, *An introduction to the theory of local zeta functions*, AMS/IP Studies in Advanced Mathematics, vol. 14 (American Mathematical Society, Providence, RI, 2000).
- Jar87 J. F. Jardine, *Simplicial presheaves*, J. Pure Appl. Algebra **47** (1987), 35–87.

- Jar02 J. F. Jardine, *Cubical homotopy theory: a beginning*, Preprints of the Newton Institute, NI02030-NST, 2002.
- Kah13 B. Kahn, *The full faithfulness conjectures in characteristic p* , in *Autour des motifs (II)- École d'été franco-asiatique de géométrie algébrique et de théorie des nombres. Asian-French summer school on algebraic geometry and number theory*, Panor. Synthèses, vol. 41 (Soc. Math. France, Paris, 2013).
- KL16 K. S. Kedlaya and R. Liu, *Relative p -adic Hodge theory, II: imperfect period rings*, Preprint (2016), [arXiv:1602.06899](https://arxiv.org/abs/1602.06899) [math.NT].
- Kle89 E. Kleinert, *Some remarks on the Chinese remainder theorem*, Arch. Math. (Basel) **52** (1989), 433–439.
- Mal09 G. Maltsiniotis, *La catégorie cubique avec connexions est une catégorie test stricte*, Homology, Homotopy Appl. **11** (2009), 309–326.
- May67 J. P. May, *Simplicial objects in algebraic topology*, Van Nostrand Mathematical Studies, vol. 11 (D. Van Nostrand, Princeton, NJ–Toronto, ON–London, 1967).
- MVW06 C. Mazza, V. Voevodsky and C. Weibel, *Lecture notes on motivic cohomology*, Clay Mathematics Monographs, vol. 2 (American Mathematical Society, Providence, RI, 2006).
- MV99 F. Morel and V. Voevodsky, *\mathbf{A}^1 -homotopy theory of schemes*, Publ. Math. Inst. Hautes Études Sci. **90** (1999), 45–143.
- Pop85 D. Popescu, *General Néron desingularization*, Nagoya Math. J. **100** (1985), 97–126.
- Pop86 D. Popescu, *General Néron desingularization and approximation*, Nagoya Math. J. **104** (1986), 85–115.
- Rio05 J. Riou, *Dualité de Spanier-Whitehead en géométrie algébrique*, C. R. Math. Acad. Sci. Paris **340** (2005), 431–436.
- Rio07 J. Riou, *Catégorie homotopique stable d'un site suspendu avec intervalle*, Bull. Soc. Math. France **135** (2007), 495–547.
- Sch12 P. Scholze, *Perfectoid spaces*, Publ. Math. Inst. Hautes Études Sci. **116** (2012), 245–313.
- Sch13 P. Scholze, *p -adic Hodge theory for rigid-analytic varieties*, Forum Math. Pi **1** (2013), e1, 77.
- Sch14 P. Scholze, *Perfectoid spaces and their applications*, in *Proceedings of the International Congress of Mathematicians—Seoul 2014, Vol. II* (Kyung Moon Sa, Seoul, 2014), 461–486.
- SGA4 M. Artin, A. Grothendieck and J.-L. Verdier (eds), *Théorie des topos et cohomologie étale des schémas*, in *Séminaire de Géométrie Algébrique du Bois-Marie 1963–1964 (SGA 4) – Tome 2*, Lecture Notes in Mathematics, vol. 270 (Springer, Berlin, 1972).
- SS00 S. Schwede and B. E. Shipley, *Algebras and modules in monoidal model categories*, Proc. Lond. Math. Soc. (3) **80** (2000), 491–511.
- SS03 S. Schwede and B. Shipley, *Equivalences of monoidal model categories*, Algebr. Geom. Topol. **3** (2003), 287–334 (electronic).
- Sta18 The Stacks Project Authors, *Stacks Project*, <http://stacks.math.columbia.edu>, 2018.
- SW13 P. Scholze and J. Weinstein, *Moduli of p -divisible groups*, Camb. J. Math. **1** (2013), 145–237.
- SV00 A. Suslin and V. Voevodsky, *Bloch-Kato conjecture and motivic cohomology with finite coefficients*, in *The arithmetic and geometry of algebraic cycles (Banff, AB, 1998)*, NATO Sci. Ser. C Math. Phys. Sci., vol. 548 (Kluwer Academic Publishers, Dordrecht, 2000), 117–189.
- Ton92 A. P. Tonks, *Cubical groups which are Kan*, J. Pure Appl. Algebra **81** (1992), 83–87.
- Vezi7 A. Vezzani, *Effective motives with and without transfers in characteristic p* , Adv. Math. **306** (2017), 852–879.
- Vezi8 A. Vezzani, *The Monsky–Washnitzer and the overconvergent realizations*, Int. Math. Res. Not. IMRN **2018** (2018), 3443–3489.
- Vezi9 A. Vezzani, *Rigid cohomology via the tilting equivalence*, J. Pure Appl. Algebra **223** (2019), 818–843.

- Voe96 V. Voevodsky, *Homology of schemes*, *Selecta Math. (N.S.)* **2** (1996), 111–153.
- Voe10 V. Voevodsky, *Cancellation theorem*, *Doc. Math.* (2010), 671–685 (Extra vol.: Andrei A. Suslin sixtieth birthday).
- Wed12 T. Wedhorn, *Adic spaces*. <https://www2.math.uni-paderborn.de/fileadmin/Mathematik/People/wedhorn/Lehre/AdicSpaces.pdf>, 2012.

Alberto Vezzani vezzani@math.univ-paris13.fr
LAGA – Université Paris 13, Sorbonne Paris Cité,
99 av. Jean-Baptiste Clément, 93430 Villetaneuse, France