

ON THE GORENSTEINNESS OF REES ALGEBRAS OVER LOCAL RINGS

SHIN IKEDA

Introduction

Let (A, m, k) be a Noetherian local ring and I an ideal of A . We set $R(I) = \bigoplus_{n \geq 0} I^n$ and call this graded A -algebra the Rees algebra of I . The importance of the Rees algebra $R(I)$ is in the fact that $\text{Proj } R(I)$ is the blowing up of $\text{Spec } A$ with center in $V(I)$. The Cohen-Macaulayness of Rees algebras was studied by many mathematicians. In [GS] S. Goto and Y. Shimoda gave a criterion for $R(m)$ to be Cohen-Macaulay under the assumption that A is Cohen-Macaulay. Their results have been generalized to $R(I)$ in [HI].

Let $\text{grade}(I) \geq 2$. The purpose of this paper is to characterize the Gorensteinness of $R(I)$ in terms of canonical modules of A and the associated graded ring $G(I) = \bigoplus_{n \geq 0} I^n/I^{n+1}$. The notion of canonical modules of local rings plays an important role in the homological theory of local rings, cf. [HK]. The canonical modules of graded rings defined over a field were introduced and studied extensively in [GW]. In Section 1 we introduce the notion of canonical modules of graded rings defined over a local ring. Our definition of canonical modules coincides with that of [GW] if the local ring is a field. In Section 2 we collect several facts about the behaviour of the local cohomology modules of Rees algebras. Section 3 will be devoted to the proof of our criterion of the Gorensteinness of $R(I)$ and to the construction of an example of a local ring (A, m, k) such that $R(m)$ is Gorenstein but A is not Cohen-Macaulay.

§ 1. Local cohomology of graded rings

In this section we give a brief summary of the theory of local cohomology and duality of graded rings.

Let $R = \bigoplus_{n \in \mathbb{Z}} R_n$ be a Noetherian graded ring and let M, N be graded

Received January 29, 1985.

R -modules. Let us denote the category of graded R -modules by $M_H(R)$. A morphism in $M_H(R)$ $f: M \rightarrow N$ is an R -linear map such that $f(M_n) \subset N_n$ for all $n \in \mathbb{Z}$. Let $n \in \mathbb{Z}$. We denote by $M(n)$ the graded R -module whose grading is defined by $M(n)_m = M_{n+m}$ for all $m \in \mathbb{Z}$. Let $\mathcal{H}om_R(M, N)_n$ be the abelian group of all homomorphisms from M into $N(n)$ in $M_H(R)$. Let $\mathcal{H}om_R(M, N) = \bigoplus_{n \in \mathbb{Z}} \mathcal{H}om_R(M, N)_n$. Then $\mathcal{H}om_R(M, N)$ is a graded R -module whose homogeneous component of degree n is $\mathcal{H}om_R(M, N)_n$. A graded R -module E is injective (resp. projective) in $M_H(R)$ if the functor $\mathcal{H}om_R(_, E)$ (resp. $\mathcal{H}om_R(E, _)$) from $M_H(R)$ into itself is an exact functor.

The tensor product $M \otimes_R N$ is a graded R -module whose n -th homogeneous component is the abelian group generated by the elements of the form $x \otimes y$ with $x \in M_i, y \in N_j$ and $i + j = n$.

The category $M_H(R)$ is an abelian category with enough injectives (cf. [Gr₁], (1, 10)). A homomorphism $f: M \rightarrow N$ in $M_H(R)$ is called essential if f is an injection and for any non-trivial graded R -submodule L of N we have $f(M) \cap L \neq 0$. The injective envelope of a graded R -module M is an injective object $\mathcal{E}_R(M)$ of $M_H(R)$ with an essential homomorphism $M \rightarrow \mathcal{E}_R(M)$ in $M_H(R)$.

The following proposition describes the structure of injective objects in $M_H(R)$.

PROPOSITION (1.1). (1) *Let M be a graded R -module. Then*

$$\text{Ass}_R(\mathcal{E}_R(M)) = \text{Ass}_R(M).$$

(2) *Let E be an injective object of $M_H(R)$. Then E is indecomposable if and only if $E = \mathcal{E}_R(R/p)(n)$ for some homogeneous prime ideal of R and for some $n \in \mathbb{Z}$.*

(3) *Every injective object of $M_H(R)$ can be decomposed into a direct sum of indecomposable injective objects of $M_H(R)$. This decomposition is unique up to isomorphism.*

Proof. This is [GW], (1.2.1).

For $i \geq 0$ the functor $\mathcal{E}_{x^i_R}(_, _)$ is defined to be the i -th derived functor of the functor $\mathcal{H}om_R(_, _)$. Suppose that M is a finitely generated graded R -module. Then $\mathcal{H}om_R(M, N) = \text{Hom}_R(M, N)$ as underlying R -modules. Hence $\mathcal{E}_{x^i_R}(M, N) = \text{Ext}_R^i(M, N)$ for all $i \geq 0$. For any $p \in \text{Spec}(R)$ and for any R -module L we define

$$\mu^i(p, L) = \dim_{k(p)} \text{Ext}_{R_p}^i(k(p), L_p),$$

where $k(p) = R_p/pR_p$, and call this number the i -th Bass number of M at p (cf. [B]).

PROPOSITION (1.2). *Let M be a graded R -module and let*

$$0 \rightarrow M \rightarrow I^0 \rightarrow I^1 \rightarrow \dots \rightarrow I^n \rightarrow I^{n+1} \rightarrow \dots$$

be a minimal injective resolution of M in $M_H(R)$. Then for any homogeneous prime ideal p and for any integer $i \geq 0$, $\mu^i(p, M)$ is equal to the number of the graded R -modules of the form $\mathcal{E}_R(R/p)(n)$ which appear in I^i as direct summands.

Proof. This is [GW], (1.2.4).

In this paper a Noetherian graded R is called defined over a local ring if R_0 is a Noetherian local ring and $R_n = 0$ for $n < 0$. If R is defined over a local ring we denote the graded ring $R \otimes_{R_0} \hat{R}_0$ by \hat{R} , where \hat{R}_0 is the completion of R_0 . In the rest of this section R denotes a graded ring defined over a local ring (R_0, m_0, k) and M denotes the maximal homogeneous ideal of R . R can be regarded as a graded R_0 -module in a natural way. Let E_{R_0} be the injective envelope of k as an R_0 -module. We denote by \mathcal{E}_{R_0} the graded R -module whose underlying R_0 -module is E_{R_0} and whose grading is given by $[\mathcal{E}_{R_0}]_0 = E_{R_0}$ and $[\mathcal{E}_{R_0}]_n = 0$ for $n \neq 0$.

DEFINITION (1.3). $\mathcal{E}_R(k) = \mathcal{H}om_{R_0}(R, \mathcal{E}_{R_0})$.

PROPOSITION (1.4). (1) $\mathcal{E}_R(k)$ is the injective envelope of R/M in $M_H(R)$.

(2) $\mathcal{H}om_R(\mathcal{E}_R(k), \mathcal{E}_R(k)) = R \otimes_{R_0} \hat{R}_0$, where \hat{R}_0 is the completion of R_0 .

Proof. (1) As in the non-graded case, in order to show that $\mathcal{E}_R(k)$ is injective in $M_H(R)$ it is enough to show that for any homogeneous ideal of R and for any integer n every homomorphism $f: I(n) \rightarrow \mathcal{E}_R(k)$ can be extended to a homomorphism $f': R(n) \rightarrow \mathcal{E}_R(k)$. Since

$$\mathcal{H}om_{R_0}(R, \mathcal{E}_{R_0}) \subset \text{Hom}_{R_0}(R, E_{R_0}) = \prod_{i \in \mathbb{Z}} \text{Hom}_{R_0}(R_i, E_{R_0}),$$

and since $\text{Hom}_{R_0}(R, E_{R_0})$ is an injective R -module f can be extended to an R -homomorphism $f'': R \rightarrow \text{Hom}_{R_0}(R, E_{R_0})$.

Let $f''(1) = (g_i)_{i \in \mathbb{Z}}$, where $g_i \in \text{Hom}_{R_0}(R_{-i}, E_{R_0})$. Since f is homogeneous for any homogeneous element $x \in I$ we have $xg_j = 0$ for $j \neq -n$. This shows that the homomorphism f' in $M_H(R)$ defined by $f'(1) = g_{-n} \in$

$\text{Hom}_{R_0}(R_n, E_{R_0})$ extends f . It is not difficult to show that $\text{Supp}(\mathcal{E}_R(k)) = M$. Moreover we have

$$\begin{aligned} \mathcal{H}om_R(R/M, \mathcal{E}_R(k)) &= \mathcal{H}om_R(R/M, \mathcal{H}om_{R_0}(R, \mathcal{E}_{R_0})) \\ &= \mathcal{H}om_{R_0}(R/M, \mathcal{E}_{R_0}) \\ &= k. \end{aligned}$$

This shows that $\mathcal{E}_R(k)$ is the injective envelope of R/M in $M_H(R)$.

$$\begin{aligned} (2) \quad \mathcal{H}om_R(\mathcal{E}_R(k), \mathcal{E}_R(k)) &= \mathcal{H}om_R(\mathcal{E}_R(k), \mathcal{H}om_{R_0}(R, \mathcal{E}_{R_0})) \\ &= \mathcal{H}om_{R_0}(\mathcal{E}_R(k), \mathcal{E}_{R_0}) \\ &= \mathcal{H}om_{R_0}(\mathcal{H}om_{R_0}(R, \mathcal{E}_{R_0}), \mathcal{E}_{R_0}) \\ &= \bigoplus_{n \in \mathbb{Z}} \text{Hom}_{R_0}(\text{Hom}_{R_0}(R_n, E_{R_0}), E_{R_0}) \\ &= \bigoplus_{n \in \mathbb{Z}} R_n \otimes_{R_0} \hat{R}_0 \\ &= R \otimes_{R_0} \hat{R}_0. \end{aligned}$$

PROPOSITION (1.5). *Let R be a graded ring defined over a complete local ring R_0 and N a graded R -module. Then, we have:*

- (1) *If N is Noetherian (resp. Artinian) $\mathcal{H}om_R(N, \mathcal{E}_R(k))$ is Artinian (resp. Noetherian).*
- (2) *If N is Noetherian or Artinian*

$$\mathcal{H}om_R(\mathcal{H}om_R(N, \mathcal{E}_R(k)), \mathcal{E}_R(k)) = N.$$

Proof. Using Proposition (1.4) this can be proved as in [M].

For every integer $i \geq 0$ we put

$$\mathcal{H}_M^i() = \varinjlim_n \mathcal{E}_{x^i_R}(R/M^n,)$$

and call it the i -th local cohomology functor, where R is a graded ring defined over a local ring and M is the maximal homogeneous ideal of R . $\mathcal{H}_M^i()$ is the i -th derived functor of $\mathcal{H}_M^0()$ (cf. [Gr₂] and [HK]).

DEFINITION (1.6). Suppose that R_0 is complete. We put

$$\mathcal{K}_R = \mathcal{H}om_R(\mathcal{H}_M^d(R), \mathcal{E}_R(k)),$$

where $d = \dim R$, and call this graded R -module the canonical module of R .

If R_0 is not complete a graded R -module \mathcal{K}_R is a canonical module of R if there is an isomorphism in $M_H(\hat{R})$ $\mathcal{K}_{\hat{R}} = \mathcal{K}_R \otimes_R \hat{R}$.

PROPOSITION (1.7). *If there is a canonical module of R it is a finitely generated R -module and unique up to isomorphisms.*

Proof. Since \hat{R} is faithfully flat over R it is sufficient to show that $\mathcal{H}_{\hat{R}}$ is finitely generated. But this follows from Proposition (1.5). For the proof of the uniqueness it is enough to show that if K and L are finitely generated graded R -modules such that $K \otimes_R \hat{R} = L \otimes_R \hat{R}$ then $K = L$. Let $f \in \mathcal{H}_{om_{\hat{R}}}(K \otimes_R \hat{R}, L \otimes_R \hat{R})_0$ be an isomorphism. Since \hat{R} is flat over R and K is finitely generated over R one gets

$$\begin{aligned} \mathcal{H}_{om_{\hat{R}}}(K \otimes_R \hat{R}, L \otimes_R \hat{R}) &= \mathcal{H}_{om_R}(K, L) \otimes_R \hat{R} \\ &= \mathcal{H}_{om_R}(K, L) \otimes_{R_0} \hat{R}_0 \end{aligned}$$

which implies that $\mathcal{H}_{om_{\hat{R}}}(K \otimes_R \hat{R}, L \otimes_R \hat{R})_0$ is the completion of $\mathcal{H}_{om_R}(K, L)_0$ since $\mathcal{H}_{om_R}(K, L)_0$ is a finitely generated R_0 -module. Let $\mathcal{H}_{om_R}(K, L)_0^\wedge$ be the m_0 -adic completion of $\mathcal{H}_{om_R}(K, L)_0$. For any integer $n > 0$ there is a homomorphism $f_n \in \mathcal{H}_{om_R}(K, L)_0$ such that $f - f_n \in m_0^n \mathcal{H}_{om_R}(K, L)_0^\wedge$. By assumption f_n induces an isomorphism $\bar{f}_n: K/m_0^n K \rightarrow L/m_0^n L$. Hence f_n is a surjective homomorphism. Since K/MK and L/ML are isomorphic there exist finitely generated graded free R -modules F and G of the same rank $\dim_k K/MK$ such that there are surjective homomorphisms in $M_H(R)$ $g: F \rightarrow K$ and $h: G \rightarrow L$. Let $S = \text{Ker}(g)$ and $T = \text{Ker}(h)$. We get a commutative diagram with exact rows

$$(I) \quad \begin{array}{ccccccc} 0 & \longrightarrow & S & \longrightarrow & F & \longrightarrow & K \longrightarrow 0 \\ & & \downarrow b_n & & \downarrow a_n & & \downarrow f_n \\ 0 & \longrightarrow & T & \longrightarrow & G & \longrightarrow & L \longrightarrow 0. \end{array}$$

a_n is an isomorphism since F and G are free R -modules of the same rank. Since \bar{f}_n is an isomorphism from (I) we get

$$T \subset b_n(S) + m_0^n G \cap T.$$

By Artin-Rees lemma there is an integer $r > 0$ such that

$$m_0^n G \cap T = m_0^{n-r} (m_0^r G \cap T) \quad \text{for } n > r.$$

Therefore we get $T \subset b_n(S) + m_0 T$ for $n > r$. By Nakayama's lemma $T = b_n(S)$. From (I) one knows that f_n is an isomorphism.

Let us recall that R is Cohen-Macaulay (resp. Gorenstein) if and only if R_M is Cohen-Macaulay (resp. Gorenstein), see [AG], [MR] and [GW].

PROPOSITION (1.8). *Let $d = \dim R$ and assume that R_0 is complete. Then R is Cohen-Macaulay if and only if for any finitely generate graded R -module N and for all $i \geq 0$ we have*

$$\mathcal{H}_{om_R}(\mathcal{H}_M^i(N), \mathcal{E}_R(k)) = \mathcal{E}_{xt_R^{d-i}}(N, \mathcal{K}_R).$$

Proof. Suppose that R is Cohen-Macaulay. We will show that the functor $T^i(\) = \mathcal{H}_{om_R}(\mathcal{H}_M^{d-i}(\), \mathcal{E}_R(k))$ is the i -th derived functor of $\mathcal{H}_{om_R}(\ , \mathcal{K}_R)$. We must show that

(1) from the short exact sequence $0 \rightarrow N' \rightarrow N \rightarrow N'' \rightarrow 0$ we obtain the long exact sequence

$$0 \rightarrow T^0(N'') \rightarrow T^0(N) \rightarrow T^0(N') \rightarrow T^1(N'') \rightarrow T^1(N) \rightarrow T^1(N') \rightarrow \dots$$

(2) $T^i(R) = 0$ for $i > 0$.

Since $\mathcal{E}_R(k)$ is an injective object in $M_H(R)$ (1) follows from the long exact sequence of the local cohomology. (2) follows from the fact that for any graded R -module N $\mathcal{H}_{om_R}(N, \mathcal{E}_R(k)) = 0$ if and only if $N = 0$. The converse is immediate.

PROPOSITION (1.9). *Suppose that R is Cohen-Macaulay. Then R is Gorenstein if and only if R has a canonical module \mathcal{K}_R and $\mathcal{K}_R = R(n)$ for some $n \in \mathbb{Z}$.*

Proof. Recall that R is Gorenstein if and only if

$$\mathcal{E}_{xt_R^i}(R/M, R) = \begin{cases} R/M & \text{for } i = \dim R \\ 0 & \text{for } i \neq \dim R. \end{cases}$$

If R is Gorenstein we have $\mathcal{H}_M^d(R) = \mathcal{E}_R(k)(n)$ for some $n \in \mathbb{Z}$. Hence $\mathcal{K}_{\hat{R}} = \hat{R}(-n)$ by Proposition (1.3). By the uniqueness of canonical modules we have $\mathcal{K}_R = R(-n)$. Conversely assume that $\mathcal{K}_R = R(-n)$ for some $n \in \mathbb{Z}$. By Proposition (1.8) we get

$$\mathcal{E}_{xt_{\hat{R}}^i}(\hat{R}/\hat{M}, \hat{R}) = \mathcal{H}_{om_{\hat{R}}}(\mathcal{H}_M^{d-i}(\hat{R}/\hat{M}, \mathcal{E}_{\hat{R}}(k))(n)$$

for all $i \geq 0$, where \hat{M} is the maximal homogeneous ideal of \hat{R} . Hence \hat{R} is Gorenstein since $\mathcal{H}_M^0(\hat{R}/\hat{M}) = \hat{R}/\hat{M}$ and $\mathcal{H}_M^i(\hat{R}/\hat{M}) = 0$ for $i > 0$. Since \hat{R} is faithfully flat over R it follows that R is Gorenstein.

Remark. Let $a = \max \{n \mid \mathcal{H}_M^d(R)_n \neq 0\}$. If R is Gorenstein we have $\mathcal{K}_R = R(a)$. In the sequel we denote this number by $a(R)$ and call it the a -invariant of R .

PROPOSITION (1.10). *Let $R \rightarrow S$ be a finite homomorphism of graded rings defined over local rings. Assume that R is Cohen-Macaulay and has a canonical module. Then*

$$\mathcal{K}_S = \mathcal{E}_{xt_R^r}(S, \mathcal{K}_R),$$

where $r = \dim R - \dim S$.

Proof. Let n_0 be the maximal ideal of S_0 and \hat{S}_0 be the n_0 -adic completion of S_0 . Since S_0 is finite over R_0 we have $\hat{S}_0 = S_0 \otimes_{R_0} \hat{R}_0$. Let N be the maximal homogeneous ideal of S and $\hat{N} = N \otimes_{R_0} \hat{R}_0$. Let $\hat{S} = S \otimes_{R_0} \hat{R}_0$. Note that $\mathcal{H}om_{\hat{R}}(\hat{S}, \mathcal{E}_{\hat{R}}(k))$ is the injective envelope of \hat{S}/\hat{N} in $M_H(\hat{S})$.

$$\begin{aligned} \mathcal{K}_{\hat{S}} &= \mathcal{H}om_{\hat{S}}(\mathcal{H}_{\hat{N}}^s(\hat{S}), \mathcal{E}_{\hat{S}}(\hat{S}/\hat{N})) \quad (s = \dim S) \\ &= \mathcal{H}om_{\hat{S}}(\mathcal{H}_{\hat{M}}^s(\hat{S}), \mathcal{H}om_{\hat{R}}(\hat{S}, \mathcal{E}_{\hat{R}}(k))) \\ &= \mathcal{H}om_{\hat{R}}(\mathcal{H}_{\hat{M}}^s(\hat{S}), \mathcal{E}_{\hat{R}}(k)) \\ &= \mathcal{E}_{xt_{\hat{R}}^r}(\hat{S}, \mathcal{K}_{\hat{R}}) \\ &= \mathcal{E}_{xt_R^r}(S, \mathcal{K}_R) \otimes_R \hat{R}. \end{aligned}$$

Since S is finite over R it follows that $\mathcal{K}_S = \mathcal{E}_{xt_R^r}(S, \mathcal{K}_R)$.

COROLLARY (1.11). *If moreover R is Gorenstein in Proposition (1.10) we get $\mathcal{K}_S = \mathcal{E}_{xt_R^n}(S, R)(n)$ for some $n \in \mathbb{Z}$.*

From Corollary (1.11) one knows that for any $p \in \text{Supp}(\mathcal{K}_S)$ $(\mathcal{K}_S)_p$ is a canonical module of the local ring S_p in the sense of [HK].

§ 2. Preliminaries

In this section we collect fundamental facts about the local cohomology of Rees algebras over Noetherian local rings.

Let (A, m, k) be a local ring and I an ideal of A . We put $R(I) = \bigoplus_{n \geq 0} I^n$ and call this graded A -algebra the Rees algebra of I . Let $I = (a_1, \dots, a_n)$. Then $R(I)$ can be identified with the subalgebra $A[a_1X, \dots, a_nX]$ of the polynomial ring $A[X]$ in one variable. Throughout this paper we use this identification without mentioning. Let $M = mR(I) + (a_1X, \dots, a_nX)R(I)$ be the maximal homogeneous ideal of $R(I)$. Let $G(I) = \bigoplus_{n \geq 0} I^n/I^{n+1}$ be the associated graded ring of I . Note that

$$G(I) = R(I)/IR(I) \quad \text{and} \quad A = R(I)/R(I)_+,$$

where $R(I)_+ = \bigoplus_{n > 0} I^n$. Let $\ell(I) = \dim R(I)/mR(I)$; we call this number the analytic spread of I . The analytic spread $\ell(I)$ of I is equal to the

minimum number of generators of a minimal reduction of I if the residue field k is infinite (cf. [NR]).

PROPOSITION (2.1). *Let (A, m, k) be a local ring and I an ideal of A with $\text{ht}(I) > 0$. If $R(I)$ is Cohen-Macaulay then*

- a) $a(G(I)) < 0$ and
- b) for $i < \dim A$ we have

$$\mathcal{H}_M^i(G(I))_n = \begin{cases} H_m^i(A) & \text{for } n = -1 \\ 0 & \text{for } n \neq -1. \end{cases}$$

Proof. For b) see the proof of [HI], Proposition (1.5). Let $J = R(I)_+$. From the exact sequences

$$0 \longrightarrow J \longrightarrow R(I) \longrightarrow A \longrightarrow 0$$

and

$$0 \longrightarrow J(1) \longrightarrow R(I) \longrightarrow G(I) \longrightarrow 0$$

we obtain the exact sequences of local cohomology

$$0 \longrightarrow H_m^d(A) \longrightarrow \mathcal{H}_M^{d+1}(J) \xrightarrow{f} \mathcal{H}_M^{d+1}(R(I)) \longrightarrow 0$$

and

$$0 \longrightarrow \mathcal{H}_M^d(G(I)) \longrightarrow \mathcal{H}_M^{d+1}(J)(1) \xrightarrow{g} \mathcal{H}_M^{d+1}(R(I)) \longrightarrow 0,$$

where $d = \dim A$. From this one gets the isomorphisms

$$f_n: \mathcal{H}_M^{d+1}(J)_n \longrightarrow \mathcal{H}_M^{d+1}(R(I))_n \quad \text{for } n \neq 0$$

and surjective homomorphisms

$$g_n: \mathcal{H}_M^{d+1}(J)_n \longrightarrow \mathcal{H}_M^{d+1}(R(I))_{n-1} \quad \text{for all } n.$$

Since $\mathcal{H}_M^{d+1}(J)$ and $\mathcal{H}_M^{d+1}(R(I))$ are Artinian $R(I)$ -modules their homogeneous components of sufficiently large degree are zero. By an easy diagram chase we know that $\mathcal{H}_M^{d+1}(J)_n = 0$ for $n \geq 1$ and $\mathcal{H}_M^{d+1}(R(I))_n = 0$ for $n \geq 0$. Now it is easy to see that $a(G(I)) < 0$.

COROLLARY (2.2). *Let A and I be the same as in Proposition (2.2). Then $\mathcal{H}_M^{d+1}(R(I))_n = 0$ for $n \geq 0$.*

Proof. This follows from the proof of Proposition (2.1).

If, in particular, $I = m$ we have the following result.

PROPOSITION (2.3). *If $d = \dim A > 0$ the following conditions are*

equivalent.

- (1) $R(m)$ is Cohen-Macaulay.
- (2) a) $a(G(m)) < 0$ and
 b) for $i < d$ we have

$$\mathcal{H}_M^i(G(m))_n = \begin{cases} H_m^i(A) & \text{for } n = -1 \\ 0 & \text{for } n \neq -1. \end{cases}$$

In this case A and $G(m)$ are Buchsbaum.

Proof. See [I].

For the technical simplicity in the rest of this paper we assume that every local ring has an infinite residue field.

LEMMA (2.4). *Let A and I be the same as above and let q be a minimal reduction of I . We put $r(q) = \min \{r \in \mathbb{Z} \mid I^{r+1} = qI^r\}$. If $\text{ht}(I) = \ell(I)$ we have $r(q) \geq a(G(I)) + \text{ht}(I)$.*

Proof. See [HI], Lemma (2.3).

LEMMA (2.5). *Let A and I be the same as above. We put*

$$n_i = \max \{n \in \mathbb{Z} \mid \mathcal{H}_M^i(G(I))_n \neq 0\} \quad \text{for } 0 \leq i \leq d = \dim A.$$

If I is m -primary we have $r(q) \leq \max_i \{n_i + i\}$ for any minimal reduction q of I .

Proof. Let $x \in A$. We denote by x^* the initial form of x with respect to \bar{I} . Let $q = (a_1, \dots, a_d)$ and $q^* = (a_1^*, \dots, a_d^*)$. Then

$$r(q) = \max \{r \in \mathbb{Z} \mid (G(I)/q^*)_r \neq 0\}.$$

If $\dim G(I) = 0$ the assertion is clear. Let $\dim G(I) > 0$. Since the residue field is infinite we may assume that $l_{a(i)}((0 : a_1^*)) < \infty$. From the exact sequences

$$0 \longrightarrow G(I)/(0 : a_1^*)(-1) \longrightarrow G(I) \longrightarrow G(I)/a_1^*G(I) \longrightarrow 0$$

and

$$0 \longrightarrow (0 : a_1^*) \longrightarrow G(I) \longrightarrow G(I)/(0 : a_1^*) \longrightarrow 0$$

we get the exact sequence

$$\mathcal{H}_M^i(G(I)) \longrightarrow \mathcal{H}_M^i(G(I)/a_1^*G(I)) \longrightarrow \mathcal{H}_M^{i+1}(G(I))(-1)$$

for $0 \leq i \leq d$. Let $n'_i = \max \{n \in \mathbb{Z} \mid \mathcal{H}_M^i(G(I)/a_1^*G(I))_n \neq 0\}$. Then $n'_i \leq \max \{n_i, n_{i+1} + 1\}$. By induction we have $r(q) \leq \max \{n_i + i\}$.

§3. The Gorensteinness of Rees algebras

This section is devoted to the proof of the following theorem.

THEOREM (3.1). *Let (A, m, k) be a local ring and I an ideal of A . Suppose that $R(I)$ is Cohen-Macaulay and $\text{grade}(I) \geq 2$. Then the following conditions are equivalent.*

- (1) $R(I)$ is Gorenstein.
- (2) $K_A = A$ and $\mathcal{H}_{G(I)} = G(I)(-2)$.

Remark. Since A and $G(I)$ are homomorphic images of $R(I)$, A and $G(I)$ have canonical modules if $R(I)$ is Gorenstein.

We need several preliminaries to prove this theorem.

LEMMA (3.2). *Let A be a local ring which has a canonical module K_A . Then the following conditions are equivalent.*

- (1) A satisfies (S_2) .
- (2) \hat{A} satisfies (S_2) .
- (3) $\text{Hom}_A(K_A, K_A) = A$.

Proof. See [A], (4.4) and (4.5).

LEMMA (3.3). *Let A and I be the same as in Theorem (3.1). Let $a \in I - I^2$ be an element whose initial form in $G(I)$ is a non zero-divisor. We put $\bar{R} = R(I)/(a, aX)$. If $R(I)$ is Gorenstein and $\text{grade}(I) \geq 2$ we have $\mathcal{H}_M^{d-1}(\bar{R}) = 0$, where $\dim A = d$.*

Proof. Let $R = R(I)$ and $G = G(I)$. Since a is a non zero divisor, by Propositions (1.8) and (1.9), it is enough to show that $\mathcal{E}_{\text{xt}_{R/aR}^1}(\bar{R}, R/aR) = 0$. Let $I = (a_1, \dots, a_n)$. Then we have the exact sequence

$$(R/aR)^n(-1) \xrightarrow{\begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix}} R/aR(-1) \xrightarrow{aX} R/aR \longrightarrow \bar{R} \longrightarrow 0.$$

Applying the functor $\mathcal{H}om_{R/aR}(\ , R/aR)$ to this sequence, we see

$$\mathcal{E}_{\text{xt}_{R/aR}^1}(\bar{R}, R/aR) = (aR : IR)/(a, aX).$$

Let $f^m \in (aR : IR)$, where $f \in I^m$ and $m \geq 0$. Then we have

$$f \in (aA : I) \cap (I^{m+1} : I) \subset (aA : I) \cap (I^{m+1} : a).$$

Since $\text{grade}(I) \geq 2$ we have $(aA : I) = aA$. That a^* is a non zero-divisor of $G(I)$ is equivalent to that $(I^m : a) = I^{m-1}$ for all $m > 0$. Hence we

have $(I^{m+1} : a) = I^m$. Therefore $f \in I^m \cap aA = aI^{m-1}$. This means $(aR : IR) = (a, aX)$, which completes the proof.

LEMMA (3.4). *Let A and I be the same as in Theorem (3.1). Assume that $R(I)$ is Cohen-Macaulay and $\mathcal{H}_{G(I)} = G(I)(-2)$. Then*

$$\mathcal{H}_{om_R(I)}(k, \mathcal{H}_M^{d+1}(R(I)))_n = 0 \quad \text{for } n \neq -1,$$

where $d = \dim A$ and $k = R(I)/M$.

Proof. Let R and G be as in the proof of Lemma (3.3). Put $J = \bigoplus_{n>0} R_n$. Then we get the exact sequence (cf. the proof of Proposition (2.1))

$$(I) \quad 0 \longrightarrow \mathcal{H}_{om_R}(k, H_m^d(A)) \longrightarrow \mathcal{H}_{om_R}(k, \mathcal{H}_M^{d+1}(J)) \xrightarrow{f} \mathcal{H}_{om_R}(k, \mathcal{H}_M^{d+1}(R)) \longrightarrow \mathcal{E}xt_R^1(k, H_m^d(A))$$

and

$$(II) \quad 0 \longrightarrow \mathcal{H}_{om_R}(k, \mathcal{H}_M^d(G)) \longrightarrow \mathcal{H}_{om_R}(k, \mathcal{H}_M^{d+1}(J))(1) \xrightarrow{g} \mathcal{H}_{om_R}(k, \mathcal{H}_M^{d+1}(R)) \longrightarrow \mathcal{E}xt_R^1(k, \mathcal{H}_M^d(G)).$$

Since $\mathcal{H}_{om_R}(k, H_m^d(A))$ is concentrated in degree 0 and since $\mathcal{E}xt_R^1(k, H_m^d(A))_n = 0$ for $n \leq -2$ from (I) we get isomorphisms

$$f_n : \mathcal{H}_{om_R}(k, \mathcal{H}_M^{d+1}(J))_n \longrightarrow \mathcal{H}_{om_R}(k, \mathcal{H}_M^{d+1}(R))_n$$

for $n \leq -2$. By assumption $\mathcal{H}_{om_R}(k, \mathcal{H}_M^d(G))_n = 0$ for $n \neq -2$. (II) yields injective homomorphisms

$$g_n : \mathcal{H}_{om_R}(k, \mathcal{H}_M^{d+1}(J))_n \longrightarrow \mathcal{H}_{om_R}(k, \mathcal{H}_M^{d+1}(R))_{n-1}$$

for $n \leq -2$. Since $\mathcal{H}_M^{d+1}(J)$ and $\mathcal{H}_M^{d+1}(R)$ are Artinian

$$\mathcal{H}_{om_R}(k, \mathcal{H}_M^{d+1}(J))_n = \mathcal{H}_{om_R}(k, \mathcal{H}_M^{d+1}(R))_n = 0 \quad \text{for } n \ll 0.$$

Now, it is easy to see that

$$\mathcal{H}_{om_R}(k, \mathcal{H}_M^{d+1}(R))_n = 0$$

for $n \leq -2$. On the other hand, by Corollary (2.2) we have $\mathcal{H}_M^{d+1}(R)_n = 0$ for $n \geq 0$. This completes the proof.

LEMMA (3.5). *Let (A, m, k) be a local ring and I an ideal of A such that $R(I)$ is Cohen-Macaulay. Suppose that $\text{grade}(I) \geq n > 0$. Then A and $G(I)$ satisfy (S_n) .*

Proof. We may assume that A is complete. Let $G = G(I)$. Let B be a Gorenstein local ring such that A is a homomorphic image of B and $d = \dim A = \dim B$. Let n be the maximal ideal of B . By the local duality we have

$$\text{Ext}_B^i(A, B) = \text{Hom}_B(H_n^{d-i}(A), E_B(B/n)) \quad \text{for } i \geq 0,$$

where $E_B(B/n)$ is the injective envelope of B/n as B -module. By Proposition (2.1) we see that $\text{Ext}_B^i(A, B)$ is annihilated by I for $i > 0$. Let $p \in \text{Spec}(A)$ and P be the inverse image of p in B . Then if $p \not\supseteq I$ we have

$$\text{Ext}_{B_P}^i(A_p, B_P) = 0 \quad \text{for } i > 0.$$

Hence A_p is Cohen-Macaulay. If $p \supseteq I$ we have $\text{depth } A_p \geq n$ by assumption. Therefore A satisfies (S_n) .

To prove the assertion on G we use induction on $\dim A/I$. Let $\dim A/I = 0$. By Proposition (2.1) we know that $l_G(\mathcal{H}_M^i(G)) < \infty$ for $i < d$ and $\text{depth } G_N \geq n$, where N is the maximal homogeneous ideal of G . Hence G satisfies (S_n) because G_Q is Cohen-Macaulay for $Q \in \text{Spec}(G) - \{N\}$. Let $\dim A/I > 0$. Note that G can be written as a homomorphic image of a Gorenstein graded ring of the same dimension. By Proposition (1.8) we see that G_p is Cohen-Macaulay if $p \not\supseteq G_+$, where $G_+ = \bigoplus_{n>0} G_n$. Assume that $p \supseteq G_+$ and $p \neq N$. Then $p \cap A/I = P/I$ for some $P \in \text{Spec}(A) - \{m\}$. Since $R(I)_P$ is Cohen-Macaulay and $\dim A/I > \dim A_P/IA_P$ one knows that G_P satisfies (S_n) by induction on $\dim A/I$.

Proof of Theorem (3.1). First we show that if $\text{ht}(I) > 0$ and $R(I)$ is Cohen-Macaulay then there is an element $a \in I - I^2$ whose initial form in $G(I)$ is a non zero-divisor. Since $\text{ht}(IR(I)) > 0$ one can choose an element $b \in I$ which is a non zero-divisor on $R(I)$. Noting that $R(I)/IR(I) + IXR(I) = A/I$, we have $\text{ht}(IR(I) + IXR(I)) = \dim R(I) - \dim A/I = d + 1 - \dim A/I \geq 2$. Since the residue field of A is infinite we can choose an element $c + aX$ of $IR(I) + IXR(I)$ such that $b, c + aX$ is an $R(I)$ -sequence and $a \in I - I^2$. Since b is also a non zero-divisor on A one can easily verify that $(bR(I) : bX) = IR(I)$. This implies that there exists an exact sequence

$$0 \longrightarrow G(I)(-1) \longrightarrow R(I)/bR(I) \longrightarrow R(I)/(b, bX)R(I) \longrightarrow 0.$$

By the choice of $c + aX$ we see that $c + aX$ is a non zero-divisor on $G(I)$. The canonical image of $c + aX$ in $G(I) = R(I)/IR(I)$ is nothing but

the initial form of a because $c \in I$. Therefore the initial form a^* of a in $G(I)$ is a non zero-divisor on $G(I)$.

(1) \Rightarrow (2): Let $R = R(I)$ and $G = G(I)$. Let $a \in I - I^2$ be as above. Since a is a non zero-divisor on A there are two exact sequences

$$(\#) \quad 0 \longrightarrow A \longrightarrow R/aXR \longrightarrow R/(a, ax) \longrightarrow 0$$

and

$$(\#\#) \quad 0 \longrightarrow G(-1) \longrightarrow R/aR \longrightarrow R/(a, aX) \longrightarrow 0.$$

These exact sequences induce the exact sequences by Lemma (3.3)

$$(+)\quad 0 \longrightarrow \mathcal{H}_M^d(A) \longrightarrow \mathcal{H}_M^d(R/aXR) \longrightarrow \mathcal{H}_M^d(R/(a, aX)) \longrightarrow 0$$

and

$$(++)\quad 0 \longrightarrow \mathcal{H}_M^d(G)(-1) \longrightarrow \mathcal{H}_M^d(R/aR) \longrightarrow \mathcal{H}_M^d(R/(a, aX)) \longrightarrow 0,$$

where $d = \dim A$ and M is the maximal homogeneous ideal of R as before. Since R is Gorenstein $\mathcal{K}_R = R(n)$ for some $n \in \mathbb{Z}$. Since aX is a non zero divisor of degree 1 we have $\mathcal{K}_{R/aXR} = R/aXR(n + 1)$. From the exact sequence (+) we know that $n = -1$ and $K_A = A/J$ for some ideal J of A . From (++) we have $\mathcal{K}_G = G/L(-2)$ for some homogeneous ideal L of G . By Lemma (3.5) A and G satisfy (S_2) , hence by Lemma (3.2) we have $J = 0$ and $L = 0$.

(2) \Rightarrow (1): From the exact sequences (#) and (\#\#) we get two injections $\mathcal{H}_M^{d-1}(R/(a, aX)) \rightarrow \mathcal{H}_M^d(A)$ and $\mathcal{H}_M^{d-1}(R/(a, aX)) \rightarrow \mathcal{H}_M^d(G)(-1)$ since R is Cohen-Macaulay. From the first one we know that $\mathcal{H}_M^{d-1}(R/(a, aX))$ is concentrated in degree 0. The assumption $\mathcal{K}_G = G(-2)$ shows that $\mathcal{H}_M^d(G)_n = 0$ for $n \geq -1$. From the second injection we see that $\mathcal{H}_M^{d-1}(R/(a, aX)) = 0$. Hence we have the exact sequences (+) and (++) . By Lemma (3.4) we know that $\mathcal{H}_{om_R}(k, \mathcal{H}_M^d(R/aXR))$ is concentrated in degree 0. By (+) we get

$$\mathcal{H}_{om_R}(k, \mathcal{H}_M^d(R/aXR)) = \text{Hom}_A(k, H_m^d(A))$$

since $\mathcal{H}_M^d(R/aR)_n = \mathcal{H}_M^d(R/(a, aX))_n = 0$ for $n \geq 0$ by Corollary (2.2). By the assumption $K_A = A$ we have $\text{Hom}_A(k, H_m^d(A)) = k$. This shows that R is Gorenstein.

Let I be an ideal of a local ring and q a minimal reduction of I . We put $r(q) = \min\{r \mid I^{r+1} = qI^r\}$. We call $r(q)$ the reduction exponent of q .

COROLLARY (3.6) *Let A be a local ring and I an ideal of A such that $\text{ht}(I) = \ell(I) > 0$ and $R(I)$ is Cohen-Macaulay. Then we have:*

(1) *Suppose that $a(G(I)) \geq -2$. Then we have $r(q) = \text{ht}(I) - 1$ or $\text{ht}(I) - 2$ for any minimal reduction q of I .*

(2) *Suppose moreover that $\text{grade}(I) \geq 2$ and $R(I)$ is Gorenstein. Then for any minimal reduction q of I we have $r(q) = \text{ht}(I) - 2$ if and only if $\text{depth } A \geq \dim A/I + 2$.*

Proof. (1) By induction on $\dim A/I$. If $\dim A/I = 0$ this follows from Lemmas (2.4) and (2.5). Let $\dim A/I > 0$. Choose an element $b \in A$ whose image in A/I is a part of system of parameters of A/I . By Proposition (1.5) b is a non zero-divisor on $G(I)$ and $R(I)$ and we have $R(I)/bR(I) = R(I(A/bA))$ and $G(I) = G(I(A/bA))$. It is easy to see that the ideal $I(A/bA)$ in A/bA satisfies the same assumption on I . By induction hypothesis we have $r(q(A/bA)) = \text{ht}(I) - 1$ or $\text{ht}(I) - 2$. By Nakayama's lemma we have $r(q(A/bA)) = r(q)$.

(2) First we assume that $\text{depth } A \geq \dim A/I + 2$. If $\dim A/I = 0$ we have $r(q) = \text{ht}(I) - 2$ by Proposition (2.1), Lemmas (2.4) and (2.5). We proceed by induction on $\dim A/I$. Let $\dim A/I > 0$. Then by assumption $\text{depth } A \geq 3$. Let a_1, a_2 be a regular sequence in I . One can choose an element $b \in m$ so that a_1, a_2, b is a regular sequence and the image of b in A/I is a part of system of parameters of A/I . Then $\text{grade}(I(A/bA)) \geq 2$ and $R(I(A/bA))$ is Gorenstein. Since $\text{depth } A/bA \geq \dim A/(b, I) + 2$ we have $r(q) = \text{ht}(I) - 2$ by induction hypothesis.

Conversely assume that $r(q) = \text{ht}(I) - 2$. Let $b_1, \dots, b_s \in m$ be a system of parameters of A/I . We set $\bar{A} = A/(b_1, \dots, b_s)$. Since b_1, \dots, b_s is a regular sequence we have only to show that $\text{depth } \bar{A} \geq 2$. Let $\bar{I} = I\bar{A}$ and $\bar{q} = q\bar{A}$. Since $r(\bar{q}) = \text{ht}(\bar{I}) - 2$ we see that $\mathcal{H}_M^{h_n}(G(\bar{I})) = 0$ for $n \geq -1$ by Lemma (2.4), where $h = \text{ht}(I)$. By [HI], Proposition (1.5) we know that b_1, \dots, b_s is a $G(I)$ -sequence. Let $q_i = (b_1, \dots, b_i)$ for $1 \leq i \leq s$. Then we see that $G(I)/q_i G(I) = G(I(A/q_i))$. We set $G_i = G(I)/q_i G(I)$ for $1 \leq i \leq s$. Then we have an exact sequence

$$\mathcal{H}_M^{d-i}(G_{i-1}) \xrightarrow{b_i} \mathcal{H}_M^{d-i}(G_{i-1}) \longrightarrow \mathcal{H}_M^{d-i}(G_i) \longrightarrow \mathcal{H}_M^{d-i+1}(G_{i-1}).$$

Since $r(q(A/q_i)) = \text{ht}(I) - 2$ we know that $\mathcal{H}_M^{d-i}(G_i) = 0$ for $n \geq -1$ by Lemma (2.4). By Proposition (2.1) we see that $H_m^{d-i}(A/q_{i-1}) = b_i H_m^{d-i}(A/q_{i-1})$ for $1 \leq i \leq s$.

This implies that $K_{A/q_i} = A/q_i$ for $0 \leq i \leq s$, where $q_0 = 0$. In particular, $K_{\bar{A}} = \bar{A}$. One sees that $\text{depth } \bar{A} \geq 2$ by [A]. The following is a generalization of a result in [GS].

COROLLARY (3.7). *Let A be a Cohen-Macaulay local ring and I an ideal of A with $\text{ht}(I) = \ell(I) \geq 2$. Then the following conditions are equivalent.*

- (1) $R(I)$ is Gorenstein.
- (2) $G(I)$ is Gorenstein and $a(G(I)) = -2$.
- (3) $G(I)$ is Gorenstein and there exists a minimal reduction q of I such that $r(q) = \text{ht}(I) - 2$.

In this case A is Gorenstein.

Proof. This follows from Theorem (3.1), Corollary (3.6) and the fact that the Gorensteinness of $G(I)$ implies that of A .

COROLLARY (3.8). *Let A and I be the same as in Corollary (3.6). Suppose that*

- (1) $R(I)$ is Gorenstein,
- (2) $l_A(H_m^i(A)) < \infty$ for $i < d = \dim A$ and
- (3) $2\text{ht}(I) \leq \dim A$.

Then A is Gorenstein.

Proof. By Corollary (3.7) it is sufficient to prove that A is Cohen-Macaulay. Let b_1, \dots, b_s be a system of parameters of A/I . We put $q_i = (b_1, \dots, b_i)$ and $G_i = G(I)/q_i G(I)$ for $1 \leq i \leq s$. Let $a_1, \dots, a_n, h = \text{ht}(I)$, be a minimal generators of a minimal reduction of I . Then $a_1, \dots, a_n, b_1, \dots, b_s$ is a system of parameters of A . Since $l_A(H_m^i(A)) < \infty$ for $i < d$ we know that if $t \leq \text{depth } A$ then any t elements of a system of parameters of A form a regular sequence by [CST], (3.3). Hence we have $\text{grade}(I(A/q_i)) \geq 2$ for $1 \leq i < s$ by [HI], Proposition (1.5). We are going to show that $H_m^{d-s+i}(A/q_{s-j}) = 0$ for $2 \leq j \leq s - 2$ and $1 \leq i \leq j - 1$ by induction on j . Let $j = 2$. From the exact sequence

$$\mathcal{H}_M^{d-s+1}(G_{s-2}) \xrightarrow{b_{s-1}} \mathcal{H}_M^{d-s+1}(G_{s-2}) \longrightarrow \mathcal{H}_M^{d-s+1}(G_{s-1})$$

we get $H_m^{d-s+1}(A/q_{s-2}) = b_{s-1}H_m^{d-s+1}(A/q_{s-2})$ by Theorem (3.1) and Proposition (2.1) since $R(I(A/q_{s-1}))$ is Gorenstein. By the assumption (2) we get $H_m^{d-s+1}(A/q_{s-2}) = 0$. Let us assume that our assertion is true for $j < s - 2$ and we will prove that the assertion is true for $j + 1$. Since b_{s-j} is a

non zero-divisor on A/q_{s-j-i} we obtain the exact sequence

$$H_m^{d-s+i}(A/q_{s-j-1}) \xrightarrow{b_{s-j}} H_m^{d-s+i}(A/q_{s-j-1}) \longrightarrow H_m^{d-s+i}(A/q_{s-j})$$

for $i > 0$. By the induction hypothesis $H_m^{d-s+i}(A/q_{s-j}) = 0$ for $1 \leq i \leq j - 1$. Therefore $H_m^{d-s+i}(A/q_{s-j-1}) = 0$ for $1 \leq i \leq j - 1$ by assumption (2). It remains to prove that $H_m^{d-s+j}(A/q_{s-j-1}) = 0$. But this can be proved by the same method used for $j = 2$. Hence, in particular, we get $H_m^i(A) = 0$ for $h + 1 \leq i \leq d$. By assumption (3) we get $\text{depth } A \geq \dim A/I + 1 \geq h + 1$ cf. [HI]. Therefore A is Cohen-Macaulay.

§ 4. Example

In this section we construct a local ring (A, m, k) such that $R(m)$ is Gorenstein but A is not Cohen-Macaulay. For a local ring A we denote the multiplicity of A by $e(A)$.

LEMMA (4.1). *Let (A, m, k) be a local ring with $\dim A = 3$ and $q = (a_1, a_2, a_3)$ be a minimal reduction of m . Let*

$$I = ((a_1, a_2) : a_3) + ((a_2, a_3) : a_1) + ((a_1, a_3) : a_2) + m^2.$$

Then $R(m)$ is Cohen-Macaulay if and only if $m^3 = qm^2$ and $l_A(I/m^2) = 3(l_A(A/q) - e(A)) + 3$.

Proof. See [I₂], Theorem 5.

LEMMA (4.2). *Let A be the same as in Lemma (4.1). Suppose that $R(m)$ is Cohen-Macaulay and A is not Cohen-Macaulay. If $l_A(m/m^2) = 6$ we have*

- (1) A is a Buchsbaum ring with $\text{depth } A = 2$ and $l_A(H_m^2(A)) = 1$,
- (2) $m^2 = qm$ for any minimal reduction q of m and
- (3) $e(A) = 3$.

Proof. See [I₂], Corollary 11.

EXAMPLE (1). Let k be a field and X_i, Y_i ($1 \leq i \leq 3$) be indeterminates over k . We put

$$A = k[[X_1, X_2, X_3, Y_1, Y_2, Y_3]]/(X_1Y_1 + X_2Y_2 + X_3Y_3, (Y_1, Y_2, Y_3)^2).$$

Then A is not Cohen-Macaulay but $R(m)$ is Cohen-Macaulay. By Lemma (4.2) $e(A) = 3$ (cf. [I₁] and [I₂]).

EXAMPLE (2). Let k be a field of $\text{ch}(k) = 2$ and $X_1, X_2, X_3, Y_1, Y_2, Y_3, Y_4$ indeterminates over k . Let

$$A = k[[X_1, X_2, X_3, Y_1, Y_2, Y_3, Y_4]]/J$$

$$= k[[x_1, x_2, x_3, y_1, \dots, y_4]]$$

where J is the ideal generated by $X_1Y_1 + X_2Y_2 + X_3Y_3, Y_1^2, Y_2^2, Y_3^2, Y_4^2, Y_1Y_2, Y_2Y_4, Y_3Y_4, Y_1Y_2 - X_3Y_4, Y_2Y_3 - X_1Y_4$ and $Y_1Y_3 - X_2Y_4$.

Then A is not Cohen-Macaulay but $R(m)$ is Gorenstein.

To prove this we need the following lemma.

LEMMA (4.3). $(0 : y_i) = (y_1, \dots, y_i)$

Proof. Let $R = k[[X_1, X_2, X_3, Y_1, Y_2, Y_3, Y_4]]$ and let $f \in (J : Y_i)$. Since

$$(J : Y_i) = (X_1Y_1 + X_2Y_2 + X_3Y_3, Y_1Y_2 - X_3Y_4, Y_2Y_3 - X_1Y_4, Y_1Y_3 - X_2Y_4) : Y_i + (Y_1, \dots, Y_i)$$

we may assume that f belongs to the first ideal on the right side. Let us write

$$fY_i = (g_1 + g'_1Y_4)(X_1Y_1 + X_2Y_2 + X_3Y_3) + (g_2 + g'_2Y_4)(Y_1Y_2 - X_3Y_4) + (g_3 + g'_3Y_4)(Y_2Y_3 - X_1Y_4) + (g_4 + g'_4Y_4)(Y_1Y_3 - X_2Y_4),$$

where $g_i \in k[[X_1, X_2, X_3, Y_1, Y_2, Y_3]]$. From this we see that

$$(I) \quad g_1(X_1Y_1 + X_2Y_2 + X_3Y_3) + g_2Y_1Y_2 + g_3Y_2Y_3 + g_4Y_1Y_3 = 0$$

and

$$(II) \quad f \equiv -g_2X_3 - g_3X_1 - g_4X_4 \pmod{(Y_1, \dots, Y_i)}.$$

From (I) we have

$$Y_1(g_1X_1 + g_2Y_2 + g_4Y_3) + Y_2(g_1X_2 + g_3Y_3) = 0.$$

Since Y_1, Y_2 is a regular sequence in R we have

$$g_1X_1 + g_2Y_2 + g_4Y_3 = hY_2$$

$$g_1X_2 + g_3Y_3 = -hY_1$$

for some $h \in R$. Since X_1, Y_2, Y_3 and X_2, Y_1, Y_3 are regular sequences in R there are elements $a_1, a_2, a_3, b_1, b_2, b_3$ of R such that

$$(III) \quad (g_1, g_2 - h, g_4) = (X_1, Y_2, Y_3) \begin{pmatrix} 0 & a_1 & a_2 \\ -a_1 & 0 & a_3 \\ -a_2 & -a_3 & 0 \end{pmatrix}$$

$$(IV) \quad (g_1, h, g_3) = (X_2, Y_1, Y_3) \begin{pmatrix} 0 & b_1 & b_2 \\ -b_1 & 0 & b_3 \\ -b_2 & -b_3 & 0 \end{pmatrix}.$$

Hence

$$g_1 = -a_1 Y_2 - a_2 Y_3 = -b_1 Y_1 - b_2 Y_3.$$

Since $\text{ch}(k) = 2$ we have

$$\left. \begin{matrix} a_2 + b_2 \equiv 0 \\ a_1 \equiv 0 \\ b_1 \equiv 0 \end{matrix} \right\} \text{mod } (Y_1, \dots, Y_4).$$

By (II), (III) and (IV) we obtain

$$\begin{aligned} f &\equiv a_1 X_1 X_3 + b_1 X_2 X_3 + (a_2 + b_2) X_1 X_2 \\ &\equiv 0 \text{ mod } (Y_1, \dots, Y_4). \end{aligned}$$

Proof of Example (2). By Lemma (4.3) we have $(0 : y_4) = (y_1, \dots, y_4)$. From the exact sequence

$$0 \longrightarrow A/(0 : y_4) \longrightarrow A \longrightarrow A/y_4 A \longrightarrow 0$$

we get $e(A) = e(A/(0 : y_4)) + e(A/y_4 A)$. Since $A/y_4 A$ is isomorphic to the local ring in Example (1) and since $A/(0 : y_4)$ is a regular local ring we have $e(A) = 4$. It is easy to see that x_1, x_2, x_3 is a system of parameters of A and that $((x_1, x_2) : x_3) = (x_1, x_2, y_3)$, $((x_2, x_3) : x_1) = (x_2, x_3, y_1)$ and $((x_1, x_3) : x_2) = (x_1, x_3, y_2)$. This shows that A is not Cohen-Macaulay. It is easy to verify that $m^2 = (x_1, x_2, x_3)m$. By Lemma (4.1) we see that $R(m)$ is Cohen-Macaulay. By Theorem (3.1) it is enough to show that $\mathcal{K}_{G(m)} = G(m)(-2)$. Since A is defined by homogeneous polynomials

$$G(m) = k[X_1, X_2, X_3, Y_1, Y_2, Y_3, Y_4]/J^* = k[x_1, x_2, \dots, y_4],$$

where J^* is generated by the polynomials generating J . Let $S = k[x_1, x_2, x_3]$. Then S is a polynomial ring with $\dim S = 3$ and $G(m)$ is generated by $1, y_1, y_2, y_3, y_4$ as an S -module. Since $G(m)$ has rank 4 as an S -module and $\text{depth } G(m) = 2$ we get a finite free resolution of $G(m)$ as S -module

$$0 \longrightarrow S(-2) \xrightarrow{[0, x_1, x_2, x_3, 0]} S \oplus S^4(-1) \xrightarrow{d} G(m) \longrightarrow 0,$$

where d is given by $d(e_0) = 1$ and $d(e_i) = y_i$ for $1 \leq i \leq 4$, with suitable free basis e_0, e_1, \dots, e_4 of $S \oplus S^4(-1)$ with $\deg(e_0) = 0$ and $\deg(e_i) = 1$ for

$1 \leq i \leq 4$. By Corollary (1.11) $\mathcal{H}_{G(m)} = \mathcal{H}_{om_S}(G(m), S(-3))$.

The $G(m)$ -structure of $\mathcal{H}_{G(m)}$ is given by

$$(xf)(y) = f(xy) \quad \text{for } f \in \mathcal{H}_{om_S}(G(m), S(-3)) \text{ and } x, y \in G(m).$$

$\mathcal{H}_{G(m)}$ is generated by $e_0^*, x_2e_3^* - x_3e_2^*, x_3e_1^* - x_1e_3^*, x_1e_2^* - x_2e_1^*$ and e_4^* as an S -module, where e_i^* is the dual base of e_i with $\deg(e_0^*) = 3$ and $\deg(e_i^*) = 2$ for $1 \leq i \leq 4$. Using the fact that $\text{ch}(k) = 2$ we can easily verify the following relations as $G(m)$ -module.

$$\begin{aligned} y_4e_4^* &= e_0^* \\ y_1e_4^* &= x_2e_3^* - x_3e_2^* \\ y_2e_4^* &= x_3e_1^* - x_1e_3^* \\ y_3e_4^* &= x_1e_2^* - x_2e_1^* \end{aligned}$$

Hence $\mathcal{H}_{G(m)} = G(m)(-2)$ and hence $R(m)$ is Gorenstein by Theorem (3.1).

EXAMPLE (3). Let A be same as in Example (2). We put $B = A[[T_1, \dots, T_n]]$, where T_1, \dots, T_n are indeterminates over A . Let $I = mB$. Then $R(I) = R(m) \otimes_A B$ is Gorenstein since B is faithfully flat over A . If $n \geq 3$ we have $2 \text{ ht}(I) = 6 \leq \dim B$. But B is not Gorenstein.

Remark. a) If in Example (2) $\text{ch}(k) \neq 2$ A is not Buchsbaum. This can be seen as follows. If A is Buchsbaum we have

$$\begin{aligned} e(A) &= l_A(A/(x_1, x_2, x_3)) - l_A((x_1, x_2) : x_3/(x_1, x_2)) \\ &= 5 - 1 = 4. \end{aligned}$$

On the other hand one can easily see $(0 : y_4) \supset (y_1, \dots, y_4, x_1x_2)$ and $\dim A/(0 : y_4) < 3$. This implies $e(A) = e(A/y_4A) = 3$, a contradiction.

b) Example (3) shows that Corollary (3.8) is false without any restriction on the local cohomology modules of A .

REFERENCES

[A] Y. Aoyama, Some basic results on canonical modules, *J. Math. Kyoto Univ.*, **23** (1983), 85–94.
 [AG] Y. Aoyama and S. Goto, On the type of graded Cohen-Macaulay rings, *J. Math. Kyoto Univ.*, **15** (1975), 275–284.
 [B] H. Bass, On the ubiquity of Gorenstein rings, *Math. Z.*, **82** (1963), 8–28.
 [Gr₁] A. Grothendieck, Sur quelque points d’algebre homologique, *Tohoku Math. J.*, **IX** (1957), 119–221.
 [Gr₂] —, Local cohomology, *Lect. Notes in Math.*, **41**, Berlin-Heidelberg-New York, 1967.

- [GS] S. Goto and Y. Shimoda, On the Rees algebras of Cohen-Macaulay local rings, Commutative algebra (analytic methods), Lecture Notes in Pure and Applied Mathematics, **68** (1982), 201–231.
- [GW] S. Goto and K. Watanabe, On graded rings I, *J. Math. Soc. Japan*, **30** (1978), 179–213.
- [HI] M. Herrmann and S. Ikeda, Remarks on lifting of Cohen-Macaulay property, *Nagoya Math. J.*, **92** (1983), 121–132.
- [HK] J. Herzog and E. Kunz, Der kanonische Modul eines Cohen-Macaulay Rings, Springer Lect. Notes in Math., **238** (1971).
- [I₁] S. Ikeda, The Cohen-Macaulayness of the Rees algebras of local rings, *Nagoya Math. J.*, **89** (1983), 47–63.
- [I₂] —, Remarks on Rees algebras and graded rings with multiplicity 3, Preprint.
- [M] E. Matlis, Injective modules over Noetherian rings, *Pacific J. Math.*, **8** (1958), 511–528.
- [MR] J. Matijevic and P. Roberts, A conjecture of Nagata on graded Cohen-Macaulay rings, *J. Math. Kyoto Univ.*, **14** (1974) 125–128.
- [NR] D. G. Northcott and D. Rees, Reductions of ideals in local rings, *Proc. Camb. Phil. Soc.*, **50** (1954), 145–158.
- [CST] N. T. Cuong, P. Schenzel and N. V. Trung, Verallgemeinerte Cohen-Macaulay-Moduln, *Math. Nachr.*, **85** (1978), 57–73.