

peripheral neuropathy symptoms (pins and needles and numbness), dizziness, anosmia, symptoms of depression, anxiety and fatigue. Patients complain of reduced quality of life and impairment on daily functioning. Although the burden of disease is high there is until now very few data available, the etiopathology is still unknown and treatment strategies are not established.

Objectives: The objective of this study is to gather standardized data of patients with long-covid syndrome who suffer from neuropsychiatric symptoms in order to better understand the complexity of this syndrome.

Methods: Patients were referred from the long-covid outpatient unit of the internal medicine department to our specialized outpatient unit, so that the previous infection was confirmed. A standardized psychiatric interview and a thorough neuropsychological assessment was conducted.

Results: We will present preliminary data on psychiatric symptoms, neuropsychology and quality of life with patients with long-covid syndrome.

Conclusions: Potential treatment strategies to improve psychiatric and neurocognitive symptoms as well as improvement of quality of life will be discussed.

Disclosure: Daniela Roesch Ely and Matthias Weisbrod have a contract with Schuhfried GmbH (development of neurocognitive batteries and training programs)

Keywords: Long COVID; psychiatry; Neuropsychology

EPV0515

Suicide presentation and the risk at the time of the mandatory quarantine for the SARS-Cov-2 pandemic: medico-legal and forensic aspects

C. Scalise¹, M.A. Sacco¹, A. Zibetti¹, P. De Fazio², P. Ricci¹ and I. Aquila^{1*}

¹Institute of Legal Medicine, Medical And Surgical Science, Catanzaro, Italy and ²Institute of Psychiatry, Medical And Surgical Sciences, Catanzaro, Italy

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1356

Introduction: COVID-19 pandemic is the most important health emergency of the 21st century. Since the high number of infected people and as there is still no specific therapy worldwide, the pandemic has been countered through the application of prevention measures based on social distancing and home isolation. These elements are known risk factors for the development of various psychiatric conditions. From a forensic point of view, these pathologies are related to a high suicide rate.

Objectives: It is no coincidence that during the previous pandemics that have occurred in history there has been a significant increase in suicides. By this work, we therefore want to highlight the psychological consequences of a pandemic and the importance of preventive strategies.

Methods: It is important to focus not only on physical well-being but also on the psychological aspects that the pandemic produces in the daily life of each individual

Results: If the infecting agent causes the death of millions of people around the world, the socio-economic context that is created indirectly determines as many deaths.

Conclusions: Therefore it is necessary to underline how it is advisable to implement preventive measures in order to significantly

reduce deaths from suicide, a problem with an important impact in the social and forensic fields.

Disclosure: No significant relationships.

Keywords: Suicide; covid 19; emergency

EPV0519

Burnout syndrome in healthcare professionals almost two years after the declaration of the Covid-19 pandemic

M. Solis*, A. Jurado Arevalo and E. Blázquez Garcia

Universitary Hospital of Jaén, Psychiatric, Jaén, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1357

Introduction: The coronavirus disease 2019 (COVID-19) pandemic has caused major sanitary crisis worldwide. Frontline healthcare workers face many difficulties, such as: direct exposure to patients with high viral load, physical exhaustion, reorganization of workspaces, face the unusually high number of deaths among patients, colleagues or relatives and ethical issues in a tense health system.

Objectives: Provide up-to-date information of Burnout syndrome associated with exposure of healthcare workers to the COVID-19 pandemic, after almost 20 months of the declaration of pandemic by the World Health Organization.

Methods: A cross-sectional study was carried out that included 84 healthcare workers from Spain in October 2021, through an anonymous, voluntary and multiple response type online survey which included questions about sociodemographic aspects and the Maslach burnout inventory

Results: 62% were doctors and 29% were nurses. 70% work on the front line of Covid-19. 38% report not having been able to enjoy their vacations when they wanted. 8% admit to having had suicidal ideas. Almost 52% admit low personal fulfillment, 38.6% admit a high depersonalization count, and 45% report high emotional exhaustion. Of the total sample, 17 respondents have burnout syndrome.

Conclusions: It is necessary create strategies to promote mental well-being in health professionals exposed to COVID-19 after 20 months of active work. Protecting and identifying health care professionals who could be at high risk for developing a mental health pathology or detecting Burnout syndrome in them should be the priority of public health post pandemic.

Disclosure: No significant relationships.

Keywords: Healthcare professionals; burnout; mental health; Covid-19

EPV0520

Covid-19: knowledge, risk perception, trust and vaccination readiness among German medical students

F. Baessler

Centre for Psychosocial Medicine, Department Of General Internal And Psychosomatic Medicine, Heidelberg, Germany

doi: 10.1192/j.eurpsy.2022.1358

Introduction: The Covid-19 pandemic has highlighted the urgency for innovative vaccine strategies since the best of vaccines cannot be useful if people do not accept vaccinations. The current situation suggests that vaccinology has been ignored in the medical curriculum and needs more representation in teaching.