

Fruit and vegetable intakes, sources and contribution to total diet in very young children (1–4 years): the Irish National Pre-School Nutrition Survey

Laura O'Connor^{1,2}, Janette Walton^{1*} and Albert Flynn¹

¹School of Food and Nutritional Sciences, University College Cork, Cork, Republic of Ireland

²Food and Nutrition Research Cluster, Holling's Faculty, Manchester Metropolitan University, Manchester M15 6BH, UK

(Submitted 5 January 2016 – Final revision received 4 March 2016 – Accepted 11 March 2016 – First published online 22 April 2016)

Abstract

Although the importance of fruit and vegetable (F&V) intakes in the prevention of chronic diseases is well established, there are limited data on intakes in very young children. This study estimates F&V intakes and sources and the contribution to the total diet using data from the National Pre-School Nutrition Survey, a nationally representative sample (n 500) of Irish children aged 1–4 years. A 4-d weighed food record was used to collect food intake data. Of 1652 food codes consumed, 740 had a fruit/vegetable component. The percentage of edible fruits and/or vegetables in each food code was calculated. Intakes (g/d), sources (g/d) and the contribution of F&V to the weight of the total diet (%) were estimated, split by age. All children consumed F&V. Intakes of total fruits, in particular fruit juice, increased with age. The contribution to total fruit intake was discrete fruits (47–56% range across age), 100% fruit juice, smoothies and purees (32–45%) as well as fruits in composite dishes (7–13%). Total vegetable intake comprised of discrete vegetables (48–62% range across age) and vegetables in composite dishes (38–52%). F&V contributed on average 20% (15% fruit; 5% vegetables) to the weight of the total diet and was <10% in sixty-one children (12%). F&V contributed 50% of vitamin C, 53% of carotene, 34% of dietary fibre and 42% of non-milk sugar intakes from the total diet. F&V are important components of the diet of Irish pre-school children; however, some aspects of F&V intake patterns could be improved in this age group.

Key words: Children: Fruits: Vegetables: Juices: Dietary intakes: Surveys: Composite dishes

Early childhood is a pivotal time in the development of food preferences and as such an opportunity to foster the development of healthy eating practices. Fruit and vegetable intakes play an important role in the prevention of chronic diseases⁽¹⁾ and obesity⁽²⁾, and to halt the escalating prevalence of both, fruit and vegetable consumption should be encouraged from an early age. There are, however, limited data on fruit and vegetable intakes in very young children.

The beneficial role of both quantity and variety of fruit and vegetable intake has been acknowledged in disease prevention in adults⁽³⁾. There are also unresolved questions as to the health outcomes associated with fruit juice intake^(4–6) and concerns about sugar intake including that from fruits^(7,8). Therefore, it is important to characterise fruit and vegetable intake by identifying the dietary sources and estimating the contribution to nutrient intakes.

At present, there are no established quantitative guidelines for fruit and vegetable intakes for very young children in Ireland or in many other countries. However, there are number of evidence-based resources available, including 'The Infant and Toddler Forum'⁽⁹⁾. It is generally acknowledged that because of the volume that a young child can consume, 400 g/d or 5 (80 g) portions/d, the guideline for those aged ≥ 5 years in Ireland^(10,11) and the World Health Organization population goal⁽¹⁾, is not appropriate. Evaluating the proportion of the diet

that is comprised of fruits and vegetables in very young children could inform whether public health intervention such as the development of an age-specific guideline is necessary. The aim of this study was to estimate fruit and vegetable intakes and identify the sources and contribution to the total diet in Irish pre-school children aged 1–4 years using data from the National Pre-School Nutrition Survey (NPNS).

Methods

Survey design and population

The NPNS, a nationally representative dietary survey, was carried out by the Irish Universities Nutrition Alliance between October 2010 and September 2011 to establish a database of habitual food and drink consumption in a representative sample of Irish children aged 1–4 years.

In total, 500 (boys 251, girls 249) pre-school children, aged 12–59 months inclusive, were recruited from a database of children compiled by 'eumom' (an Irish parenting resource; www.eumom.ie) or from randomly selected childcare facilities in selected locations. Although this facilitated a representative sample of the population of the Republic of Ireland with regard to age, sex and residential location, the sample contained a higher proportion of children of professional workers and

* Corresponding author: J. Walton, email janette.walton@ucc.ie

a lower proportion of children of skilled manual workers than the general population⁽¹²⁾.

Children and their families were visited in their own home by a research nutritionist; 4-d weighed food records were used to collect food intake data. Training in completion of the food record was given to the primary caregivers, and a brief set of instructions on a laminated sheet was provided to accompany the child and food record when in the care of others. The food records benefited from a large amount of researcher–participant interaction, allowing for detailed training of the participants and clarification of recorded data where necessary. All participants completed the records over a continuous 4-d period, including at least 1 weekend day. Food intake data were converted to nutrient intakes using UK and Irish food composition data^(13,14).

This study was conducted according to the guidelines laid down in the Declaration of Helsinki, and ethics approval was obtained from the Clinical Research Ethics Committee of the Cork Teaching Hospitals, University College Cork. Written informed consent was obtained from parents/guardians. A detailed description of the survey methodology is available at www.iuna.net.

Fruit and vegetable intake

In all, 1652 different food codes were consumed on the NPNS, of these 740 had a fruit and/or vegetable component. These included discrete fruits and vegetables and fruits and vegetables contained in composite foods and dishes. The percentage of fruits and/or vegetables in each of these food codes was estimated using standard recipes from the UK food composition database⁽¹⁴⁾, as well as from participants' food diaries and manufacturers' product information. Calculations were included to remove inedible or unconsumed portions – for example, cores and uneaten peel. The effects of concentration, for example, in the case of tomato purée, were also accounted for.

Vegetables included the edible parts of plants commonly consumed as vegetables; foods used as vegetables, such as green pulses and sprouts and fresh sweet corn; botanical fruits used as vegetables, such as tomatoes, peppers or cucumbers; and mushrooms and seaweed. Cereals, potatoes and other tubers, as well as dry pulses, were not considered as vegetables. Fruits included the edible part of all fruits that were fresh, canned, frozen and dried, unless they were classified as vegetables. Fruits with high energy content, such as avocados and olives, were included as fruit. In calculating the fruit and/or vegetable proportion of composite foods and dishes, all dishes/foods with a fruit and/or vegetable component regardless of nutritional profile of the dish/food were included.

Statistical analysis

Intakes (g/d) and sources (g/d) were described for the total population and for consumers only using means, medians, standard deviations and percentage of consumers. Intakes from major sources were stratified by age, and the association with age was assessed using ANOVA. Associations were considered significant at $P < 0.05$.

Intakes and sources were further disaggregated by age and are presented for the total population and for consumers only (online Supplementary Table S1 and S2).

The contribution of fruit and vegetable intake to the weight of the total diet (total weight of all food and beverages in the diet) was estimated separately as the contribution of all sources (g), discrete vegetables (g), discrete fruits (g), vegetables in composite foods and dishes (g) and fruits in composite foods and dishes (g) to the total weight of the diet (g). These were expressed as percentages and reported for the total population and by age.

We estimated the proportion of children with contributions of fruits and vegetables to the weight of the total diet at four levels: <10 , ≥ 10 and <20 , ≥ 20 and <30 and >30 %. As the weight of fruit juice contributes disproportionately to the weight of total fruits and vegetables, the proportion of children in each category of percentage contribution was calculated both including fruits and vegetables from all sources and limiting the contribution of 100% fruit juice to total fruit and vegetable intake to 1 portion/d (50–120 ml)⁽¹⁵⁾. As food and beverage intake data were weighed and subsequently reported in grams, a 1:1 conversion of millilitres to grams was assumed.

Using age-appropriate portion sizes for individual fresh fruits and vegetables, dried fruits and fruit juices (derived from medians of intakes, mostly weighed) (range 14–124 g)⁽¹⁵⁾, we estimated the means and standard deviations of daily intake of portions of fruits and vegetables from all sources of fruits and vegetables, and separately limiting the contribution of 100% juice to 1 portion. We reported these for the total population and by age.

The contribution fruit and vegetable intakes made to the intake of key nutrients was estimated as mean daily intake and percentage contribution of total dietary intake for total sugar, non-milk sugars (NMS), dietary fibre, K, folate, vitamin C and carotene.

Results

The daily intakes of fruits and vegetables in the total population and in consumers only, from all sources and disaggregated by source, are displayed in Table 1. All pre-school children consumed fruits and vegetables during the four survey days. Mean daily intake of fruits and vegetables from all sources was 247 (SD 124) g/d. There were more consumers of fruits (98%) than of vegetables (90%). The largest proportion (70%) of total fruit and vegetable intake (247 g/d) came from discrete fruit intake (171 g/d).

There were no differences in intakes by sex ($P = 0.303$). Intakes of total fruit were positively associated with age ($P < 0.001$), but there was no significant association between vegetable intake and age ($P = 0.184$) (Table 2). Higher intakes of total fruit in older children were largely explained by higher intakes of 100% fruit juice, smoothies and purées (Table 2). This was driven by a 3-fold higher intake of 100% fruit juice in those aged 4 years (77 (SD 99) g/d) *v.* those aged 1 year (23 (SD 50) g/d) (online Supplementary Table S1).

The sources of total fruit intake were discrete fruit excluding 100% fruit juice, smoothies and purees (48–56% contribution

Table 1. Daily intake (g/d) of fruits and vegetables in Irish pre-school children aged 1–4 years (*n* 500) (Mean values, medians, percentages and standard deviations)

	Total population			%	Consumers only		
	Mean	Median	SD		Mean	Median	SD
All sources	247	231	124	100	247	231	124
Discrete vegetables	32	24	30	89	35	28	30
Peas, bean and lentils	8	1	13	52	16	11	14
Onions, peppers, squashes and other vegetables	7	0	12	50	14	10	14
Carrots	7	0	11	50	14	10	13
Green vegetables (including green beans)	5	0	10	39	13	9	12
Salad vegetables	2	0	7	19	13	11	10
Tinned and jarred vegetables	1	0	4	10	11	8	9
Sweet potatoes	1	0	6	3	29	22	25
Fresh herbs	0	0	0	2	1	1	1
Discrete fruit	171	147	117	98	175	150	115
Fruit juices (100% juice)	51	8	78	52	98	71	84
Bananas	30	24	30	71	42	37	28
Fruit purees and smoothies (100% fruit)	22	0	44	31	69	50	53
Apples	21	12	28	62	34	26	30
Citrus fruits	11	0	22	36	31	23	29
Berries	11	0	22	37	29	21	28
Grapes	10	0	18	38	25	19	21
Kiwi, melons, pineapples, plums and other fruit	8	0	20	25	31	21	31
Pears	5	0	14	19	26	20	20
Dried fruit	3	0	6	32	10	8	7
Tinned fruit	1	0	3	6	11	11	8
Vegetables in composite foods and dishes	26	18	25	90	29	22	24
Meat and meat products/dishes	13	7	17	67	19	14	18
Soups, sauces and miscellaneous foods	5	0	12	32	16	11	18
Vegetable dishes	3	0	10	20	17	13	17
Grains, rice, pasta and savouries	3	0	7	36	9	5	9
Fish and fish dishes	1	0	4	8	10	8	8
Breads and rolls	0	0	1	7	3	2	4
Eggs and egg dishes	0	0	1	1	8	7	5
Potato dishes	0	0	0	2	2	1	2
Fruits in composite foods and dishes	19	9	37	98	19	9	37
Beverages	11	2	36	64	17	5	44
Fruited yogurt	4	3	4	9	4	3	4
Confectionery and preserves	1	0	3	50	3	2	4
Fruit dishes	1	0	8	3	37	29	37
Ice-creams and chilled desserts	1	0	3	9	7	5	7
Biscuits, cakes and pastries	1	0	2	21	3	1	3
Breakfast cereals	0	0	2	13	3	1	5
Sauces	0	0	1	1	7	6	8

to total fruit, range of contribution across age) of which bananas (14–19%) and apples (7–13%) contributed the most; 100% fruit juice, smoothies and purees (32–45%); and fruits in composite foods and dishes (7–13%) of which beverages contributed the most (48–63%). Total vegetable intake comprised of discrete vegetables (48–62% contribution to total vegetables, range of contribution across age) of which peas, beans and lentils (11–16%) and carrots (9–15%) contributed the most and vegetables in composite foods and dishes (38–52%) of which meat-based dishes (15–30%) contributed the most (Table 2 and further disaggregation in the online Supplementary Table S1).

Trends observed for intakes and sources by age in the total population were similar when examined in consumers only (online Supplementary Table S2).

Fruits and vegetables contributed to 20% (fruit: 15%, vegetables: 5%) of the weight of the total diet (Fig. 1). This did not vary significantly with age. The ratio of the contribution of discrete fruits and vegetables:fruits and vegetables in composite

foods and dishes varied with age, with older children consuming more discrete fruits and vegetables (Fig. 1).

Of the 500 children, sixty-one had fruit and vegetable intakes that contributed <10% of the weight of the total diet, 224 had intakes that contributed ≥10 and <20%, 148 had intakes that contributed ≥20 and <30% and sixty-seven had intakes that contributed ≥30%. Limiting the contribution of 100% fruit juice to total fruit and vegetable intake to 1 portion/d most affected the number of children categorised as having contributions ≥30%. After applying the limitation, sixty-four children had fruit and vegetable intakes that contributed <10% of the weight of the total diet, 243 had intakes that contributed ≥10 and <20%, 151 had intakes that contributed ≥20 and <30% and forty-two had intakes that contributed ≥30%.

The mean daily intake in the total population was 4.7 (SD 2.2) portions of fruits and vegetables and 4.5 (SD 2.0) portions when the contribution of juice was limited to 1 portion/d. Mean daily intake of portions differed by age ($P \leq 0.001$ when including all juice and when limiting juice). Children aged 1 year consumed



Table 2. Daily intakes and sources of fruits and vegetables in Irish pre-school children aged 1–4 years by age (n 500) (MDI and percentage contribution)

	Age 1 year (n 126)		Age 2 years (n 124)		Age 3 years (n 126)		Age 4 years (n 124)		P*
	MDI (g/d)	% contribution	MDI (g/d)	% contribution	MDI (g/d)	% contribution	MDI (g/d)	% contribution	
Total vegetables		63		53		53		61	0.184
Discrete vegetables	30	48	28	53	31	58	38	62	0.065
Vegetables in composite foods and dishes	33	52	25	47	22	42	23	38	0.001
Total fruit		151		185		209		213	0.000
Discrete fruit	84	56	102	55	105	50	102	48	0.000
100% fruit juice, smoothie, purée	48	32	61	33	86	41	96	45	0.002
Fruit in composite foods and dishes	19	12	22	12	18	9	15	7	0.504

* As calculated using ANOVA.

the most portions per day (mean 5.5 (SD 2.6) when including all juice and 5.3 (SD 2.3) when limiting juice) and children aged 4 years consumed the least (mean 4.3 (SD 1.8) when including all juice and 4.1 (SD 1.7) when limiting juice).

Fruit and vegetable intakes contributed to 32% (24 g/d) of total sugar, 42% (24 g/d) of NMS, 34% (4.1 g/d) of dietary fibre, 25% (449 mg/d) of K, 18% (30 µg/d) of folate, 50% (43 mg/d) of vitamin C and 53% (1225 µg/d) of carotene daily intakes (Table 3).

Discussion


Summary of findings

In this nationally representative survey of pre-school children aged 1–4 years, fruits and vegetables were important foods, contributing 20% of the weight of the total diet and were consumed by all children during the 4 d surveyed. Discrete fruit intake was the largest contributor to total fruit and vegetable intakes. Total fruit intake was higher in older children, largely driven by higher intakes of 100% fruit juice. Vegetable intakes contributed to only 5% of the total weight of the diet, and 10% of children consumed no vegetables during the four survey days. Composite foods and dishes were important sources of vegetables particularly in younger children in whom they contributed to 52% of total vegetable intake. Mean portion intake was 4.7 portions/d. In all, 12% of children had what could be considered a very low (<10%) contribution of fruit and vegetable intake to the weight of the total diet. Fruits and vegetables are an important source of nutrients, particularly vitamin C (50% of mean daily intake), carotene (53%) and dietary fibre (34%). They also contributed to 42% of NMS daily intakes.

Results in context

To compare our findings directly with those from other studies is difficult as we have included fruits and vegetables in composite dishes, excluded inedible portions and included calculations for concentration and cooking losses. These are methodologies not routinely included in studies as they require a detailed level of data collection that is often not available. Some general comparisons can be made with surveys from the UK and USA.

In the UK National Diet and Nutrition Survey, mean intakes in 1.5–3-year-olds were 106 g/d for fruit, 114 g/d for fruit juice, 22 g/d for salad and raw vegetables and 48 g/d for cooked vegetables including those from composite dishes, totalling a crude mean of 290 g/d⁽¹⁶⁾. This is similar to the mean intakes of total fruits and vegetables in the present study of 247 g/d. The Feeding Infants and Toddlers Study (FITS) reported that a substantial proportion of US children do not consume any fruits or vegetables in a given day⁽¹⁷⁾. In contrast, all children in the present study consumed fruit and/or vegetables. This difference may be explained by methodological difference as FITS is based on one 24-h recall and only included discrete fruit and vegetable intakes.



Disaggregated by age

	Age 1 year (n 126)	Age 2 years (n 124)	Age 3 years (n 126)	Age 4 years (n 124)
	% of total diet			
Total fruit and vegetables	17.8	19.1	21.4	21.3
Discrete vegetables	2.5	2.2	2.5	2.9
Discrete fruit	11.0	13.1	15.5	15.5
Vegetables in composite foods and dishes	2.8	2.0	1.8	1.8
Fruit in composite foods and dishes	1.5	1.8	1.6	1.1
Remainder of diet	82.2	80.9	78.6	78.7

Fig. 1. Percentage contribution of fruit and vegetable intake to the weight of the total diet in the total population (n 500).

Contribution of composite foods and dishes

The importance of composite foods and dishes in the estimation of fruit and vegetable intakes has been highlighted previously, although in adults^(18,19). Our study has shown that inclusion of the contribution of composite foods and dishes is particularly important for estimating vegetable intakes in this age group, as 38–52% (range across age) of their vegetable intake was from this source. The contribution of composite foods and dishes should be included in estimating intakes for other analyses. If it is not feasible to disaggregate composite food codes, the percentages reported in this study could be applied. Without accounting for this important source, intakes may be underestimated by as much as 52% in this age group. Thus, composite foods and dishes should also be considered if generating guidelines for consumption.

Fruit and vegetable dietary patterns and guidelines

It is difficult to interpret what contribution to diet fruit and vegetable intake should make in very young children. At present, Irish food-based dietary guidelines pertaining to fruit and vegetable intake are for those aged 5 years and above^(10,11). Other countries similarly have a gap between breast-/bottle-feeding and weaning advice for infants and the availability of food-based dietary guidelines. For example, the *eatwell* plate in the UK is also aimed at those over the age of 5 years⁽²⁰⁾. Some countries have established guidelines, including Australia, who recommend two to three servings of vegetables and legumes and 0.5 servings of fruits for boys and girls aged 1–2 years and 2.5 and one servings, respectively, for 2–3-year-olds⁽²¹⁾.

To inform whether the development of age-specific fruit and vegetable guidelines or public health interventions for fruit and vegetable intake in pre-school children is necessary, we carried out a number of observational analyses. Areas we noted for improvement included the contribution of fruit juice to total

fruit intake, the proportionally lower vegetable intakes than fruit intakes, the very low intakes of fruits and vegetables (<10% weight) and 0 g/d intakes of vegetables in a sub-group of the population.

Pre-school age is an important time for establishing good dietary behaviours to carry forward into later life⁽²²⁾. Previous evaluation of discrete fruits and vegetables in Irish school-age children has shown that pre-school children have a higher intake and less reliance on fruit juice than older children and teenagers⁽²³⁾. Our more detailed observations compounded this finding and also provided further insight as to the evolution of dietary patterns in this age group. The greater dependence on fruit juice as a source of fruits with increasing age noted for older children was already evident in pre-school children. In pre-school children, the total fruit and vegetable intake excluding fruit juice increased with age as did the percentage of the total diet comprised of fruits and vegetables. This suggests that the decline in intake of fruits and vegetables coincides with starting school, marking this as a point for targeted intervention. The mean intake of the pre-school population was 4.7 portions/d. This is encouraging as currently the Irish child and teenager⁽²³⁾ as well as the adult⁽²⁴⁾ populations need to double their intakes to meet the 5-a-day fruit and vegetable intake recommendations⁽¹⁾.

Fruit juice

There is mixed evidence for the nutritional benefits of consuming fruit juice. In this nationally representative survey of very young children, 100% fruit juice was a large contributor to vitamin C intakes, second only to discrete fruit intakes (data not shown). Fruit juice has also been highlighted as the largest contributor to vitamin C intake in older UK children diets⁽¹⁶⁾. However, fruit juice is also a source of free sugar, and limiting the consumption of free sugars is recommended for optimum health^(25,26). The benefits of replacing fruit juice with whole fruit

Table 3. Contribution of fruit and vegetable intake to daily intake of key nutrients in Irish pre-school children aged 1–4 years by age (*n* 500)
(Mean values and standard deviations)

	MDI		% contribution to total dietary intake	
	Mean	SD	Mean	SD
Total sugars (g/d)	24.0	13.9	31.5	14.8
Non-milk sugars (g/d)	24.0	13.8	42.4	17.7
Dietary fibre (g/d)	4.1	2.4	33.9	14.6
K (mg/d)	449	243	25	12
Folate (µg/d)	30	23	18	12
Vitamin C (mg/d)	43	35	50	26
Carotene (µg/d)	1225	1668	53	33

in terms of increasing fibre intake and decreasing energy intake have also been shown using data from children in the USA⁽²⁷⁾.

The association of fruit juice intake with health outcomes is also as of yet inconclusive. For example, higher intake of 100% fruit juice has been associated with higher risk of incident type 2 diabetes⁽⁵⁾ but a null association has also been reported⁽²⁸⁾. The associations of fruit juice intake and the risk of the metabolic syndrome, obesity⁽²⁹⁾ and blood pressure⁽³⁰⁾ have also been investigated, but there is currently insufficient evidence from which to draw conclusions.

In all, 100% fruit juice was a large contributor to total fruit intakes in this age group and its contribution increased substantially with age. However, even in the oldest children, the mean intake was 77 g/d. Intakes at this level are unlikely to negatively impact on health outcomes, although the rapid increase in consumption of 100% fruit juice with age may be of concern. Until a definitive conclusion is made as to the role of fruit juice in the diet, communication of the nutritional benefits of whole fruit instead of fruit juice consumption in this age group would be pragmatic.

Reporting bias

The data are self-reported, and are thus susceptible to reporting bias. Under-reporting food and energy intakes is of particular concern in this context as there are noted implications of under-reporting for the development of food-based dietary guidelines⁽³¹⁾. However, as the focus in these analyses is fruit and vegetable intakes, perceived as ‘healthy’ foods, the social biases associated with food reporting are more likely to bias towards over-reporting not under-reporting. The high level of researcher–participant interaction may have eliminated much misreporting due to forgetting foods, poor or incorrect descriptions of foods and through encouraging compliance. The use of weighed records likely reduced inaccurate estimation of portion sizes. However, researcher–participant interaction and weighing increases participant burden, which can introduce its own biases.

Strengths and limitations

The main strengths of this study are the national representativeness of the sample, the detailed prospective dietary intake data and the comprehensiveness of the estimation of fruit

and vegetable intakes. A limitation of this study is that we used the weight of all food and beverages for the weight of the total diet. This may have led to children with high intakes of liquid-like foods and beverages having lower estimated percentage contribution of fruits and vegetables to the total diet. However, as there was no difference in the weight of the total diet across the four groups of level of contribution of fruits and vegetables, this was considered a justified approach.

Conclusions

Fruits and vegetables are important components of the diet of Irish pre-school children. They were eaten by all children, mean intakes were >4 portions/d, they contributed 20% of the weight of the total pre-school diet and were important sources of dietary fibre, carotene and vitamin C. Some aspects of fruit and vegetable intakes in pre-school children could be improved and would benefit from targeted public health interventions. These include low vegetable intakes overall, increasing dependence on fruit juice with age and very low intakes of both fruits and vegetables in a sub-group of the population.

Acknowledgements

The authors thank the study participants and their families for contributing their time to the survey. The authors also thank the field research staff for collecting the data.

The project was funded by the Irish Government, Department of Agriculture, Food and the Marine under the ‘Food for Health Research Initiative’ 2007–2012.

L. O’C. conceived the research questions, generated the fruit and vegetable intake data, analysed the data and wrote the manuscript. J. W. contributed to the contents and writing of the manuscript and is the study coordinator. A. F. gave critical input to the manuscript and is a principal investigator of the National Pre-School Nutrition Survey. All the authors approved the final version of the manuscript.

The authors have no conflicts of interest or any additional funding to declare.

Supplementary material

For supplementary material/s referred to in this article, please visit <http://dx.doi.org/10.1017/S0007114516001422>

References

1. World Health Organization/Food and Agriculture Organization (2003) *Diet, Nutrition and the Prevention of Chronic Disease: Report of Joint WHO/FAO Expert Consultation. World Health Organisation Technical Report Series*. Geneva: WHO/FAO.
2. Mytton OT, Nnoaham K, Eyles H, *et al.* (2014) Systematic review and meta-analysis of the effect of increased vegetable and fruit consumption on body weight and energy intake. *BMC Public Health* **14**, 886.
3. Cooper AJ, Sharp SJ, Lentjes MA, *et al.* (2012) A prospective study of the association between quantity and variety of fruit and vegetable intake and incident type 2 diabetes. *Diabetes Care* **35**, 1293–1300.
4. Xi B, Li S, Liu Z, *et al.* (2014) Intake of fruit juice and incidence of type 2 diabetes: a systematic review and meta-analysis. *PLOS ONE* **9**, e93471.
5. Muraki I, Imamura F, Manson JE, *et al.* (2013) Fruit consumption and risk of type 2 diabetes: results from three prospective longitudinal cohort studies. *BMJ* **347**, f5001.
6. Pan A, Malik VS, Hao T, *et al.* (2013) Changes in water and beverage intake and long-term weight changes: results from three prospective cohort studies. *Int J Obes (Lond)* **37**, 1378–1385.
7. Te Morenga L, Mallard S & Mann J (2013) Dietary sugars and body weight: systematic review and meta-analyses of randomised controlled trials and cohort studies. *BMJ* **346**, e7492.
8. Te Morenga LA, Howatson AJ, Jones RM, *et al.* (2014) Dietary sugars and cardiometabolic risk: systematic review and meta-analyses of randomized controlled trials of the effects on blood pressure and lipids. *Am J Clin Nutr* **100**, 65–79.
9. Infant and Toddler Forum (2016) Healthy Eating for Toddlers. www.infantandtoddlerforum.org (accessed March 2016).
10. Flynn MA, O'Brien CM, Faulkner G, *et al.* (2012) Revision of food-based dietary guidelines for Ireland, phase 1: evaluation of Ireland's food guide. *Public Health Nutr* **15**, 518–526.
11. Flynn MA, O'Brien CM, Ross V, *et al.* (2012) Revision of food-based dietary guidelines for Ireland, phase 2: recommendations for healthy eating and affordability. *Public Health Nutr* **15**, 527–537.
12. Central Statistics Office (2007) *Census 2006 Principal Demographic Results*. Dublin: Central Statistics Office.
13. Black IJ, Ireland J, Moller A, *et al.* (2011) Development of an on-line Irish food composition database for nutrients. *J Food Compos Anal* **24**, 1017–1023.
14. Food Standards Agency (2002) *McCance and Widdowson's The Composition of Foods*, 6th ed. Cambridge: Food Standards Agency, Royal Society of Chemistry.
15. Lyons J, Walton J & Flynn A (2013) Development of an online database of typical food portion sizes in Irish population groups. *J Nutr Sci* **2**, e25.
16. Public Health England (2014) National Diet and Nutrition Survey. Results from years 1, 2, 3 and 4 (combined) of the Rolling Programme (2008/2009–2011/2012). <https://www.gov.uk/government/statistics/national-diet-and-nutrition-survey-results-from-years-1-to-4-combined-of-the-rolling-programme-for-2008-and-2009-to-2011-and-2012> (accessed April 2016).
17. Siega-Riz AM, Deming DM, Reidy KC, *et al.* (2010) Food consumption patterns of infants and toddlers: where are we now? *J Am Diet Assoc* **110**, S38–S51.
18. O'Brien MM, Kiely M, Galvin M, *et al.* (2003) The importance of composite foods for estimates of vegetable and fruit intakes. *Public Health Nutr* **6**, 711–726.
19. Fitt E, Mak TN, Stephen AM, *et al.* (2010) Disaggregating composite food codes in the UK National Diet and Nutrition Survey food composition databank. *Eur J Clin Nutr* **64**, Suppl. 3, S32–S36.
20. Food Standards Agency (2007) FSA nutrient and food based guidelines for UK institutions. <http://www.food.gov.uk/sites/default/files/multimedia/pdfs/nutrientinstitution.pdf> (accessed April 2016).
21. Nutrition Australia (2013) Australian Dietary Guidelines: Recommended daily intakes. <http://www.nutritionaustralia.org/national/resource/australian-dietary-guidelines-recommended-daily-intakes#toddlers> (accessed March 2016).
22. Kudlova E & Schneidrova D (2012) Dietary patterns and their changes in early childhood. *Cent Eur J Public Health* **20**, 126–134.
23. Walton J, McNulty BA, Nugent AP, *et al.* (2014) Diet, lifestyle and body weight in Irish children: findings from Irish Universities Nutrition Alliance national surveys. *Proc Nutr Soc* **73**, 190–200.
24. Irish Universities Nutrition Alliance (2011) *National Adult Nutrition Survey Summary Report*. Dublin: IUNA. <http://www.iuna.net/wp-content/uploads/2010/12/National-Adult-Nutrition-Survey-Summary-Report-March-2011.pdf> (accessed April 2016).
25. World Health Organization (2014) Draft guideline: sugars intake for adults and children. http://www.who.int/nutrition/sugars_public_consultation/en/ (accessed April 2016).
26. The Scientific Advisory Committee on Nutrition (2014) Scientific consultation: draft SACN carbohydrates and health report – June 2014. http://www.sacn.gov.uk/reports_position_statements/reports/scientific_consultation_draft_sacn_carbohydrates_and_health_report_-_june_2014.html (accessed August 2014).
27. Monsivais P & Rehm CD (2012) Potential nutritional and economic effects of replacing juice with fruit in the diets of children in the United States. *Arch Pediatr Adolesc Med* **166**, 459–464.
28. O'Connor L, Imamura F, Lentjes MA, *et al.* (2015) Prospective associations and population impact of sweet beverage intake and type 2 diabetes, and effects of substitutions with alternative beverages. *Diabetologia* **58**, 1474–1483.
29. Pereira MA & Fulgoni VL 3rd (2010) Consumption of 100% fruit juice and risk of obesity and metabolic syndrome: findings from the national health and nutrition examination survey 1999–2004. *J Am Coll Nutr* **29**, 625–629.
30. Oude Griep LM, Stamler J, Chan Q, *et al.* (2013) Association of raw fruit and fruit juice consumption with blood pressure: the INTERMAP Study. *Am J Clin Nutr* **97**, 1083–1091.
31. Becker W & Welten D (2001) Under-reporting in dietary surveys – implications for development of food-based dietary guidelines. *Public Health Nutr* **4**, 683–687.