

# Invited review: Tannins as a potential alternative to antibiotics to prevent coliform diarrhea in weaned pigs

M. Girard<sup>†</sup> and G. Bee

Swine Research Unit, Agroscope, 1725 Posieux, Switzerland

(Received 24 August 2018; Accepted 13 August 2019; First published online 1 October 2019)

---

*In addition to a multifactorial etiology of nutritional, social and environmental stressors, post-weaning diarrhea (PWD) in pigs is often related to infection with specific pathogens such as enterotoxigenic Escherichia coli (ETEC). In swine farming operations, the incidence of PWD is a global concern and is associated with an unbalanced gut status, resulting in poor performance and high antimicrobial consumption via prophylaxis and metaphylaxis. Increases in antimicrobial resistance are reinforcing an already-urgent need for sustainable, alternative solutions for maintaining optimal gut health in livestock. Tannin-rich plants and extracts contain bioactive compounds that could be of great interest in this respect. This review describes how the use of tannins around weaning could be beneficial for pigs, with special emphasis on the reduction of ETEC-related PWD. An overview of the broad chemical diversity of tannins is presented together with their physicochemical and biological properties, as well as how they may be metabolized in the digestive tract. The pharmacological effects exerted by tannins are summarized; more precisely, the possible mechanisms by which tannins can disrupt the different steps of the pathogenesis of ETEC-related PWD are highlighted. The factors affecting the bioactivity of tannins are also discussed, shedding light on the importance of chemical structure among different tannins.*

---

**Keywords:** bioactive compounds, *Escherichia coli*, piglet, nutrition, weaning

## Implications

Antimicrobials are widely used worldwide in swine production to prevent the occurrence of post-weaning diarrhea. The present review explores the wide-ranging effects of tannins as a promising alternative to antimicrobials. The data presented herein could help mitigate post-weaning diarrhea problems for pig producers, including those in organic farming operations. For researchers, this review highlights potential areas for further investigation.

## Introduction

Early and abrupt weaning as practiced in pig farms is a critical period of dietary transition and environmental and social upheaval for piglets, and is associated with numerous physiological, immunological and microbiological changes to the gastrointestinal tract. During this stressful period, piglets are more susceptible to infection by enteropathogens, the most widespread being enterotoxigenic *Escherichia coli* (ETEC). The combination of gastrointestinal disturbances occurring around

weaning often leads to post-weaning diarrhea (PWD). Across Europe, ETEC is detected in 60% of PWD-affected farms (Luppi *et al.*, 2016). The number seems to be lower in Switzerland, with 37% of pig husbandry operations reporting problems with PWD (Hartmann, 2016). The worsened productive performances, that is, the high morbidity and the depressed growth rate, as well as the increased medical costs related to PWD, result in substantial economic losses to farmers (Fairbrother *et al.*, 2005). Excluding treatment costs, Sjölund *et al.* (2014) estimated annual costs for low-grade PWD of 40 euros per sow.

ETEC colonizes the gastrointestinal tract by binding to the apical side of enterocytes via F4 (or K88) or F18 fimbrial adhesion to receptors; once attached, the bacteria secrete the enterotoxins heat-labile toxin (LT) and heat-stable toxins a and b (STa and STb). These toxins trigger an intracellular signaling cascade resulting in the hypersecretion of electrolytes and water into the intestinal lumen, leading to watery diarrhea (Heo *et al.*, 2013). Lekagul *et al.* (2019) reported that the prophylactic use of in-feed antimicrobials, although strongly discouraged, remains a common practice for coping with PWD in some countries.

---

<sup>†</sup> E-mail: [marion.girard@agroscope.admin.ch](mailto:marion.girard@agroscope.admin.ch)

The extensive use of in-feed antimicrobials has been associated with an increasing incidence of antimicrobial resistance, with 70% of the pathogenic *E. coli* isolated from Swiss pig farms displaying resistance to at least one antimicrobial (Brand *et al.*, 2017). Across Europe, the situation is similar. In 2017, 61% of indicator *E. coli* from the fecal content of fattening pigs exhibited resistance to one or more antimicrobials, with some countries in southern Europe (e.g. Spain, Italy and Portugal) having >90% of resistant isolates (European Food Safety Authority and European Centre for Disease Prevention Control, 2019). This increasing incidence of resistances gives rise to animal and public health concerns and consequently increases the political pressure to reduce antimicrobial use in pig production and generally in livestock. Although considerable efforts have been achieved in recent years to reduce antimicrobial use in Europe, solutions are still needed (European Medicines Agency, 2018). The development of alternative approaches that could contribute to optimal gut health is a prerequisite to minimize the prophylactic use of antimicrobials and could ultimately lead to the improved productivity, health and welfare of pigs, as well as ensure the long-term sustainability of pig production. The two recent decisions of the European Union (EU) to ban within the next 5 years the use of zinc oxide, added at medicinal levels in piglet feed, and the prophylactic use of antimicrobials to prevent PWD, emphasize the urgent need for alternatives. Recent, well-supported research delineates many PWD management strategies, including genetic selection for ETEC-resistant piglets devoid of F4-receptors, vaccination and nutritional approaches (Fairbrother *et al.*, 2005; Heo *et al.*, 2013). Nutritional approaches, such as the management of protein sources and levels in the diet and the use of feed additives (prebiotics, probiotics and organic acids), seem to be the most cost-effective solutions currently used, although the effectiveness of some of these strategies remains somewhat doubtful (Lauridsen *et al.*, 2017). For several years, a growing interest in the use of bioactive compounds from plants as alternatives to antibiotics has emerged, in concurrence with a drive toward sustainable production strategies and consumer acceptability. Bioactive compounds are secondary plant metabolites that elicit pharmacological or toxicological effects on organisms (Bernhoft, 2010). Polyphenols are one of the largest groups of bioactive compounds in the plant kingdom, with more than 8000 known phenolic structures. Among polyphenols, tannins are widely studied for their antiviral, insecticidal, nematocidal, antifungal, antibacterial and antioxidant properties and might therefore represent one solution to tackle PWD (Barbehenn and Constabel, 2011; Hoste *et al.*, 2015; Smeriglio *et al.*, 2017).

This review aims to provide an overview of the large diversity of tannins and their possible contributions to the reduction of coliform PWD and improvement in and maintenance of gut health. A definition of tannin chemical structures and the wide range of their effects on bacteria, with a special emphasis on ETEC-related PWD, is explored from existing *in vitro* and *in vivo* studies. The factors affecting the bioactivity of tannins are also discussed.

## Tannins, a diverse family of bioactive compounds

### *Definition, localization and roles in plants*

Tannins are a heterogeneous group of astringent polyphenolic biomolecules that can interact with and precipitate macromolecules, such as proteins, gelatins, polysaccharides and alkaloids. Plants synthesize these compounds in mixtures that cover a wide range of molecular weights, and up to 20 000 Da have been reported (Khanbabaee and van Ree, 2001). Commonly, tannins are classified into three categories: condensed tannins (CTs), hydrolyzable tannins (HTs) and complex tannins.

Many tannin compounds are found in flowers, leaves, seeds, fruits, roots and tree bark samples (Sieniawska and Baj, 2017). The CTs appear to be stored inside tannosomes, a chloroplast-derived organelle enclosed within tonoplasts inside plant vacuoles. In these tannosomes, tannins do not interact with proteins and consequently do not interfere with plant metabolism (Brillouet *et al.*, 2013). It has been shown that HTs may be synthesized and deposited in chloroplasts but also in cell walls or the intercellular space (Grundhöfer and Gross, 2001).

The role of CTs is to defend the plant against predation by herbivore and insects primarily by reducing plant palatability, particularly in young leaves (Barbehenn and Constabel, 2011). In roots, CTs are a chemical barrier against penetration and colonization by pathogens; in seeds, they maintain plant dormancy and have bactericidal properties (Constabel *et al.*, 2014).

### *Classifications of tannins*

The most widespread CTs, also known as proanthocyanidins, are oligomers (2 to 10 monomers) or polymers (>10 monomers) of flavan-3-ol units. When subjected to acidic alcohol treatment, CTs degrade to anthocyanidins, the pink-purple pigments responsible for flower coloring (hence the name). Common flavan-3-ols are [epi]catechin, [epi]afzelechin, [epi]gallocatechin, [epi]fisetinidol and [epi]robinetinidol, in accordance with the position of their –OH and –H groups, which give rise to procyanidins, prodelphinidins, profisetinidins and prorobinetinidins (Figure 1). For instance, procyanidins contain catechin or epicatechin, and prodelphinidins contain gallocatechin or epigallocatechin (Figure 1). In plants, a mixture of the aforementioned classes is often present and varies according to growing conditions (growth stage, temperature, etc.) and mode of conservation (fresh, wilted, etc.) (Girard, 2016). The size of the CT molecule is characterized by its degree of polymerization. In addition to differences in their chemical groups, flavan-3-ols differ in their stereoisomerism: catechin and gallocatechin have a *trans* configuration, whereas epicatechin and epigallocatechin have a *cis* configuration at the C2–C3 bond. The interflavan linkages between flavan-3-ols units can be either A-type or B-type (Girard, 2016). A-type linkages comprise C4β–C8 bonds and C2β–O–C7 or C2β–O–C5 ether bonds. B-type linkages consist of C4β–C8 or C4β–C6 linkages (Figure 1).




Figure 1 Condensed tannins consisting of typical flavan-3-ol subunits and two different linkage types.






Figure 2 Chemical structure of: (a) a gallotannin molecule with depsidic bonds and (b) gallic acid.

The two main classes of HTs are the relatively rare gallotannins and the widespread ellagitannins. Simple gallic acid derivatives can be considered a third subclass of HTs (Salminen and Karonen, 2011). Simple gallic acid derivatives and gallotannins contain gallic acid substituents that are esterified with a polyol residue, usually a D-glucose. However, glucitol, fructose, shikimic acid, xylose, hamamelose, saccharose, quercitol or quinic acid have also been identified as polyol residues in certain plants such as maple,

chestnut, oak and witch hazel (Smeriglio *et al.*, 2017). In the first step of biosynthesis, galloylation reactions with 1-O-galloyl- $\beta$ -D-glucose ( $\beta$ -glucogallin) yield di-, tri-, tetra-, and pentagalloylglucose (1,2,3,4,6-penta-O-galloyl- $\beta$ -D-glucopyranose) that correspond to simple gallic acid derivatives (Figure 2). In the second step of galloylation, depsidic binding (*meta*- or *para*-depside) yields, among others, hexa-, hepta-, octa- and -galloylglucose; in fact, up to 12 units of gallic acids can be esterified on a single glucose



**Figure 3** Chemical structure of: (a) an ellagitannin (casuarictin) with two hexahydroxydiphenyl units and (b) ellagic acid.



**Figure 4** Chemical structure of camelliatannin E, a complex tannin.

moiety (Figure 2) (Sieniawska and Baj, 2017; Smeriglio *et al.*, 2017). Hydrolysis of gallotannins with strong acids generates the core polyol and gallic acid. Ellagitannins are produced through gallotannin oxidation by intermolecular carbon-carbon coupling between at least two galloyl units, forming hexahydroxydiphenyl (HHDP; Figure 3a) or its derivatives, such as dehydrohexahydroxydiphenyl units. In aqueous solutions, HHDP spontaneously lactonizes and releases ellagic acid (Figure 3b). Ellagitannins can be in monomeric, dimeric, oligomeric or C-glycosidic forms, and almost 500 different ellagitannin molecules have been isolated and identified to

date (Smeriglio *et al.*, 2017). Complex tannins contain a carbon-carbon link between the central carbohydrate and a flavan-3-ol unit (e.g. epicatechin) plus galloyl and HHDP groups, such as camelliatannins (Figure 4; Hatano *et al.*, 1995).

#### *Chemical and biological properties*

Owing to their hydroxyl and phenolic groups, tannins interact in a number of ways with other molecules as reviewed by Girard (2016). These interactions are based on hydrogen bonding or hydrophobic interactions. They can also interact with molecules to form covalent bonds. Such bonds can occur

under oxidative conditions, at high temperatures, at high pH, in ultraviolet radiation, in autoxidation or in the presence of catalytic enzymes, such as polyphenol oxidase, which produces reactive quinones (Barbehenn and Constabel, 2011; Constabel *et al.*, 2014).

Originally, tannins were used in tanning processes to convert animal hides into leather through the ability of tannins to adhere to proteins or, more precisely, to collagen. One of the main physico-chemical characteristics of tannins is their ability to interact strongly with proteins including enzymes. The maximum interaction between CTs and protein occurs when the solution pH is near the isoelectric pH of the protein; thus, for instance, greater affinities for CTs are reported at pH 4.9 for globular proteins with acidic isoelectric points, whereas at pH 7.8, basic proteins have a higher affinity for CTs (Hagerman and Butler, 1981). The interaction of tannins with salivary proteins and taste receptors, particularly bitter receptors, generates the astringent taste appreciated in wine and beer (Soares *et al.*, 2013). In addition to proteins, various carbohydrates originating from the cell wall, such as pectin, cellulose or dietary fibers, can interact with tannins (Jakobek, 2015). The detailed interactions between lipids and polyphenols (including tannins) have been described in a review of Jakobek (2015). In a previous study, He *et al.* (2006) showed a strong binding affinity between the gallic acid derivative pentagalloylglucose and phospholipids, which are constituents of the cell membranes. The interaction between HTs and phospholipids relies on the hydrophobic association between the galloyl groups of gallotannins and the hydrophobic hydrocarbon chains of phospholipids (He *et al.*, 2006; Yu *et al.*, 2011). Because of the orthohydroxyl groups present on their B-rings (catechol or pyrogallol), tannins can chelate and thereby sequester metal ions, particularly cations such as calcium, iron, magnesium, manganese and copper (Oladoja *et al.*, 2011).

The aforementioned properties confer tannins to several biological applications, and their antiviral, insecticidal, nematocidal, antifungal, antibacterial and antioxidant activities have been well established (Barbehenn and Constabel, 2011; Hoste *et al.*, 2015; Smeriglio *et al.*, 2017). In particular, their potential antioxidant and anti-inflammatory effects on piglets would be interesting to help these animals to better cope with the ETEC-related diarrhea. However, our review only focuses on the effect of tannins on the pathogenic and commensal gut bacteria of piglets.

#### *Tannin absorption and metabolism in the gastro-intestinal tract*

Studies investigating the pharmacokinetics of tannins in animals are scarce. In the stomach, CTs remain relatively stable. The cleavage of CTs into monomers, such as flavan-3-ols, may occur, but it seems to be highly dependent on the solubility of CTs in this gastric juice and on the pH of the gastric juice (Zhang *et al.*, 2016). Flavan-3-ols are often acylated by gallic acid (galloylation) and small amounts could be passively absorbed without being metabolized in the small intestine, along with CT dimers or trimers (Marín *et al.*, 2015). Through the action of sulfotransferases, uridine-5'-diphosphate-glucuronosyl-transferases and catechol-*O*-methyltransferases,

the main metabolites of flavan-3-ols or CT dimers in the plasma are sulfated, glucuronidated and methylated derivatives (Zhang *et al.*, 2016; Mena *et al.*, 2019). However, oligomers larger than trimers and polymers cannot be absorbed via passive paracellular transport in their native form, and most ingested CTs reach the colon intact (Deprez *et al.*, 2001). The colonic microbiota catabolize CTs to produce hydroxyphenolic acids: phenylvaleric, phenylpropionic and phenylacetic acids, as well as phenylvalerolactones and benzoic acids (Marín *et al.*, 2015). A part of these bioactive catabolites can be absorbed by the colonocytes. Subsequently, they are conjugated by specific hepatic enzymes to produce conjugated derivatives that are further eliminated in the urine (Mena *et al.*, 2019).

The polymers of HTs cannot be absorbed in their native form. Several types of bacteria, such as lactic acid bacteria, possess tannase activity (tannin acyl hydrolase), in which they hydrolyze the ester bonds of gallotannins and ellagitannins (although not the carbon–carbon bonds of ellagitannins; Mingshu *et al.*, 2006). Hydrolysis of ester and depside linkages of gallotannins by intestinal enzymes and colonic microbiota yields the core polyol (glucose) and gallic acid. The resultant aglycone (gallic acid) is metabolized to pyrogallol and phloroglucinol and, ultimately, to acetate and butyrate (Krumholz *et al.*, 1987). *Eubacterium oxidoreducans* and *Coprococcus* sp., which are present in the rumen and the distal portion of the monogastric intestine, are involved in this reaction (Krumholz *et al.*, 1987).

Due to their carbon–carbon bonds, ellagitannins are more difficult to degrade than gallotannins. Ellagitannins are resistant to the action of lactase-phlorizin hydrolase and  $\beta$ -glucosidase, present in the small intestine, and consequently are preferentially cleaved by colon microbiota (Marín *et al.*, 2015). Hydrolysis of ellagitannins yields ellagic acid following lactonization. Whether ellagitannins can efficiently release ellagic acid under gastrointestinal physiological conditions without the involvement of specific gut microbiota remains unclear. In pigs fed acorns (rich in ellagitannins), ellagic acid has been shown to gradually be metabolized to urolithins throughout the intestine – to urolithins D and C in the jejunum and urolithins B and A in more distal parts of the intestinal tract (Espín *et al.*, 2007). Owing to the presence of ellagic acid metabolites in the bile and urine, as well as their absence in intestinal tissues, ellagic acid has also been suggested to be possibly absorbed in the stomach (Espín *et al.*, 2007).

#### **Effects of tannins on bacteria**


##### *Pathogenesis of enterotoxigenic Escherichia coli infection*

Diarrhea is a clinical symptom of an imbalance between absorption and secretion in the intestine, resulting in the hypersecretion of chloride and bicarbonate ions together with osmosis-driven water migration into the lumen. The main pathotype causing coliform PWD is ETEC. The pathogenicity of ETEC is due to their adhesins and

enterotoxins, which impair enterocyte function and are primarily responsible for their virulence. After being ingested by ETEC-susceptible piglets, ETEC colonize the gastrointestinal tract by attachment to microvilli or mucus coating the epithelium via their fimbrial adhesins, enabling them to avoid elimination by peristalsis. Fimbriae are long, thin protein appendages present in high numbers (100 to 300 per bacterium) on the bacterial surface. The common types of fimbriae found in ETEC in piglets suffering from PWD are F18 and F4. The F18 fimbriae have two antigenic variants, F18ab and F18ac, whereas F4 fimbriae possess three antigenic variants, F4ab, F4ac and F4ad. These variants differ slightly in amino acid composition; variants F4ab and F4ad are made up of 264 amino acids, whereas variant F4ac is made up of 262 amino acids. Fimbriae primarily attach to specific receptors located on the apical side of enterocytes in the jejunum and ileum, but they may also bind to non-specific receptors in the mucus that coats the epithelium. The F4ad fimbriae are more likely to bind glycolipids, while the F18, F4ab and F4ac fimbriae preferentially bind to glycoproteins (Nagy and Fekete, 2005). Once attached to the epithelium, ETEC replicate and secrete two types of enterotoxins, LT and ST ones, which are extracellular proteins and peptides. The pathogenesis of ETEC is summarized in Figure 5. The LT toxin has a high molecular weight (88 kDa, 343 amino acids), and its structure is similar to that of the *Vibrio cholerae* toxin. The LT toxin consists of six subunits: one A subunit with enzymatic activity and five B subunits (pentamer) involved in the adhesion of LT to the GM1 (monosialotetrahexoxylganglioside) receptor

present on the surface of enterocytes. Once bound, LT is internalized by endocytosis and transported to the Golgi apparatus for disassembly. The A subunit migrates to the endoplasmic reticulum where it is cleaved to A1 (an NAD-dependent ADP-ribosyl transferase) and A2 (peptide) subunits. The activated A1 subunit transfers an ADP-ribosyl from NAD to the regulatory G<sub>s</sub> protein α subunit located in the basolateral membrane. This results in the permanent activation of the adenylyl cyclase cascade and the increased intracellular concentrations of cyclic adenosine monophosphate (cAMP). The accumulation of cAMP activates protein kinase A (PKA), which phosphorylates the cystic fibrosis transmembrane regulator (CFTR) and a Na<sup>+</sup>/H<sup>+</sup> exchanger (Dubreuil *et al.*, 2016). This process leads to the secretion of chloride and carbonate ions and to the inhibition of Na<sup>+</sup> reabsorption. Water is then osmotically drawn into the intestinal lumen, resulting in watery diarrhea (Dubreuil *et al.*, 2016).

Enterotoxigenic *E. coli* isolated from pigs produce two types of ST toxins: STa (18 amino acids and approximately 2 kDa) and STb (48 amino acids and 5.2 kDa). Both toxins remain active even after incubation at 100°C for 30 min. The STa mechanism is similar to that of LT. On the apical side of enterocytes, STa binds to guanylate cyclase-C, which synthesizes cyclic guanosine monophosphate (cGMP) from guanosine triphosphate. The accumulation of cGMP in the enterocytes has two effects: first, it activates a cGMP-dependent protein kinase II (PKII), inducing phosphorylation of the CFTR and secretion of chloride and carbonate ions; second, it inhibits phosphodiesterase 3, resulting in an increase



**Figure 5** Pathogenesis of enterotoxigenic *Escherichia coli*. Signal cascade following toxin production by enterotoxigenic *Escherichia coli*, which results in the activation of ion channels and the disruption of tight junctions leading to electrolyte and water hypersecretion in the lumen. AA = arachidonic acid; AC = adenylate cyclase; ARF = ADP-ribosylation factor; CaCC = calcium-activated chloride channel; CFTR = cystic fibrosis transmembrane regulator; GC-C = guanylate cyclase C; GM1 = monosialotetrahexoxylganglioside; G = protein; Gsα = Gs protein α; LT = heat-labile toxin; NHE3 = Na<sup>+</sup>/H<sup>+</sup> exchanger 3; P = phosphorylation; PDE3 = phosphodiesterase 3; PGE<sub>2</sub> = prostaglandin E<sub>2</sub>; PL A<sub>2</sub>-PL C = phospholipases A<sub>2</sub> and C; PKA = protein kinase A; PKC = protein kinase C; PKII = protein kinase II; ST = heat-stable toxin.

in cAMP and the activation of PKA (Dubreuil *et al.*, 2016). As in the LT signal cascade, CFTR and the Na<sup>+</sup>/H<sup>+</sup> exchanger are phosphorylated, leading to a hypersecretion of electrolytes and water into the lumen.

The signal transduction pathway of STb does not involve cyclic nucleotides as secondary messengers. Research suggests that STb binds to a sulfatide receptor and is thereby internalized (Labrie *et al.*, 2002). The G protein cascade is then activated leading to an increase in the concentration of intracellular calcium, which activates a number of enzymes. The first is calmodulin-dependent PKII, which opens specific chloride channels and activates PKC; this in turn phosphorylates CFTR and inhibits Na<sup>+</sup> uptake. The second and third activated enzymes are phospholipases A2 and C, which catalyze the release of arachidonic acid from membrane phospholipids as well as the formation of prostaglandin E2 and serotonin, known as secretory agents, from enterochromaffin cells (Peterson and Whipp, 1995; Dubreuil *et al.*, 2016). The release of these agents has been suggested to be a potential mechanism of effect for STb on the enteric nervous system (Peterson and Whipp, 1995).

Finally, ETEC has been shown to induce pro-inflammatory responses with the production of cytokines in epithelial intestinal cells (Devriendt *et al.*, 2010). However, the possible effects of tannins on inflammatory responses are not discussed in the present review (Williams *et al.*, 2017).

#### *In vitro tannin inhibition of the growth of some bacteria*

The bactericidal and bacteriostatic effects of CTs and HTs have been well documented (Smeriglio *et al.*, 2017). Table 1 presents tannin-containing plants with antimicrobial properties. Taguri *et al.* (2004) noted that of the 10 polyphenols they tested, the 2 flavan-3-ols, epigallocatechin and epigallocatechin gallate and the 2 tannins, castalagin and prodelphinidins, demonstrated the greatest antimicrobial activity against all strains of *E. coli*, particularly on ETEC strains. The A-type CTs from cranberries possess antibacterial activity against several pathogenic bacteria, including *Helicobacter pylori*, *Salmonella*, *Staphylococcus aureus*, *E. coli*, and *Campylobacter* (Côté *et al.*, 2010). The addition of 10 mg/ml HTs from chestnut wood extract had a slight bacteriostatic effect on the growth of ETEC F4ac, whereas 20 and 40 mg/ml exhibited marked bacteriostatic effects (Pradervand N, personal communication). Jelager *et al.* (1998) observed that the antibacterial properties of tannins from Mauritian medicinal plants were lost when tannins were removed from the extract by precipitation with gelatin prior to testing. This finding confirmed that tannins were responsible for the observed antibacterial effects. Tannins may inhibit bacterial growth by destabilization and permeabilization of the cytoplasmic membrane and by inhibition of extracellular microbial enzymes, either by direct action against microbial metabolism and/or sequestration of the substrates required for microbial growth (Min *et al.*, 2008). For instance, tannins have been shown to chelate mineral macronutrients, such as iron or zinc, that are required for the growth of many bacterial species (Chung *et al.*, 1998). The addition of iron to

the medium restored HT-inhibited bacterial growth (Engels *et al.*, 2009).

#### *Inhibition of bacterial adhesion to the intestinal epithelium and biofilm formation*

Microorganism adhesion to the intestinal epithelium is a prerequisite step for bacterial colonization and biofilm formation. Cranberry juice containing A-type CTs has been reported to inhibit the adhesion of uropathogenic *E. coli*, which are responsible for urinary tract infections. The catabolites of cranberry CTs in the colon, the hydroxy-phenyl- $\gamma$ -valerolactones, have been proposed as plausible candidates to exert anti-adhesive activity in the bladder (Mena *et al.*, 2019). Liu *et al.* (2006) noted that cranberry tannins decreased the adhesion forces between bacterial and epithelial cells and altered the conformation of surface macromolecules on *E. coli*, leading to a 60% decrease in the average P-fimbriae length (from 148 to 53 nm). A few studies focused on the effects of tannins on the intestinal epithelium adhesion of ETEC strains. Coddens *et al.* (2017) found that 10  $\mu$ g of cranberry extract was sufficient to reduce the *in vitro* adhesion of verotoxigenic *E. coli* F18, whereas a greater amount (100  $\mu$ g) was necessary to strongly inhibit ETEC F4 adhesion. These results were confirmed in ligated loop experiments on pigs in which the immunochimistry results clearly demonstrated that pre-incubation of F4 or F18 fimbriae with cranberry extract abolished the binding of fimbriae to the intestinal epithelial brush border. However, the authors demonstrated that this binding inhibition was not caused by an inhibition of bacterial growth. Similarly, Verhelst *et al.* (2010) observed that at a concentration of 150  $\mu$ g/ml, three tannin-rich extracts reduced ETEC F4ac binding to epithelial brush border tissues isolated from pigs. The three tannin-rich extracts contained pentagalloylglucose, a mixture of medium and high molecular weight HTs and procyanidins and flavan-3-ols of cocoa beans, respectively.

Some tannins can inhibit biofilm formation. Bacteria form biofilms following adhesion to a surface. Bacterial biofilm formation is now believed to play an important role in intestinal colonization (Hancock *et al.*, 2010). Owing to its capacity to autoaggregate and form biofilms, a bacterial afimbrial adhesin involved in diffuse adherence (AIDA) associated with some diarrheagenic *E. coli* strains was shown to promote bacteria-to-bacteria adherence. A mutation of the genes coding for AIDA in an AIDA-positive ETEC resulted in an inability to aggregate and induce biofilms in the large intestine as well as to induce diarrhea (Ravi *et al.*, 2007). The biofilm facilitates the exchange of plasmids for horizontal gene transfer, concentrates bacterial enzymes that inactivate antibiotics and decreases the penetration of antimicrobials. Ellagitannins (mainly punicalagin) and ellagic acid from *Punica granatum L.* (pomegranate) extract inhibited the formation of biofilms in *Staphylococcus aureus*, methicillin-resistant *S. aureus* (MRSA), *E. coli* and *Candida albicans* (Bakkiyaraj *et al.*, 2013). In the same experiment, a minimum inhibitory concentration of 250  $\mu$ g/ml of pomegranate extract was necessary to inhibit *E. coli* growth, but only 150  $\mu$ g/ml of the same extract was necessary to fully inhibit biofilm

**Table 1** Tannin-containing plants with antimicrobial properties

Plant source	Plant part	Tannins (CTs or HTs)	Tannin or metabolite type	<i>In vitro</i> and/or <i>in vivo</i>	Bacteria	Reference
Pomegranate ( <i>Punica granatum</i> L.)	Fruit peel	HTs	Ellagitannins (punicalagin) Ellagic acid (metabolite)	<i>In vitro</i>	Food-borne pathogens: <i>Escherichia coli</i> , <i>Listeria monocytogenes</i> , <i>Staphylococcus aureus</i> , <i>Yersinia enterocolitica</i>	Al-Zoreky (2009)
Chestnut ( <i>Castanea sativa</i> )	Wood	HTs	Gallotannins Ellagitannins	<i>In vitro</i>	Poultry pathogens: <i>Campylobacter jejuni</i> , <i>Clostridium perfringens</i> type A, <i>Escherichia coli</i> , <i>Pasteurella multocida</i> , <i>Salmonella enteritidis</i> , <i>Salmonella gallinarum</i> , <i>Salmonella typhimurium</i> , <i>Salmonella virchow</i> , <i>Staphylococcus aureus</i>	Graziani et al. (2006)
Japanese rose ( <i>Rosa rugosa</i> )	Petals	HTs	Ellagitannins (tellimagrandin II, rugosin A and D)	<i>In vitro</i>	Intestinal bacteria: <i>Bacillus cereus</i> , <i>Escherichia coli</i> , <i>Salmonella</i> sp., <i>Staphylococcus aureus</i>	Kamijo et al. (2008)
Chestnut ( <i>Castanea sativa</i> )	Not specified	HTs	Gallotannins Ellagitannins	<i>In vitro</i> / <i>in vivo</i>	<i>Escherichia coli</i> O157:H7 ( <i>in vitro</i> ) and generic fecal <i>Escherichia coli</i> ( <i>in vivo</i> )	Min et al. (2007)
Mimosa ( <i>Acacia mearnsii</i> )	Not specified	CTs	Procyanidins Prodelfinidins Prorobinetinidins			
Sumac ( <i>Rhus copallina</i> )	Leaves	CTs : HTs (17% : 83%)	Gallotannins Ellagitannins	<i>In vitro</i>	<i>Escherichia coli</i> , <i>Klebsiella pneumoniae</i> , <i>Staphylococcus aureus</i>	Min et al. (2008)
Shinnery oak ( <i>Quercus havardii</i> )	Leaves	CTs : HTs (29% : 71%)	Catechin (flavan-3-ol) Gallotannins Ellagitannins			
Quebracho extract ( <i>Schinopsis</i> spp.)	Not specified	CTs : HTs (98.5% : 1.5%)	Profisetinidins Prorobinetinidins			
Japanese chestnut ( <i>Castanea crenata</i> )	Wood	HTs	Castalagin	<i>In vitro</i>	<i>Escherichia coli</i> (non-pathogenic <i>E. coli</i> , enterohemorrhagic <i>E. coli</i> , enteroinvasive <i>E. coli</i> , enterotoxigenic <i>E. coli</i> ), <i>Salmonella</i> , <i>Staphylococcus aureus</i>	Taguri et al. (2004)
Woodland elaeocarpus ( <i>Elaeocarpus sylvestris</i> )	Bark	CTs	Prodelfinidins			
Loquat ( <i>Eriobotrya japonica</i> )	Seeds	CTs	Procyanidins			
Mimosa ( <i>Acacia mearnsii</i> )	Bark	CTs	Procyanidins Prodelfinidins Prorobinetinidins Profisetinidins	<i>In vitro</i>	<i>Bacillus subtilis</i> , <i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i> , <i>Shigella dysenteriae</i> , <i>Staphylococcus aureus</i>	Yao et al. (2006)

CTs = condensed tannins; HTs = hydrolyzable tannins.

formation. Similarly, a concentration of 8 mg/ml of gallic acid, a metabolite of gallotannin catabolism, inhibited *E. coli* biofilm formation at 25°C and 37°C (Shao et al., 2015).

#### Inhibition of enterotoxin production and activities

Some tannins have the ability to inhibit bacterial enterotoxins or interrupt the signal transduction pathway of enterotoxins

by inhibiting ionic channels. Elizondo et al. (2010) showed that CTs from quebracho, containing polymers of profisetinidins and prorobinetinidins, and HTs from chestnut, containing gallotannins and ellagitannins (mainly vesicalagin and castalagin), were able to reduce the  $\alpha$  toxin lecithinase activity and  $\epsilon$  toxin cytotoxicity of *C. perfringens*. Similarly, 0.05% (v/v) of methanol extract from pomegranate extract containing



ellagitannins (mainly punicalagin) and ellagic acid drastically reduced the production of staphylococcal enterotoxin A by *S. aureus* (Braga *et al.*, 2005). Similar results were observed for ETEC strains. Chen *et al.* (2006) showed that Galla Chinensis extract, which is rich in gallotannins and gallic acid, inhibited the binding of the B subunit of LT to its GM1 receptor, which may explain the antidiarrheal properties of this extract. Likewise, 750 µg/ml of two HT extracts, one containing the gallic acid derivative pentagalloylglucose and the other containing a mixture of medium and high molecular weight HTs, clearly reduced the binding affinity of LT for GM1, although the same concentration of a procyanidin CT extract had no inhibitory effect (Verhelst *et al.*, 2010). However, the authors did not mention the tannin content of each extract. Certain specific tannins and flavan-3-ols are inhibitors of the ion channels involved in the development of diarrhea. The flavan-3-ols, epigallocatechin-gallate and epicatechin-gallate, from green tea (*Camellia sinensis*) and tannic acid (a mixture of gallotannins) inhibited calcium-activated Cl<sup>-</sup> channels by up to 50%, thereby reducing intestinal Cl<sup>-</sup> secretion. Epicatechin- or catechin-containing compounds are present in green tea and in grape seeds, but they have no documented effect on the aforementioned channels (Namkung *et al.*, 2010). Schuier *et al.* (2005) demonstrated that procyanidins of cocoa beans partially blocked intestinal CFTR activity in a human colon epithelial cell line, which could explain their antisecretory properties.

#### *Tannin effect on commensal microbiota and coliform diarrhea development*

Several *in vitro* studies have shown that some tannins can have a negative effect on the growth of pathogenic bacteria and no or positive effects on the growth of commensal, non-pathogenic bacteria. For instance, ellagitannins isolated from *Rosa rugosa* petals (tellimagrandins II, rugosin D) showed antibacterial activities against *E. coli*, *S. aureus*, *Bacillus cereus* and *Salmonella* sp., but they had little or no effect against *Bifidobacterium breve* or *Lactobacillus salivarius* (Kamijo *et al.*, 2008). Similarly, tannic acid, some berries rich in CTs (mainly procyanidins) and gallotannins from *Mangifera indica* L. (mango) did not inhibit the growth of probiotic lactic acid bacteria, such as *Bifidobacterium* spp. and *Lactobacillus* spp. (Chung *et al.*, 1998; Puupponen-Pimiä *et al.*, 2001; Engels *et al.*, 2009). The beneficial effects of tannins in reducing the growth of pathogenic bacteria and promoting that of beneficial bacteria have also been reported in *in vivo* trials (Biagi *et al.*, 2010; Brus *et al.*, 2013). Likely owing to their bactericidal and bacteriostatic properties, tannins have been found to modify the microbiota of pigs (Tretola M., personal communication). In a previous experiment, the addition of chestnut wood extract containing HTs to the feed of healthy pigs during fattening resulted in a reduction in the total *E. coli* number, an increase in the total lactic acid bacteria count and a greater average daily gain (ADG) (Brus *et al.*, 2013). Supplementation with chestnut wood extract for 1 month following weaning tended to increase the viable counts of *Lactobacilli* in the jejunum, although the bacterial counts in the cecum were not affected

(Biagi *et al.*, 2010). Tannins were found to modify the microbiota of piglets artificially infected with ETEC. The inclusion of CT extract from grape seeds (10 g/kg in feed; B-type CT) and from cranberries (10 g/kg in feed + 1 g/l in water; A-type CT) in their rations significantly decreased ETEC F4 and verotoxinogenic F18 *E. coli* shedding, respectively, and decreased the incidence of diarrhea (Verhelst *et al.*, 2014; Coddens *et al.*, 2017). Although the severity of diarrhea in infected piglets was reduced upon 10 g/kg feed supplementation with chestnut wood extract rich in HTs, no reduction in ETEC F4 shedding in the feces was observed at 4 days post-infection compared with the infected piglets that did not receive HT supplementation (Girard *et al.*, 2018). Nevertheless, increasing the percentage of chestnut wood extract to 20 g/kg in the weaning diet of F4-susceptible piglets did result in lower ETEC F4 shedding in the feces at 3 days post-infection, a 40 g/day greater ADG and a marked decrease in the number of piglets with diarrhea (Girard M. and Hu D., unpublished observation). In addition, the inclusion of 20 g/kg of chestnut wood extract in the diets of piglets infected or not with ETEC F4 decreased the relative abundance of ETEC F4 and some *Clostridium* spp. in the jejunum without lowering the abundance of *Lactobacillus* spp. (Girard *et al.*, 2019). In the aforementioned studies, the chestnut wood extract contained 45 mg/g gallotannins, 9 mg/g ellagitannins and 3.7 mg/g gallic acid.

#### **Modifying factors of the bioactivity of tannins**

The wide range of effects that tannins exert on pathogenic bacteria and their toxins is based on their ability to interact not only with proteins (including enzymes) but also with carbohydrates, lipids and metal ions. However, the bioactivity of tannins against bacteria varies according to the prevailing physico-chemical conditions and to factors related to bacteria and/or to the tannin molecules themselves.

#### *Factors related to bacteria: type, concentration and time of exposure*

An important factor is the target bacteria species. As previously mentioned, some CTs and HTs can inhibit the growth of some species but do not affect the growth of probiotic lactic acid bacteria (Kamijo *et al.*, 2008). Unlike other bacteria, some probiotic bacteria of the genera *Bifidobacterium* and *Lactobacillus* do not require heme-containing enzymes for metabolism and are not sensitive to the ion chelation induced by tannins. This may be why the growth of probiotic bacteria is not affected by tannins (Marín *et al.*, 2015). In studies in which tannins influenced bacterial growth, researchers also demonstrated that Gram-positive bacteria, such as *S. aureus*, were more susceptible to tannins than Gram-negative bacteria, such as *E. coli*, were (Taguri *et al.*, 2004; Min *et al.*, 2008). The different susceptibilities between Gram-positive and Gram-negative bacteria might be related to their cell wall structures. Gram-positive bacteria have a single lipid bilayer surrounded by a thick porous layer of peptidoglycans, whereas Gram-negative bacteria have an inner lipid bilayer, a thin layer

of peptidoglycans and an outer asymmetrical lipid bilayer on which lipopolysaccharides are anchored (Barer, 2012). Compared with the double lipid bilayer, the single lipid bilayer might therefore be more easily destabilized by some tannins. The effects of tannins are directly associated with the concentration of bacteria and the duration of tannin exposure. Graziani *et al.* (2006) found an inhibitory capacity against  $1.8 \times 10^6$  cfu/ml of *E. coli* using 1.5 g/kg of HTs from chestnut wood extract, but no inhibition was observed against  $1.2 \times 10^9$  cfu/ml of *E. coli* at 1.0 and 2.5 g/kg of this same extract. Elizondo *et al.* (2010) observed a positive correlation between the time of exposure and the bactericidal effect of chestnut HTs against *C. perfringens*. Of note, the effectiveness of tannins in interfering with bacteria can be modified by tannin interaction with other compounds. Verhelst *et al.* (2010) reported that the inclusion of bovine serum albumin (BSA) reversed the inhibitory effect of LT on its GM1 receptor, likely because tannins preferentially bind to BSA. Ropiak *et al.* (2017) noted that the affinity of CTs for gelatin was greater than that for BSA, likely because CTs have high affinities for proline-rich proteins, such as gelatin that offers more binding sites (Hagerman and Butler, 1981). Similarly, He *et al.* (2006) clearly demonstrated that pentagalloylglucose formed stronger hydrophobic associations with amino acids containing aromatic groups and aliphatic side chains. These findings reveal that the extent of tannin-protein interaction strongly depends on the amino acid profile.

#### *Associated factors: tannin dose, type and chemical structure*

The beneficial effects of tannins depend primarily on the dose of tannins in the media or the diet. Both CTs from quebracho and HTs from chestnut inhibited the growth rate and toxic effects of *C. perfringens* in a dose-dependent manner (Elizondo *et al.*, 2010). Chen *et al.* (2006) also presented a clear dose-dependent inhibition of LT binding to GM1 with increasing concentrations of *Galla Chinensis* extract containing gallotannins. However, several studies highlighted the fact that the type of tannins is also important (Digrak *et al.*, 1999; Elizondo *et al.*, 2010). Elizondo *et al.* (2010) found that 8 mg/ml of HTs of chestnut wood extract (gallotannins and ellagitannins) had strong bactericidal properties, whereas 7.5 mg/ml of CTs from quebracho, containing polymers of prorobinetidinins and profisetidinins, had no effect. Despite a lower tannin content, CTs from *Acacia mollissima* (mimosa bark), containing polymers of prorobinetidinins and profisetidinins, possessed greater antibacterial activity than did ellagitannins (mainly castalagin and vescalagin) from the gallnuts of *Quercus macrolepis* (valonia oak) and gallotannins from *Quercus infectoria* (Digrak *et al.*, 1999). At the same concentration, Yu *et al.* (2011) showed that pentagalloylglucose, a nonpolar gallic acid derivative, disordered the acyl chains of the lipid bilayers compared with a very polar CT trimer of catechins. The authors suggested that the polarity of the tannin was inversely related to the strength of interaction.

The method of tannin extraction can also affect the bioactivity of tannins because different solvents extract different

types of tannin molecules (Chen *et al.*, 2006; Yao *et al.*, 2006; Unaeze *et al.*, 2017). In a previous experiment, Chen *et al.* (2006) compared the ability of several soluble fractions of *Galla Chinensis* extract to inhibit the binding of LT to GM1. At 500 µg/ml, an ethyl acetate soluble fraction of *Galla Chinensis* extract completely inhibited the binding of LT to GM1, whereas a 50% inhibition was observed for the butan-1-ol fraction, and no inhibition was found for aqueous soluble fractions. These differences might be directly associated with the chemical properties of tannins, such as the molecular weight of the tannin molecule, or the degree of galloylation of the gallotannin polymers, and tannin composition. Yao *et al.* (2006) confirmed the importance of molecular weight in relation to biological activity (interaction with proteins, enzyme inhibition and antimicrobial potency) by comparing different fractions of CTs from *Acacia mearnsii* (black wattle). The average molecular weights of water, ethyl acetate and ether extracts were 2050, 980 and 440 Da, respectively. The authors found that water fraction with high molecular weight had the strongest capacity to interact with proteins, whereas the ether extract with low molecular weight displayed greater antibacterial potency. Ethyl acetate extract, which was of medium molecular weight, had both a strong capacity to inhibit enzyme activity and strong antibacterial properties. Baert *et al.* (2016) also demonstrated that tellimagrandin I-based oligomeric ellagitannins isolated from *Epilobium angustifolium* (rosebay willow herb) flowers decreased gas production and total volatile fatty acid concentration proportionately to their degree of oligomerization. Because gas, such as methane, and volatile fatty acids are produced by the rumen microbiota, the authors hypothesized that these ellagitannins may have a direct inhibition of the methanogenic microbiota and/or could bind to dietary fibers, the substrate required for gas production. Similar conclusions were drawn by Ropiak *et al.* (2017), who showed that the greater the mean degree of polymerization or the average molecular weight, the fewer the moles of CTs are needed to precipitate the same amount of BSA or gelatin. Therefore, CT size is an important factor to consider in CT-protein interaction. In the case of ETEC-related diarrhea, testing whether small CTs would differently affect ETEC growth, biofilm formation, ETEC adhesion to the mucosa and ST and LT enterotoxin activities compared with larger CTs would be worthwhile. Finally, the chemical composition of tannins is another important component relating to tannin bioactivity. Among the diverse family of tannins, each plant has a unique chemical composition. The CTs from *Schinopsis lorentzii* (quebracho) and *Acacia mollissima* (mimosa bark) are polymers with different profisetidin : prorobinetidin ratios, whereas the CTs from temperate forage legumes are polymers with different percentages of procyanidins and prodelphinidins. Prodelphinidins have more hydroxyl groups than procyanidins, which generally indicates greater reactivity toward protein. Greater molar proportions of prodelphinidins have been shown to decrease the apparent  $\alpha$ -helix content and increase the apparent  $\beta$ -sheet content of BSA, which

indicates a conformational change in the tertiary structure of the protein (Ropiak *et al.*, 2017). The ability of such tannins to affect protein structure could be of interest to inactivate some proteins involved in ETEC pathogenesis, such as fimbriae, enterotoxins (ST and LT) and their respective receptors in the enterocytes. The antimicrobial activity of flavan-3-ols and CTs can be modified by the presence of galloyl groups as opposed to hydroxyl groups at the C3 position of the C ring (Taguri *et al.*, 2004). Similarly, comparing the binding properties of gallotannins and ellagitannins, Deaville *et al.* (2007) highlighted the importance of the conformational flexibility of tannin molecules. Their HHDP groups render ellagitannins less flexible than gallotannins. The authors demonstrated that gallotannins and pentagalloylglucose bind with equal strength to gelatin and BSA, whereas less-flexible ellagitannins bind more strongly to flexible proteins such as gelatin, and weakly to BSA. Therefore, gallotannins may be more interesting than ellagitannins to interact with a larger range of proteins involved in ETEC pathogenicity, while ensuring that the dose is not deleterious to the animal ingesting them.

The aforementioned results demonstrated that differences in the chemical structure of tannins, such as tannin size (degree of polymerization, galloylation or oligomerization) and composition, lead to differences in tannin properties, such as the flexibility, the hydrophobicity and the polarity of tannin molecules. In turn, this results in varying affinities for proteins and lipids, which ultimately disrupt ETEC pathogenicity. Moreover, owing to the broad diversity of tannins, concluding which ones are the most promising in reducing the development of coliform PWD is very difficult at present because both types of tannins, CTs and HTs, can affect ETEC pathogenicity. In addition, numerous *in vivo* experiments use commercial extracts that are mixtures of tannins, and very often, no information about the detailed tannin composition is given. Finding a type of tannin or a combination of tannins that would affect, at the same time, bacterial growth, bacterial attachment to the mucosa and enterotoxin activities introduces new challenges. Designing innovative experiments that can establish the types of tannins or tannin-containing feeds that are most effective and at which dose would help to tackle these challenges. This can be achieved through a careful choice of tannin-containing plants or the selection or breeding of the required plant models. Nevertheless, some gallotannins or gallic acid derivatives, such as pentagalloylglucose (940 Da), seem promising in concomitantly disrupting lipid membranes and in interacting strongly with different proteins. Their ability to reduce the binding of ETEC to brush borders and to disrupt the binding between the LT toxin and its receptor GM1 has proven this (Verhelst *et al.*, 2010).

## Conclusion

Tannins offer great potential for the prevention of microbial infections and the reduction of coliform PWD. With their numerous hydroxyl and phenolic groups, tannins exhibit several biological properties with a wide range of effects

on bacteria and on enterotoxin production and their activities. They directly act on pathogenic bacteria to inhibit or slow down growth by preventing bacterial adhesion to the intestinal epithelium and by inhibiting the bacterial enterotoxins and channels involved in the secretion of electrolytes and water into the lumen. In addition, some tannins seem to have prebiotic effects in the gut, together with antioxidant and anti-inflammatory effects, which were not discussed in the present review but may help piglets better cope with ETEC-related diarrhea. *In vitro* results are supported by *in vivo* experiments showing that the inclusion of tannins in the diet can decrease the severity and occurrence of diarrhea. This can engender improved health status and growth performance, as well as promote pig welfare via improved appetite. Tannins may counteract the intestinal dysbiosis that occurs at weaning, promoting and maintaining optimal gut health.

Disparities in terms of efficacy and specificity exist among the different types of tannins present in a plant or extract, as well as differences attributed to their concentration and chemical structure. Further investigations are needed to determine the optimal combinations of tannins and the most suitable and cost-effective manner with which to deliver them. Certain discrepancies between *in vitro* and *in vivo* results could be explained by the metabolism of tannins along the intestinal tract. Whether and which tannins and their metabolites remain active in the intestine warrant further research. Their antibacterial properties and the possibility for bacteria to develop resistances to tannins should be investigated, as well as how this risk could be minimized.

## Acknowledgements

The authors gratefully acknowledge the Pig Commission of the European Association of Animal Production (EAAP) for the opportunity to present this review. This work refers in part to the literature review of the doctoral thesis of Marion Girard funded by the European Marie Curie Initial Training Network ('LegumePlus'; PITNGA-2011-289377).

## Declaration of interest

The authors declare that they have no competing interests.

## Ethics statement

None.

## Software and data repository resources

None.

## References

- Al-Zoreky NS 2009. Antimicrobial activity of pomegranate (*Punica granatum* L.) fruit peels. *International Journal of Food Microbiology* 134, 244–248.
- Baert N, Pellikaan WF, Karonen M and Salminen JP 2016. A study of the structure-activity relationship of oligomeric ellagitannins on ruminal fermentation *in vitro*. *Journal of Dairy Science* 99, 8041–8052.

- Bakkiyaraj D, Nandhini JR, Malathy B and Pandian SK 2013. The anti-biofilm potential of pomegranate (*Punica granatum L.*) extract against human bacterial and fungal pathogens. *Biofouling* 29, 929–937.
- Barbehenn RV and Constabel CP 2011. Tannins in plant–herbivore interactions. *Phytochemistry* 72, 1551–1565.
- Barer MR 2012. Morphology and nature of micro-organisms. In *Medical microbiology*, 18th edition (ed. D Greenwood, M Barer, R Slack and W Irving), Chapter 2, pp. 9–23. Churchill Livingstone, Edinburgh, UK.
- Bernhoft A 2010. A brief review on bioactive compounds in plants. In *Bioactive compounds in plants – benefits and risks for man and animals* (ed. A Bernhoft), pp 11–17. The Norwegian Academy of Science and Letters, Oslo, Norway.
- Biagi G, Cipollini I, Paulicks BR and Roth FX 2010. Effect of tannins on growth performance and intestinal ecosystem in weaned piglets. *Archives of Animal Nutrition* 64, 121–135.
- Braga LC, Shupp JW, Cummings C, Jett M, Takahashi JA, Carmo LS, Chartone-Souza E and Nascimento AMA 2005. Pomegranate extract inhibits *Staphylococcus aureus* growth and subsequent enterotoxin production. *Journal of Ethnopharmacology* 96, 335–339.
- Brand P, Gobeli S and Perreten V 2017. Pathotyping and antibiotic resistance of porcine enterovirulent *Escherichia coli* strains from Switzerland (2014–2015). *Schweizer Archiv für Tierheilkunde* 159, 373–380.
- Brillouet JM, Romieu C, Schoefs B, Solymosi K, Cheynier V, Fulcrand H, Verdeil JL and Conéjéro G 2013. The tannosome is an organelle forming condensed tannins in the chlorophyllous organs of Tracheophyta. *Annals of Botany* 112, 1003–1014.
- Brus M, Dolinšek MJ, Cencič A and Škorjanc D 2013. Effect of chestnut (*Castanea sativa* Mill.) wood tannins and organic acids on growth performance and faecal microbiota of pigs from 23 to 127 days of age. *Bulgarian Journal of Agricultural Science* 19, 841–847.
- Chen JC, Ho TY, Chang YS, Wu SL and Hsiang CY 2006. Anti-diarrheal effect of *Galla chinensis* on the *Escherichia coli* heat-labile enterotoxin and ganglioside interaction. *Journal of Ethnopharmacology* 103, 385–391.
- Chung KT, Wong TY, Wei CI, Huang YW and Lin Y 1998. Tannins and human health: a review. *Critical Reviews in Food Science and Nutrition* 38, 421–464.
- Coddens A, Loos M, Vanrompay D, Remon JP and Cox E 2017. Cranberry extract inhibits *in vitro* adhesion of F4 and F18+ *Escherichia coli* to pig intestinal epithelium and reduces *in vivo* excretion of pigs orally challenged with F18+ verotoxigenic *E. coli*. *Veterinary Microbiology* 202, 64–71.
- Constabel CP, Yoshida K and Walker V 2014. Diverse ecological roles of plant tannins: plant defense and beyond. *Recent Advances in Polyphenol Research* 4, 115–142.
- Côté J, Caillet S, Doyon G, Sylvain JF and Lacroix M 2010. Bioactive compounds in cranberries and their biological properties. *Critical Reviews in Food Science and Nutrition* 50, 666–679.
- Deaville ER, Green RJ, Mueller-Harvey I, Willoughby I and Frazier RA 2007. Hydrolyzable tannin structures influence relative globular and random coil protein binding strengths. *Journal of Agricultural and Food Chemistry* 55, 4554–4561.
- Deprez S, Mila I, Huneau JF, Tome D and Scalbert A 2001. Transport of proanthocyanidin dimer, trimer, and polymer across monolayers of human intestinal epithelial Caco-2 cells. *Antioxidants and Redox Signaling* 3, 957–967.
- Devriendt B, Stuyven E, Verdonck F, Goddeeris BM and Cox E 2010. Enterotoxigenic *Escherichia coli* (K88) induce proinflammatory responses in porcine intestinal epithelial cells. *Developmental and Comparative Immunology* 34, 1175–1182.
- Digrak M, Alma MH, İlçim A and Sen S 1999. Antibacterial and antifungal effects of various commercial plant extracts. *Pharmaceutical Biology* 37, 216–220.
- Dubreuil JD, Isaacson RE and Schifferli DM 2016. Animal Enterotoxigenic *Escherichia coli*. *EcoSal Plus* 7, <https://doi.org/10.1128/ecosalplus.ESP-0006-2016>.
- Elizondo AM, Mercado EC, Rabinovitz BC and Fernandez-Miyakawa ME 2010. Effect of tannins on the *in vitro* growth of *Clostridium perfringens*. *Veterinary Microbiology* 145, 308–314.
- Engels C, Knödler M, Zhao YY, Carle R, Gänzle MG and Schieber A 2009. Antimicrobial activity of gallotannins isolated from Mango (*Mangifera indica L.*) kernels. *Journal of Agricultural and Food Chemistry* 57, 7712–7718.
- Espín JC, Gonzalez-Barrio R, Cerda B, Lopez-Bote C, Rey AI and Tomas-Barberan FA 2007. Iberian pig as a model to clarify obscure points in the bioavailability and metabolism of ellagitannins in humans. *Journal of Agricultural and Food Chemistry* 55, 10476–10485.
- European Food Safety Authority and European Centre for Disease Prevention Control 2019. The European Union summary report on antimicrobial resistance in zoonotic and indicator bacteria from humans, animals and food in 2017. *EFSA Journal* 17, e05598.
- European Medicines Agency 2018. Sales of veterinary antimicrobial agents in 30 European countries in 2016: trends from 2010 to 2016, Eighth European Surveillance of Veterinary Antimicrobial Consumption (ESVAC) report. Retrieved on 17 July 2019 from [https://www.ema.europa.eu/en/documents/report/sales-veterinary-antimicrobial-agents-30-european-countries-2016-trends-2010-2016-eighth-esvac\\_en.pdf](https://www.ema.europa.eu/en/documents/report/sales-veterinary-antimicrobial-agents-30-european-countries-2016-trends-2010-2016-eighth-esvac_en.pdf)
- Fairbrother JM, Nadeau E and Gyles CL 2005. *Escherichia coli* in postweaning diarrhea in pigs: an update on bacterial types, pathogenesis, and prevention strategies. *Animal Health Research Reviews* 6, 17–39.
- Girard M 2016. Bioactive compounds in forage legumes: structural changes during conservation, their fate along the digestive tract and their potential to impact ruminant products. Doctoral thesis, ETH Zurich, Zurich, Switzerland.
- Girard M, Pradervand N, Silacci P and Bee G 2019. L'extrait de châtaignier réduit les diarrhées de post-sevrage et modifie la population bactérienne du jéjunum. In *Proceedings of the 51st Journées de la recherche porcine*, 5–6 February 2019, Paris, France, pp. 101–102.
- Girard M, Thanner S, Pradervand N, Hu D, Ollagnier C and Bee G 2018. Hydrolysable chestnut tannins for reduction of postweaning diarrhea: efficacy on an experimental ETEC F4 model. *PLoS ONE* 13, e0197878.
- Graziani R, Tosi G and Denti R 2006. *In vitro* antimicrobial activity of SILVA FEED ENC® on bacterial strains of poultry origin. In *Proceedings of the 12th European Poultry Conference*, 10–14 September 2006, Verona, Italy, pp. 328.
- Grundhöfer P and Gross GG 2001. Immunocytochemical studies on the origin and deposition sites of hydrolyzable tannins. *Plant Science* 160, 987–995.
- Hagerman AE and Butler LG 1981. The specificity of proanthocyanidin-protein interactions. *The Journal of Biological Chemistry* 256, 4494–4497.
- Hancock V, Dahl M and Klemm P 2010. Probiotic *Escherichia coli* strain Nissle 1917 outcompetes intestinal pathogens during biofilm formation. *Journal of Medical Microbiology* 59, 392–399.
- Hartmann S 2016. Antibiotikainsatz und Tierbehandlungsindex in Schweizer Ferkelerzeugungsbetrieben. Doctoral thesis, Vetsuisse-Faculty Zurich University, Zurich, Switzerland.
- Hatano T, Han L, Taniguchi S, Shingu T, Okuda T and Yoshida T 1995. Tannins and related polyphenols of theaceous plants. VIII. Camelliatannins C and E, new complex tannins from *Camellia japonica* Leaves. *Chemical and Pharmaceutical Bulletin* 43, 1629–1633.
- He Q, Shi B and Yao K 2006. Interactions of gallotannins with proteins, amino acids, phospholipids and sugars. *Food Chemistry* 95, 250–254.
- Heo JM, Opapeju FO, Pluske JR, Kim JC, Hampson DJ and Nyachoti CM 2013. Gastrointestinal health and function in weaned pigs: a review of feeding strategies to control post-weaning diarrhoea without using in-feed antimicrobial compounds. *Journal of Animal Physiology and Animal Nutrition* 97, 207–237.
- Hoste H, Torres-Acosta JF, Sandoval-Castro CA, Mueller-Harvey I, Sotiraki S, Louvandini H, Thamsborg SM and Terrill TH 2015. Tannin containing legumes as a model for nutraceuticals against digestive parasites in livestock. *Veterinary Parasitology* 212, 5–17.
- Jakobek L 2015. Interactions of polyphenols with carbohydrates, lipids and proteins. *Food Chemistry* 175, 556–567.
- Jelager L, Gurib-Fakim A and Adrsen A 1998. Antibacterial and antifungal activity of medicinal plants of Mauritius. *Pharmaceutical Biology* 36, 153–161.
- Kamijo M, Kanazawa T, Funaki M, Nishizawa M and Yamagishi T 2008. Effects of *Rosa rugosa* petals on intestinal bacteria. *Bioscience Biotechnology and Biochemistry* 72, 773–777.
- Khanbabaee K and van Ree T 2001. Tannins: classification and definition. *Natural Product Reports* 18, 641–649.
- Krumholz LR, Crawford RL, Hemling ME and Bryant MP 1987. Metabolism of gallate and phloroglucinol in *Eubacterium oxidoreducens* via 3-hydroxy-5-oxohexanoate. *Journal of Bacteriology* 169, 1886–1890.
- Labrie V, Harel J and Dubreuil JD 2002. *Escherichia coli* heat-stable enterotoxin b (STb) *in vivo* internalization within rat intestinal epithelial cells. *Veterinary Research* 33, 223–228.

- Lauridsen C, Højberg O, Kongsted H and Canibe N 2017. A critical review on alternatives to antibiotics and pharmacological zinc for prevention of diarrhoea in pigs post-weaning. DCA – Danish Centre For Food and Agriculture, 56pp. Retrieved on 17 July 2018 from <https://www.ft.dk/samling/20161/almdele/MOF/bilag/492/1769796.pdf>
- Lekagul A, Tangcharoensathien V and Yeung S 2019. Patterns of antibiotic use in global pig production: a systematic review. *Veterinary and Animal Science* 7, 100058.
- Liu Y, Black MA, Caron L and Camesano TA 2006. Role of cranberry juice on molecular-scale surface characteristics and adhesion behavior of *Escherichia coli*. *Biotechnology and Bioengineering* 93, 297–305.
- Luppi A, Gibellini M, Gin T, Vangroenweghe F, Vandebroucke V, Bauerfeind R, Bonilauri P, Labarque G and Hidalgo Á 2016. Prevalence of virulence factors in enterotoxigenic *Escherichia coli* isolated from pigs with post-weaning diarrhoea in Europe. *Porcine Health Management* 2, 20.
- Marín L, Miguélez EM, Villar CJ and Lombó F 2015. Bioavailability of dietary polyphenols and gut microbiota metabolism: antimicrobial properties. *BioMed Research International* 2015, 905215.
- Mena P, Bresciani L, Brindani N, Ludwig IA, Pereira-Caro G, Angelino D, Llorach R, Calani L, Brighenti F, Clifford MN, Gill CIR, Crozier A, Curti C and Del Rio D 2019. Phenyl- $\gamma$ -valerolactones and phenylvaleric acids, the main colonic metabolites of flavan-3-ols: synthesis, analysis, bioavailability, and bioactivity. *Natural Product Reports* 36, 714–752.
- Min BR, Pinchak WE, Anderson RC and Callaway TR 2007. Effect of tannins on the *in vitro* growth of *Escherichia coli* O157: H7 and *in vivo* growth of generic *Escherichia coli* excreted from steers. *Journal of Food Protection* 70, 543–550.
- Min BR, Pinchak WE, Merkel R, Walker S, Tomita G and Anderson RC 2008. Comparative antimicrobial activity of tannin extract from perennial plants on mastitis pathogens. *Scientific Research and Essay* 3, 66–73.
- Mingshu L, Kai Y, Qiang H and Dongying J 2006. Biodegradation of gallotannins and ellagitannins. *Journal of Basic Microbiology* 46, 68–84.
- Nagy B and Fekete PZ 2005. Enterotoxigenic *Escherichia coli* in veterinary medicine. *International Journal of Medical Microbiology* 295, 443–454.
- Namkung W, Thiagarajah JR, Phuan PW and Verkman AS 2010. Inhibition of Ca<sup>2+</sup>-activated Cl<sup>-</sup> channels by gallotannins as a possible molecular basis for health benefits of red wine and green tea. *FASEB Journal* 24, 4178–4186.
- Oladoja NA, Alliu YB, Ofomaja AE and Unuabonah IE 2011. Synchronous attenuation of metal ions and colour in aqua stream using tannin–alum synergy. *Desalination* 271, 34–40.
- Peterson JW and Whipp SC 1995. Comparison of the mechanisms of action of cholera toxin and the heat-stable enterotoxins of *Escherichia coli*. *Infection and Immunity* 63, 1452–1461.
- Puupponen-Pimiä R, Nohynek L, Meier C, Kähkönen M, Heinonen M, Hopia A and Oksman-Caldentey KM 2001. Antimicrobial properties of phenolic compounds from berries. *Journal of Applied Microbiology* 90, 494–507.
- Ravi M, Ngeleka M, Kim SH, Gyles C, Berthiaume F, Mourez M, Middleton D and Simko E 2007. Contribution of AIDA-I to the pathogenicity of a porcine diarrheagenic *Escherichia coli* and to intestinal colonization through biofilm formation in pigs. *Veterinary Microbiology* 120, 308–319.
- Ropiak HM, Lachmann P, Ramsay A, Green RJ and Mueller-Harvey I 2017. Identification of structural features of condensed tannins that affect protein aggregation. *PLoS ONE* 12, e0170768.
- Salminen J-P and Karonen M 2011. Chemical ecology of tannins and other phenolics: we need a change in approach. *Functional Ecology* 25, 325–338.
- Schuijer M, Sies H, Illek B and Fischer H 2005. Cocoa-related flavonoids inhibit CFTR-mediated chloride transport across T84 human colon epithelia. *Journal of Nutrition* 135, 2320–2325.
- Shao D, Li J, Li J, Tang R, Liu L, Shi J, Huang Q and Yang H 2015. Inhibition of gallic acid on the growth and biofilm formation of *Escherichia coli* and *Streptococcus mutans*. *Journal of Food Science* 80, M1299–M1305.
- Sieniawska E and Baj T 2017. Tannins. In *Pharmacognosy: fundamentals, applications and strategies* (ed. S Badal and R Delgoda), Chapter 10, pp. 199–232. Academic Press, Boston, MA, USA.
- Sjölund M, Zoric M and Wallgren P 2014. Financial impact of disease on pig production. Part III. Gastrointestinal disorders. In *Proceedings of the 6th European symposium of Porcine Health Management, 7–9 May 2014, Sorrento*, pp. 189.
- Smeriglio A, Barreca D, Bellocco E and Trombetta D 2017. Proanthocyanidins and hydrolysable tannins: occurrence, dietary intake and pharmacological effects. *British Journal of Pharmacology* 174, 1244–1262.
- Soares S, Kohl S, Thalman S, Mateus N, Meyerhof W and De Freitas V 2013. Different phenolic compounds activate distinct human bitter taste receptors. *Journal of Agricultural and Food Chemistry* 61, 1525–1533.
- Taguri T, Tanaka T and Kouno I 2004. Antimicrobial activity of 10 different plant polyphenols against bacteria causing food-borne disease. *Biological and Pharmaceutical Bulletin* 27, 1965–1969.
- Unaëze BC, Ilo CE, Ekwuatu C, Orabueze I and Obi E 2017. Anti-diarrhoeal effects of three Nigerian medicinal plant extracts on *E. coli*-induced diarrhea. *International Journal of Biological and Chemical Sciences* 11, 414–419.
- Verhelst R, Schroyen M, Buys N and Niewold T 2010. The effects of plant polyphenols on enterotoxigenic *Escherichia coli* adhesion and toxin binding. *Livestock Science* 133, 101–103.
- Verhelst R, Schroyen M, Buys N and Niewold T 2014. Dietary polyphenols reduce diarrhea in enterotoxigenic *Escherichia coli* (ETEC) infected post-weaning piglets. *Livestock Science* 160, 138–140.
- Williams AR, Klaver EJ, Laan LC, Ramsay A, Frygas C, Difborg R, Kringel H, Reed JD, Mueller-Harvey I, Skov S, van Die I and Thamsborg SM 2017. Co-operative suppression of inflammatory responses in human dendritic cells by plant proanthocyanidins and products from the parasitic nematode *Trichuris suis*. *Immunology* 150, 312–328.
- Yao K, He Q, Ying Jia D and Shi B 2006. The potential of wattle tannin extracts for fine use. *Natural Product Research* 20, 271–278.
- Yu X, Chu S, Hagerman AE and Lorigan GA 2011. Probing the interaction of polyphenols with lipid bilayers by solid-state NMR spectroscopy. *Journal of Agricultural and Food Chemistry* 59, 6783–6789.
- Zhang L, Wang Y, Li D, Ho CT, Li J and Wan X 2016. The absorption, distribution, metabolism and excretion of procyanidins. *Food and Function* 7, 1273–1281.