



On Greenberg–Benois \mathcal{L} -invariants and Fontaine–Mazur \mathcal{L} -invariants

Ju-Feng Wu

Abstract. We prove a comparison theorem between Greenberg–Benois \mathcal{L} -invariants and Fontaine–Mazur \mathcal{L} -invariants. Such a comparison theorem supplies an affirmative answer to a speculation of Besser–de Shalit.

1 Introduction

Let f be a cuspidal normalized newform of weight $2k$ and level $\Gamma_0(pN)$, where p is a prime number and $N \in \mathbf{Z}_{>0}$ such that $p \nmid N$. Consider the complex L -function attached to f

$$L(f, s) = \sum_{n=1}^{\infty} a_n n^{-s}.$$

One way to study $L(f, s)$ is via p -adic method. That is, one can associate f with a p -adic L -function $L_p(f, s)$, which p -adically interpolates the algebraic part of the special values $L(f, j)$ for $1 \leq j \leq 2k - 1$. In particular, the interpolation property at $s = k$ is given by the formula

$$L_p(f, k) = \left(1 - \frac{p^{k-1}}{a_p}\right) \frac{L(f, k)}{\Omega_f},$$

where Ω_f is the Deligne period of f at k ([6]).

Suppose moreover that $a_p = p^{k-1}$; the formula above shows that $L_p(f, s)$ vanishes at $s = k$. In the case when $k = 1$, Mazur–Tate–Teitelbaum conjectured in [15] that there exists an invariant $\mathcal{L}(f)$ such that

$$\frac{d}{ds} L_p(f, s)|_{s=k} = \mathcal{L}(f) \frac{L(f, k)}{\Omega_f}.$$

This conjecture is known as the *trivial zero conjecture* and has been proven by Greenberg–Stevens in [9]. Moreover, for higher weights, various generalizations of the invariant $\mathcal{L}(f)$ has been proposed. The following is an incomplete list:

- In [8], R. Greenberg constructed the \mathcal{L} -invariants for Galois representations that are ordinary at p and suggested a generalization of the trivial zero conjecture.

Received by the editors December 27, 2023; revised June 11, 2024; accepted September 10, 2024.

AMS subject classification: 11F80, 11R34, 11S25, 11R23.

Keywords: Galois representations, p -adic Hodge theory, \mathcal{L} -invariants.



- In [14], Fontaine–Mazur defined the \mathcal{L} -invariant by studying the semistable module (à la Fontaine) associated with a p -adic representation.
- In [5], R. Coleman proposed a construction of \mathcal{L} -invariants as an application of his p -adic integration theory.
- In [18], J. Teitelbaum proposed a construction of \mathcal{L} -invariants by applying the p -adic integration theory to p -adically uniformized Shimura curve.

All these \mathcal{L} -invariants are known to be equal: Coleman–Iovita compared the second and the third in [4]; Iovita–Spieß compared the second and the fourth in [13]; and the comparison between the first and the second is a special case of [2, Proposition 2.3.7].

It is a natural question to ask whether one can establish a similar philosophy for higher rank automorphic forms. Let us mention the following generalizations in our consideration:

- In [2], D. Benois generalized Greeberg’s construction to Galois representations of $\text{Gal}_{\mathbb{Q}}$ that satisfies some reasonable conditions. He also stated a trivial zero conjecture in such a generality ([*op. cit.*, pp. 1579]).
- In [1], Besser–de Shalit generalized both the Fontaine–Mazur \mathcal{L} -invariants and Coleman (or Teitelbaum) \mathcal{L} -invariants by studying the p -adic cohomology groups of p -adically uniformized Shimura varieties. It is conjectured in *loc. cit.* that these two constructions give rise to the same \mathcal{L} -invariants (or \mathcal{L} -operators as called in *loc. cit.*). Authors of *loc. cit.* also speculated that the existence of a *trivial zero conjecture* for these two \mathcal{L} -invariants. However, they were not able to provide an explicit statement.

This article concerns the comparison between Benois’s \mathcal{L} -invariants and the Fontaine–Mazur type \mathcal{L} -invariants of Besser–de Shalit. To explain our result, let us fix some notations: Let F be a number field such that for every prime ideal $\mathfrak{p} \subset \mathcal{O}_F$ sitting above p , the maximal unramified extension of $\mathbb{Q}_{\mathfrak{p}}$ in $F_{\mathfrak{p}}$ is $\mathbb{Q}_{\mathfrak{p}}$ itself; let E be a large enough value field that is a finite extension over $\mathbb{Q}_{\mathfrak{p}}$. Suppose

$$\rho : \text{Gal}_F \rightarrow \text{GL}_n(E)$$

is a Galois representation that is semistable at places above p . We further assume that ρ satisfies the assumptions in §5.1. In particular, we assume the Frobenius eigenvalues on the associated semistable modules are given by p^m, \dots, p^{m-n+1} (for some suitable $m \in \mathbb{Z}$ that is independent of the prime ideals sitting above p) and the monodromy is maximal. We remark in the beginning that these assumptions are required so that we can perform the following two constructions:

- Following the suggestion in [17] (see also [10]), one can consider the induction $\text{Ind}_F^{\mathbb{Q}} \rho$. Part of the assumptions then allows us to attach the \mathcal{L} -invariant in Benois’s style to $\text{Ind}_F^{\mathbb{Q}} \rho(m)$. This resulting \mathcal{L} -invariant is denoted by $\mathcal{L}_{\text{GB}}(\rho(m))$, where the subscript GB stands for “Greenberg–Benois”. We refer the readers to §3 for the construction of \mathcal{L}_{GB} .
- We realized that the generalization of Fontaine–Mazur \mathcal{L} -invariants suggested by Besser–de Shalit can be translated to the world of semistable modules of a local Galois representation. The other part of the assumptions in §5.1 then allow us to attach \mathcal{L} -invariants of Fontaine–Mazur style to each local Galois representation

$\rho_{\mathfrak{p}} = \rho|_{\text{Gal}_{F_{\mathfrak{p}}}}$ for every prime \mathfrak{p} above p . We term such \mathcal{L} -invariants $\mathcal{L}_{\text{FM}}(\rho_{\mathfrak{p}})$, where the subscript FM stands for “Fontaine–Mazur”. We refer the readers to §4 for the construction of \mathcal{L}_{FM} .

Our main result reads as follows.

Theorem 5.4 *We have an equality*

$$\mathcal{L}_{\text{GB}}(\rho(m)) = \prod_{\mathfrak{p}|p} -\mathcal{L}_{\text{FM}}(\rho_{\mathfrak{p}}),$$

where the index set runs through all prime ideals in \mathcal{O}_F sitting above p

Since there is a well-stated trivial zero conjecture for $\mathcal{L}_{\text{GB}}(\rho(m))$ in [2], our result immediately supplies an affirmative answer to Besser–de Shalit’s speculation of the relationship between their \mathcal{L} -invariants and p -adic L -functions.

To close this introduction, let us mention that the generalization of \mathcal{L} -invariants à la Coleman (or Teitelbaum) suggested by Besser–de Shalit replaces Coleman’s integration theory with Besser’s theory of finite polynomial cohomology. Although they only consider the case for the trivial coefficient (so that we can only see automorphic forms of weight associated with the differential 1-forms), one can hope a generalization for nontrivial coefficients by using *finite polynomial cohomology with coefficients* ([12]). We wish to come back to this in future projects and hopefully to compare this type of \mathcal{L} -invariants with $\mathcal{L}_{\text{FM}}(\rho_{\mathfrak{p}})$ as suggested in [1].

Notations

- Through out this article, we fix a prime number p .
- Given a field F , we fix a separable closure \bar{F} and denote by $\text{Gal}_F = \text{Gal}(\bar{F}/F)$ its absolute Galois group.

2 Preliminaries on (φ, Γ) -modules

2.1 General (φ, Γ) -modules

Fix a compatible system of primitive p -power roots of unity $(\zeta_{p^n})_{n \in \mathbb{Z}_{\geq 0}}$ in $\bar{\mathbf{Q}}_p$. Given a finite extension K of \mathbf{Q}_p , consider $K(\zeta_{p^\infty}) = \bigcup_{n \in \mathbb{Z}_{\geq 0}} K(\zeta_{p^n})$ and denote by $\Gamma = \Gamma_K$ the Galois group $\text{Gal}(K(\zeta_{p^\infty})/K)$. Moreover, for any $r \in [0, 1)$, let

$$\mathcal{R}_K^r := \left\{ f = \sum_{i \in \mathbb{Z}} a_i T^i : \begin{array}{l} a_i \in K^{\text{unr}} \cap K(\zeta_{p^\infty}) \\ f \text{ is holomorphic on the annulus } r \leq |T| < 1 \end{array} \right\}$$

and

$$\mathcal{R}_K := \bigcup_{r \in [0, 1)} \mathcal{R}_K^r,$$

where K^{unr} is the maximal unramified extension of K in $\bar{\mathbf{Q}}_p$ and the infinite union is taken with respect to the inclusions $\mathcal{R}_K^r \hookrightarrow \mathcal{R}_K^{r'}$ for $r \leq r' < 1$. We call the ring \mathcal{R}_K the

Robba ring over K . It carries a φ -action and a Γ -action via the formula

$$\varphi(T) = (1 + T)^p - 1 \quad \text{and} \quad \gamma(T) = (1 + T)^{\chi_{\text{cyc}}(\gamma)} - 1 \text{ for any } \gamma \in \Gamma,$$

where χ_{cyc} is the cyclotomic character.

In what follows, we shall consider a more generalized version of \mathcal{R}_K . Let E be a finite extension of \mathbf{Q}_p . We denote by $\mathcal{R}_{K,E} := \mathcal{R}_K \otimes_{\mathbf{Q}_p} E$ and call it the *Robba ring over K with coefficients in E* . We linearize the actions of φ and Γ on $\mathcal{R}_{K,E}$ via $\varphi \otimes \text{id}$ and $\gamma \otimes \text{id}$, respectively. In what follows, we often assume E is large enough so that $K \subset E$.

By a (φ, Γ) -module over $\mathcal{R}_{K,E}$, we mean a finite free $\mathcal{R}_{K,E}$ -module D together with a φ -semilinear endomorphism φ_D and a semilinear action by Γ , which commute with each other, such that the induced map

$$\varphi_D : \varphi^* D = D \otimes_{\varphi} \mathcal{R}_{K,E} \rightarrow D$$

is an isomorphism. We shall denote by $\mathbf{Mod}_{\mathcal{R}_{K,E}}^{(\varphi, \Gamma)}$ the category of (φ, Γ) -modules over $\mathcal{R}_{K,E}$.

Let $\mathbf{Rep}_K(E)$ the category of Galois representations of Gal_K with coefficients in E . Then, by [2, Proposition 1.1.4], there is a fully faithful functor

$$\mathbf{D}_{\text{rig}}^\dagger : \mathbf{Rep}_K(E) \rightarrow \mathbf{Mod}_{\mathcal{R}_{K,E}}^{(\varphi, \Gamma)}.$$

Moreover, by letting $\mathbf{Mod}_{K,E}^{(\varphi, N)}$ (resp., $\mathbf{Mod}_{K,E}^\varphi$) the category of (φ, N) -modules (resp., φ -modules) over $K_0 = K \cap \mathbf{Q}_p^{\text{unr}}$ with coefficients in E , there is a functor (see, for example, [2, §1.2.3])

$$\mathcal{D}_{\text{st}} : \mathbf{Mod}_{\mathcal{R}_{K,E}}^{(\varphi, \Gamma)} \rightarrow \mathbf{Mod}_{K,E}^{(\varphi, N)} \quad (\text{resp., } \mathcal{D}_{\text{cris}} : \mathbf{Mod}_{\mathcal{R}_{K,E}}^{(\varphi, \Gamma)} \rightarrow \mathbf{Mod}_{K,E}^\varphi)$$

such that if $\rho \in \mathbf{Rep}_K(E)$ is semistable (resp., crystalline), then ([3, Théorème 0.2])

$$\mathcal{D}_{\text{st}}(\mathbf{D}_{\text{rig}}^\dagger(\rho)) = \mathbf{D}_{\text{st}}(\rho) \quad (\text{resp., } \mathcal{D}_{\text{cris}}(\mathbf{D}_{\text{rig}}^\dagger(\rho)) = \mathbf{D}_{\text{cris}}(\rho)).$$

Here, \mathbf{D}_{st} (resp., \mathbf{D}_{cris}) is Fontaine’s semistable (resp., crystalline) functor ([7, 3]), assigning a Galois representation in $\mathbf{Rep}_K(E)$ a (φ, N) -module (resp., φ -module) over K_0 with coefficients in E .

Now, let D be a (φ, Γ) -module over $\mathcal{R}_{K,E}$. Recall the cohomology of D is defined by the cohomology of the *Herr complex*

$$0 \rightarrow D \xrightarrow{x \mapsto ((\varphi_D - 1)x, (\gamma - 1)x)} D \oplus D \xrightarrow{(x, y) \mapsto (\gamma - 1)x - (\varphi_D - 1)y} D \rightarrow 0,$$

where γ is a (fixed) topological generator of Γ . Note that, given $\alpha = (x, y) \in D \oplus D$ such that $(\gamma - 1)x - (\varphi_D - 1)y = 0$, there is an extension

$$0 \rightarrow D \rightarrow D_\alpha \rightarrow \mathcal{R}_{K,E} \rightarrow 0$$

defined by

$$(2.1) \quad D_\alpha = D \oplus \mathcal{R}_{K,E}e, \quad (\varphi_{D_\alpha} - 1)e = x, \quad (\gamma - 1)e = y.$$

It turns out that such an assignment gives rise to an isomorphism

$$H^1(D) \cong \text{Ext}_{(\varphi, \Gamma)}^1(\mathcal{R}_{K,E}, D).$$

Furthermore, we write $H_{\text{st}}^1(D)$ (resp., $H_f^1(D)$) the subspace of $H^1(D)$, consisting of those semistable (resp., crystalline) extensions D_α (i.e., those satisfy $\text{rank}_{K_0 \otimes_{\mathbb{Q}_p} E} \mathcal{D}_{\text{st}}(D_\alpha) = \text{rank}_{K_0 \otimes_{\mathbb{Q}_p} E} \mathcal{D}_{\text{st}}(D) + 1$ (resp., $\text{rank}_{K_0 \otimes_{\mathbb{Q}_p} E} \mathcal{D}_{\text{cris}}(D_\alpha) = \text{rank}_{K_0 \otimes_{\mathbb{Q}_p} E} \mathcal{D}_{\text{cris}}(D) + 1$). According to [2, Proposition 1.4.2], if $\rho \in \mathbf{Rep}_K(E)$, then

$$H_{\text{st}}^1(\mathbf{D}_{\text{rig}}^\dagger(\rho)) \cong H_{\text{st}}^1(K, \rho) \quad (\text{resp., } H_f^1(\mathbf{D}_{\text{rig}}^\dagger(\rho)) \cong H_f^1(K, \rho)),$$

where

$$H_{\text{st}}^1(K, \rho) = \ker \left(H^1(K, \rho) \rightarrow H^1(K, \rho \otimes_{\mathbb{Q}_p} \mathbf{B}_{\text{st}}) \right) \\ (\text{resp., } H_f^1(K, \rho) = \ker \left(H^1(K, \rho) \rightarrow H^1(K, \rho \otimes_{\mathbb{Q}_p} \mathbf{B}_{\text{cris}}) \right))^{11}$$

is the usual local Bloch–Kato Selmer group.

To conclude our discussion for general (φ, Γ) -modules, we mention that if D is semistable,² then $H_{\text{st}}^1(D)$ and $H_f^1(D)$ can be computed by complexes C_{st}^\bullet and C_{cris}^\bullet , respectively ([2, Proposition 1.4.4]). Here,

$$C_{\text{st}}^\bullet(D) = \left[\begin{array}{c} \mathcal{D}_{\text{st}}(D) \xrightarrow{a \mapsto (a, (\varphi-1)a, N(a))} \frac{\mathcal{D}_{\text{st}}(D)}{\text{Fil}_{\text{dR}}^0 \mathcal{D}_{\text{st}}(D)} \oplus \mathcal{D}_{\text{st}}(D) \oplus \mathcal{D}_{\text{st}}(D) \\ \downarrow (a, b, c) \mapsto N(b) - (p\varphi-1)c \\ \mathcal{D}_{\text{st}}(D) \end{array} \right]$$

and

$$C_{\text{cris}}^\bullet(D) = \left[\mathcal{D}_{\text{cris}}(D) \xrightarrow{a \mapsto (a, (\varphi-1)a)} \frac{\mathcal{D}_{\text{cris}}(D)}{\text{Fil}_{\text{dR}}^0 \mathcal{D}_{\text{cris}}(D)} \oplus \mathcal{D}_{\text{cris}}(D) \right].$$

2.2 (φ, Γ) -modules of rank 1

Recall that (φ, Γ) -modules of rank 1 can be understood via continuous characters. More precisely, given a continuous character $\delta : K^\times \rightarrow E^\times$ and fixing a uniformizer $\omega \in K$, we can write $\delta = \delta' \delta''$ with $\delta'|_{\mathcal{O}_K^\times} = \delta|_{\mathcal{O}_K^\times}$, $\delta'(\omega) = 1$ and $\delta''(\omega) = \delta(\omega)$, $\delta''|_{\mathcal{O}_K^\times} = 1$. By local class field theory, δ' defines a unique one-dimensional Galois representation $\chi_{\delta'}$; that is,

$$\chi_{\delta'} : \text{Gal}_K \xrightarrow{\text{local Artin map}} \widehat{K^\times} \cong \mathcal{O}_K^\times \times \widehat{\mathbf{Z}} \xrightarrow{(a,b) \mapsto \delta'(a)} E^\times,^3$$

which admits its associated (φ, Γ) -module $\mathbf{D}_{\text{rig}}^\dagger(\chi_{\delta'})$. However, we define $\mathcal{R}_{K,E}(\delta'') = \mathcal{R}_{K,E} e_{\delta''}$ such that $\varphi(e_{\delta''}) = \delta(\omega) e_{\delta''}$ and $\gamma(e_{\delta''}) = e_{\delta''}$. Then, the (φ, Γ) -module associated with δ is defined to be

$$\mathcal{R}_{K,E}(\delta) := \mathcal{R}_{K,E}(\delta'') \otimes_{\mathcal{R}_{K,E}} \mathbf{D}_{\text{rig}}^\dagger(\chi_{\delta'}).$$

¹Here, \mathbf{B}_{st} and \mathbf{B}_{cris} are, respectively, Fontaine’s semistable and crystalline period rings.

²In fact, the condition can be loosened to being *potentially semistable*, but we do not need such a generality here.

³Here, $\widehat{K^\times}$ is the profinite completion of K^\times . Note that the isomorphism $\widehat{K^\times} \cong \mathcal{O}_K^\times \times \widehat{\mathbf{Z}}$ depends on the choice of ω , which is fixed.

In particular, the cyclotomic character $\text{Gal}_K \rightarrow \mathcal{O}_E^\times$ has the associated (φ, Γ) -module $\mathbf{D}_{\text{rig}}^\dagger(\chi_{\text{cyc}})$. By [16, Lemma 2.13], we know that

$$\mathbf{D}_{\text{rig}}^\dagger(\chi_{\text{cyc}}) = \mathcal{R}_{K,E}(\text{Nm}_{\mathbf{Q}_p}^K(z) | \text{Nm}_{\mathbf{Q}_p}^K(z)|),$$

where $\text{Nm}_{\mathbf{Q}_p}^K$ is the norm function from K to \mathbf{Q}_p .

Lemma 2.1. *Let $\delta : K^\times \rightarrow E^\times$ be the character*

$$\delta(z) = \left(\prod_{\sigma: K \rightarrow \overline{\mathbf{Q}_p}} \sigma(z)^{m_\sigma} \right) \left| \text{Nm}_{\mathbf{Q}_p}^K(z) \right|$$

such that all $m_\sigma \geq 1$.

- If $(\mathcal{D}_{\text{st}}(\mathcal{R}_{K,E}(\delta)))^\vee(\chi_{\text{cyc}})^{\varphi=1}$ is nonzero, then the inclusion $H_{\text{st}}^1(\mathcal{R}_{K,E}(\delta)) \hookrightarrow H^1(\mathcal{R}_{K,E}(\delta))$ is an isomorphism.
- If $(\mathcal{D}_{\text{st}}(\mathcal{R}_{K,E}(\delta)))^\vee(\chi_{\text{cyc}})^{\varphi=1} = 0$, then the inclusion $H_f^1(\mathcal{R}_{K,E}(\delta)) \hookrightarrow H_{\text{st}}^1(\mathcal{R}_{K,E}(\delta))$ is an isomorphism.

Proof By [2, Corollary 1.4.5], we have the formula

$$\dim_E H_{\text{st}}^1(\mathcal{R}_{K,E}(\delta)) - \dim_E H_f^1(\mathcal{R}_{K,E}(\delta)) = \dim_E (\mathcal{D}_{\text{st}}(\mathcal{R}_{K,E}(\delta)))^\vee(\chi_{\text{cyc}})^{\varphi=1}.$$

Applying [17, Proposition 2.1 & Lemma 2.3], we know that

$$\dim_E H_{\text{st}}^1(\mathcal{R}_{K,E}(\delta)) \leq [K : \mathbf{Q}_p] + 1 \quad \text{and} \quad \dim_E H_f^1(\mathcal{R}_{K,E}(\delta)) = [K : \mathbf{Q}_p].$$

The lemma then follows easily. ■

Suppose $\delta : K^\times \rightarrow E^\times$ is a continuous character as in Lemma 2.1. Suppose $\mathcal{R}_{K,E}(\delta)$ is semistable and so $\text{rank}_{K_0 \otimes_{\mathbf{Q}_p} E} \mathcal{D}_{\text{st}}(\mathcal{R}_{K,E}(\delta)) = 1$. We fix a $K_0 \otimes_{\mathbf{Q}_p} E$ -basis ν_δ for $\mathcal{D}_{\text{st}}(\mathcal{R}_{K,E}(\delta))$ and define

$$\beta_\delta^* = -\text{cl}(0, 0, \nu_\delta), \quad \alpha_\delta^* = \text{cl}(\nu_\delta, 0, 0) \in H^1(C_{\text{st}}^\bullet(\mathcal{R}_{K,E}(\delta))) = H_{\text{st}}^1(\mathcal{R}_{K,E}(\delta)).^4$$

Lemma 2.2. *Suppose $\eta : K^\times \rightarrow E^\times$ is a continuous character of the form $\eta(z) = \prod_{\sigma: K \rightarrow \overline{\mathbf{Q}_p}} \sigma(z)^{n_\sigma}$ with all $n_\sigma \leq 0$. Suppose*

$$0 \rightarrow \mathcal{R}_{K,E}(\delta) \rightarrow D \rightarrow \mathcal{R}_{K,E}(\eta) \rightarrow 0$$

is a semistable extension (in the sense of §2.1). Then,

$$\text{image}(\partial : H^0(\mathcal{R}_{K,E}(\eta)) \rightarrow H^1(\mathcal{R}_{K,E}(\delta))) \subset H_{\text{st}}^1(\mathcal{R}_{K,E}(\delta)).$$

Moreover, there exists a unique $\mathcal{L}(D) \in E$ such that

$$\beta_\delta^* + \mathcal{L}(D)\alpha_\delta^* \in \text{image } \partial.$$

⁴We use such notations due to [2, Theorem 1.5.7].

Proof First of all, it follows from [2, Proposition 1.2.7] that $\mathcal{R}_{K,E}(\eta)$ is also semistable. Hence, by applying [op. cit., Proposition 1.4.4], we know that

$$H^0(\mathcal{R}_{K,E}(\eta)) = H_{\text{st}}^0(\mathcal{R}_{K,E}).$$

Taking the cohomology of the short exact sequence in the lemma, we have a commutative diagram

$$\begin{array}{ccc} H^0(\mathcal{R}_{K,E}(\eta)) & \xrightarrow{\partial} & H^1(\mathcal{R}_{K,E}(\delta)) \\ \parallel & & \uparrow \\ H_{\text{st}}^0(\mathcal{R}_{K,E}(\eta)) & \xrightarrow{\partial} & H_{\text{st}}^1(\mathcal{R}_{K,E}(\delta)) \end{array},$$

which shows the first claim.

Since $\mathcal{R}_{K,E}(\eta)$ is semistable and it is of rank 1 over $\mathcal{R}_{K,E}$, it is crystalline and $\mathcal{D}_{\text{st}}(\mathcal{R}_{K,E}(\eta)) = \mathcal{D}_{\text{cris}}(\mathcal{R}_{K,E}(\eta))$. This is because the monodromy operator is nilpotent. We consider the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{D}_{\text{st}}(\mathcal{R}_{K,E}(\delta)) & \longrightarrow & \mathcal{D}_{\text{st}}(D) & \longrightarrow & \mathcal{D}_{\text{st}}(\mathcal{R}_{K,E}(\eta)) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \frac{\mathcal{D}_{\text{st}}(\mathcal{R}_{K,E}(\delta))}{\text{Fil}_{\text{dR}}^0 \mathcal{D}_{\text{st}}(\mathcal{R}_{K,E}(\delta))} \oplus \mathcal{D}_{\text{st}}(\mathcal{R}_{K,E}(\delta))^2 & \longrightarrow & \frac{\mathcal{D}_{\text{st}}(D)}{\text{Fil}_{\text{dR}}^0 \mathcal{D}_{\text{st}}(D)} \oplus \mathcal{D}_{\text{st}}(D)^2 & \longrightarrow & \frac{\mathcal{D}_{\text{st}}(\mathcal{R}_{K,E}(\eta))}{\text{Fil}_{\text{dR}}^0 \mathcal{D}_{\text{st}}(\mathcal{R}_{K,E}(\eta))} \oplus \mathcal{D}_{\text{st}}(\mathcal{R}_{K,E}(\eta))^2 \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathcal{D}_{\text{st}}(\mathcal{R}_{K,E}(\delta)) & \longrightarrow & \mathcal{D}_{\text{st}}(D) & \longrightarrow & \mathcal{D}_{\text{st}}(\mathcal{R}_{K,E}(\eta)) \longrightarrow 0 \end{array}$$

induced by the short exact sequence in the lemma, where the rows are exact and the columns are the semistable complexes. Let v_η be the element in $\mathcal{D}_{\text{st}}(\mathcal{R}_{K,E}(\eta))$ that gives rise to the basis of $H^0(\mathcal{R}_{K,E}(\eta))$ as in [17, Proposition 2.1]. In particular, $v_\eta \in \text{Fil}_{\text{dR}}^0 \mathcal{D}_{\text{st}}(\mathcal{R}_{K,E}(\eta))$ and $\varphi(v_\eta) = v_\eta$. Using the relation $N\varphi = p\varphi N$, one deduces that 1 and p^{-1} are Frobenius eigenvalues of $\mathcal{D}_{\text{st}}(D)$. We choose a lift $\tilde{v}_\eta \in \mathcal{D}_{\text{st}}(D)$ such that $\varphi(\tilde{v}_\eta) = \tilde{v}_\eta$. This then implies that $N(\tilde{v}_\eta)$ has Frobenius eigenvalue p^{-1} . The commutativity of the diagram then yields

$$\begin{array}{ccc} \tilde{v}_\eta & \longmapsto & v_\eta \\ \downarrow & & \downarrow \\ (\tilde{v}_\eta, 0, N(\tilde{v}_\eta)) & \longmapsto & 0 \end{array}.$$

Applying the exactness of the middle row, we see that

$$(\tilde{v}_\eta, 0, N(\tilde{v}_\eta)) = a(v_\delta, 0, 0) - b(0, 0, v_\delta).$$

Since $N(\tilde{v}_\eta)$ is a basis for the Frobenius eigensubspace of $\mathcal{D}_{\text{st}}(D)$ on which φ acts via p^{-1} , we see that b is invertible. We then conclude that

$$\partial : H^0(\mathcal{R}_{K,E}(\eta)) \rightarrow H_{\text{st}}^1(\mathcal{R}_{K,E}(\delta)), \quad \text{cl}(v_\eta) \mapsto a\alpha_\delta^* + b\beta_\delta^*.$$

Therefore, $\mathcal{L}(D) := a/b$. ■

3 Greenberg–Benois \mathcal{L} -invariants

In this section, we first discuss the construction of Greenberg–Benois \mathcal{L} -invariant over \mathbf{Q} in §3.1 by summarizing Benoist’s construction in [2, §2] (in particular [*op. cit.*, (26)]). Then we follow the strategy in [17], generalizing Benoist’s construction to general number fields by considering inductions of Galois representations (§3.2).

3.1 Greenberg–Benois \mathcal{L} -invariants over \mathbf{Q}

To define Greenberg–Benois \mathcal{L} -invariants over \mathbf{Q} , we start with a Galois representation

$$\rho : \text{Gal}_{\mathbf{Q}} \rightarrow \text{GL}_n(E),$$

which is unramified outside a finite set of places. We denote by

$$S = \{\ell : \rho|_{\text{Gal}_{\mathbf{Q}_\ell}} \text{ is ramified}\} \cup \{p, \infty\}$$

and let \mathbf{Q}_S be the maximal extension of \mathbf{Q} that is unramified outside S .

Recall the Bloch–Kato Selmer group associated with ρ : Given $v \in S$, define the local Selmer groups

$$H_f^1(\mathbf{Q}_v, \rho) = \begin{cases} \ker(H^1(\mathbf{Q}_\ell, \rho) \rightarrow H^1(I_\ell, \rho)), & \text{if } v = \ell \nmid p\infty, \\ \ker(H^1(\mathbf{Q}_p, \rho) \rightarrow H^1(\mathbf{Q}_p, \rho \otimes_{\mathbf{Q}_p} \mathbf{B}_{\text{cris}})), & \text{if } v = p, \\ H^1(\mathbf{R}, \rho), & \text{if } v = \infty, \end{cases}$$

where I_ℓ stands for the inertia group at ℓ . Then, the Bloch–Kato Selmer group associated with ρ is defined to be

$$H_f^1(\mathbf{Q}, \rho) := \ker\left(H^1(\text{Gal}_{\mathbf{Q}_S}, \rho) \rightarrow \bigoplus_{v \in S} \frac{H^1(\mathbf{Q}_v, \rho)}{H_f^1(\mathbf{Q}_v, \rho)}\right).$$

Let $\rho_p := \rho|_{\text{Gal}_{\mathbf{Q}_p}}$. We follow [2, §2.1.2, 2.1.4] and proceed with the following conditions:

- (B1) The local representation ρ_p is semistable with Hodge–Tate weights $k_1 \leq k_2 \leq \dots \leq k_n$, giving rise to the *de Rham filtration* $\text{Fil}_{\text{dR}}^\bullet \mathbf{D}_{\text{st}}(\rho)$.
- (B2) The Frobenius action on $\mathbf{D}_{\text{st}}(\rho_p)$ is semisimple at 1 and p^{-1} .
- (GB1) $H_f^1(\mathbf{Q}, \rho) = 0 = H_f^1(\mathbf{Q}, \rho^\vee(1))$.⁵
- (GB2) $H^0(\text{Gal}_{\mathbf{Q}_S}, \rho) = 0 = H^0(\text{Gal}_{\mathbf{Q}_S}, \rho^\vee(1))$.
- (GB3) The associated (φ, Γ) -module $\mathbf{D}_{\text{rig}}^\dagger(\rho_p)$ has no saturated subquotient⁶ isomorphic to $U_{k,m}$ with $k \geq 1$ and $m \geq 0$ ([2, §2.1.2]), where $U_{k,m}$ is the unique crystalline (φ, Γ) -module sitting in a nonsplit short exact sequence

$$0 \rightarrow \mathcal{R}_{\mathbf{Q}_p, E}(|z|z^k) \rightarrow U_{k,m} \rightarrow \mathcal{R}_{\mathbf{Q}_p, E}(z^{-m}) \rightarrow 0.$$

⁵Here, by confusing ρ with its underlying vector space, $\rho^\vee = \text{Hom}(\rho, E)$ is the dual representation of ρ and $\rho^\vee(1)$ is the twist of ρ^\vee by the cyclotomic character.

⁶Here, by “saturated”, we mean the following: A *saturated* (φ, Γ) -submodule of a (φ, Γ) -module is a (φ, Γ) -submodule that has a torsion-free quotient. A *saturated* subquotient is a subquotient arising from saturated (φ, Γ) -submodules; in particular, a saturated subquotient is torsion-free.

Given a regular submodule $D \subset \mathbf{D}_{\text{st}}(\rho_p)$ (i.e., a (φ, N) -submodule such that $\mathbf{D}_{\text{st}}(\rho_p) = D \oplus \text{Fil}_{\text{dR}}^0 \mathbf{D}_{\text{st}}(\rho_p)$), Benois defines a five-step filtration

$$(3.1) \quad D_i^{\text{GB}} := \begin{cases} 0, & i = -2, \\ (1 - p^{-1}\varphi^{-1})D + N(D^{\varphi=1}), & i = -1, \\ D, & i = 0, \\ D + \mathbf{D}_{\text{st}}(\rho_p)^{\varphi=1} \cap N^{-1}(D^{\varphi=p^{-1}}), & i = 1, \\ \mathbf{D}_{\text{st}}(\rho_p), & i = 2. \end{cases}$$

Such a filtration then yields a filtration on $\mathbf{D}_{\text{rig}}^\dagger(\rho_p)$ by

$$\text{Fil}_i^{\text{GB}} \mathbf{D}_{\text{rig}}^\dagger(\rho_p) = \mathbf{D}_{\text{rig}}^\dagger(\rho_p) \cap (D_i^{\text{GB}} \otimes_{\mathbf{Q}_p} \mathcal{R}_{\mathbf{Q}_p, E}[1/t]),$$

where $t = \log(1 + T) \in \mathcal{R}_{\mathbf{Q}_p, E}$.

Using this filtration, we define the exceptional subquotient

$$W := \text{Fil}_1^{\text{GB}} \mathbf{D}_{\text{rig}}^\dagger(\rho_p) / \text{Fil}_{-1}^{\text{GB}} \mathbf{D}_{\text{rig}}^\dagger(\rho_p).$$

By [2, Proposition 2.1.7], we have

$$\begin{aligned} W &\cong W_0 \oplus W_1 \oplus M && \text{rank } W_0 = \dim_E H^0(W^\vee(1)), \\ \text{Gr}_0^{\text{GB}} \mathbf{D}_{\text{rig}}^\dagger(\rho_p) &\cong W_0 \oplus M_0 && \text{with } \text{rank } W_1 = \dim_E H^0(W), \\ \text{Gr}_1^{\text{GB}} \mathbf{D}_{\text{rig}}^\dagger(\rho_p) &\cong W_1 \oplus M_1 && \text{rank } M_0 = \text{rank } M_1, \end{aligned}$$

where $M, M_0,$ and M_1 sit inside a short exact sequence

$$0 \rightarrow M_0 \rightarrow M \rightarrow M_1 \rightarrow 0.$$

Moreover, one has

$$\begin{aligned} H^1(W) &= \text{coker} \left(H^1(\text{Fil}_{-1}^{\text{GB}} \mathbf{D}_{\text{rig}}^\dagger(\rho_p)) \rightarrow H^1(\text{Fil}_1^{\text{GB}} \mathbf{D}_{\text{rig}}^\dagger(\rho_p)) \right), \\ H_f^1(W) &= \text{coker} \left(H_f^1(\text{Fil}_{-1}^{\text{GB}} \mathbf{D}_{\text{rig}}^\dagger(\rho_p)) \rightarrow H_f^1(\text{Fil}_1^{\text{GB}} \mathbf{D}_{\text{rig}}^\dagger(\rho_p)) \right), \end{aligned}$$

and $\dim_E H^1(W)/H_f^1(W) = e_D = \text{rank } M_0 + \text{rank } W_0 + \text{rank } W_1$ ([2, §2.2.1]).

Under the assumption (GB1) and (GB2), one applies Poitou–Tate exact sequence and deduces an isomorphism

$$H^1(\text{Gal}_{\mathbf{Q}_s}, \rho) \cong \bigoplus_{v \in S} \frac{H^1(\mathbf{Q}_v, \rho)}{H_f^1(\mathbf{Q}_v, \rho)}.$$

Note that the latter space contains an e_D -dimensional subspace $\frac{H^1(W)}{H_f^1(W)} \cong \frac{H^1(\text{Fil}_1^{\text{GB}} \mathbf{D}_{\text{rig}}^\dagger(\rho_p))}{H_f^1(\mathbf{Q}_p, \rho)}$. We then define $H^1(D, \rho)$ to be the image of $\frac{H^1(W)}{H_f^1(W)}$ in $H^1(\text{Gal}_{\mathbf{Q}_s}, \rho)$.

To define the \mathcal{L} -invariant, we further assume that

(GB4) $W_0 = 0$ and the Hodge–Tate weights for $\text{Gr}_1^{\text{GB}} \mathbf{D}_{\text{rig}}^\dagger(\rho_p)$ are positive (see [2, Proposition 1.5.9]).

Benois shows that there is a decomposition ([2, §2.1.9] (see also the discussion in [11, §1.2]))

$$H^1(\mathrm{Gr}_1^{\mathrm{GB}} \mathbf{D}_{\mathrm{rig}}^\dagger(\rho_p)) \cong H_f^1(\mathrm{Gr}_1^{\mathrm{GB}} \mathbf{D}_{\mathrm{rig}}^\dagger(\rho_p)) \oplus H_c^1(\mathrm{Gr}_1^{\mathrm{GB}} \mathbf{D}_{\mathrm{rig}}^\dagger(\rho_p))$$

and isomorphisms

$$H_f^1(\mathrm{Gr}_1^{\mathrm{GB}} \mathbf{D}_{\mathrm{rig}}^\dagger(\rho_p)) \cong \mathcal{D}_{\mathrm{cris}}(\mathrm{Gr}_1^{\mathrm{GB}} \mathbf{D}_{\mathrm{rig}}^\dagger(\rho_p)) \cong H_c^1(\mathrm{Gr}_1^{\mathrm{GB}} \mathbf{D}_{\mathrm{rig}}^\dagger(\rho_p)).$$

There are natural morphisms $\rho_{D,?} : H^1(D, \rho) \rightarrow \mathcal{D}_{\mathrm{cris}}(\mathrm{Gr}_1^{\mathrm{GB}} \mathbf{D}_{\mathrm{rig}}^\dagger(\rho_p))$ (for $? \in \{f, c\}$) making the diagram

$$\begin{array}{ccc} \mathcal{D}_{\mathrm{cris}}(\mathrm{Gr}_1^{\mathrm{GB}} \mathbf{D}_{\mathrm{rig}}^\dagger(\rho_p)) & \xrightarrow{\cong} & H_f^1(\mathrm{Gr}_1^{\mathrm{GB}} \mathbf{D}_{\mathrm{rig}}^\dagger(\rho_p)) \\ \rho_{D,f} \uparrow & & \uparrow \\ H^1(D, \rho) & \longrightarrow & H^1(\mathrm{Gr}_1^{\mathrm{GB}} \mathbf{D}_{\mathrm{rig}}^\dagger(\rho_p)) \\ \rho_{D,c} \downarrow & & \downarrow \\ \mathcal{D}_{\mathrm{cris}}(\mathrm{Gr}_1^{\mathrm{GB}} \mathbf{D}_{\mathrm{rig}}^\dagger(\rho_p)) & \xrightarrow{\cong} & H_c^1(\mathrm{Gr}_1^{\mathrm{GB}} \mathbf{D}_{\mathrm{rig}}^\dagger(\rho_p)) \end{array}$$

commutative. Under the assumption of (GB4), Benois shows that $\rho_{D,c}$ is an isomorphism, and so one can define the *Greenberg–Benois \mathcal{L} -invariant* attached to ρ (with respect to D) as

$$\mathcal{L}_{\mathrm{GB}}(\rho) = \mathcal{L}_{\mathrm{GB}}(\rho, D) := \det(\rho_{D,f} \circ \rho_{D,c}^{-1}) \in E.$$

3.2 Greenberg–Benois \mathcal{L} -invariants over general number fields

To define the Greenberg–Benois \mathcal{L} -invariants over general number fields, we follow the idea in [17] (see also [10]) and consider the induction of a Galois representation. More precisely, let F be a number field and suppose we are given a Galois representation

$$\rho : \mathrm{Gal}_F \rightarrow \mathrm{GL}_n(E),$$

where E is (again) a finite extension of \mathbf{Q}_p . We shall consider the induction $\mathrm{Ind}_F^{\mathbf{Q}} \rho$ and define S similarly as before.

Assume the following conditions hold for ρ :

- (B1) For each place $\mathfrak{p}|p$ in F , $\rho_{\mathfrak{p}} = \rho|_{\mathrm{Gal}_{F_{\mathfrak{p}}}}$ is semistable with Hodge–Tate weights $k_{\mathfrak{p},\sigma,1} \leq k_{\mathfrak{p},\sigma,2} \leq \dots \leq k_{\mathfrak{p},\sigma,n}$, where $\sigma : F_{\mathfrak{p}} \hookrightarrow \overline{\mathbf{Q}_p}$.
- (B2) For each place $\mathfrak{p}|p$ in F , the Frobenius action on $\mathbf{D}_{\mathrm{st}}(\rho_{\mathfrak{p}})$ is semistable at 1 and p^{-1} .
- (GB1) $H_f^1(\mathbf{Q}, \mathrm{Ind}_F^{\mathbf{Q}} \rho) = 0 = H_f^1(\mathbf{Q}, \mathrm{Ind}_F^{\mathbf{Q}} \rho^\vee(1))$.
- (GB2) $H^0(\mathrm{Gal}_{\mathbf{Q}_S}, \rho) = 0 = H^0(\mathrm{Gal}_{\mathbf{Q}_S}, \rho^\vee(1))$.
- (GB3) The associated (φ, Γ) -module $\mathbf{D}_{\mathrm{rig}}^\dagger((\mathrm{Ind}_F^{\mathbf{Q}} \rho)_p) = \bigoplus_{\mathfrak{p}|p} \mathbf{D}_{\mathrm{rig}}^\dagger(\rho_{\mathfrak{p}})$ has no saturated subquotient isomorphic to $U_{k,m}$ with $k \geq 1$ and $m \geq 0$ ([2, §2.1.2]).

For every $\mathfrak{p}|p$, choose a regular submodule $D_{\mathfrak{p}} \subset \mathbf{D}_{\text{st}}(\rho_{\mathfrak{p}})$. Then, $D := \bigoplus_{\mathfrak{p}|p} D_{\mathfrak{p}} \subset \bigoplus_{\mathfrak{p}|p} \mathbf{D}_{\text{st}}(\rho_{\mathfrak{p}}) = \mathbf{D}_{\text{rig}}^{\dagger}((\text{Ind}_F^{\mathbf{Q}} \rho)_{\rho})$ is a regular submodule.⁷ Moreover, if W_0, M_0, M_1 (resp., $W_{\mathfrak{p},0}, M_{\mathfrak{p},0}, M_{\mathfrak{p},1}$) are the corresponding subquotients of $\mathbf{D}_{\text{rig}}^{\dagger}((\text{Ind}_F^{\mathbf{Q}} \rho)_{\rho})$ (resp., $\mathbf{D}_{\text{rig}}^{\dagger}(\rho_{\mathfrak{p}})$) with respect to D (resp., $D_{\mathfrak{p}}$), then we have decompositions

$$W_0 = \bigoplus_{\mathfrak{p}|p} W_{\mathfrak{p},0}, \quad M_0 = \bigoplus_{\mathfrak{p}|p} M_{\mathfrak{p},0}, \quad M_1 = \bigoplus_{\mathfrak{p}|p} M_{\mathfrak{p},1}.$$

Hence, by assuming

(GB4) $W_{\mathfrak{p},0} = 0$ for every $\mathfrak{p}|p$ and the Hodge–Tate weights for $\text{Gr}_1^{\text{GB}} \mathbf{D}_{\text{rig}}^{\dagger}((\text{Ind}_F^{\mathbf{Q}} \rho)_{\rho})$ are all positive,

we may then follow the same recipe and define the *Greenberg–Benois \mathcal{L} -invariant* attached to ρ (with respect to $\{D_{\mathfrak{p}}\}_{\mathfrak{p}|p}$)

$$\mathcal{L}_{\text{GB}}(\rho) = \mathcal{L}_{\text{GB}}(\rho, \{D_{\mathfrak{p}}\}_{\mathfrak{p}|p}) := \mathcal{L}_{\text{GB}}(\text{Ind}_F^{\mathbf{Q}} \rho, D) \in E.$$

4 Fontaine–Mazur \mathcal{L} -invariants

To define the Fontaine–Mazur \mathcal{L} -invariants, we fix a finite extension K over \mathbf{Q}_p . We shall be considering Galois representations

$$\rho : \text{Gal}_K \rightarrow \text{GL}_n(E),$$

where E is (again) a finite extension of \mathbf{Q}_p . In what follows, we consider the (φ, N) -module $\mathbf{D}_{\text{st}}(\rho)$ associated with ρ . Note that if K_0 is the maximal unramified extension of \mathbf{Q}_p in K , then $\mathbf{D}_{\text{st}}(\rho)$ is a priori a K_0 -vector space. However, we shall linearize everything by base change to E .

Let q be the order of the residue field of K . We further assume ρ enjoys the following properties:

- (B1) The representation ρ is semistable with Hodge–Tate weights $k_{\sigma,1} \leq k_{\sigma,2} \leq \dots \leq k_{\sigma,n-1} \leq k_{\sigma,n}$, where $\sigma : K \hookrightarrow \overline{\mathbf{Q}_p}$. The Hodge–Tate weights give rise to the de Rham filtration $\text{Fil}_{\text{dR}}^{\bullet} \mathbf{D}_{\text{st}}(\rho) = [\text{Fil}_{\text{dR}}^{k_{\sigma,1}} \mathbf{D}_{\text{st}}(\rho) \supset \text{Fil}_{\text{dR}}^{k_{\sigma,2}} \mathbf{D}_{\text{st}}(\rho) \supset \dots \supset \text{Fil}_{\text{dR}}^{k_{\sigma,n}} \mathbf{D}_{\text{st}}(\rho)]$.
- (B2) The linearized Frobenius eigenvalues on $\mathbf{D}_{\text{st}}(\rho)$ are q^m, \dots, q^{m-n+1} .
- (FM1) Let $D_{(\varphi,N)}^{(i)}$ be the eigenspace in $\mathbf{D}_{\text{st}}(\rho)$ on which the Frobenius acts via q^{m-i} , and we assume that the induced monodromy operator N on $D_{(\varphi,N)}^{(i)}$ gives an isomorphism

$$N : D_{(\varphi,N)}^{(i)} \rightarrow D_{(\varphi,N)}^{(i+1)}.$$

⁷Note that $\mathbf{D}_{\text{st}}(\text{Ind}_{F_{\mathfrak{p}}}^{\mathbf{Q}_p} \rho_{\mathfrak{p}})$ is nothing but $\mathbf{D}_{\text{st}}(\rho_{\mathfrak{p}})$ (a priori a $K_0 \otimes_{\mathbf{Q}_p} E$ -module) viewing as a E -vector space.

(FM2) Define Frobenius filtration $\text{Fil}_\bullet^\varphi \mathbf{D}_{\text{st}}(\rho)$ by $\text{Fil}_j^\varphi \mathbf{D}_{\text{st}}(\rho) := \sum_{i>n-1-j} D_{(\varphi,N)}^{(i)}$ and assume the orthogonality

$$\mathbf{D}_{\text{st}}(\rho) = \text{Fil}_{\text{dR}}^{k_\bullet, i} \mathbf{D}_{\text{st}}(\rho) \oplus \text{Fil}_i^\varphi \mathbf{D}_{\text{st}}(\rho).$$

Lemma 4.1. *Keep the notations and the assumptions as above. We abuse the notation and denote by $\text{Gr}_{\text{dR}}^{n-1} \mathbf{D}_{\text{st}}(\rho) := \text{Fil}_{\text{dR}}^{k_\bullet, n-1} \mathbf{D}_{\text{st}}(\rho) / \text{Fil}_{\text{dR}}^{k_\bullet, n} \mathbf{D}_{\text{st}}(\rho)$. Then, we have an inclusion*

$$\text{Gr}_{\text{dR}}^{n-1} \mathbf{D}_{\text{st}}(\rho) \hookrightarrow D_{(\varphi,N)}^{(0)} \oplus D_{(\varphi,N)}^{(1)}.$$

Proof Indeed, we have a sequence of identifications

$$\begin{aligned} D_{(\varphi,N)}^{(0)} \oplus D_{(\varphi,N)}^{(1)} &= \frac{\text{Fil}_n^\varphi \mathbf{D}_{\text{st}}(\rho)}{\text{Fil}_{n-2}^\varphi \mathbf{D}_{\text{st}}(\rho)} \\ &= \frac{\text{Fil}_n^\varphi \mathbf{D}_{\text{st}}(\rho) \oplus \text{Fil}_{\text{dR}}^{k_\bullet, n-2} \mathbf{D}_{\text{st}}(\rho)}{\text{Fil}_{n-2}^\varphi \mathbf{D}_{\text{st}}(\rho) \oplus \text{Fil}_{\text{dR}}^{k_\bullet, n-2} \mathbf{D}_{\text{st}}(\rho)} \\ &= \frac{\text{Fil}_n^\varphi \mathbf{D}_{\text{st}}(\rho) \oplus \text{Fil}_{\text{dR}}^{k_\bullet, n-2} \mathbf{D}_{\text{st}}(\rho)}{\mathbf{D}_{\text{st}}(\rho)} \\ &= \frac{\text{Fil}_n^\varphi \mathbf{D}_{\text{st}}(\rho) \oplus \text{Fil}_{\text{dR}}^{k_\bullet, n-2} \mathbf{D}_{\text{st}}(\rho)}{\text{Fil}_n^\varphi \mathbf{D}_{\text{st}}(\rho) \oplus \text{Fil}_{\text{dR}}^{k_\bullet, n} \mathbf{D}_{\text{st}}(\rho)} \\ &= \frac{\text{Fil}_{\text{dR}}^{k_\bullet, n-2} \mathbf{D}_{\text{st}}(\rho)}{\text{Fil}_{\text{dR}}^{k_\bullet, n} \mathbf{D}_{\text{st}}(\rho)}, \end{aligned}$$

where the third and the fourth identifications follow from the orthogonality assumption. ■

Lemma 4.2. *For every i , we have*

$$\text{rank}_{K_0 \otimes_{\mathbb{Q}_p} E} D_{(\varphi,N)}^{(i)} = 1.$$

Moreover, $m < k_{\sigma,n}$ for every $\sigma : K \hookrightarrow \overline{\mathbb{Q}_p}$.

Proof Consider the twisted Galois representation $\rho(m)$. One can similarly define the Frobenius filtration $\text{Fil}_\bullet^\varphi \mathbf{D}_{\text{st}}(\rho(m))$, and we denote by $D_{(\varphi,N)}^{(i)}(m)$ the graded pieces. Since each $\text{Fil}_i^\varphi \mathbf{D}_{\text{st}}(\rho(m))$ is a (φ, N) -module, [2, Proposition 1.2.7 (ii)] implies that we have an associated filtration $\text{Fil}_\bullet \mathbf{D}_{\text{rig}}^\dagger(\rho(m))$ such that $\mathcal{D}_{\text{st}}(\text{Fil}_\bullet \mathbf{D}_{\text{rig}}^\dagger(\rho(m))) = \text{Fil}_i^\varphi \mathbf{D}_{\text{st}}(\rho(m))$.

Consider $\text{Gr}_n \mathbf{D}_{\text{rig}}^\dagger(\rho(m))$. One sees that

$$\mathcal{D}_{\text{st}}(\text{Gr}_n \mathbf{D}_{\text{rig}}^\dagger(\rho(m))) = D_{(\varphi,N)}^{(0)}(m),$$

on which the semistable Frobenius acts via 1. Hence, by [17, Proposition 2.4], $\text{Gr}_n \mathbf{D}_{\text{rig}}^\dagger(\rho(m))$ is crystalline and

$$\mathrm{Gr}_n \mathbf{D}_{\mathrm{rig}}^\dagger(\rho(m)) \cong \mathcal{R}_{K,E}(\delta) \quad \text{with} \quad \delta(z) = \prod_{\sigma:K \rightarrow \overline{\mathbf{Q}}_p} \sigma(z)^{-k_{\sigma,n}+m}.$$

This shows that $\mathrm{rank}_{K_0 \otimes_{\mathbf{Q}_p} E} D_{(\varphi,N)}^{(0)}(m) = 1$. Using the formula in *loc. cit.*, one also sees that $k_{\sigma,n} > m$.

Since $\mathrm{rank}_{K_0 \otimes_{\mathbf{Q}_p} E} D_{(\varphi,N)}^{(0)}(m) = 1$, we see that $\mathrm{rank}_{K_0 \otimes_{\mathbf{Q}_p} E} D_{(\varphi,N)}^{(0)} = 1$. The result then can be concluded by applying (FM1). ■

Thanks to Lemma 4.1 and Lemma 4.2, we can now define the *Fontaine–Mazur \mathcal{L} -invariant*. Let v_0 be a $K_0 \otimes_{\mathbf{Q}_p} E$ -basis for $D_{(\varphi,N)}^{(0)}$ and let $v_1 := Nv_0$, which is a $K_0 \otimes_{\mathbf{Q}_p} E$ -basis for $D_{(\varphi,N)}^{(1)}$. The *Fontaine–Mazur \mathcal{L} -invariant* attached to ρ is then defined to be $\mathcal{L}_{\mathrm{FM}}(\rho) \in K_0 \otimes_{\mathbf{Q}_p} E$ such that

$$v_0 - \mathcal{L}_{\mathrm{FM}}(\rho)v_1 \in \mathrm{Gr}_{\mathrm{dR}}^{n-1} \mathbf{D}_{\mathrm{st}}(\rho).$$

Remark 4.3. In fact, if we write $\mathrm{Gr}_{\mathrm{dR}}^i \mathbf{D}_{\mathrm{st}}(\rho) := \mathrm{Fil}_{\mathrm{dR}}^{k_i} \mathbf{D}_{\mathrm{st}}(\rho) / \mathrm{Fil}_{\mathrm{dR}}^{k_{i+1}} \mathbf{D}_{\mathrm{st}}(\rho)$, then a similar argument as in Lemma 4.1 shows that

$$\mathrm{Gr}_{\mathrm{dR}}^{n-i} \mathbf{D}_{\mathrm{st}}(\rho) \hookrightarrow D_{(\varphi,N)}^{(i-1)} \oplus D_{(\varphi,N)}^{(i)}.$$

By using this inclusion, one can similarly define the *i -th Fontaine–Mazur \mathcal{L} -operator* attached to ρ to be $\mathcal{L}_{\mathrm{FM}}^{(i)}(\rho) \in K_0 \otimes_{\mathbf{Q}_p} E$ such that $v_{i-1} - \mathcal{L}_{\mathrm{FM}}^{(i)}(\rho)v_i \in \mathrm{Gr}_{\mathrm{dR}}^{n-i} \mathbf{D}_{\mathrm{st}}(\rho)$, where $v_j = N^j v$. Such a strategy was taken in [1]. However, it is believed that $\mathcal{L}_{\mathrm{FM}}^{(0)}(\rho) = \mathcal{L}_{\mathrm{FM}}(\rho)$ should determine all the other $\mathcal{L}_{\mathrm{FM}}^{(i)}(\rho)$'s (see, for example, [op. cit., §4.3.2]). Hence, we focus on $\mathcal{L}_{\mathrm{FM}}(\rho)$. Moreover, one shall see, in what follows, that it is $\mathcal{L}_{\mathrm{FM}}(\rho)$ we can relate to Greenberg–Benois \mathcal{L} -invariants.

5 Comparing the two \mathcal{L} -invariants

The aim of this section is to prove the comparison theorem (Theorem 5.4). However, as aforementioned, to define \mathcal{L} -invariants, there are some constraints one needs to put on the Galois representations. For reader's convenience, we collect all the assumptions in §5.1 and briefly discuss a folklore about these assumptions.

5.1 Assumptions on the Galois representation

Let F be a number field and let E be a finite extension of \mathbf{Q}_p such that, for every prime ideal \mathfrak{p} in \mathcal{O}_F sitting above p , $F_{\mathfrak{p}} \subset E$. Write $F_{\mathfrak{p},0}$ for the maximal unramified extension of \mathbf{Q}_p in $F_{\mathfrak{p}}$; we further assume that $F_{\mathfrak{p},0} = \mathbf{Q}_p$ for every \mathfrak{p} . Suppose we are given a Galois representation

$$\rho : \mathrm{Gal}_F \rightarrow \mathrm{GL}_n(E)$$

that is unramified outside a finite set of places. Let S be the set of places in F such that ρ ramifies. We make the following assumptions:

- (I) Basic assumptions:
 - (B1) For any prime ideal $\mathfrak{p} \subset \mathcal{O}_F$ sitting above p , $\rho_{\mathfrak{p}} := \rho|_{\text{Gal}_{F_{\mathfrak{p}}}}$ is semistable with Hodge–Tate weights $0 \leq k_{\mathfrak{p},\sigma,1} \leq k_{\mathfrak{p},\sigma,2} \leq \dots \leq k_{\mathfrak{p},\sigma,n-1} \leq k_{\mathfrak{p},\sigma,n}$, where $\sigma : F_{\mathfrak{p}} \hookrightarrow \mathbf{Q}_p$.
 - (B2) For any prime ideal $\mathfrak{p} \subset \mathcal{O}_F$ sitting above p , the Frobenius eigenvalues on $\mathbf{D}_{\text{st}}(\rho_{\mathfrak{p}})$ are p^m, \dots, p^{m-n+1} such that $k_{\mathfrak{p},\sigma,n} > m > k_{\mathfrak{p},\sigma,n-1}$, where the first inequality is always guaranteed by Lemma 4.2.⁸
- (II) Fontaine–Mazur assumptions:
 - (FM1) For any $\mathfrak{p}|p$, let $D_{\mathfrak{p},(\varphi,N)}^{(i)}$ be the eigenspace in $\mathbf{D}_{\text{st}}(\rho_{\mathfrak{p}})$ on which the Frobenius acts via p^{m-i} . We assume that the induced monodromy operator N on $D_{\mathfrak{p},(\varphi,N)}^{(i)}$ gives an isomorphism

$$N : D_{\mathfrak{p},(\varphi,N)}^{(i)} \rightarrow D_{\mathfrak{p},(\varphi,N)}^{(i+1)}.$$
 - (FM2) Define $\text{Fil}_j^{\varphi} \mathbf{D}_{\text{st}}(\rho_{\mathfrak{p}}) := \sum_{i>n-1-j} D_{\mathfrak{p},(\varphi,N)}^{(i)}$, and we call the ascending filtration $\text{Fil}_{\bullet}^{\varphi} \mathbf{D}_{\text{st}}(\rho_{\mathfrak{p}})$ the *Frobenius filtration* on $\mathbf{D}_{\text{st}}(\rho_{\mathfrak{p}})$. We assume the orthogonality

$$\mathbf{D}_{\text{st}}(\rho_{\mathfrak{p}}) = \text{Fil}_{\text{dR}}^{k_{\mathfrak{p},\bullet,i}} \mathbf{D}_{\text{st}}(\rho_{\mathfrak{p}}) \oplus \text{Fil}_i^{\varphi} \mathbf{D}_{\text{st}}(\rho_{\mathfrak{p}}).$$
- (III) Greenberg–Benois assumptions:
 - (GB1) Vanishing of the Bloch–Kato Selmer groups

$$H_f^1(\mathbf{Q}, \text{Ind}_F^{\mathbf{Q}} \rho(m)) = H_f^1(\mathbf{Q}, \text{Ind}_F^{\mathbf{Q}} \rho^{\vee}(1-m)) = 0.$$
 - (GB2) Vanishing of the zero-degree Galois cohomology

$$H^0(\text{Gal}_{\mathbf{Q}_S}, \text{Ind}_F^{\mathbf{Q}} \rho(m)) = H^0(\text{Gal}_{\mathbf{Q}_S}, \text{Ind}_F^{\mathbf{Q}} \rho^{\vee}(1-m)) = 0.$$
 - (GB3) The associated (φ, Γ) -module $\mathbf{D}_{\text{rig}}^{\dagger}((\text{Ind}_F^{\mathbf{Q}} \rho(m))_{\mathfrak{p}}) = \bigoplus_{\mathfrak{p}|p} \mathbf{D}_{\text{rig}}^{\dagger}(\rho_{\mathfrak{p}}(m))$ does not admit a subquotient of the form $U_{k,r}$ with $k \geq 1$ and $r \geq 0$ ([2, §2.1.2]).
 - (GB4) For any $\mathfrak{p}|p$, the space $W_{\mathfrak{p},0}$ for $\rho_{\mathfrak{p}}(m)$ vanishes (see [2, Proposition 2.1.7] or [17, pp. 1238]).

Remark 5.1. For every $\mathfrak{p}|p$, the Frobenius filtration $\text{Fil}_{\bullet}^{\varphi} \mathbf{D}_{\text{st}}(\rho_{\mathfrak{p}})$ defines a filtration $\text{Fil}_{\bullet} \mathbf{D}_{\text{rig}}^{\dagger}(\rho)$ (similar as in the proof of Lemma 4.2). One observes that the graded pieces $\text{Gr}_i \mathbf{D}_{\text{rig}}^{\dagger}(\rho)$ of this filtration are all of rank 1 over $\mathcal{R}_{F_{\mathfrak{p}},E}$ (by [2, Proposition 1.2.7 (ii)]). In particular, $\text{Fil}_{\bullet} \mathbf{D}_{\text{rig}}^{\dagger}(\rho)$ is a triangulation of $\mathbf{D}_{\text{rig}}^{\dagger}(\rho)$. In fact, we have the following description for the graded pieces:

$$\text{Gr}_i \mathbf{D}_{\text{rig}}^{\dagger}(\rho) = \mathcal{R}_{F_{\mathfrak{p}},E}(\delta_{n-i}),$$

⁸Here, m does not depend on \mathfrak{p} and σ .

where

$$\delta_{n-i}(z) = \left(\prod_{\sigma: F_p \rightarrow \overline{\mathbf{Q}}_p} \sigma(z)^{-k_{p,\sigma,i}} \right) |\mathrm{Nm}_{\mathbf{Q}_p} z|^{-(m-n+i)}.$$

Remark 5.2. For every $p|p$, since $k_{p,\sigma,n} > m > k_{p,\sigma,n-1}$, we see that $\mathrm{Fil}_{\mathrm{dR}}^{k_{p,\sigma,n-1}-m} \mathbf{D}_{\mathrm{st}}(\rho_p(m)) = \mathrm{Fil}_{\mathrm{dR}}^0 \mathbf{D}_{\mathrm{st}}(\rho_p(m))$. Moreover, the orthogonality condition (FM2) implies that $\mathrm{Fil}_{n-1}^\varphi \mathbf{D}_{\mathrm{st}}(\rho_p(m))$ is a regular (φ, N) -submodule of $\mathbf{D}_{\mathrm{st}}(\rho_p(m))$. Hence, in what follows, we naturally work with $D_p := \mathrm{Fil}_{n-1}^\varphi \mathbf{D}_{\mathrm{st}}(\rho_p(m)) \subset \mathbf{D}_{\mathrm{st}}(\rho_p(m))$ and $D = \bigoplus_{p|p} D_p \subset \bigoplus_{p|p} \mathbf{D}_{\mathrm{st}}(\rho_p(m))$. Moreover, in our situation, we shall see in the proof (e.g., (5.2)) that the corresponding $\mathrm{Gr}_1^{\mathrm{GB}} \mathbf{D}_{\mathrm{rig}}^\dagger((\mathrm{Ind}_F^{\mathbf{Q}} \rho)_p)$ has positive Hodge–Tate weights, and so we remove such assumption in (GB4).

Remark 5.3. We have many assumptions on our Galois representation ρ . On the one hand, one sees that they are necessary in order to attach both $\mathcal{L}_{\mathrm{GB}}$ and $\mathcal{L}_{\mathrm{FM}}$ to it. On the other hand, we remark that it is a folklore that they shall appear as Galois representations for automorphic forms of unitary groups whose corresponding Shimura varieties can be p -adically uniformized by Drinfeld’s upper-half spaces. For example, we are requiring maximal monodromy on our Galois representations. Such a phenomenon is expected to appear for the Galois representations attached to unitary automorphic representations that are Steinberg at p .

5.2 The main theorem

Theorem 5.4. *Keep the notations and assumptions as above. We have an equality*

$$\mathcal{L}_{\mathrm{GB}}(\rho(m)) = \prod_{p|p} -\mathcal{L}_{\mathrm{FM}}(\rho_p).$$

Proof The proof of the theorem is similar to the proof of [2, Proposition 2.3.7], which relies on the following three steps:

Step 1. Fontaine–Mazur \mathcal{L} -invariants and cohomology of (φ, Γ) -modules.

Consider the triangulation $\mathrm{Fil}_\bullet \mathbf{D}_{\mathrm{rig}}^\dagger(\rho)$ in Remark 5.1. We define

$$\widetilde{W}_p := \mathrm{Fil}_n \mathbf{D}_{\mathrm{rig}}^\dagger(\rho) / \mathrm{Fil}_{n-2} \mathbf{D}_{\mathrm{rig}}^\dagger(\rho).$$

Hence, \widetilde{W}_p sits inside the short exact sequence

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathrm{Gr}_{n-1} \mathbf{D}_{\mathrm{rig}}^\dagger(\rho) & \longrightarrow & \widetilde{W}_p & \longrightarrow & \mathrm{Gr}_n \mathbf{D}_{\mathrm{rig}}^\dagger(\rho) \longrightarrow 0 \\ & & \downarrow \cong & & & & \downarrow \cong \\ & & \mathcal{R}_{F_p, E}(\delta_{p,1}) & & & & \mathcal{R}_{F_p, E}(\delta_{p,0}) \end{array}$$

for $\delta_{p,i} : F_p^\times \rightarrow E^\times$ described as in Remark 5.1.

As a result, \widetilde{W}_p defines a class

$$\begin{aligned} \text{cl}(\widetilde{W}_p) &\in \text{Ext}_{(\varphi, \Gamma)}(\mathcal{R}_{F_p, E}(\delta_{p,0}), \mathcal{R}_{F_p, E}(\delta_{p,1})) \\ &\cong \text{Ext}_{(\varphi, \Gamma)}(\mathcal{R}, \mathcal{R}(\delta_{p,1}\delta_{p,0}^{-1})) \\ &\cong H^1(\mathcal{R}_{F_p, E}(\delta_{p,1}\delta_{p,0}^{-1})). \end{aligned}$$

However, by construction, we know that \widetilde{W}_p is semistable (since $\mathcal{D}_{\text{st}}(\widetilde{W}_p) = D_{p,(\varphi, N)}^{(0)} \oplus D_{p,(\varphi, N)}^{(1)}$), and so $\text{cl}(\widetilde{W}_p) \in H_{\text{st}}^1(\mathcal{R}_{F_p, E}(\delta_{p,1}\delta_{p,0}^{-1}))$. Recall that $H_{\text{st}}^1(\mathcal{R}_{F_p, E}(\delta_{p,1}\delta_{p,0}^{-1}))$ can be computed via the complex $C_{\text{st}}^\bullet(\mathcal{R}_{F_p, E}(\delta_{p,1}\delta_{p,0}^{-1}))$ with

$$\begin{array}{ccc} \mathcal{D}_{\text{st}}(\mathcal{R}_{F_p, E}(\delta_{p,1}\delta_{p,0}^{-1})) & \rightarrow & \frac{\mathcal{D}_{\text{st}}(\mathcal{R}_{F_p, E}(\delta_{p,1}\delta_{p,0}^{-1}))}{\text{Fil}_{\text{dR}}^0 \mathcal{D}_{\text{st}}(\mathcal{R}_{F_p, E}(\delta_{p,1}\delta_{p,0}^{-1}))} \oplus \mathcal{D}_{\text{st}}(\mathcal{R}_{F_p, E}(\delta_{p,1}\delta_{p,0}^{-1})) \oplus \mathcal{D}_{\text{st}}(\mathcal{R}_{F_p, E}(\delta_{p,1}\delta_{p,0}^{-1})) \\ & & \downarrow \\ & & \mathcal{D}_{\text{st}}(\mathcal{R}_{F_p, E}(\delta_{p,1}\delta_{p,0}^{-1})) \end{array},$$

where the first map is given by $a \mapsto (a \bmod \text{Fil}_{\text{dR}}^0 \mathcal{D}_{\text{st}}(\mathcal{R}_{F_p, E}(\delta_{p,1}\delta_{p,0}^{-1})), (\varphi - 1)a, Na)$, while the second arrow is defined by $(a, b, c) \mapsto Nb - (p\varphi - 1)c$.

Now, choose a basis $v_{p,0} \in D_{p,(\varphi, N)}^{(0)}$ over $F_{p,0} \otimes_{\mathbb{Q}_p} E$ and let $v_{p,1} := Nv_{p,0}$, which is a $F_{p,0} \otimes_{\mathbb{Q}_p} E$ -basis for $D_{p,(\varphi, N)}^{(1)}$. We again denote by $v_{p,i}$ for the image of $v_{p,i}$ in $\mathcal{D}_{\text{st}}(\widetilde{W}_p(\delta_{p,0}^{-1}))$. By the proof of [2, Proposition 1.4.4 (ii)], we know that the class $\text{cl}(\widetilde{W}_p)$ in $H^1(C_{\text{st}}^\bullet(\mathcal{R}_{F_p, E}(\delta_{p,1}\delta_{p,0}^{-1})))$ is given by

$$\text{cl}(a, b, c) = \text{cl}(a, (\varphi - 1)v_{p,0}, Nv_{p,0}) = \text{cl}(a, 0, v_{p,1}),$$

where $a \in \mathcal{D}_{\text{st}}(\widetilde{W}_p(\delta_{p,0}^{-1}))$ such that $v_{p,0} + a \in \text{Fil}_{\text{dR}}^0 \mathcal{D}_{\text{st}}(\widetilde{W}_p(\delta_{p,0}^{-1}))$. After untwisting, a defines an element, still denoted by $a \in \mathcal{D}_{\text{st}}(\widetilde{W}_p)$ such that $v_{p,0} + a \in \text{Fil}_{\text{dR}}^{k_{p, \bullet, n}} \mathcal{D}_{\text{st}}(\widetilde{W}_p)$. However, by construction,

$$\text{Fil}_{\text{dR}}^{k_{p, \bullet, n}} \mathcal{D}_{\text{st}}(\widetilde{W}_p) = \text{Gr}_{\text{dR}}^{n-1} \mathbf{D}_{\text{st}}(\rho) \text{ (notation as in Lemma 4.1).}$$

Hence, we conclude that

$$(5.1) \quad \text{cl}(\widetilde{W}_p) = \text{cl}(-\mathcal{L}_{\text{FM}}(\rho_p)v_{p,1}, 0, v_{p,1}) \in H^1(C_{\text{st}}^\bullet(\mathcal{R}_{F_p, E}(\delta_{p,1}\delta_{p,0}^{-1}))).$$

Step 2. Computing $\mathcal{L}_{\text{GB}}(\rho)$.

Next, we would also like to compute the Greenberg–Benois \mathcal{L} -invariant $\mathcal{L}_{\text{GB}}(\rho)$ via cohomology of (φ, Γ) -modules. As before, because of the decomposition $\mathbf{D}_{\text{rig}}^\dagger((\text{Ind}_F^{\mathbb{Q}} \rho)_p) = \bigoplus_{p|p} \mathbf{D}_{\text{rig}}^\dagger(\text{Ind}_{F_p}^{\mathbb{Q}_p} \rho_p)$, we can study each p individually. Hence, fix $p|p$. Computing the five-step filtration (3.1) explicitly, we have

$$D_{p,i}^{\text{GB}} = \begin{cases} 0, & i = -2, \\ \text{Fil}_{n-2}^\varphi \mathbf{D}_{\text{st}}(\rho_p(m)), & i = -1, \\ D_p, & i = 0 \\ \mathbf{D}_{\text{st}}(\rho_p(m)), & i = 1, \\ \mathbf{D}_{\text{st}}(\rho_p(m)), & i = 2, \end{cases}$$

which gives rise to a five-step filtration $\text{Fil}_{\bullet}^{\text{GB}} \mathbf{D}_{\text{rig}}^\dagger(\rho_p(m))$.

Let us simplify the notation and write

$$W_p = \text{Fil}_1^{\text{GB}} \mathbf{D}_{\text{rig}}^\dagger(\rho_p(m)) / \text{Fil}_{-1}^{\text{GB}} \mathbf{D}_{\text{rig}}^\dagger(\rho_p(m)).$$

Similar as before, we see that W_p sits inside the short exact sequence

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \text{Gr}_0^{\text{GB}} \mathbf{D}_{\text{rig}}^\dagger(\rho_p(m)) & \longrightarrow & W_p & \longrightarrow & \text{Gr}_1^{\text{GB}} \mathbf{D}_{\text{rig}}^\dagger(\rho_p(m)) \longrightarrow 0 \\
 & & \downarrow \cong & & & & \downarrow \cong \\
 & & \mathcal{R}_{F_p, E}(\delta'_{p,1}) & & & & \mathcal{R}_{F_p, E}(\delta'_{p,0})
 \end{array} \tag{5.2}$$

where

$$\delta'_{p, n-i} = \delta_{p, n-i} \left(\text{Nm}_{\mathbb{Q}_p}^{F_p} z \right)^m |\text{Nm}_{\mathbb{Q}_p}^{F_p} z|^m = \left(\prod_{\sigma: F_p \hookrightarrow \overline{\mathbb{Q}_p}} \sigma(z)^{-k_{p, \sigma, i+m}} \right) |\text{Nm}_{\mathbb{Q}_p}^{F_p} z|^{n-i}.$$

By taking cohomology, we have the connecting homomorphism

$$\partial : H^0(\mathcal{R}_{F_p, E}(\delta'_{p,0})) \rightarrow H^1(\mathcal{R}_{F_p, E}(\delta'_{p,1})) = H_{\text{st}}^1(\mathcal{R}_{F_p, E}(\delta'_{p,1})),$$

where the equation follows from Lemma 2.1. Denoted by $\alpha_{\delta'_{p,1}}^*$ and $\beta_{\delta'_{p,1}}^*$ the two classes in $H^1(\mathcal{R}_{F_p, E}(\delta'_{p,1}))$ in Lemma 2.2. We know from *loc. cit.* that ∂ gives rise to a unique number $\mathcal{L}(W_p) \in E$ such that

$$\beta_{\delta'_{p,1}}^* + \mathcal{L}(W_p) \alpha_{\delta'_{p,1}}^* \in \text{image } \partial.$$

We claim that

$$\mathcal{L}_{\text{GB}}(\rho(m)) = \prod_{p|\rho} \mathcal{L}(W_p). \tag{5.3}$$

Note that, in the definition of $\mathcal{L}_{\text{GB}}(\rho(m))$, one studies the cohomology of $\text{Gr}_1^{\text{GB}} \mathbf{D}_{\text{rig}}^\dagger(\rho_p(m))$. However, we are now having cohomology classes in $H^1(\text{Gr}_0^{\text{GB}} \mathbf{D}_{\text{rig}}^\dagger(\rho_p(m)))$. To resolve this, we look at the short exact sequence

$$\begin{array}{ccccccc}
 0 & \rightarrow & (\text{Gr}_1^{\text{GB}} \mathbf{D}_{\text{rig}}(\rho_p(m)))^\vee & (\chi_{\text{cyc}}) & \rightarrow & W_p^\vee & (\chi_{\text{cyc}}) & \rightarrow & (\text{Gr}_0^{\text{GB}} \mathbf{D}_{\text{rig}}(\rho_p(m)))^\vee & (\chi_{\text{cyc}}) & \rightarrow & 0 \\
 & & \downarrow \cong & & & & & & \downarrow \cong & & & & \\
 & & \mathcal{R}_{F_p, E}(\kappa_{p,0}) & & & & & & \mathcal{R}_{F_p, E}(\kappa_{p,1}) & & & &
 \end{array}$$

By [2, Proposition 2.2.4], the Greenberg–Benois \mathcal{L} -invariant computed by this exact sequence (at each p) is the same as $\mathcal{L}_{\text{GB}}(\rho(m))$. Here,

$$\kappa_{p,i}(z) = \left(\prod_{\sigma: F_p \hookrightarrow \overline{\mathbb{Q}_p}} \sigma(z)^{k_{p, \sigma, n-i-m+1}} \right) |\text{Nm}_{\mathbb{Q}_p}^{F_p} z|^{1-i},$$

and we want to compute $\mathcal{L}_{\text{GB}}(\rho(m))$ using the cohomology of $\mathcal{R}_{F_p, E}(\kappa_{p,1})$.

By (B2), we have $k_{p,\sigma,n-1} - m + 1 \leq 0$ and we let $u_p := \min_{\sigma} \{k_{p,\sigma,n-1} - m + 1\}$. By [17, (2.8)], there is an injection

$$H^1(\mathcal{R}_{Q_p,E}(z^{u_p})) \hookrightarrow H^1(\mathcal{R}_{F_p,E}(\kappa_{p,1})), \quad \begin{matrix} x_{u_p} \mapsto x_{k_{p,\bullet,n-1-m+1}} \\ y_{u_p} \mapsto y_{k_{p,\bullet,n-1-m+1}} \end{matrix},$$

where x_{u_p} , $x_{k_{p,\bullet,n-1-m+1}}$, y_{u_p} , and $y_{k_{p,\bullet,n-1-m+1}}$ are as defined in *loc. cit.*^{9, 10} By the discussion on [17, pp. 1238], we have a commutative diagram

$$(5.4) \quad \begin{array}{ccccc} & & H^1\left(\bigoplus_{p|p} D_p^{\vee}(1-m), \text{Ind}_F^Q \rho^{\vee}(1-m)\right) & & \\ & \swarrow \iota_f & \downarrow & \searrow \iota_c & \\ \bigoplus_{p|p} Ex_{k_{p,\bullet,n-1-m+1}} & \hookrightarrow & \bigoplus_{p|p} H^1(\mathcal{R}_{F_p,E}(\kappa_{p,1})) & \longleftarrow & \bigoplus_{p|p} Ey_{k_{p,\bullet,n-1-m+1}} \end{array}$$

where ι_c is an isomorphism. Moreover, [op. cit., Corollary 3.9] yields that

$$\mathcal{L}_{GB}(\rho(m)) = \det(\iota_f \circ \iota_c^{-1}).$$

In particular, if $\mathcal{L}_p \in E$ such that

$$\mathcal{L}_p x_{k_{p,\bullet,n-1-m+1}} + y_{k_{p,\bullet,n-1-m+1}} \in \text{image} \left(H^1\left(\bigoplus_{p|p} D_p^{\vee}(1-m), \text{Ind}_F^Q \rho^{\vee}(1-m)\right) \rightarrow H^1(\mathcal{R}_{F_p,E}(\kappa_{p,1})) \right),$$

then

$$\mathcal{L}_{GB}(\rho(m)) = \prod_{p|p} \mathcal{L}_p.$$

By definition, $H^1\left(\bigoplus_{p|p} D_p^{\vee}(1-m), \text{Ind}_F^Q \rho^{\vee}(1-m)\right) \cong \bigoplus_{p|p} \frac{H^1(W_p^{\vee}(\chi_{\text{cyc}}))}{H_f^1(W_p^{\vee}(\chi_{\text{cyc}}))}$. The vertical morphism in (5.4) is compatible with the natural morphism

$$H^1(W_p^{\vee}(\chi_{\text{cyc}})) \rightarrow H^1(\mathcal{R}_{F_p,E}(\kappa_{p,1})),$$

⁹For the convenience of the readers, we briefly recall the definitions of x_{u_p} and y_{u_p} . The definitions for $x_{k_{p,\bullet,n-1-m+1}}$ and $y_{k_{p,\bullet,n-1-m+1}}$ are similar; we refer the readers to [17, pp. 1233, 1234] for the precise definitions. Given $\alpha = (a, b) \in \mathcal{R}_{Q_p,E}(z^{u_p})^{\oplus 2}$, one can define an extension D_{α} as in (2.1), which defines a class $\text{cl}(a, b) \in H^1(\mathcal{R}_{Q_p,E}(z^{u_p}))$. We simplify the notation and write e for the basis for $\mathcal{R}_{Q_p,E}(z^{u_p})$. Then $x_{u_p} := \text{cl}(t^{-u_p} e, 0)$ and $y_{u_p} = \log \chi_{\text{cyc}}(\gamma) \text{cl}(0, t^{-u_p} e)$, where recall γ is a (fixed) topological generator for $\Gamma = \text{Gal}(\mathbb{Q}_p(\zeta_{p^{\infty}})/\mathbb{Q}_p)$ and $t = \log[\varepsilon] \in B_{\text{dR}}^+$. Here, $\varepsilon \in \mathcal{O}_{C_p}^b = \varprojlim_{a \rightarrow a^p} \mathcal{O}_{C_p}$ is defined by the fixed compatible system of primitive p -power roots of unity (see the beginning of §2.1), and we implicitly use the fact that certain subring of \mathcal{R}_{Q_p} can be embedded into B_{dR}^+ (see [2, §1.2.2]).

¹⁰This injection comes from a natural injection $\mathcal{R}_{Q_p,E}(z^{u_p}) \hookrightarrow \text{Ind}_{\Gamma_p}^{\Gamma} \mathcal{R}_{F_p,E}(\kappa_{p,1})$. By duality, we have a natural projection $\text{Ind}_{\Gamma_p}^{\Gamma} \mathcal{R}_{F_p,E}(\delta'_{p,1}) \twoheadrightarrow \mathcal{R}_{Q_p,R}(z^{-u_p+1}|z|)$ as well as $H^1(\mathcal{R}_{F_p,E}(\delta'_{p,1})) \twoheadrightarrow H^1(\mathcal{R}_{Q_p,E}(z^{-u_p+1}|z|))$. According to [2, Theorem 1.5.7], x_{u_p} (resp. y_{u_p}) is dual to $\beta_{-u_p+1}^*$ (resp. $-\alpha_{-u_p+1}^*$). Thus, we may choose $v_{\delta'_{p,1}} \in \mathcal{D}_{\text{st}}(\mathcal{R}_{F_p,E}(\delta'_{p,1}))$ such that the corresponding $\alpha_{\delta'_{p,1}}^* = \text{cl}(v_{\delta'_{p,1}}, 0, 0) \mapsto \alpha_{-u_p+1}^*$ and $\beta_{\delta'_{p,1}}^* = -\text{cl}(0, 0, v_{\delta'_{p,1}}) \mapsto \beta_{-u_p+1}^*$.

induced from the short exact sequence (5.2). Note that the exact sequence

$$H^1(W_p^\vee(\chi_{\text{cyc}})) \rightarrow H^1(\mathcal{R}_{F_p,E}(\kappa_{p,1})) \xrightarrow{\partial} H^2(\mathcal{R}_{F_p,E}(\kappa_{p,0}))$$

is dual to the exact sequence

$$H^0(\mathcal{R}_{F_p,E}(\delta'_{p,0})) \xrightarrow{\partial} H^1(\mathcal{R}_{F_p,E}(\delta'_{p,1})) \rightarrow H^1(W_p).$$

We have $\partial(\mathcal{L}_p x_{k_p, \bullet, n-1-m+1} + y_{k_p, \bullet, n-1-m+1}) = 0 \in H^2(\mathcal{R}_{F_p,E}(\kappa_{p,0}))$ and $\beta_{\delta'_{p,1}}^* + \mathcal{L}(W_p)\alpha_{\delta'_{p,1}}^* = 0 \in H^1(W_p)$. Moreover, using the relation between $x_{k_p, \bullet, n-1-m+1}$ (resp., $y_{k_p, \bullet, n-1-m+1}$) and $\beta_{\delta'_{p,1}}^*$ (resp., $-\alpha_{\delta'_{p,1}}^*$), one sees that

$$\mathcal{L}_p = \mathcal{L}(W_p),$$

which concludes our claim.

Step 3. Conclusion.

By construction, W_p defines a class (see (5.2))

$$\begin{aligned} \text{cl}(W_p) &\in \text{Ext}_{(\varphi, \Gamma)}^1(\mathcal{R}_{F_p,E}(\delta'_{p,0}), \mathcal{R}_{F_p,E}(\delta'_{p,1})) \\ &\cong \text{Ext}_{(\varphi, \Gamma)}^1(\mathcal{R}_{F_p,E}, \mathcal{R}_{F_p,E}(\delta'_{p,1}\delta_{p,0}^{-1})) \\ &\cong H^1(\mathcal{R}_{F_p,E}(\delta_{p,1}\delta_{p,0}^{-1})). \end{aligned}$$

Note that, as classes in $H^1(\mathcal{R}_{F_p,E}(\delta_{p,1}\delta_{p,0}^{-1}))$, we have

$$\text{cl}(W_p) = \text{cl}(\tilde{W}_p).$$

Unwinding everything, we have

$$c \text{cl}(\mathcal{L}(W_p)v_{p,1}, 0, -v_{p,1}) = \text{cl}(W_p) = \text{cl}(\tilde{W}_p) \stackrel{(5.1)}{=} \text{cl}(-\mathcal{L}_{\text{FM}}(\rho_p)v_{p,1}, 0, v_{p,1})$$

for some $c \in E$. In particular, we conclude that

$$\mathcal{L}_{\text{FM}}(\rho_p) = -\mathcal{L}(W_p)$$

and so,

$$\mathcal{L}_{\text{GB}}(\rho(m)) = \prod_{p|m} -\mathcal{L}_{\text{FM}}(\rho_p)$$

by (5.3). ■

Acknowledgements This paper grew from a working group with Ting-Han Huang, Martí Roset Julià, and Giovanni Rosso, and I would like to thank them for interesting discussions. I especially thank Giovanni Rosso for valuable feedback regarding the early draft of this paper. I also thank David Loeffler for pointing out a mistake in an early version of this paper. I would also like to thank Muhammad Manji for interesting discussions on (φ, Γ) -modules. Part of the work was done when I was visiting National Center for Theoretical Sciences in Taipei; I would like to thank the hospitality of the institute and Ming-Lun Hsieh. Finally, I thank the anonymous referees for their corrections and valuable suggestions, which helped to improve the

exposition of the article. This work is supported by the ERC Consolidator grant “Shimura varieties and the BSD conjecture” and the Irish Research Council under grant number IRCLA/2023/849 (HighCritical).

References

- [1] A. Besser and E. de Shalit, \mathcal{L} -invariants of p -adically uniformized varieties. *Ann. Math. Qué.* 40(2016), 29–54.
- [2] D. Benois, *A generalization of Greenberg’s \mathcal{L} -invariant*. *Amer. J. Math.* 133(2011), no. 6, 1573–1632.
- [3] L. Berger, *Représentations p -adiques et équations différentielles*. *Invent. Math.* 148(2002), no. 2, 219–284.
- [4] R. Coleman and A. Iovita, *Hidden structures on semistable curves*. *Astérisque* 331(2010), 179–254.
- [5] R. Coleman, *A p -adic Shimura isomorphism and p -adic periods of modular forms*. In p -adic monodromy and the Birch and Swinnerton-Dyer conjecture (Boston, MA, 1991), *Contemp. Math.*, Vol. 165, Amer. Math. Soc., Providence, RI, 1994, 21–51.
- [6] P. Deligne, *Valeurs de fonctions L et périodes d’intégrales*. In *Automorphic forms, representations and L -functions* (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 2, Proc. Sympos. Pure Math., XXXIII, with an appendix by N. Koblitz and A. Ogus, Amer. Math. Soc., Providence, RI, 1979, 313–346.
- [7] J.-M. Fontaine, *Le corps des périodes p -adiques*. In 223, with an appendix by Pierre Colmez, *Périodes p -adiques* (Bures-sur-Yvette, 1988), 1994, 59–111.
- [8] R. Greenberg, *Trivial zeros of p -adic L -functions*. In p -adic monodromy and the Birch and Swinnerton-Dyer conjecture (Boston, MA, 1991), *Contemp. Math.*, Vol. 165, Amer. Math. Soc., Providence, RI, 1994, 149–174.
- [9] R. Greenberg and G. Stevens, *On the conjecture of Mazur, Tate, and Teitelbaum*. In p -adic monodromy and the Birch and Swinnerton-Dyer conjecture (Boston, MA, 1991), *Contemp. Math.*, Vol. 165, Amer. Math. Soc., Providence, RI, 1994, 183–211.
- [10] H. Hida, *On a generalization of the conjecture of Mazur–Tate–Teitelbaum*. *Int. Math. Res. Not. IMRN* 23(2007), Art. ID rnm102, 49.
- [11] R. Harron and A. Jorza, *On symmetric power \mathcal{L} -invariants of Iwahori level Hilbert modular forms*. *Amer. J. Math.* 139(2017), no. 6, 1605–1647. issn: 0002-9327,1080-6377.
- [12] T.-H. Huang and Wu J.-F. *Finite polynomial cohomology with coefficients*. Preprint, 2024, <https://arxiv.org/abs/2209.10289>.
- [13] A. Iovita and M. Spieß, *Derivatives of p -adic L -functions, Heegner cycles and monodromy modules attached to modular forms*. *Invent. Math.* 154(2003), no. 2, 333–384.
- [14] B. Mazur, *On monodromy invariants occurring in global arithmetic, and Fontaine’s theory*. In p -adic monodromy and the Birch and Swinnerton-Dyer conjecture (Boston, MA, 1991), *Contemp. Math.*, Vol. 165, Amer. Math. Soc., Providence, RI, 1994, 1–20.
- [15] B. Mazur, J. Tate and J. Teitelbaum, *On p -adic analogues of the conjectures of Birch and Swinnerton-Dyer*. *Invent. Math.* 84(1986), no. 1, 1–48.
- [16] K. Nakamura, *Classification of two-dimensional split trianguline representations of p -adic fields*. *Compos. Math.* 145(2009), no. 4, 865–914.
- [17] G. Rosso, *\mathcal{L} -Invariant for Siegel–Hilbert forms*. *Doc. Math.* 20(2015), 1227–1253.
- [18] J. Teitelbaum, *Values of p -adic L -functions and a p -adic Poisson kernel*. *Invent. Math.* 101(1990), no. 2, 395–410.

UCD School of Mathematics and Statistics, University College Dublin, Belfield, Dublin 4, Ireland, D04 V1W8
 e-mail: ju-feng.wu@ucd.ie