

A note on the countable chain condition and sigma-finiteness of measures

K.P.S. Bhaskara Rao and M. Bhaskara Rao

The objectives of this paper are the following:

- (1) to show that a theorem of Ficker is incorrect;
- (2) to show that a stronger version of Ficker's Theorem is valid for a certain class of measures;
- (3) characterize all σ -algebras on which every measure is a countable sum of finite measures.

1. Introduction, notation and definitions

A measure μ is an extended real valued, nonnegative, countably additive function defined either on a σ -algebra A of subsets of a set X or on a boolean σ -algebra B vanishing at the empty set \emptyset or the zero element of B . Ficker [1, p. 242] proved the following theorem.

THEOREM (*). *Let μ be a measure on a σ -algebra A of X and N denote the collection of all sets in A of μ -measure zero. Then $A - N$ satisfies countable chain condition (CCC) if and only if μ can be written as a countable sum of finite measures.*

We give an example to show that this Theorem (*) is incorrect.

2. Example

Let B be a boolean σ -algebra satisfying CCC such that there is no strictly positive, finite measure on B . For example, one can take the

Received 16 November 1971. The authors are grateful to Dr Ashok Maitra for useful discussions they had with him on this paper.

boolean σ -algebra of all Borel subsets of the real line modulo first category Borel sets. Let X be the Stone space of B , \mathcal{A} the Baire σ -algebra on X and I the collection of all first category Baire subsets of X . By Loomis' Theorem (see, for example, [2, p. 102]), the quotient boolean σ -algebra \mathcal{A}/I and B are σ -isomorphic. Since I is a σ -ideal, the function μ defined by the formula, $\mu(A) = 0$ if $A \in I$, $\mu(A) = \infty$ if $A \in \mathcal{A} - I$, is a measure on \mathcal{A} . Note that \mathcal{A}/I satisfies CCC and so $\mathcal{A} - I$ satisfies CCC. If Ficker's Theorem (*) were to be true, we can write μ as a countable sum of finite measures on \mathcal{A} which implies that μ is equivalent to a finite measure λ on \mathcal{A} . Since I is precisely the collection of all λ -null sets, we have a strictly positive finite measure on \mathcal{A}/I . But this is a contradiction.

3. Semi-finite measures

A measure μ on a σ -algebra \mathcal{A} of X is said to be semi-finite if $F \in \mathcal{A}$, $\mu(F) = \infty$ implies there exists $E \in \mathcal{A}$ such that E is contained in F and $0 < \mu(E) < \infty$. For a measure μ on \mathcal{A} , there are two definitions of μ -atoms.

(I) A set A in \mathcal{A} is said to be a μ -atom if

(i) $\mu(A) > 0$ and

(ii) $B \in \mathcal{A}$, B contained in A implies $\mu(B) = 0$ or $\mu(B) = \mu(A)$.

(II) A set A in \mathcal{A} is said to be a μ -atom if

(i) $\mu(A) > 0$ and

(ii) $B \in \mathcal{A}$, B contained in A implies $\mu(B) = 0$ or $\mu(A-B) = 0$.

These definitions are not equivalent. It is easy to construct an example. However, when μ is semi-finite these two definitions are equivalent. Ficker [1] adopted definition (II) in the course of his proof of the Theorem (*). Under this definition, his Lemma 3 [1, p. 239] is not correct. However, if μ is semi-finite all his proofs are valid and hence for such a class of measures his Theorem (*) is true.

Here we prove a stronger version of his Theorem (*) directly for semi-finite measures.

THEOREM 1. *Let μ be a semi-finite measure on a σ -algebra \mathcal{A} of*

X . Let N denote the collection of sets of μ -measure zero. Then $A - N$ satisfies CCC if and only if μ is σ -finite.

Proof. If μ is σ -finite, it is obvious that $A - N$ satisfies CCC. Conversely, if $\mu(X) < \infty$, there is nothing to prove. If $\mu(X) = \infty$, choose A_1 in A such that $0 < \mu(A_1) < \infty$. Choose A_2 in A such that A_2 is contained in $X - A_1$ and $0 < \mu(A_2) < \infty$. Thus we can find a sequence of disjoint sets A_1, A_2, \dots in A such that each $A_i \in A - N$ and $\mu(A_i) < \infty$. If $\mu\left(X - \bigcup_{i \geq 1} A_i\right) < \infty$, then we have a decomposition of X which implies that μ is σ -finite. If $\mu\left(X - \bigcup_{i \geq 1} A_i\right) = \infty$, choose A_ω in A such that A_ω is contained in $X - \bigcup_{i \geq 1} A_i$ and $0 < \mu(A_\omega) < \infty$, where ω is the first countable ordinal. Continue this process. Since $A - N$ satisfies CCC, there exists a countable ordinal α such that $\mu\left(X - \bigcup_{\beta < \alpha} A_\beta\right) < \infty$. This implies that μ is σ -finite.

4. Some characterizations

Let A be a σ -algebra on a set X . A set A in A is said to be an atom of A if

- (i) $A \neq \emptyset$ and
- (ii) B in A , B contained in A implies $B = \emptyset$ or $B = A$.

A σ -algebra A on X is said to be atomless if there are no atoms of A .

The following result is known. See Remark 11 of [3, p. 203]. For completeness sake, we give a proof of this result.

PROPOSITION. *Let A be a σ -algebra on a set X . A is atomless if and only if every nonempty set in A contains \aleph_1 disjoint nonempty sets in A .*

Proof. Let A in A be nonempty. Fix $x \in A$. Find A_1 in A such that $x \notin A_1$, $A_1 \neq \emptyset$ and A_1 is contained in A . Choose A_2 in A such that $x \notin A_2$, $A_2 \neq \emptyset$ and A_2 is contained in $A - A_1$.

Continuing this process, we obtain a family $A_\alpha : \alpha < \Omega$ of nonempty disjoint sets contained in A , where Ω is the first uncountable ordinal. The converse part is trivial.

THEOREM 2. *Let A be a σ -algebra on a set X . The following statements are all true:*

- (i) *A satisfies CCC if and only if A is isomorphic to the power set, that is, the class of all subsets, of some countable (finite or infinite) set;*
- (ii) *there exists a strictly positive finite measure on A if and only if A is isomorphic to the power set of some countable set;*
- (iii) *every measure on A can be written as a countable sum of finite measures if and only if A is isomorphic to the power set of some countable set;*
- (iv) *every measure on A is equivalent to a finite measure if and only if A is isomorphic to the power set of some countable set.*

Proof. A proof of (i) can be obtained using the Proposition proved earlier. Since A satisfies CCC, the number of atoms of A is countable. From X remove all atoms of A . In view of the Proposition the remaining part is empty. The proofs of (ii), (iii) and (iv) are easy.

Professor Ashok Maitra suggested an alternative proof of (i). Since A satisfies CCC, it is complete as a boolean algebra. For x in X , the infimum of all sets in A containing x is an atom of A . This implies that A is atomic. Again by CCC, the number of atoms of A is at most countable.

References

- [1] V. Ficker, "On the equivalence of a countable disjoint class of sets of positive measure and a weaker condition than total σ -finiteness of measures", *Bull. Austral. Math. Soc.* 1 (1969), 237-243.

- [2] Paul R. Halmos, *Lectures on Boolean algebras* (Van Nostrand, Princeton, New Jersey; Toronto; New York; London; 1963).
- [3] B.V. Rao, "On Borel structures", *Colloq. Math.* 21 (1970), 199-204.

Research and Training School,
Indian Statistical Institute,
Calcutta,
India.