

Invited Review

Cite this article: Crippa BL, de Matos LG, Souza FN and Silva NCC (2024). Non-*aureus* staphylococci and mammaliococci (NASM): their role in bovine mastitis and One Health. *Journal of Dairy Research* **91**, 44–56. <https://doi.org/10.1017/S0022029924000165>

Received: 25 April 2023

Revised: 19 January 2024

Accepted: 22 January 2024

First published online: 8 April 2024

Keywords:

Food safety; mastitis; milk; non-*aureus* staphylococci and Mammaliococci (NASM); One Health; public health

Corresponding author:

Bruna Lourenço Crippa;
Email: lourencobruna@yahoo.com.br

Non-*aureus* staphylococci and mammaliococci (NASM): their role in bovine mastitis and One Health

Bruna Lourenço Crippa¹, Luiz Gustavo de Matos^{1,2}, Fernando Nogueira Souza^{3,4} and Nathália Cristina Cirone Silva¹

¹Department of Food Science and Nutrition, School of Food Engineering (FEA), University of Campinas (UNICAMP), Campinas, SP, 13083-862, Brazil; ²Department of Veterinary Medicine, Università degli Studi di Milano, Lodi, Lombardia, Italy; ³Department of Clinical Science, Faculty of Veterinary Medicine and Animal Sciences, University of São Paulo, São Paulo, 05508-270, Brazil and ⁴Department of Veterinary Medicine, School of Food Engineering (FEA), University of Campinas (UNICAMP), Campinas, SP, 13083-862, Brazil

Abstract

Non-*aureus* staphylococci (NAS) are gaining importance in mastitis and public health, and some NAS have been reclassified as mammaliococci (NASM). Bovine milk production has a major influence on the world economy, being an essential source of income for small, medium and large producers, and bovine mastitis caused by NASM can cause an economic impact. Mastitis generates financial losses due to reduced revenue, increased veterinary costs and expenses associated with animal slaughter. However, it is also a public health issue involving animal health and welfare, human health and the ecosystem. Furthermore, it is an increasingly common infection caused by NASM, including antimicrobial-resistant strains. Despite all these adverse effects that NASM can cause, some studies also point to its protective role against mastitis. Therefore, this review article addresses the negative and positive aspects that NASM can cause in bovine mastitis, the virulence of the disease and resistance factors that make it difficult to treat and, through the One Health approach, presents a holistic view of how mastitis caused by NASM can affect both animal and human health at one and the same time.

Introduction

Milk and milk products are important global dietary products consumed by more than 6 billion people worldwide. In 2021, recorded milk consumption was 928 million tons, making the dairy industry a very profitable market (Food and Agriculture Organization of the United Nations, 2021). However, an infection of the mammary gland caused mainly by bacteria, mastitis, is a major problem affecting animal welfare, productivity and the economy, especially in dairy production, which can lead to losses for the dairy industry (Abera *et al.*, 2013).

Among the main microorganisms involved in mastitis, we can mention *Staphylococcus* sp., of which *S. aureus* is the most significant mastitis pathogen, whilst the NAS (non-*aureus* staphylococci) are regarded as minor mastitis pathogens. Among all NAS found among cattle, *S. haemolyticus*, *S. chromogenes*, *S. epidermidis*, *S. warneri*, *S. cohnii*, *S. simulans*, *S. hominis*, *S. capitis*, and *S. xylosum* are the most prevalent species (Traversari *et al.*, 2019). Besides NAS, it is important to highlight the new genus reclassified from *Staphylococcus*, the *Mammaliococcus*. A study published by Madhaiyan *et al.* (2020) suggests the reassignment of five species of *Staphylococcus* (*S. sciuri*, *S. fleurettii*, *S. lentus*, *S. stepanovicii* and *S. vitulinus*) to this novel genus, with *Mammaliococcus sciuri* as the type species. By whole genome sequences, these mammaliococci species are as distant from staphylococci (64.2–67.3% amino acid identity – AAI) as *Macroccus* is distant from *Staphylococcus* (61.1–64.3%). Non-*aureus* staphylococci and mammaliococci (NASM) have become a concern among dairy producers, as they are the most common microorganisms isolated from aseptically collected quarter milk samples (Fergestad *et al.*, 2021).

According to the World Health Organization (WHO, 2024) 420 000 lives are lost due to food poisoning, and *Staphylococcus* spp. are characterized as important agents that can cause foodborne diseases. Poisoning occurs due to ingesting enterotoxins produced in food, including those produced by NASM, and symptoms include vomiting, diarrhea and cramps (da Silva Cândido *et al.*, 2020). But there are other problems as well. Political agendas, legislation, development of therapies and educational initiatives are essential to mitigate the increasing rate of antibiotic resistance amongst pathogenic organisms. In addition to bacteria having the ability to form biofilms, other genes are also involved in antibiotic resistance, and the process of becoming resistant can also occur through mutation at the genetic level (Zaman *et al.*, 2017). In addition, it has been suggested that NASM might act as a potential reservoir for resistance

genes, which can be transferred and integrated into the genome of *S. aureus* (Vitali *et al.*, 2014). Despite being under debate, it is essential to highlight that there are species of *Mammaliicoccus* that are suggested to be the evolutionary origin of the *mecA* gene in *S. aureus* (Lakhundi and Zhang, 2018).

For the most part, NASM can be regarded as minor pathogens that do have the ability to cause mastitis. However, some studies have demonstrated that NAS can reduce infection by *S. aureus* and other pathogens. Therefore, this review article aims to address the negative and positive aspects that NASM can have in bovine mastitis, in addition to addressing the importance of adopting the concept of One Health.

Non-aureus staphylococci

Differences among and within bovine NAS species have been reported regarding epidemiology and ecology (Souza *et al.*, 2016), antimicrobial resistance (Fergestad *et al.*, 2021), virulence (Wuytack *et al.*, 2020), potential protective traits (Toledo-Silva *et al.*, 2021a, 2021b), host interaction (Piccart *et al.*, 2016) and impact on udder health and milk yield (Valckenier *et al.*, 2019, 2020, 2021) as recently reviewed by De Buck *et al.* (2021). Non-aureus staphylococci can be present in a broad range of ecological habitats, which includes bovine teat canals (Traversari *et al.*, 2019) and teat apices (Adkins *et al.*, 2018) as well as the farm environment including bulk tank milk, bedding (Adkins *et al.*, 2022) and feces (Wuytack *et al.*, 2020). *S. chromogenes* and *S. epidermidis* are predominantly isolated from milk. Therefore, they are considered host-adapted species, whereas *S. equorum* and *S. devriesei* are mostly found in the environment and are classified as environmental NAS species (Adkins *et al.*, 2022). Other species, such as *S. haemolyticus*, originate from various (extra)mammary sites and so are labeled opportunistic (Adkins *et al.*, 2022). Unfortunately, multiple studies contradict this current species-stratification (Thorberg *et al.*, 2009; Supré *et al.*, 2011; De Visscher *et al.*, 2014; Vanderhaeghen *et al.*, 2015), and these confounding observations could be attributed to strain differences within NAS species. Wuytack *et al.* (2019), for example, suggested in a recent study on strain distribution of NAS in multiple bovine-associated habitats that different NAS strains within a species appear to be habitat-specific or site-specific.

After several taxonomic reclassifications, introducing the coagulase test enabled differentiation between CoNS and CoPS (Fairbrother, 1940). Non-aureus staphylococci is a group of Gram-positive bacteria that were before known as coagulase-negative staphylococci (CoNS); they are usually found in the normal flora of humans and different animals, and some species are recognized as human and facultative animal pathogens (Belhout *et al.*, 2022). For a long time, only *S. aureus* was considered to be harmful to its host, however, as a consequence of patient- and procedure-related changes, NAS now includes a large number of nosocomial microorganisms, with *S. epidermidis* and *S. haemolyticus* being the most significant ones (Pereira-Ribeiro *et al.*, 2019).

Some of the pathogenicity of NAS strains is probably due to their ability to adhere to surfaces and form biofilm on polymer surfaces of medical devices used during the hospitalization process in hospitals (Le *et al.*, 2019). In addition, NAS were also recognized to be capable of 'hiding' from the immune system by being internalized by nonprofessional phagocytes, a mechanism that is mediated by the AtlE gene that helps the

microorganism not only bind but also be phagocyted by human endothelial cells (Christner *et al.*, 2010). Besides that, NAS showed antimicrobial resistance and carried multiple resistance genes, which can be shared with other species (Lee *et al.*, 2020).

NAS brings issues not only to human health but also to animals. They are often found as the majority in bovine intramammary infection, especially in dairy heifers, and several studies in different areas of the globe have found NAS in different proportions in bovine udder, mainly in non-bred or first lactation animals (De Buck *et al.*, 2021). This could have happened because the NAS group was considered to comprise minor pathogens and so were typically not treated. However, grouping these species contributed to a better understanding of their distribution and importance to the animal's health care (Condas *et al.*, 2017).

Despite NAS being a large group, we can highlight some individual species. *S. epidermidis* is recognized as a pathogen and is the most important cause of nosocomial infection (Hugh and Ellis, 1968). This species can also adapt quickly to extreme conditions. For example, to survive in a place with a high salt concentration, the species has a different number of ion/proton exchangers and systems for osmo protectants (Gill *et al.*, 2005), and carries many genes that allow it to survive in these severe situations (Otto, 2009). Also, it is known that *S. epidermidis* is capable of producing an extracellular polysaccharide adhesive, termed polysaccharide intercellular adhesin (PIA) or polymeric N-acetyl-glucosamine, by *ica* operon-encoded enzymes. This is a significant virulence factor of the specie, helping them to colonize medical devices and leading to device-related infections (Le *et al.*, 2019). Besides this, they can also produce different toxins (Namvar *et al.*, 2014). However, studies suggest that this bacterium has evolved to play an important role in maintaining the microbiomes of humans and other mammals (Otto, 2009).

Due to its potential to produce biofilm, *S. haemolyticus* is also a critical species for nosocomial infections in humans, especially from blood infections. However, there is not as much information for them as for *S. epidermidis* (Barros *et al.*, 2015). When the resistance to β -lactam antibiotics is considered, *S. aureus* and *S. haemolyticus* present enough similarity to say that they have the exact same resistance mechanism (Czekaj *et al.*, 2015).

In several countries, *S. chromogenes* has been identified as the most prevalent species in milk samples from healthy cows and those with subclinical mastitis and clinical mastitis (Valckenier *et al.*, 2020). *S. chromogenes* could overcome the physical barrier of the udder and form biofilms. In addition, this species is more adapted to the cow's mammary gland and may become a microbial reservoir and possible source of infection (Tomazi *et al.*, 2015; Isaac *et al.*, 2017). Molecular studies have also shown that *S. chromogenes* has several antibiotic resistance genes, in addition to having this resistance tested and confirmed in phenotypic tests (Qu *et al.*, 2019).

Mammaliicoccus

Mammaliicoccus are Gram-positive, non-motile, non-spore-forming, paired cocci cells with aerobic to facultative anaerobic growth. They are catalase-positive and possess variable oxidase activities. The G + C content varies from 31.6 to 35.7% in genomes that range from 2.44 to 2.81 Mbp. Based on the analyses of 16S rRNA gene sequences, five species (*M. sciuri*, *M. fleurettii*, *M. lentus*, *M. stepanovicii*, and *M. vitulinus*), are now in this novel genus (Madhaiyan *et al.*, 2020) and are as distant from *Staphylococcus* species as *Macrococcus* (another genus) is from

Staphylococcus. Recognized to colonize humans and animals, they are also considered potentially opportunist pathogens capable of affecting the immune system (Beuckelaere *et al.*, 2021). Like the NAS, this group may also cause mastitis in different species, and/or may serve to transmit resistance genes to other more pathogenic species, as suggested by the study of Lienen *et al.* (2022) where *M. sciuri*, *M. lentus*, *M. fleurettii* and *M. vitulinus* were isolated from milk and swab samples collected from German dairy cows.

M. sciuri (previously *Staphylococcus sciuri*) is usually found on the skin of wild and domestic animals. Based on their ribotype pattern, the species was divided into three different subspecies called *S. sciuri* subsp. *sciuri*, *S. sciuri* subsp. *carnaticus* and *S. sciuri* subsp. *rodentium* (Nemeghaire *et al.*, 2014). In addition, some studies suggest this species as the origin of the *mecA* gene, which is located on a mobile genetic element called staphylococcal cassette chromosome *mec* (SCC*mec*), and is the gene responsible for methicillin-resistance in *S. aureus* (MRSA) or more generally in *Staphylococcus* spp (MRSS: Couto *et al.*, 1996; Fuda *et al.*, 2007). Whilst being characterized as a typical colonizer of a wide range of animals, this species has also been associated with various health problems including skin disease in goats and pigs (Nemeghaire *et al.*, 2014). Despite displaying various virulence factors (lipolytic, proteolytic and hemolytic), these bacteria are not related to infection in humans, except after previous infection due to a pathogen (Nemeghaire *et al.*, 2014).

M. lentus (previously *S. lentus*) was first characterized as a subspecies of *S. sciuri*. However, after analysis of DNA-DNA hybridization, it was placed into a new species (Schleifer *et al.*, 1983). It was originally isolated from poultry and other animals and can produce enterotoxins, but only to a minor extent (less than 100 ng/ml supernatant: Irlinger, 2008). Guerre *et al.* (1999) showed that the enterotoxigen strain of *M. lentus* inoculated in milk for the production of Camembert-type cheeses does not produce enterotoxins despite high levels of the microorganisms.

For *M. fleurettii* (previously *S. fleurettii*), it was also possible to identify enterotoxin production, and some studies suggest that this species, rather than *S. sciuri*, is the origin of the *mecA* gene in MRSA. However, the exact origin is still debated (Lakhundi and Zhang, 2018). In this context, *M. stepanovicii* was also reported with the presence of the gene (Loncaric *et al.*, 2013). Table 1 provides a complete list of NASM and the different samples from which they were isolated.

NASM as a cause of mastitis

NAS forms a heterogeneous group of more than 50 species, of which approximately ten are associated with intramammary infections in dairy cattle and are abundantly present in bulk milk (Vanderhaeghen *et al.*, 2015). As other species of *Staphylococci* become more commonly associated with the disease, it is of interest to learn more about the distribution of NAS on dairy farms. They have many virulence factors, and mastitis control is complicated because this group contains many species (Thorberg *et al.*, 2009).

The main ways to diagnose subclinical mastitis from NAS are from the results of somatic cell count (SCC) and culture. Tomazi *et al.* (2015) observed that mammary quarters infected by NAS had elevated SCC (> 306 000 vs. < 65 000 in contralateral uninfected quarters), without any affect on milk yield or contents of fat, crude protein, casein, lactose, total solids, and solids-not-fat.

Although considered as pathogens with a lower clinical incidence, NAS infections have been identified as relatively common (Persson Waller *et al.*, 2011; Gao *et al.*, 2017; Heikkilä *et al.*, 2018), clinical cases being due mainly to *S. chromogenes*, *S. simulans* and *S. haemolyticus* (Persson Waller *et al.*, 2011). In subclinical mastitis cases, NAS are the second most commonly identified pathogen after *S. aureus* in Swedish dairy cows, *S. epidermidis* followed by *S. chromogenes*, *S. simulans*, and *S. haemolyticus* being the most common (Persson Waller *et al.*, 2011). Another study isolated *S. chromogenes*, *S. fleurettii*, *S. haemolyticus*, *S. sciuri*, *S. simulans*, *S. succinus* and *S. xylosus* from mastitic bovine milk (Khazandi *et al.*, 2018).

The main pathogenic factors of NASM

According to Vanderhaeghen *et al.* (2014), there are three virulence factor groups involved in the pathogenesis of staphylococcal infections: secreted proteins (e.g., superantigens, cytotoxins and tissue-degrading enzymes), cell surface-bound proteins (e.g. microbial surface components recognizing adhesive matrix molecules: MSCRAMM) and cell wall components, with the polysaccharide capsule and lipoteichoic being examples of those. Naushad *et al.* (2019), using whole genome sequencing analyzed the distribution of 191 virulence factors in 441 genomes of 25 NAS species; the authors reported virulence factors in functional categories such as toxins ($n = 93$), iron metabolism ($n = 29$), adherence ($n = 28$), exoenzymes ($n = 21$) and immune evasion ($n = 20$). Another study by Wuytack *et al.* (2020) compared the presence of virulence genes (*agrA*, *bap*, and *cap5H* and the *mecA* methicillin resistance gene) of NAS isolated from the milk samples of healthy quarters and quarters with clinical mastitis. Based on this comparison, they suggested that NAS has the potential to cause mild clinical mastitis.

Enterotoxins

Staphylococcal poisoning is widespread and mainly transmitted by food products, as it can affect healthy people. NAS is also able to produce enterotoxins, which can be one of the causes of many food diseases (da Silva Cândido *et al.*, 2020). Enterotoxins belong to a group of extracellular single-chain proteins that are water-soluble and resistant to proteolytic enzyme action in the digestive system. This means that they remain active after ingestion and can compromise human health and, consequently, the public health system. These proteins are also thermostable and resist thermal treatments such as pasteurization (Vanderhaeghen *et al.*, 2014). There are more than 25 types of staphylococcal enterotoxins (SE) and enterotoxin-like (SEL) toxins; the former have an emetic effect in primates, whilst the latter do not. The most known of the enterotoxins are the classical serotypes (SEA through SEE: Bergdoll *et al.*, 1971), followed by the non-classical SEG through SET as well as SEY and more recently SE02 (Su and Lee Wong, 1995; Munson *et al.*, 1998; Ono *et al.*, 2008, 2015, 2019; Omoe *et al.*, 2013; Suzuki *et al.*, 2020). The SELs identified until today are SELJ and then SELU through SELX, SELZ and SEL 01 (Zhang *et al.*, 1998; Letertre *et al.*, 2003; Thomas *et al.*, 2006; Wilson *et al.*, 2011; Spoor *et al.*, 2015; Hisatsune *et al.*, 2017). NASM are capable of producing the majority of these (Rodrigues *et al.*, 2017; da Silva Cândido *et al.*, 2020). However, this has largely been ignored by authorities as NAS can establish a commensal relationship with humans and

Table 1. NASM isolated from different samples (human, animal, environment, and food) and different countries

Specie	Prevalence (%)	Sample collection	Number of samples	Presence of enterotoxin	Place	Reference
<i>S. arlettae</i>	12	Subclinical mastitis milk samples	209	<i>seb</i> (16.6%)	China	Xu <i>et al.</i> (2015)
<i>S. xyloso</i>	12			No presence		
<i>M. sciuri</i>	12			<i>sei</i> (8.3%)		
<i>S. chromogenes</i>	49	Intramammary infection	5,434	Not analyzed	Canada	Condas <i>et al.</i> (2017)
<i>S. simulans</i>	17					
<i>S. xyloso</i>	12					
<i>S. equorum</i>	7	Rectal feces	80	Not analyzed	Belgium	Wuytack <i>et al.</i> (2020)
<i>S. chromogenes</i>	29	Quarters from 82 heifers	324	Not analyzed		Valckenier <i>et al.</i> (2020)
<i>S. xyloso</i>	9					
<i>S. vitulinus</i>	9					
<i>S. chromogenes</i>	49	82 dairy cows	328	Not analyzed	Finland	Taponen <i>et al.</i> (2008)
<i>S. simulans</i>	23					
<i>S. warneri</i>	5					
<i>S. chromogenes</i>	47	Composite milk samples	441	Not analyzed	Argentina	Raspanti <i>et al.</i> (2016)
<i>S. haemolyticus</i>	32					
<i>S. warner</i>	7					
<i>S. chromogenes</i>	30	Quarter milk samples	4220	Not analyzed	Netherlands	Sampimon <i>et al.</i> (2009)
<i>S. epidermidis</i>	13					
<i>S. capitis</i>	11					
<i>S. warneri</i>	37	Udder quarters	46	Not analyzed	Poland	Malinowski <i>et al.</i> (2006)
<i>S. chromogenes</i>	33					
<i>S. xyloso</i>	23					
<i>S. chromogenes</i>	48	Milk samples	604	Not analyzed	USA	Jenkins <i>et al.</i> (2019)
<i>S. haemolyticus</i>	18					
<i>S. simulans</i>	7					
<i>S. epidermidis</i>	75	Human Nasal swab	176	Not analyzed	Germany	Marincola <i>et al.</i> (2021)
<i>S. hominis</i>	15					
<i>S. warneri/ S. pasteurii</i>	7.1					
<i>S. epidermidis</i>	57.9	Prosthetic joint infection	113	Not analyzed	USA	Tande <i>et al.</i> (2014)
<i>S. capitis/caprae</i>	5.3					
<i>S. lugdunensis</i>	7.9					

(Continued)

Table 1. (Continued.)

Specie	Prevalence (%)	Sample collection	Number of samples	Presence of enterotoxin	Place	Reference
<i>S. epidermidis</i>	83.3	Prosthetic joint infection	6	Not analyzed	Poland	Bogut et al. (2014)
<i>S. warneri</i>	16.7					
<i>S. epidermidis</i>	56.7	Blood, pus, wound swab, drain fluid, tracheal aspirate, peritoneal fluid, and pleural fluid	120	Not analyzed	India	Bora et al. (2018)
<i>S. haemolyticus</i>	21.7					
<i>S. lugdunensis</i>	11.7					
<i>S. epidermidis</i> ,	80.7	Human nasal swab among patients undergoing haemodialysis	71	Not analyzed	Poland	Kozioł-Montewka et al. (2006)
<i>S. lugdunensis</i>	7.7					
<i>S. haemolyticus</i>	3.8					
<i>S. warneri</i>	3.8					
<i>S. capitis</i>	3.8					
<i>M. sciuri</i>	58.8	Environment	17	Not analyzed	German	Schnitt et al. (2021)
<i>S. lentus</i>	17.6					
<i>S. fleurettii</i>	5.8					
<i>S. epidermidis</i>	5.8					
<i>S. haemolyticus</i>	5.8					
<i>S. cohnii</i>	5.8					
<i>S. haemolyticus</i>	1	Ready-to-eat raw fish	200	Not analyzed	Japan	Hammad et al. (2012)
<i>S. haemolyticus</i>	1					
<i>S. pasteurii</i>	1					
<i>S. intermedius</i>	25	Meat	286	<i>sec</i> (12.5%)	Jordan	Al-Tarazi et al. (2009)
<i>S. warneri</i>	12.5			No presence		
<i>S. lugdenesis</i>	12.5			No presence		
<i>S. epidermidis</i>	6.3%			<i>sec</i> (66.6%)		
<i>M. sciuri</i>	3.1			<i>sec</i> (100%)		
<i>S. intermedius</i>	3.1			No presence		
<i>S. haemolyticus</i>	29.0	Bicycles/ Environment	79	Not analyzed	China	Xu et al. (2019)
<i>S. hominis</i>	29.0					
<i>S. epidermidis</i>	14.0					

et al. (2020), this was their study's most frequently detected gene, followed by *eno* and *jbe*, then *altE*. Srednik *et al.* (2017) showed in their research that NAS isolated from cows with mastitis can produce biofilm in vitro, and they also showed that 90% of the isolates tested positive for one of the above genes.

Zuniga *et al.* (2015) also studied the components on microbial surfaces that recognize the adhesive matrix and found the presence of the *can* gene in 17% of isolates, and the *eno* gene at 82.1% was the most prevalent gene among all. The *ebpS* gene was positive in 17.9%, and there were 72.6% strains positive for the *fnbA* gene and 5.7% for the *fnbB* gene. The *fib* gene was among the most prevalent, present in 76/106 (71.6%), and the *bap* gene was in 60/106 (56%). Darwish and Asfour (2013) identified the presence of the *eno* gene in 92.6% of all 68 NAS isolates. Simojoki *et al.* (2012) also showed a prevalence of the *eno* gene, which was positive in 75% of the 84 studied strains, and the genes *bbp* (1.2%), *cflA* (4.8%), *cflB* (3.6%), *ebpS* (4.8%), *fib* (4.8%), *fnbA* (3.6%), and *fnbB* (2.4%) were also detected. The *cflA* gene is involved with adhesion and immune evasion whilst the *ebpS* gene is responsible for encoding elastin-binding proteins present in the extracellular matrix. Fibrinogen-binding proteins are encoded by the *fib* gene, and fibronectin-binding protein A and fibronectin-binding protein B are encoded by the *fnbA* and *fnbB* genes respectively (Zuniga *et al.*, 2015).

Resistance to antibiotics

Biofilm formation is also connected to antibiotic resistance. According to Srednik *et al.* (2017), NAS isolates growing within biofilms are less susceptible to antimicrobials commonly used on farms. Antibiotics are manufactured at an estimated scale of about 100 000 tons annually worldwide, and their use profoundly impacts bacterial life on Earth. More pathogenic strains have become antibiotic-resistant, and some have become resistant to multiple antibiotics, known as multidrug resistance (Nikaido, 2009). El-Seedy *et al.* (2017) tested how biofilm formation could affect antibiotic resistance, and a total of 95 NAS were identified and tested for biofilm formation, of which 95.8% were resistant to ampicillin, 77.9% were resistant to cefoxitin, 35.8% were resistant to cefuroxime and 31.6% were resistant to amoxicillin.

Besides biofilm formation, there are also genes involved in antibiotic resistance. Xu *et al.* (2015) studied the presence of some of these genes in 76 NAS isolates. They reported the following results: the most prevalent gene was *linA*, which gives resistance to lincosamides, followed by *tetK*, which is responsible for resistance to tetracycline. Conferring resistance to penicillin, the gene *blaZ* was found in 30.3%. Also present were the genes *aacA* - *aphD* which provide resistance to aminoglycoside. Another gene studied was *msrA/B*, which was positive in 17.1 and 19.7% of isolates, respectively, conferring resistance to macrolide and erythromycin. The gene *mecA* is one of the most studied genes, and it is responsible for methicillin resistance and was positive in 17.1% of isolates; the genes *ermB/C* that confer resistance to erythromycin and clindamycin were present in 9.2 and 13.2% respectively whilst the genes *aac(6')/aph(2')*, responsible for resistance to streptomycin, were present in 10.5% of the strains (Kumar *et al.*, 2010).

Another study identified 109 NAS, all of which were positive for the *blaZ* gene, and 73% of isolates had the *mecA* gene, both of which resist β -lactam groups. The genes that grant resistance to tetracycline *tetK*, *tetL* and *tetM* all showed a high presence, and other genes identified in a significant amount of isolates

were the MLSb resistance genes, *ermB*, *inuA*, *msrA*, and *mphC*. In addition, *dfrG*, *aacA-aphD* and *aphA3* were present in 25, 24, and 43 NAS isolates (Qu *et al.*, 2019). Ahmed *et al.* (2020) found NASM isolates exhibited a high prevalence of the resistance genes *mecA*, *blaZ* and *tetK*. Approximately half of the isolates harbored the *aac-aphD*, *erm(C)*, and *erm(B)* genes.

Klibi *et al.* (2018) studied 83 isolates; sixty-eight of these strains were identified as NAS. In the study, 29.41% showed oxacillin and cefoxitin resistance whilst some strains were *mecA* positive. Eleven of those twenty methicillin resistant non-*aureus* staphylococci (MRNAS) showed erythromycin-resistance and were positive for *ermB*, *ermT*, *ermC*, *mphC* or *msrA*. Four strains were *tetK* positive, which were resistant to tetracycline. Silva *et al.* (2014) also studied MRNAS, and they reported 26 strains with oxacillin and cefoxitin resistance that were positive for the *mecA* gene. In the same study, the SCC*mec* gene, a mobile genetic element that carries the central determinant for broad-spectrum beta-lactam resistance encoded by the *mecA* gene, was analyzed; it was present in 9 out of 26 strains. MRNAS isolates were also found to display additional resistance to tetracycline, streptomycin, tobramycin and gentamicin, and harbored the genes: *tet(K)*, *str*, *ant(4')-I*, and *aac(6')/aph(2')*.

Multidrug resistance

Studies suggest that the frequent use of drugs in veterinary practices, such as penicillin and tetracycline, especially among cattle, has increased the resistance of microorganisms over the years. The Food and Drug Administration (FDA) and the European Medicine Agency (EMA) show that tetracycline, penicillin, and sulfonamides are the classes of antibiotics most sold in the USA and Europe (Dorneles *et al.*, 2019). Several studies in the literature point to multidrug-resistant NAS and these are enumerated in Table 2. In addition, NAS are known to be more resistant to antibiotics when compared to *S. aureus*, and this resistance can be to different drugs and contribute to a low cure rate. The resistance can be due to NAS, typically on the cow's body, being more exposed to antimicrobials (Khazandi *et al.*, 2018).

NASM involvement in mastitis prevention

While studies show NASM as the cause of mastitis, some studies have demonstrated that NAS can reduce infection by *S. aureus* and other pathogens (Brouillette *et al.*, 2022). In another study by Toledo-Silva *et al.*, (2021a, 2021b), it was observed that NASM suppressed the biofilm dispersion of a strain of *S. aureus*. Some studies also point to a possible protective role of NASM, as described by Piepers *et al.* (2013), who in comparing infected heifers with uninfected heifers, observed that heifers infected with NASM had fewer cases of clinical mastitis throughout lactation.

Some gram-positive bacteria can produce bacteriocins, inhibiting the growth of other bacterial strains. Bacteriocins produced by NASM include epidermin, hyacin 3682, nukacin ISK-1, nukacin 3299 and nukacin KQU-131. *Staphylococcus epidermidis* produces Tu 3298, *Staphylococcus hyicus* 3682, *Staphylococcus warneri* ISK-1, *Staphylococcus simulans* 3299, and *Staphylococcus hominis* KQU-131 (Yong *et al.*, 2019), so it can be seen that there is potential for NASM to negatively affect the growth of pathogenic bacteria. In this context, Ferronato *et al.* (2019) used the cross-streaking method to test growth inhibition of mastitis-causing pathogens. Using this method, it is possible to determine if there was partial inhibition, total inhibition (no colony was

Table 2. Number of multidrug resistant non-*aureus* Staphylococci; antibiotic groups they have resistance to; percentage of resistant strains for each drug or group.

Number of MDR-NAS	Antibiotic resistance	% of MDR	Reference
73/152	Penicillin + ampicillin + erythromycin	51%	Phophi <i>et al.</i> (2019)
60/107	Enrofloxacin + oxacillin + penicillin + tetracycline	56%	Cheng <i>et al.</i> (2019)
14/37	Oxacillin + chloramphenicol + novobiocin + tetracycline	38%	Khazandi <i>et al.</i> (2018)
3/8	Ampicillin + clindamycin + oxacillin erythromycin + gentamycin + penicillin + sulfonamide + trimethoprim/sulfamethoxazole + tetracycline	37.5%	Dorneles <i>et al.</i> (2019)
2/8	Ampicillin + penicillin + trimethoprim/sulfamethoxazole	25%	
1/8	Ampicillin + gentamycin + oxacillin + penicillin + sulfonamide + trimethoprim/sulfamethoxazole + tetracycline	12.5%	
2/8	Ampicillin + gentamycin + oxacillin + penicillin + trimethoprim/sulfamethoxazole + tetracycline	25%	
18/170	Same group of antibiotics (beta-lactam or MLS compounds)	10.6%	Sampimon <i>et al.</i> (2011)
18/170	Different groups	10.6%	

observed), or no growth inhibition (same size and number of colonies as on the positive control plate). It was reported that all of the 19 NAS studied could inhibit growth partially, and in the case of *S. chromogenes* there was total inhibition. Another study showed the capacity of 38 strains of NAS to inhibit *Listeria innocua* growth, with the strain that most displayed this ability being *S. quorum* (Braem *et al.*, 2014).

De Vlieghe *et al.* (2004) also studied how *S. chromogenes* could inhibit the growth of some bacteria, such as *S. aureus*, *S. dysgalactiae* and *S. uberis*, and the result was positive. The NAS studied inhibited the growth of these strains but not that of *E. coli*, which is typically expected, as the inhibitory effect of bacteriocins is usually more intense against phylogenetically related bacterial species (Ferronato *et al.*, 2019). Furthermore, the same authors in a previous study suggested that colonization of the teat apex by *S. chromogenes* in pre-calving heifers protected the udder against increased SCC in the early postpartum period (De Vlieghe *et al.*, 2003).

According to Isaac *et al.* (2017), NASM is a group of bacteria classified as either minor mastitis pathogens or commensal microbiota. A study showed that *S. epidermidis*, *S. simulans*, *S. hominis*, *S. saprophyticus*, and *S. arlettae* isolated from milk were able to produce antimicrobial compounds to inhibit the growth of some mastitis pathogens, including *S. aureus* (Nascimento *et al.*, 2005). Beuckelaere *et al.* (2021) observed a probiotic effect of *S. chromogenes* when colonizing dry quarters with this microorganism. As a result, the authors observed a shift to the Th1 response in late pregnancy and early lactation, increasing IgG2 concentration. In a study with a murine mastitis model, the authors observed that *S. aureus* intramammary colonization is reduced by exoproducts from *S. chromogenes* and *S. simulans* (Brouillette *et al.*, 2022). Some NAS species (*S. chromogenes*, *S. simulans* and *S. epidermidis*) inhibited the growth of *S. aureus* and downregulated the expression of rnaIII, important molecule of the QS system (Toledo-Silva *et al.*, 2021a, 2021b). In addition, other studies point out that NASM protects the mammary quarters against IMI caused by primary pathogens (De Buck *et al.*, 2021). Other characteristics of NASM, to cause or prevent mastitis, could be explained by differences among the species and intraspecies once the host determines the infection, the virulence of microorganisms and biofilm formation capacity (Brouillette *et al.*, 2022). However, much still needs to be understood about these rather complex interactions within the microbiota.

One health and public health

The One Health approach conceptualizes that we cannot truly understand human, animal and environmental health by only addressing each in isolation. Because of this, we must understand the connections of these areas using holistic interpretations to face public health challenges. In other words, One Health focuses equally or more on the relationships between the factors in the system rather than on the individual-level factors themselves (Dalton *et al.*, 2020).

We can observe in Tables 1 and 2 NASM being isolated from food, clinical infections and environment, as well as their resistance to antibiotics. Several NASM strains from these studies (Kozioł-Montewka *et al.*, 2006; Hammad *et al.*, 2012; Xu *et al.*, 2019; Schnitt *et al.*, 2021) have been confirmed as MRNAS. Also, the topics about virulence and part of Table 1 showed the capacity of these species to produce toxins and form biofilm. All these components place NASM as a concern for public health.

Advances in molecular techniques allow us to relate strains from community and clinical infections, which helps us understand transmission among humans, animals and environments. The species most studied in *Staphylococcus* spp. is *S. aureus* and many studies have observed clones common in clinical infections, in animals or in the environment (Silva *et al.*, 2013, 2014; Aires-de-Sousa, 2016; Zhou *et al.*, 2018). But, we also observe the same route for NAS in studies that demonstrate the presence of identical clones in hospitals and communities (Martínez-Meléndez *et al.*, 2016; Xu *et al.*, 2019; Pinheiro-Hubinger *et al.*, 2021). So, the prevalence of NASM in the dairy chain and the increasing prevalence of resistance genes raises further concern. The prevalence of antimicrobial-resistant pathogens has increased while the approval of new drugs has decreased, which is a public health challenge. However, the number of antimicrobials used in veterinary medicine is immense and should be discussed since they select resistant microorganisms. Animals are the reservoirs of these organisms and can transmit them through food (Gohl *et al.*, 2016; FDA, 2018).

According to the FDA, each year in the United States at least 2.8 million antibiotic-resistant infections occur, and more than 35 000 people die as a result. Therefore, agencies around the world are trying to minimize the harm caused by the incorrect use of antibiotics, for example, using them for purposes for which they are not intended (to treat diseases caused by viruses, fungi or yeasts), in addition to the frequent use of broad-spectrum

antibiotics (Hildreth *et al.*, 2009). The FDA and the European Food Safety Authority (EFSA) are two organizations that have been playing a crucial role in the dissemination of vital information to preserve the effectiveness of currently available antimicrobial drugs and promote the development of new medical products that can help reduce the emergence and spread of antimicrobial resistant bacteria (EFSA, *nd*; FDA, 2018).

Stevens *et al.* (2018) analyzed the consumption of antimicrobials in dairy herds and its association with the diameters of the antimicrobial inhibition zone of NAS isolated from subclinical mastitis. They reported a high incidence of antimicrobial treatment for fourth generation cephalosporins, a class of antibiotics important for human health. Strains of MRNAS were identified in milk samples with mastitis in a study in Korea, and one of the strains persisted on one of the analyzed farms for more than two years (Kim *et al.*, 2019). The role of NAS as a cause of disease in humans has been described. Diseases such as endocarditis, tract infections, catheter-related sepsis, endophthalmitis, surgical site infections, peritonitis in patients with continuous ambulatory peritoneal dialysis and foreign body infections were due to infections caused by NAS (Piette and Verschraegen, 2009). Approximately 55–75% of nosocomial isolates are resistant to methicillin, and NASM is responsible for about 30% of all nosocomial bloodstream infections (Piette and Verschraegen, 2009).

In the context of One Health and mastitis, there is a solid recommendation for reducing antimicrobial use. Taking into account that milk with antibiotic residues does not enter the supply chain, as it is withdrawn at source, this recommendation emphasizes the need to periodically search for antimicrobial residues in raw milk on farm. Combined with microbiome analysis and research on antimicrobial residues, we have the somatic cell count, which acts as an indicator of the health of the mammary gland, helping to reveal the situation of the dairy herd within the context of health (Gohl *et al.*, 2016). When milk contains residues, this indicates that it comes from unhealthy animals, so it can cause food-borne pathogens and be a potential allergen for consumers. This concern is even more significant in developing countries, where there is no supervision on using antimicrobials in animals or the opportunity to test milk and meat for drug residues. The presence of adulterating antimicrobials in milk can have a profound impact on the health of babies and children in regions that already face high rates of diarrheal diseases and malnutrition. Several studies have shown a relationship between the intestinal microbiome, health status and disease. Therefore, ensuring that milk does not contain antimicrobial residues significantly impacts public health and producers' economic livelihood (Mohan *et al.*, 2008; Gordon *et al.*, 2012; Garcia *et al.*, 2019). Since NASM is a large group that contains species with different levels of pathogenicity, virulence, resistance profile and host adaptation, analyses of the genetic profile of these species are necessary to understand the role of each NAS species in bovine mastitis (De Buck *et al.*, 2021). Thus, the One Health approach is also required when we are specifically talking about NAS.

This article gathers evidence that NASM has the potential to cause mastitis, and that all of this can cause public health problems since there is a lack of specific studies and strategies to combat these pathogens. Even with this evidence, NASM is still overlooked as causing disease in both humans and animals. With all these points raised, we must apply a One Health approach to address this critical public health problem, including implementing good antimicrobial administration practices in

human and veterinary health settings to delay the development of resistance and prolong the life of useful antimicrobials (FDA, 2018).

In conclusion, since it is an important food at all stages of life, milk consumption increases each year along with population growth. We know that NASM mastitis can harm the health and welfare of animals and, consequently, human health, either by food poisoning or by the recurrent use of antimicrobials to treat the disease, which can leave residues in the milk that will be consumed in different ways. Studies related to NASM mastitis are extremely important to learn more about the role of these microorganisms in the disease and how their virulence factors can make treatment difficult. Questions that need more concrete answers are why some NASM strains that cause mastitis appear to have a protective role against other microorganisms and why some strains can harm milk production and animals.

In addition, the increased resistance of these microorganisms to antimicrobials is worrying, requiring the implementation of residue monitoring in milk and studies of NASM antimicrobial resistance used to combat mastitis. Thus, a holistic vision that encompasses safer food and solutions to improve animal, human and ecosystem health seems to be the best alternative.

Acknowledgements. This work was carried out with the support of the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES) – Finance Code 001 and São Paulo Research Foundation – FAPESP, grant #2019/17308-4.

References

- Abera M, Demie B, Aragaw K, Regassa F and Regassa A (2013) Isolation and identification of *Staphylococcus aureus* from bovine mastitic milk and their drug resistance patterns in Adama town Ethiopia. *Journal of Veterinary Medicine and Animal Health* **1**, 19–23.
- Adkins PRF, Dufour S, Spain JN, Calcutt MJ, Reilly TJ, Stewart GC and Middleton JR (2018) Cross-sectional study to identify staphylococcal species isolated from teat and inguinal skin of different-aged dairy heifers. *Journal of Dairy Science* **101**, 3213–3225.
- Adkins PRF, Placheta LM, Borchers MR, Bewley JM and Middleton JR (2022) Distribution of staphylococcal and mammaliococcal species from compost-bedded pack or sand-bedded freestall dairy farms. *Journal of Dairy Science* **105**, 6261–6270.
- Ahmed W, Neubauer H, Tomaso H, El Hofy FI, Monecke S, Abdeltawab AA and Hotzel H (2020) Characterization of staphylococci and streptococci isolated from milk of bovines with mastitis in Egypt. *Pathogens (Basel, Switzerland)* **9**, 1–17.
- Aires-de-Sousa, M (2016) Methicillin-resistant *Staphylococcus aureus* among animals: current overview. *Clinical Microbiology and Infection* **23**, 373–380.
- Al-Tarazi YH, Albetar MA and Alaboudi AR (2009) Biotyping and enterotoxigenicity of *Staphylococci* isolated from fresh and frozen meat marketed in Jordan. *Food Research International* **42** 374–379.
- Barros EM, Lemos M, Souto-Pradón T and Giambiagi-deMarval M (2015) Phenotypic and genotypic characterization of biofilm formation in *Staphylococcus haemolyticus*. *Current Microbiology* **70**, 829–834.
- Belhout C, Elgroud R and Butaye P (2022) Methicillin-resistant *Staphylococcus aureus* (MRSA) and other methicillin-resistant *Staphylococci* and *Mammaliococcus* (MRNaS) associated with animals and food products in Arab countries: a review. *Veterinary Sciences* **9**, 317–317.
- Bergdoll MS, Borja CR, Robbins RN and Weiss KF (1971) Identification of enterotoxin E. *Infection and Immunity* **4**, 593–595.
- Beuckelaere L, De Visscher A, Souza FN, Meyer E, Haesebrouck F, Piepers S and De Vlieghe S (2021) Colonization and local host response following intramammary *Staphylococcus chromogenes* challenge in dry cows. *Veterinary Research* **52**, 137.

- Bogut A, Niedźwiadek J, Koziol-Montewka M, Strzelec-Nowak D, Blacha J, Mazurkiewicz T, Marczyński W and Plewik D (2014) Characterization of *Staphylococcus epidermidis* and *Staphylococcus warneri* small-colony variants associated with prosthetic-joint infections. *Journal of Medical Microbiology* **63**, 176–185.
- Bora P, Datta P, Gupta V, Singhal L and Chander J (2018) Characterization and antimicrobial susceptibility of coagulase-negative staphylococci isolated from clinical samples. *Journal of Laboratory Physicians* **10**, 414–419.
- Braem G, Stijlemans B, Van Haken W, De Vliegher S, De Vuyst L and Leroy F (2014) Antibacterial activities of coagulase-negative staphylococci from bovine teat apex skin and their inhibitory effect on mastitis-related pathogens. *Journal of Applied Microbiology* **116**, 1084–1093.
- Brouillette E, Goetz C, Droppa-Almeida D, Chamberland S, Jacques M and Malouin F (2022) Secondary *Staphylococcus aureus* intramammary colonization is reduced by non-*aureus* staphylococci exoproducts. *Microbes and Infection* **24**, 104879.
- Cheng J, Qu W, Barkema HW, Nobrega DB, Gao J, Liu G, De Buck J, Kastelic JP, Sun H and Han B (2019) Antimicrobial resistance profiles of 5 common bovine mastitis pathogens in large Chinese dairy herds. *Journal of Dairy Science* **102**, 2416–2426.
- Christner M, Franke GC, Schommer NN, Wendt U, Wegert K, Pehle P, Kroll G, Schulze C, Buck F, MacK D, Aepfelbacher M and Rohde H (2010) The giant extracellular matrix-binding protein of *Staphylococcus epidermidis* mediates biofilm accumulation and attachment to fibronectin. *Molecular Microbiology* **75**, 187–207.
- Condas LAZ, De Buck J, Nobrega DB, Carson DA, Roy JP, Keefe GP, DeVries TJ, Middleton JR, Dufour S and Barkema HW (2017) Distribution of non-*aureus* staphylococci species in udder quarters with low and high somatic cell count and clinical mastitis. *Journal of Dairy Science* **100**, 5613–5627.
- Couto I, De Lencastre H, Severina E, Kloos W, Webster JA, Hubner RJ, Sanches IS and Tomasz A (1996) Ubiquitous presence of a *mecA* homologue in natural isolates of *Staphylococcus sciuri*. *Microbial Drug Resistance* **2**, 377–391.
- Czekaj T, Ciszewski M and Szcwyczyk EM (2015) *Staphylococcus haemolyticus* – an emerging threat in the twilight of the antibiotics age. *Microbiology (Reading England)* **161**, 2061–2068.
- da Silva Cândido T, da Silva AC, de Matos LG, da Silva do Nascimento M, Camargo CH, Cobo Zanella R, Mores Rall VL and Silva NCC (2020) Enterotoxigenic potential and molecular typing of *Staphylococcus* sp isolated from organic and conventional fresh minas cheese in the state of São Paulo Brazil. *International Dairy Journal* **102** <https://doi.org/10.1016/j.idairyj.2019.104605>.
- Dalton KR, Rock C, Carroll KC and Davis MF (2020) One health in hospitals: how understanding the dynamics of people animals and the hospital built-environment can be used to better inform interventions for antimicrobial-resistant gram-positive infections. *Antimicrobial Resistance and Infection Control* **9**, 1–17.
- Darwish SF and Asfour HAE (2013) Investigation of biofilm forming ability in staphylococci causing bovine mastitis using phenotypic and genotypic assays. *The Scientific World Journal* **2013**, 378492.
- De Buck J, Ha V, Naushad S, Nobrega DB, Luby C, Middleton JR, De Vliegher S and Barkema HW (2021) Non-*aureus* staphylococci and bovine udder health: current understanding and knowledge gaps. *Frontiers in Veterinary Science* **8**, 360.
- de Freitas Guimarães F, Nóbrega DB, Richini-Pereira VB, Marson PM, de Figueiredo Pantoja JC and Langoni H (2013) Enterotoxin genes in coagulase-negative and coagulase-positive staphylococci isolated from bovine milk. *Journal of Dairy Science* **96**, 2866–2872.
- De Visscher A, Supré K, Haesebrouck F, Zadoks RN, Piessens V, Van Coillie E, Piepers S and De Vliegher S (2014) Further evidence for the existence of environmental and host-associated species of coagulase-negative staphylococci in dairy cattle. *Veterinary Microbiology* **172**, 466–474.
- De Vliegher S, Laevens H, Devriese LA, Opsomer G, Leroy JLM, Barkema HW and De Kruijff A (2003) Prepartum teat apex colonization with *Staphylococcus chromogenes* in dairy heifers is associated with low somatic cell count in early lactation. *Veterinary Microbiology* **92**, 245–252.
- De Vliegher S, Opsomer G, Vanrolleghem A, Devriese LA, Sampimon OC, Sol J, Barkema HW, Haesebrouck F and De Kruijff A (2004) In vitro growth inhibition of major mastitis pathogens by *Staphylococcus chromogenes* originating from teat apices of dairy heifers. *Veterinary Microbiology* **101**, 215–221.
- Dorneles EMS, Fonseca MDAM, Abreu JAP, Lage AP, Brito MAVP, Pereira CR, Brandão HM, Guimarães AS and Heinemann MB (2019) Genetic diversity and antimicrobial resistance in *Staphylococcus aureus* and coagulase-negative *Staphylococcus* isolates from bovine mastitis in Minas Gerais Brazil. *Microbiology Open* **8**, 1–7.
- Efsa H (nd) Antimicrobial resistance 2 How EFSA contributes to EU-wide cooperation 3 EFSA is working with key EU actors to reduce antimicrobial resistance 2–3.
- El-Seedy FA, Radwan IH, Hassan W and Shehata A (2017) Coagulase Negative Staphylococci as an emerging cause of bovine mastitis: prevalence antimicrobial resistance and biofilm formation. *Journal of Veterinary Medical Research* **24**, 1–1.
- Fairbrother RW (1940) Coagulase production as a criterion for the classification of the staphylococci. *The Journal of Pathology and Bacteriology* **50**, 83–88.
- Fergstad ME, Touzain F, De Vliegher S, De Visscher A, Thiry D, Ngassam Tchamba C, Mainil JGL, Abee-Lund T, Blanchard Y and Wasteson Y (2021) Whole genome sequencing of staphylococci isolated from bovine milk samples. *Frontiers in Microbiology* **12**, 3976.
- Ferronato JA, Souza FN, Della Libera AMMP, De Vliegher S, De Visscher A, Piepers S, Blagitz MG and Heinemann MB (2019) Inhibition of the growth of major mastitis-causing pathogens by non-*aureus* *Staphylococcus* isolates using the cross-streaking method. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia* **71**, 1745–1749.
- Food and Agriculture Organization of the United Nations (2021) Dairy market review. *Food and Agriculture Organization of the United Nations April* 1–13.
- Food and Drug Administration (FDA) (2018) Supporting antimicrobial stewardship in veterinary settings 1–21. Available at <https://www.fda.gov/downloads/AnimalVeterinary/SafetyHealth/AntimicrobialResistance/UCM620420.pdf#0Ahttps://www.vma.org/KB/Resources/Reference/Pages/One-Health94.aspx>
- Fuda C, Suvorov M, Shi Q, Heseck D, Lee M and Mobashery S (2007) Shared functional attributes between the *mecA* gene product of *Staphylococcus sciuri* and penicillin-binding protein 2a of methicillin-resistant *Staphylococcus aureus*. *Biochemistry* **46**, 8050–8057.
- Gao J, Barkema HW, Zhang L, Liu G, Deng Z, Cai L, Shan R, Zhang S, Zou J, Kastelic JP and Han B (2017) Incidence of clinical mastitis and distribution of pathogens on large Chinese dairy farms. *Journal of Dairy Science* **100**, 4797–4806.
- Garcia SN, Osburn BI and Cullor JS (2019) A One Health perspective on dairy production and dairy food safety. *One Health (Amsterdam, Netherlands)* **7**, 100086.
- Gill SR, Fouts DE, Archer GL, Mongodin EF, DeBoy RT, Ravel J, Paulsen IT, Kolony JF, Brinkac L, Beanan M, Dodson RJ, Daugherty SC, Madupu R, Angiuoli SV, Durkin AS, Haft DH, Vamathevan J, Khouri H, Utterback T and Fraser CM (2005) Insights on evolution of virulence and resistance from the complete genome analysis of an early methicillin-resistant *Staphylococcus aureus* strain and a biofilm-producing methicillin-resistant *Staphylococcus epidermidis* strain. *Journal of Bacteriology* **187**, 2426–2438.
- Gohl DM, Vangay P, Garbe J, MacLean A, Hauge A, Becker A, Gould TJ, Clayton JB, Johnson TJ, Hunter R, Knights D and Beckman KB (2016) Systematic improvement of amplicon marker gene methods for increased accuracy in microbiome studies. *Nature Biotechnology* **34**, 942–949.
- Gordon JL, Dewey KG, Mills DA and Medzhitov RM (2012) The human gut microbiota and undernutrition. *Science Translational Medicine* **4**, 2–7.
- Guerre P, Bailly JD, Benard G and Burgat V (1999) Risk linked to an enterotoxigenic strain of *Staphylococcus lentus* during the manufacture and ripening of raw goats' milk Camembert-type cheeses. *Revue de Médecine Vétérinaire* **150**, 703–708.
- Hammad AM, Watanabe W, Fujii T and Shimamoto T (2012) Occurrence and characteristics of methicillin-resistant and -susceptible *Staphylococcus*

- aureus* and methicillin-resistant coagulase-negative staphylococci from Japanese retail ready-to-eat raw fish. *International Journal of Food Microbiology* **156**, 286–289.
- Heikkilä AM, Liski E, Pyörälä S and Taponen S** (2018) Pathogen-specific production losses in bovine mastitis. *Journal of Dairy Science* **101**, 9493–9504.
- Hildreth CJ, Burke AE and Glass RM** (2009) Inappropriate use of antibiotics. *JAMA – Journal of the American Medical Association* **302**, 816.
- Hisatsune J, Hagiya H, Shiota S and Sugai M** (2017) Complete genome sequence of systemically disseminated sequence type 8 staphylococcal cassette chromosome mec type IV1 community-acquired methicillin-resistant *Staphylococcus aureus* *Genome Announcements* **5**, e00852-17.
- Hugh R and Ellis MA** (1968) The neotype strain for *Staphylococcus epidermidis* (Winslow and Winslow 1908) Evans 1916. *International Journal of Systematic Bacteriology* **18**, 231–239.
- Irlinger F** (2008) Safety assessment of dairy microorganisms: coagulase-negative staphylococci. *International Journal of Food Microbiology* **126**, 302–310.
- Isaac P, Bohl LP, Bresler ML, Orellano MS, Conesa A, Ferrero MA and Porporatto C** (2017) Commensal coagulase-negative *Staphylococcus* from the udder of healthy cows inhibits biofilm formation of mastitis-related pathogens. *Veterinary Microbiology* **207**, 259–266.
- Jenkins SN, Okello E, Rossitto PV, Lehenbauer TW, Champagne J, Penedo MCT, Arruda AG, Godden S, Rapnicki P, Gorden PJ, Timms LL and Aly SS** (2019) Molecular epidemiology of coagulase-negative *Staphylococcus* species isolated at different lactation stages from dairy cattle in the United States. *PeerJ* **2019**, e6749.
- Khazandi M, Al-Farha AAB, Coombs GW, O’Dea M, Pang S, Trott DJ, Aviles RR, Hemmatzadeh F, Venter H, Ogunniyi AD, Hoare A, Abraham S and Petrovski KR** (2018) Genomic characterization of coagulase-negative staphylococci including methicillin-resistant *Staphylococcus sciuri* causing bovine mastitis. *Veterinary Microbiology* **219**, 17–22.
- Kim SJ, Moon DC, Park SC, Kang HY, Na SH and Lim SK** (2019) Antimicrobial resistance and genetic characterization of coagulase-negative staphylococci from bovine mastitis milk samples in Korea. *Journal of Dairy Science* **102**, 11439–11448.
- Klibi A, Maaroufi A, Torres C and Jouini A** (2018) Detection and characterization of methicillin-resistant and susceptible coagulase-negative staphylococci in milk from cows with clinical mastitis in Tunisia. *International Journal of Antimicrobial Agents* **52**, 930–935.
- Kozioł-Montewka M, Szczepanik A, Baranowicz I, Józwiak L, Książek A and Kaczor D** (2006) The investigation of *Staphylococcus aureus* and coagulase-negative staphylococci nasal carriage among patients undergoing haemodialysis. *Microbiological Research* **161**, 281–287.
- Kumar R, Yadav BR and Singh RS** (2010) Genetic determinants of antibiotic resistance in *Staphylococcus aureus* isolates from milk of mastitic crossbred cattle. *Current Microbiology* **60**, 379–386.
- Lakhundi S and Zhang K** (2018) Methicillin-resistant *Staphylococcus aureus*: molecular characterization evolution and epidemiology. *Clinical Microbiology Reviews* **31**, e00020-18.
- Le KY, Villaruz AE, Zheng Y, He L, Fisher EL, Nguyen TH, Ho TV, Yeh AJ, Joo HS, Cheung GYC and Otto M** (2019) Role of phenol-soluble modulins in *Staphylococcus epidermidis* biofilm formation and infection of indwelling medical devices. *Journal of Molecular Biology* **431**, 3015–3027.
- Lee SI, Kim S, Do Park JH and Yang SJ** (2020) Species distribution antimicrobial resistance and enterotoxigenicity of non-*aureus* staphylococci in retail chicken meat. *Antibiotics* **9**, 809.
- Letertre C, Perelle S, Dilasser F and Fach P** (2003) Identification of a new putative enterotoxin SEU encoded by the *egc* cluster of *Staphylococcus aureus*. *Journal of Applied Microbiology* **95**, 38–43.
- Lienen T, Schnitt A, Hammerl JA, Maurischat S and Tenhagen BA** (2022) *Mammaliococcus* spp from German dairy farms exhibit a wide range of antimicrobial resistance genes and non-wildtype phenotypes to several antibiotic classes. *Biology* **11**, 152.
- Loncaric I, Kübber-Heiss A, Posautz A, Stalder GL, Hoffmann D, Rosengarten R and Walzer C** (2013) Characterization of methicillin-resistant *Staphylococcus* spp carrying the *mecC* gene isolated from wildlife. *The Journal of Antimicrobial Chemotherapy* **68**, 2222–2225.
- Madhaiyan M, Wirth JS and Saravanan VS** (2020) Phylogenomic analyses of the staphylococcaceae family suggest the reclassification of five species within the genus staphylococcus as heterotypic synonyms the promotion of five subspecies to novel species the taxonomic reassignment of five staphylococcus species to *mammaliococcus* gen Nov and the formal assignment of *nosocomiicoccus* to the family staphylococcaceae. *International Journal of Systematic and Evolutionary Microbiology* **70**, 926–936.
- Malinowski E, Lassa H, Khlossowska A, Smulski S, Markiewicz H and Kaczmarowski M** (2006) Etiological agents of dairy cows’ mastitis in west-ern part of Poland. *Polish Journal of Veterinary Sciences* **9**, 191–194.
- Marincola G, Liong O, Schoen C, Abouelfetouh A, Hamdy A, Wencker FDR, Marciniak T, Becker K, Köck R and Ziebuhr W** (2021) Antimicrobial resistance profiles of coagulase-negative staphylococci in community-based healthy individuals in Germany. *Frontiers in Public Health* **9**, 684456.
- Martínez-Meléndez A, Morfín-Otero R, Villarreal-Treviño L, Camacho-Ortiz A, González-González G, Llaca-Díaz J, Rodríguez-Noriega E and Garza-González E** (2016) Molecular epidemiology of coagulase-negative bloodstream isolates: detection of *Staphylococcus epidermidis* ST2 ST7 and linezolid-resistant ST23. *Brazilian Journal of Infectious Diseases* **20**, 419–428.
- Mohan R, Koebnick C, Schildt J, Mueller M, Radke M and Blaut M** (2008) Effects of *Bifidobacterium lactis* Bb12 supplementation on body weight fecal pH acetate lactate calprotectin and IgA in preterm infants. *Pediatric Research* **64**, 418–422.
- Munson SH, Tremaine MT, Betley MJ and Welch RA** (1998) Identification and characterization of staphylococcal enterotoxin types G and I from *Staphylococcus aureus*. *Infection and Immunity* **66**, 3337–3348.
- Namvar AE, Bastarahang S, Abbasi N, Ghehi GS, Farhadbakhtarian S, Arezi P, Hosseini M, Baravati SZ, Jokar Z and Chermahin SG** (2014) Clinical characteristics of *Staphylococcus epidermidis*: a systematic review. *GMS Hygiene and Infection Control* **9**, 23.
- Nascimento JDS, Fagundes PC, Brito MAVDP, Netto Dos Santos KR and Bastos MDCDF** (2005) Production of bacteriocins by coagulase-negative staphylococci involved in bovine mastitis. *Veterinary Microbiology* **106**, 61–71.
- Naushad S, Naqvi SA, Nobrega D, Luby C, Kastelic JP, Barkema HW and De Buck J** (2019) Comprehensive virulence gene profiling of bovine non-*aureus* staphylococci based on whole-genome sequencing data. *MSystems* **4**, 1–27.
- Nemeghaires S, Argudín MA, Feßler AT, Hauschild T, Schwarz S and Butaye P** (2014) The ecological importance of the *Staphylococcus sciuri* species group as a reservoir for resistance and virulence genes. *Veterinary Microbiology* **171**, 342–356.
- Nikaido H** (2009) Multidrug resistance in bacteria. *Annual Review of Biochemistry* **2**, 119–146.
- Omoe K, Hu DL, Ono HK, Shimizu S, Takahashi-Omoe H, Nakane A, Uchiyama T, Shinagawa K and Imanishi K** (2013) Emetic potentials of newly identified staphylococcal enterotoxin-like toxins. *Infection and Immunity* **81**, 3627–3631.
- Ono HK, Omoe K, Imanishi K, Iwakabe Y, Hu DL, Kato H, Saito N, Nakane A, Uchiyama T and Shinagawa K** (2008) Identification and characterization of two novel staphylococcal enterotoxins types S and T. *Infection and Immunity* **76**, 4999–5005.
- Ono HK, Sato’o Y, Narita K, Naito I, Hirose S, Hisatsune J, Asano K, Hu DL, Omoe K, Sugai M and Nakane A** (2015) Identification and characterization of a novel staphylococcal emetic toxin. *Applied and Environmental Microbiology* **81**, 7034–7040.
- Ono HK, Hirose S, Narita K, Sugiyama M, Asano K, Hu DL and Nakane A** (2019) Histamine release from intestinal mast cells induced by staphylococcal enterotoxin A (SEA) evokes vomiting reflex in common marmoset. *PLOS Pathogens* **15**, e1007803.
- Otto M** (2009) *Staphylococcus epidermidis* the “accidental” pathogen. *Nature Reviews Microbiology* **7**, 555–567.
- Pereira-Ribeiro PMA, Sued-Karam BR, Faria YV, Nogueira BA, Colodette SS, Fracalanza SEL, Duarte JLMB, Júnior RH and Mattos-Guaraldi AL**

- (2019) Influence of antibiotics on biofilm formation by different clones of nosocomial *Staphylococcus haemolyticus*. *Future Microbiology* **14**, 789–799.
- Persson Waller K, Aspán A, Nyman A, Persson Y and Grönlund Andersson U** (2011) CNS species and antimicrobial resistance in clinical and subclinical bovine mastitis. *Veterinary Microbiology* **152**, 112–116.
- Phophi L, Petzer IM and Qekwana DN** (2019) Antimicrobial resistance patterns and biofilm formation of coagulase-negative *Staphylococcus* species isolated from subclinical mastitis cow milk samples submitted to the Onderstepoort Milk Laboratory. *BMC Veterinary Research* **15**, 1–9.
- Piccart K, Verbeke J, De Visscher A, Piepers S, Haesebrouck F and De Vliegheer S** (2016) Local host response following an intramammary challenge with *Staphylococcus fleurettii* and different strains of *Staphylococcus chromogenes* in dairy heifers. *Veterinary Research* **47**, 1–11.
- Piepers S, Schukken YH, Passchyn P and De Vliegheer S** (2013) The effect of intramammary infection with coagulase-negative staphylococci in early lactating heifers on milk yield throughout first lactation revisited. *Journal of Dairy Science* **96**, 5095–5105.
- Piessens V, De Vliegheer S, Verbist B, Braem G, Van Nuffel A, De Vuyst L, Heyndrickx M and Van Coillie E** (2012) Characterization of coagulase-negative *staphylococcus* species from cows' milk and environment based on *bap icaA* and *mecA* genes and phenotypic susceptibility to antimicrobials and teat dips. *Journal of Dairy Science* **95**, 7027–7038.
- Piette A and Verschraegen G** (2009) Role of coagulase-negative staphylococci in human disease. *Veterinary Microbiology* **134**, 45–54.
- Pinheiro-Hubinger L, Riboli DFM, Abraão LM, Pereira Franchi EPL and Ribeiro de Souza da Cunha MdeL** (2021) Coagulase-negative staphylococci clones are widely distributed in the hospital and community. *Pathogens (Basel, Switzerland)* **10**, 792.
- Podkowik M, Park JY, Seo KS, Bystron J and Bania J** (2013) Enterotoxigenic potential of coagulase-negative staphylococci. *International Journal of Food Microbiology* **163**, 34–40.
- Qu Y, Zhao H, Nobrega DB, Cobo ER, Han B, Zhao Z, Li S, Li M, Barkema HW and Gao J** (2019) Molecular epidemiology and distribution of antimicrobial resistance genes of *Staphylococcus* species isolated from Chinese dairy cows with clinical mastitis. *Journal of Dairy Science* **102**, 1571–1583.
- Raspanti CG, Bonetto CC, Vissio C, Pellegrino MS, Reinoso EB, Dieser SA, Bogni CI, Larriestra AJ and Odierno LM** (2016) Prevalence and antibiotic susceptibility of coagulase-negative *Staphylococcus* species from bovine subclinical mastitis in dairy herds in the central region of Argentina. *Revista Argentina de Microbiología* **48**, 50–56.
- Rodrigues MX, Silva NCC, Trevelin JH, Cruzado MMB, Mui TS, Duarte FRS, Castillo CJC, Canniatti-Brazaca SG and Porto E** (2017) Molecular characterization and antibiotic resistance of *Staphylococcus* spp isolated from cheese processing plants. *Journal of Dairy Science* **100**, 5167–5175.
- Rohde H, Burandt EC, Siemssen N, Frommelt L, Burdelski C, Wurster S, Scherpe S, Davies AP, Harris LG, Horstkotte MA, Knobloch JKM, Rangunath C, Kaplan JB and Mack D** (2007) Polysaccharide intercellular adhesion or protein factors in biofilm accumulation of *Staphylococcus epidermidis* and *Staphylococcus aureus* isolated from prosthetic hip and knee joint infections. *Biomaterials* **28**, 1711–1720.
- Rumi MV, Huguet MJ, Bentancor AB and Gentilini ER** (2013) The *icaA* gene in staphylococci from bovine mastitis. *Journal of Infection in Developing Countries* **7**, 556–560.
- Sampimon OC, Barkema HW, Berends IMGA, Sol J and Lam TJGM** (2009) Prevalence and herd-level risk factors for intramammary infection with coagulase-negative staphylococci in Dutch dairy herds. *Veterinary Microbiology* **134**, 37–44.
- Sampimon OC, Lam TJGM, Mevius DJ, Schukken YH and Zadoks RN** (2011) Antimicrobial susceptibility of coagulase-negative staphylococci isolated from bovine milk samples. *Veterinary Microbiology* **150**, 173–179.
- Schleifer KH, Geyer U, Kilpper-Bälz R and Devriese LA** (1983) Elevation of *Staphylococcus sciuri* subsp *lentus*. *Systematic and Applied Microbiology* **4**, 382–387.
- Schnitt A, Lienen T, Wichmann-Schauer H and Tenhagen BA** (2021) The occurrence of methicillin-resistant non-aureus staphylococci in samples from cows young stock and the environment on German dairy farms. *Journal of Dairy Science* **104**, 4604–4614.
- Shemesh M and Ostrov I** (2020) Role of *Bacillus* species in biofilm persistence and emerging antibiofilm strategies in the dairy industry. *Journal of the Science of Food and Agriculture* **100**, 2327–2336.
- Silva NCC, Guimarães FF, Manzi MP, Budri PE, Gómez-Sanz E, Benito D, Langoni H, Rall VLM and Torres C** (2013) Molecular characterization and clonal diversity of methicillin-susceptible *Staphylococcus aureus* in milk of cows with mastitis in Brazil. *Journal of Dairy Science* **96**, 6856–6862.
- Silva NCC, Guimarães FF, Manzi MP, Júnior AF, Gómez-Sanz E, Gómez P, Langoni H, Rall VLM and Torres C** (2014) Methicillin-resistant *Staphylococcus aureus* of lineage ST398 as cause of mastitis in cows. *Letters in Applied Microbiology* **59**, 665–669.
- Simojoki H, Hyvönen P, Plumed Ferrer C, Taponen S and Pyörälä S** (2012) Is the biofilm formation and slime producing ability of coagulase-negative staphylococci associated with the persistence and severity of intramammary infection? *Veterinary Microbiology* **158**, 344–352.
- Souza FN, Piepers S, Della Libera AMMP, Heinemann MB, Cerqueira MMOP and De Vliegheer S** (2016) Interaction between bovine-associated coagulase-negative staphylococci species and strains and bovine mammary epithelial cells reflects differences in ecology and epidemiological behavior. *Journal of Dairy Science* **99**, 2867–2874.
- Spoor LE, Richardson E, Richards AC, Wilson GJ, Mendonca C, Gupta RK, McAdam PR, Nutbeam-Tuffs S, Black NS, O'gara JP, Lee CY, Corander J and Ross Fitzgerald J** (2015) Recombination-mediated remodelling of host-pathogen interactions during *Staphylococcus aureus* niche adaptation. *Microbial Genomics* **1**, 1–1.
- Srednik ME, Tremblay YDN, Labrie J, Archambault M, Jacques M, Cirelli AF and Gentilini ER** (2017) Biofilm formation and antimicrobial resistance genes of coagulase-negative staphylococci isolated from cows with mastitis in Argentina. *FEMS Microbiology Letters* **364**, 10.1093/femsle/fnx001
- Stevens M, Piepers S, Supré K and De Vliegheer S** (2018) Antimicrobial consumption on dairy herds and its association with antimicrobial inhibition zone diameters of non-aureus staphylococci and *Staphylococcus aureus* isolated from subclinical mastitis. *Journal of Dairy Science* **101**, 3311–3322.
- Su YC and Lee Wong AC** (1995) Identification and purification of a new staphylococcal enterotoxin H. *Applied and Environmental Microbiology* **61**, 1438.
- Supré K, Haesebrouck F, Zadoks RN, Vanechoutte M, Piepers S and De Vliegheer S** (2011) Some coagulase-negative *Staphylococcus* species affect udder health more than others. *Journal of Dairy Science* **94**, 2329–2340.
- Suzuki Y, Ono HK, Shimojima Y, Kubota H, Kato R, Kakuda T, Hirose S, Hu DL, Nakane A, Takai S and Sadamasu K** (2020) A novel staphylococcal enterotoxin SE02 involved in a staphylococcal food poisoning outbreak that occurred in Tokyo in 2004. *Food Microbiology* **92**, 103588.
- Tande AJ, Osmon DR, Greenwood-Quaintance KE, Mabry TM, Hanssen AD and Patel R** (2014) Clinical characteristics and outcomes of prosthetic joint infection caused by small colony variant staphylococci. *MBio* **5**, e01910-14.
- Taponen S, Björkroth J and Pyörälä S** (2008) Coagulase-negative staphylococci isolated from bovine extramammary sites and intramammary infections in a single dairy herd. *Journal of Dairy Research* **75**(4) 422–429. <https://doi.org/10.1017/S0022029908003312>
- Thomas DY, Jarraud S, Lemerrier B, Cozon G, Echasserieau K, Etienne J, Gougeon ML, Lina G and Vandenesch F** (2006) Staphylococcal enterotoxin-like toxins U2 and V two new staphylococcal superantigens arising from recombination within the enterotoxin gene cluster. *Infection and Immunity* **74**, 4724.
- Thorberg BM, Danielsson-Tham ML, Emanuelson U and Persson Waller K** (2009) Bovine subclinical mastitis caused by different types of coagulase-negative staphylococci. *Journal of Dairy Science* **92**, 4962–4970.
- Toledo-Silva B, de Souza FN, Mertens K, Piepers S, Haesebrouck F and De Vliegheer S** (2021a) Bovine-associated non-aureus staphylococci suppress *Staphylococcus aureus* biofilm dispersal in vitro yet not through agr regulation. *Veterinary Research* **52**, 114.
- Toledo-Silva B, Nogueira De Souza F, Piepers S, Mertens K, Haesebrouck F and De Vliegheer S** (2021b) Metabolites of bovine-associated non-aureus staphylococci influence expression of *Staphylococcus aureus* agr-related genes in vitro. *Veterinary Research* **52**, 62.

- Tomazi T, Gonçalves JL, Barreiro JR, Arcari MA and dos Santos MV** (2015) Bovine subclinical intramammary infection caused by coagulase-negative staphylococci increases somatic cell count but has no effect on milk yield or composition. *Journal of Dairy Science* **98**, 3071–3078.
- Traversari J, Van Den Borne BHP, Dolder C, Thomann A, Perreten V and Bodmer M** (2019) Non-*aureus* staphylococci species in the teat canal and milk in four commercial Swiss dairy herds. *Frontiers in Veterinary Science* **6**, 11–14.
- Turchi B, Bertelloni F, Marzoli F, Cerri D, Tola S, Azara E, Longheu CM, Tassi R, Schiavo M, Cilia G and Fratini F** (2020) Coagulase-negative staphylococci from ovine milk: genotypic and phenotypic characterization of susceptibility to antibiotics disinfectants and biofilm production. *Small Ruminant Research* **183**, 106030.
- Ünal N and Çinar OD** (2012) Detection of staphylococcal enterotoxin methicillin-resistant and Panton-Valentine leukocidin genes in coagulase-negative staphylococci isolated from cows and ewes with subclinical mastitis. *Tropical Animal Health and Production* **44**, 369–375.
- Valckenier D, Piepers S, De Visscher A, Bruckmaier RM and De Vliegheer S** (2019) Effect of intramammary infection with non-*aureus* staphylococci in early lactation in dairy heifers on quarter somatic cell count and quarter milk yield during the first 4 months of lactation. *Journal of Dairy Science* **102**, 6442–6453.
- Valckenier D, Piepers S, De Visscher A and De Vliegheer S** (2020) The effect of intramammary infection in early lactation with non-*aureus* staphylococci in general and *Staphylococcus chromogenes* specifically on quarter milk somatic cell count and quarter milk yield. *Journal of Dairy Science* **103**, 768–782.
- Valckenier D, Piepers S, Schukken YH, De Visscher A, Boyen F, Haesebrouck F and De Vliegheer S** (2021) Longitudinal study on the effects of intramammary infection with non-*aureus* staphylococci on udder health and milk production in dairy heifers. *Journal of Dairy Science* **104**, 899–914.
- Vanderhaeghen W, Piepers S, Leroy F, Van Coillie E, Haesebrouck F and De Vliegheer S** (2014) Invited review: Effect, persistence, and virulence of coagulase-negative *Staphylococcus* species associated with ruminant udder health. *Journal of Dairy Science* **97**, 5275–5293.
- Vanderhaeghen W, Piepers S, Leroy F, Van Coillie E, Haesebrouck F and De Vliegheer S** (2015) Identification typing ecology and epidemiology of coagulase negative staphylococci associated with ruminants. *Veterinary Journal* **203**, 44–51.
- Vitali LA, Petrelli D, Lamikanra A, Prenna M and Akinkunmi EO** (2014) Diversity of antibiotic resistance genes and staphylococcal cassette chromosome mec elements in faecal isolates of coagulase-negative staphylococci from Nigeria. *BMC Microbiology* **14**, 106.
- Walsh EJ, Miajlovic H, Gorkun OV and Foster TJ** (2008) Identification of the *Staphylococcus aureus* MSCRAMM clumping factor B (ClfB) binding site in the α C-domain of human fibrinogen. *Microbiology (Reading, England)* **154**, 550–558.
- WHO** (2024) Estimating the burden of foodborne diseases. Available at: <https://www.who.int/activities/estimating-the-burden-of-foodborne-diseases#:~:text=Each%20year%20worldwide%2C%20unsafe%20food%20causes%20600%20million,year%2C%20and%20this%20number%20is%20likely%20an%20under%20estimation>. Accessed March 2024.
- Wilson GJ, Seo KS, Cartwright RA, Connelley T, Chuang-Smith ON, Merriman JA, Guinane CM, Park JY, Bohach GA, Schlievert PM, Morrison WI and Fitzgerald JR** (2011) A novel core genome-encoded superantigen contributes to lethality of community-associated MRSA necrotizing pneumonia. *PLoS Pathogens* **7**, e1002271.
- Wuytack A, De Visscher A, Piepers S, Boyen F, Haesebrouck F and De Vliegheer S** (2019) Non-*aureus* staphylococci in fecal samples of dairy cows: first report and phenotypic and genotypic characterization. *Journal of Dairy Science* **102**, 9345–9359.
- Wuytack A, De Visscher A, Piepers S, Boyen F, Haesebrouck F and De Vliegheer S** (2020) Distribution of non-*aureus* staphylococci from quarter milk teat apices and rectal feces of dairy cows and their virulence potential. *Journal of Dairy Science* **103**, 10658–10675.
- Xu J, Tan X, Zhang X, Xia X and Sun H** (2015) The diversities of staphylococcal species virulence and antibiotic resistance genes in the subclinical mastitis milk from a single Chinese cow herd. *Microbial Pathogenesis* **88**, 29–38.
- Xu Z, Liu S, Chen L, Liu Y, Tan L, Shen J and Zhang W** (2019) Antimicrobial resistance and molecular characterization of methicillin-resistant coagulase-negative staphylococci from public shared bicycles in Tianjin China. *Journal of Global Antimicrobial Resistance* **19**, 231–235.
- Yong YY, Dykes GA and Choo WS** (2019) Biofilm formation by staphylococci in health-related environments and recent reports on their control using natural compounds. *Critical Reviews in Microbiology* **45**, 201–222.
- Zaman S, Bin Hussain MA, Nye R, Mehta V, Mamun KT and Hossain N** (2017) A review on antibiotic resistance: alarm bells are ringing. *Cureus* **9**, e1403.
- Zhang S, Iandolo JJ and Stewart GC** (1998) The enterotoxin D plasmid of *Staphylococcus aureus* encodes a second enterotoxin determinant (sej). *FEMS Microbiology Letters* **168**, 227–233.
- Zhou W, Li X, Shi L, Wang HH and Yan H** (2018) Novel SCCmec type XII methicillin-resistant *Staphylococcus aureus* isolates identified from a swine production and processing chain. *Veterinary Microbiology* **225**, 105–113.
- Zuniga E, Melville PA, Saidenberg ABS, Laes MA, Gonsales FF, Salaberry SRS, Gregori F, Brandão PE, dos Santos FGB, Lincopan NE and Benites NR** (2015) Occurrence of genes coding for MSCRAMM and biofilm-associated protein Bap in *Staphylococcus* spp isolated from bovine subclinical mastitis and relationship with somatic cell counts. *Microbial Pathogenesis* **89**, 1–6.